

Review

Derivatization of fatty acids and its application for conjugated linoleic acid studies in ruminant meat lipids

Noelia Aldai,^{1*} Brendan E Murray,² Ana I Nájera,³ Declan J Troy² and Koldo Osoro¹

¹Servicio Regional de Investigación y Desarrollo Agroalimentario (SERIDA), Apdo 13-33300 Villaviciosa, Asturias, Spain

²Teagasc, The National Food Centre, Ashtown, Dublin 15, Ireland

³Área de Tecnología de los Alimentos, Facultad de Farmacia, Universidad del País Vasco/Euskal-Herriko Unibertsitatea. Paseo de la Universidad 7, 01006 Vitoria-Gasteiz, Spain

Abstract: Conjugated linoleic acid (c9, t11 CLA) is a dietary fatty acid produced mainly by ruminant animals and exhibits promising health-promoting biological effects. For lipid fatty acid composition analyses, including CLA, lipids must be pre-treated so that the free and esterified fatty acids (triacylglycerols, phospholipids, etc) are available for determination. The most common treatments involve fatty acid methyl ester derivatives from relatively simple chemical reactions, but this becomes complicated when esterification of CLA is involved because of potential changes in its positional and geometrical isomers by reaction with certain reagents. In this review we explain concisely the advantages and disadvantages of the most popular methods (acid- and base-catalysed methods) generally employed for total fatty acid derivatization and their determination on a gas chromatograph. Based on our experiences we put forward the (trimethylsilyl) diazomethane method as an alternative and successful approach for ruminant tissue lipid determinations.

© 2005 Society of Chemical Industry

Keywords: fatty acids; derivatization methods; conjugated linoleic acid; CLA; ruminant lipids

INTRODUCTION

In ruminant meat, as in other meats, the quality of fat is as important as its quantity. Today, health professionals emphasize the association between nutrition and a number of lifestyle diseases in man (particularly heart disease and cancer), and have focused research activities on the role of metabolically important fatty acids. Dietary advice often recommends a reduction in the fat content of the diet, maintenance of the ratio of polyunsaturated to saturated fatty acids (P:S ratio) at about 0.45 or higher and an increase in the intake of *n*-3 polyunsaturated fatty acids (PUFA) relative to *n*-6 PUFA such that the *n*-6:*n*-3 ratio is less than 4.^{1–3}

Food products from ruminants, especially milk fat, represent the major dietary sources of conjugated linoleic acids (CLA), especially with respect to the *cis*-9, *trans*-11 isomer [*cis*-9(*Z*), *trans*-11(*E*) C_{18:2}]. Sources of other conjugated fatty acids are also found in plant seed oils (C_{18:3}) and marine organisms (C_{18:4}, C_{20:4}, C_{22:6}) with only trace amounts of CLA.^{4–9} The *cis*-9, *trans*-11-octadecadienoic acid

isomer (referred to as '*rumenic acid*'), accounts for 75–90% of total CLA in ruminant fats.^{5,10} Recently, a number of health-promoting biological activities for CLA using animal and *in vitro* models have been demonstrated,^{11–16} including changes in body composition and anticarcinogenic, antiatherogenic, antidiabetogenic and immunomodulatory activities. Currently, the evidence for direct beneficial effects in humans is scarce, sometimes contradictory, but promising.^{17,18}

Conjugated linoleic acids

'Conjugated linoleic acid' often describes a generic family of positional and geometric isomers of linoleic acid (C_{18:2} c9, c12) with diverse biological activity, in which the double bonds are conjugated instead of existing in the more typical methylene-interrupted configuration. Several different isomers of CLA have been reported. Of these isomers, the c9, t11 isoform is believed to be the most common natural form of CLA with biological activity as it is incorporated into the phospholipids of cell membranes.^{5,11,15,19}

* Correspondence to: Noelia Aldai, Servicio Regional de Investigación y Desarrollo Agroalimentario (SERIDA), Apdo 13-33300 Villaviciosa, Asturias, Spain

E-mail: naldai@serida.org

(Received 3 August 2004; revised version received 24 September 2004; accepted 8 October 2004)

Published online 21 February 2005

However, some authors could not detect *c9*, *t11* CLA in polar lipids of animal muscle, but found it present in neutral lipids, related to the storage lipid fraction.²⁰

Health-promoting properties have also been proposed for the other isomers, even if they are minor in quantity (eg long-chain derivatives of C_{18:2} *t10*, *c12* appear to be strong inhibitors of prostaglandin biosynthesis¹⁶). Since there are potential health benefits arising from CLA consumption, there is considerable research effort directed to determining and quantifying these components in ruminant-derived products.

CLA (predominantly the *c9*, *t11* isomer) is formed during ruminal biohydrogenation of linoleic acid as the first intermediate. It should be noted that *Butyrivibrio fibrisolvens* is thought to be the key bacterium involved in ruminal biohydrogenation and CLA formation.²¹ Other bacteria can also convert linoleic acid into CLA, including *Lactobacillus*, *Propionibacterium* and *Bifidobacterium* spp.^{22–25} For more information in the area of biohydrogenation in the rumen, the reader is directed towards more substantial reviews.^{26–28} The next step of the pathway involves the hydrogenation of the conjugated diene resulting in *trans*-11-octadecenoic acid (also referred to as *trans*-vaccenic acid, C_{18:1} *t11*). This is the major *trans* isomer of ruminant tissues and milk fat, and further biohydrogenation results in stearic acid (C_{18:0}).²⁹ The *c9*, *t11* isomer can also be synthesized by animal tissues from C_{18:1} *t11* if the Δ^9 -desaturase enzyme is present,^{30–32} this may also occur in humans through similar desaturase enzymes.^{33–35} In this way, the concentrations of C_{18:1} *t11* isomer and *c9*, *t11* CLA that are available for absorption from the gastrointestinal tract are related to the dietary intake of unsaturated fatty acids and rumen conditions that may affect the growth and activity of the reticulo rumen micropopulation. It is unlikely that *trans* octadecenoic acid availability limits this pathway in ruminants because the concentration of *trans* octadecenoic acid in milk and fat depots is always much higher than CLA. In contrast, changes in the rumen environment (pH, substrate types and concentration, dilution rate), which can lead to changes in microbial activity, will have an effect on CLA formation.

In general, lipids obtained from animal sources have complex structures and often most of the fatty acids found in meat lipids are bound to either an alcohol (glycerol) via ester bonds (triacylglycerols, phospholipids and sterol esters) or to a long-chain base (sphingosines) via amide bonds (sphingolipids). For fatty acid composition analyses, the complex lipids must be pre-treated so that the free and esterified fatty acids are available for determination. Fatty acids are converted to more non-polar and volatile derivatives by various derivatization methods. Formation of alkyl esters (methyl, ethyl, propyl or butyl esters) is the most standard type of reaction used in analysis

of fatty acids,^{36,37} in particular is the preparation of fatty acid methyl esters. Esters formed with higher molecular weight alcohols (ethanol, propanol, isopropanol, butanol) are useful in determining very volatile short chain fatty acids as are often found in dairy products and milk. For example, derivatization with isopropanol separates the low molecular analytes from the solvent front and improves the ruggedness for determination of the C₄–C₁₀ group of fatty acids by reducing the loss of such acids during the preparation procedure itself. Short chain fatty acids are more volatile than the longer chain C₁₂ and greater fatty acids characteristic of ruminant muscle and adipose tissue. The use of isopropanol gives rise to derivatives with higher boiling points (due to longer alkyl chains), which are much easier to handle and separate on a gas chromatograph.³⁷ The major disadvantage with the use of these higher molecular weight alcohols is the lack of commercially available standards and reference mixtures. Each individual fatty acid would have to be derivatized to allow for identification via similar retention time (flame ionisation detection) or ion chromatographs (mass spectroscopy). In contrast, the availability of methyl esters is very broad indeed (Supelco, Matreya, Larodan, Aldrich, etc). The preparation of the methyl ester derivatives of fatty acids is carried out in a simple reaction, but it becomes more complicated when derivatization of conjugated fatty acids or other potentially labile fatty acids are involved (eg labile functional groups include epoxy, cyclopropane and cyclopropene rings). The presence of a conjugated bond within fatty acids makes them unsuitable for the most common techniques (acid-catalysed reactions) employed for fatty acid analysis. For example, CLA can easily be isomerized by boron trifluoride (BF₃), thus changing the profile of the original isomer content.³⁸

Gas chromatography (GC) with the use of a flame ionization detector (FID), a large split flow (>30:1) and preferably hydrogen as carrier gas at a flow rate of 1.2 ml min⁻¹ is the suggested method for the analysis and quantification of fatty acids and their isomers, particularly if high resolution, polar capillary columns are employed (eg 100 m Chrompak CP-Sil88, 100 m Supelco SP-2560, 120 m SGE BPX-70). GC exhibits rapidity, high resolution and sensitivity.³⁹ However, even under optimal chromatographic conditions, a complete separation of all CLA isomers is not yet achievable and it is recognized that a combination of methods (multicolumn silver ion HPLC coupled with GC, two-dimensional GC, GC-MS) will be required to provide the full isomeric distribution of a sample.^{5,40–43}

Derivatization methods for total fatty acid analysis by gas chromatography

Fatty acids of animal origin exist as a heterogeneous group of chemical structures. It is therefore necessary to optimize the derivatization procedure in order to obtain accurate, quantitative and qualitative results

while addressing potential procedural difficulties associated with ester preparation.⁴⁴ Common problems include:

- incomplete conversion of the fatty acids to their methyl (or indeed ethyl- or propyl-) ester homologues (often as a result of the presence of water);
- alteration of the original fatty acid profile during esterification, because of chemical mediated changes in the proportion of different positional and/or geometric isomers;
- formation of artefacts which can be wrongly identified as fatty acids (giving rise to spurious peaks);
- contamination and subsequent damage of the GC column resulting from unclean samples (carry over of methoxide, trimethylsilyl, etc).

Various methods are currently employed for the conversion of fatty acids in a complex lipid matrix to fatty acid methyl esters. In this review we will try to explain the advantages and disadvantages of some of the different methods generally used for total fatty acid determinations, particularly in regard to ruminant tissues.

There are basically two mechanisms, which include both acidic and alkaline catalysis:

- (1) Methylation following hydrolysis of the fatty acids from the complex lipids involves saponification (alkaline hydrolysis) in which the ester bond between the fatty acid and the glycerol moiety is cleaved (eg triacylglycerols and phospholipids) under conditions of elevated temperature and in the presence of a strong alkali. This is followed by methylation typically performed in the presence of an acidic catalyst in methanol.
- (2) Direct *transesterification* involving alkaline or acidic catalysts in a process where the native alcohol (ie glycerol) is displaced from the lipid by another alcohol in large molar excess under anhydrous conditions.

It is essential to note that the method for preparing methyl esters for GC analysis is critical and has to

avoid modifications to the original isomer distribution. Generally, there is no single method that works optimally in all situations and for all kinds of samples. Researchers must know the nature of their sample and select the appropriate method.⁴⁵ Some of the most frequently employed methods are described below.

Acid-catalysed methods (Fig 1)

Boron trifluoride (BF₃) in methanol

An aliquot of lipid extract is mixed with BF₃–methanol and heated to a maximum of 100 °C for 2–90 min depending on the type of lipid.⁴⁶ After cooling to ambient temperature, water and a non-polar solvent are added, vortexed, and the two phases separated by centrifugation. The upper organic phase containing the methyl esters is carefully removed to a new vial where it is removed under a stream of nitrogen gas (N₂). The remaining residue containing fatty acid methyl esters is dissolved in n-hexane prior to GC analysis.

Since its introduction as a rapid method of preparation of fatty acid methyl esters from lipids,^{46–48} BF₃ has found wide application as a good direct *transesterification* reagent (methylating the fatty acids directly from the acylglycerol parent lipid), and esterification reagent (converting free fatty acids to their methyl ester derivatives). Rapid methyl ester formation is favoured by using an excess of anhydrous methanol and it can be purchased from commercial chemical companies already prepared at the desired concentration. It also has some disadvantages: the BF₃–methanol reagent is toxic and relatively unstable on storage. There have been reports that artefacts can form^{49–51} and polyunsaturated fatty acid decomposition can occur^{52–54} if the reagent is old or allowed to become too concentrated through evaporation. Some by-product of the degradation of BF₃–methanol reagent (formation of fluoroboron compounds by reaction with atmospheric oxygen⁵⁵) is capable of catalysing the addition of methanol across a double bond and new artefactual peaks have been observed eluting late in GC chromatograms. It can also cleave the rings in cyclopropane fatty acids and




Figure 1. General mechanism of acid-catalysed esterification of fatty acids (a) and *transesterification* of lipids (b). In the presence of water esterification is incomplete and *transesterification* results in some hydrolysis of the lipid to give free fatty acids. (Adapted from Christie WW, *Advances in Lipid Methodology—Two*, ed by Christie WW, pp 69–111. Oily Press, Dundee, 1993).

react with the antioxidant butylated hydroxytoluene (BHT) to produce additional unknown peaks in chromatograms.^{55,56} BF_3 is to be avoided if it is necessary to analyse fatty acids containing unusual functional groups such as conjugated double bonds as it can alter their configuration.^{54,57} It has been reported that CLA exhibits conversion from *c*, *t* or *t*, *c* isomers to *t*, *t* isomers or unknown substances as a consequence of BF_3 methylation.³⁸ The use of the milder BCl_3 -methanol reagent in place of the BF_3 -methanol reagent has similar results.⁵⁴

Hydrochloric acid (HCl) in methanol

To the lipid aliquot is added 5% (w/v) anhydrous HCl in methanol and it is heated to 80 °C for 20 min. Then some distilled water is added and the methylated fatty acids extracted with *n*-hexane. The *n*-hexane is evaporated under a stream of N_2 and re-dissolved in an adequate volume for GC analysis.

HCl-methanol converts both acylglycerols and free fatty acids into methyl esters. It has the advantages of being clean, easily prepared and applicable to all common lipid classes, including N-acyl lipids such as sphingolipids and glycosphingolipids.⁵⁸ However, several authors^{54,57,59–61} have found that acidic conditions can alter the isomer distribution of conjugated systems, usually seen as an increase in *t*, *t* isomer content. As with other acid-catalysed methylations, allylic methoxy derivatives are formed.^{57,61,62} Lowering the methylation temperature can decrease CLA isomerization, but these milder conditions may not be sufficient to methylate all lipids.^{38,60,63}

Sulphuric acid (H_2SO_4) in methanol

To the lipid aliquot is added 0.1 M H_2SO_4 in methanol and the tubes heated to 100 °C for 30–60 min. Sodium bicarbonate solution is added to neutralize the reagent and *n*-hexane to extract the FAMES. The organic layer is carefully removed, then dried over anhydrous sodium sulphate. Successive *n*-hexane extracts are added and evaporated under a stream of N_2 . The residue is re-dissolved in *n*-hexane for GC analysis.

This is an acid-catalysed methanolysis or transesterification method, and through its application it is possible to convert both acylglycerols and free fatty acids into methyl esters. Hitchcock and Hammond⁶⁴ suggested that H_2SO_4 -methanol was a very satisfactory reagent for methylation, but it also has the same disadvantages as other acid-catalysts in changing the isomeric distribution of conjugated fatty acids. Interestingly, H_2SO_4 -methanol has been found to be the mildest of the acid-based methods, and, if the reaction conditions are precisely controlled (55 °C for 5 min), little or no isomerization or methoxy artefacts are produced for the unesterified forms of CLA. For esterified lipids, artefacts are still produced (because more severe reaction conditions are required), but at a much lower quantity than those arising from the use of BF_3 .^{45,59} Some authors showed good results using

a traditional base-catalysed methylation, followed by a mild treatment using H_2SO_4 -methanol.⁴⁵

In general, acid-catalysed methods can be used in a direct transesterification or in a two-step methylation incorporating a previous hydrolysis or saponification of the lipids to free fatty acids. These methods usually require high temperatures (60–100 °C) for completion of methylation. Attention should be paid to the methylation of CLA isomers because these methods have been reported to change the *c*, *t* and *t*, *c* configurations in favour of the *t*, *t* configuration and generate allylic methoxy artefacts.^{57,59,60} In addition, higher temperatures and/or longer incubation times generate considerable amounts of artefact. When acid-catalysed methylations are carried out at a low temperature (60 °C) with a longer reaction time, the extent of isomerization and allylic methoxy derivative formation is greatly reduced.^{38,57,65} However, under such conditions, methylation of esterified fatty acids is not always complete, resulting in incomplete quantification. In experiments carried out by some researchers, *t*, *t* isomers were increased by acid-catalysed methods (HCl-MeOH and BF_3 -MeOH) but not by base-catalysed ones (NaOMe).⁶⁰ In another study, changes in CLA isomer proportions using the H_2SO_4 method were relatively small compared with those observed for the other acid-catalysed methods (BF_3 , HCl).⁵⁹ This is in agreement with results obtained by Park *et al.*⁴⁵ In the BF_3 method there is a time-dependent change in the CLA composition where *t*, *t* isomers increase markedly with time. Unknown peaks also continuously increase during methylation, but the total level observed in the H_2SO_4 method was lower than in the other two methods. Some authors also analysed the influence of antioxidants and dimethylsulphoxide (DMSO) and dimethylformamide (DMF) on CLA isomerization when used in conjunction with the acidic-catalysts. Antioxidants (butylated hydroxytoluene, ascorbic acid, β -carotene, α -tocopherol) at a 1 mM concentration had no preventive effect on the CLA isomer distribution. However, both DMSO and DMF added at the level of one-third of the reaction mixture both exhibited a protective effect against CLA isomerization by BF_3 and H_2SO_4 . It is thought that DMSO and DMF interact with the protonated alcohol (by way of their projecting oxygen atom) to prevent that protonated alcohol reacting with the conjugated double bonds in CLA and forming methoxy derivatives that upon further reaction can form *t*, *t* isomers.⁵⁹

A combination of methods appears to be the most suitable for analysis of lipid samples containing conjugated fatty acids. Normally NaOH or KOH in methanol is used at a temperature of 60–100 °C for the initial hydrolysis of acylglycerols. Saponification using sodium hydroxide in methanol followed by esterification with BF_3 -methanol was widely used and adopted as the official method for both the American Oil Chemists' Society (AOCS)⁶⁶ and the

Official Methods of Analysis of AOAC International.⁶⁷ However, it is not suitable for CLA quantification because *cis*, *trans*-conjugated fatty acids are lost (significant increase in the *t*, *t* CLA isomer occurs) and methoxy artefacts are formed.⁵⁴

Base-catalysed methods (Fig 2)

Sodium methoxide (NaOCH₃) in methanol

A lipid aliquot (20–40 mg) is dissolved in dry toluene, 2 ml 0.5 M sodium methoxide (metallic sodium in anhydrous methanol) added and the samples heated at 60 °C for 20 min. This esterification step can also be carried out at room temperature (20 °C). A few drops of glacial acetic acid are added followed by 2 ml saturated NaCl solution to eliminate excess methoxide. With 50 ppm butylated hydroxy toluene (BHT) 2 ml n-hexane are added, and vortexed vigorously. The upper organic phase containing the extracted methyl esters is taken, dried over anhydrous sodium sulphate or 100 µl of a water scavenger such as 2, 2 dimethoxypropane added, and dried under a gentle stream of N₂ at 40 °C. The fatty acid methyl esters are resuspended in 2 ml of n-hexane with 50 ppm BHT and injected for GC analysis. Dilution of the sample in n-hexane may be necessary, depending on the GC response.

This *transesterification* method is the generally accepted method for the conversion of acylglycerides to methyl esters. Nevertheless, this technique is not suitable when large amounts of free fatty acids are present in the sample, as it is unable to methylate free fatty acids and N-linked (amide-bond) fatty acids such as those found in sphingolipids and glycosphingolipids.⁶⁸ This is a characteristic of all base-catalysed methylations. Only acid-based reagents and diazomethane can methylate free fatty acids. NaOCH₃ does not cause isomerization of conjugated double bonds, nor does it produce methoxy artefacts.^{54,57} Reaction conditions need to be strictly anhydrous as the presence of water causes saponification.⁵⁴ NaOCH₃ is a popular methylating agent in the oil industry, especially for refined oils, where the free fatty acids have already been removed during the refining stage. The removal of excess methoxide before GC analysis is also strongly recommended, as the reagent tends to interfere with analysis. An aqueous wash necessary for the removal of excess reagent can affect some fatty acids. Without due care, losses in short-chain fatty acid methyl esters can

render the technique inaccurate for the determination of methyl butyrate (C₄) and caproate (C₆), which are present in considerable amounts in dairy fat.⁶⁹

Potassium methoxide (KOCH₃) or potassium hydroxide (KOH) in methanol

The KOCH₃ reagent is used in a similar manner to NaOCH₃ described in the previous section, but is more reactive and difficult to prepare. KOH in anhydrous methanol may also be used: are the lipids dissolved in n-hexane, KOH methanolic reagent added and the samples heated at 50 °C for 10–15 min. A few drops of glacial acetic acid are added followed by water and n-hexane. The solution is vortexed vigorously and the upper organic phase taken and dried over anhydrous sodium sulphate prior to injection for GC analysis.

KOH in anhydrous methanol and KOCH₃ are other *transesterification* reagents commonly employed in the production of methyl esters from oils. In a study carried out involving NaOCH₃ and KOCH₃ reagents, it was found that KOCH₃ was more active than NaOCH₃ and produced greater conversion of cholesteryl esters into their corresponding methyl esters under comparable conditions.⁷⁰ KOCH₃ was used for quantitative determination of conjugated linoleic acid isomers in beef fat.⁷¹ An interesting point about the use of KOH in methanol is its ability to efficiently *transesterify* lipids directly from dry (lyophilized) tissue samples. Other researchers showed that concentrations of CLA were greatest after direct *transesterification* of dried muscle tissue with KOH compared with NaOCH₃.⁶¹ They further reported similar efficiencies for both reagents when *transesterifying* extracted total lipid. This is to be expected, as strong alkali is efficient at solubilizing biological tissues.

Tetramethylguanidine (TMG) in methanol

To a dried sample of lipid is added 20% TMG in methanol. The tubes are heated at 100 °C for 10 min. Saturated NaCl solution and n-hexane are added to each tube, then vortexed and centrifuged to separate the phases. After centrifugation the organic layer is removed, and the n-hexane extraction repeated. n-Hexane extracts are combined and dried under a gentle stream of N₂. The residue containing FAMES is re-dissolved in an adequate volume of solvent for GC analysis.




Figure 2. General mechanism of base-catalysed *transesterification* of lipids. An unesterified fatty acid is converted to a carboxylate ion, R-COO⁻, in a basic solution and cannot be esterified. In the presence of water, the intermediate (R''=H) will dissociate irreversibly to the free acid. (Adapted from Christie WW, *Advances in Lipid Methodology – Two*, ed by Christie WW, pp 69–111. Oily Press, Dundee, 1993).

More recently, guanidine and its alkylated derivatives in methanol have been used to catalyse the esterification of oils and fats. Although the original authors of the TMG method⁷² claimed that it worked well with acylglycerols and free fatty acids, recent results by other researchers suggest otherwise.^{45,54,57} This method is not effective in methylating free fatty acids and N-acyl lipids, and has difficulty with some esterified lipids such as phospholipids.^{45,57} In addition, TMG can give undesirable products that extract with *n*-hexane and interfere with the analyses of short-chain FAME. Therefore, this method is not suitable for samples with high acid values. For clear chromatographic resolution, excess reagent needs to be removed prior to sample injection onto the GC column. The derivatization reaction is quite rapid; the reagent is inexpensive and mild, causing no isomerization of conjugated double bonds. Conditions for TMG–methanol need to be absolutely anhydrous, otherwise, soap formation will result in the loss of fatty acids and it will not be possible to obtain accurate results.⁵⁴ Park *et al*⁴⁵ obtained good results for methylation of esterified-CLA rich lipids in egg yolk, but only if they also included an additional reaction with H₂SO₄–methanol at 55 °C for 5 min.

Other reagents are available for *transesterification* with shortened derivatization times. One of these is tetramethylammonium hydroxide in methanol, which *transesterifies* triglyceride fatty acids to methyl esters.^{73,74} This reagent minimizes the losses of C_{14:0} and C_{15:0} fatty acids and unsaturated C₁₈ isomers in the analysis of plant oils, margarine, lard and animal tissues.⁷³

In general, base-catalysed methods are mild reactions suitable for methylating the fatty acids bound up in acylglycerides, but are not suitable for samples with substantial amounts of free fatty acids. Care must be taken to exclude water from the reaction medium to prevent saponification. This hydrolysis is an unwanted and irreversible side reaction.^{52,53}

Diazoalkane (diazomethane) method (Fig 3)

Trimethylsilyldiazomethane [(CH₃)₃SiCHN₂] method

The lipids are saponified with strong base such as 5 M KOH at a temperature of 60 °C for 60 min (a high concentration of base gives more efficient saponification; the authors routinely use 6 ml of 5 M KOH in 50% methanol with samples containing approximately 20–40mg of total lipid, eg 1 g of muscle

tissue). NaOH may also be used and gives identical fatty acid profiles, but reducing the concentration of base below 5 M reduces the overall efficiency of saponification, requiring longer incubation times at 60 °C. Following saponification, the reaction mixture is diluted with 12 ml of 0.5% NaCl and a non-polar solvent (5 ml petroleum spirit) added to remove non-saponifiable lipid components (cholesterol, N-acyl lipids). The samples are vortexed and the upper solvent layer discarded. The samples are acidified using glacial acetic acid (typically 3 ml) and solvent added to partition the (protonated) free fatty acids into the solvent phase. The tubes are vortexed and the non-polar phase collected, then 100 µl of 2, 2-dimethoxypropane (a water scavenger) are added and the mixture is vortexed. The solvent is removed by drying under a stream of nitrogen (N₂) at 40 °C. For esterification, the residue is re-dissolved in 1 ml toluene–methanol (1:2 v/v) and trimethylsilyldiazomethane (TMS-DM) added in molar excess (about 120–160 µl of 2 M TMS-DM in hexane depending on fatty acid content). The diazomethane solution is yellow and this is lost as the methylation reaction proceeds at 40 °C for a maximum of 10 min. In addition, in the presence of free fatty acids there is a rapid liberation of N₂ gas as the reaction proceeds, so it is important not to cap the reaction tubes. There is no need to quench the reaction with acetic acid. Excess solvent and diazomethane are removed under a gentle stream of nitrogen at 40 °C. Finally, the residue is dissolved in 2 ml of *n*-hexane containing 50 ppm butylated hydroxy toluene (BHT), centrifuged at 20 000 × g for 5 min and 1 µl of supernatant transferred to GC for chromatographic analysis. Within-sample precision is very good with typical relative standard deviations of 5–10% for each fatty acid methyl ester (bovine muscle tissue), the higher values associated with those fatty acids in low or variable abundance such as the branched chain fatty acids. The use of an internal standard such as C_{23:0} methyl ester can give variable results (60–85% recovery), but the authors consider this due to solubility issues with the saponification solution. This serves to highlight one of the inherent difficulties with fortification of samples with analytical standards—that is, the standard may not always behave in the same way as native components under the extraction procedure. This is not of much concern




Figure 3. General mechanism of trimethylsilyldiazomethane mediated methylation of free fatty acids. The reaction must have methanol present to get good yields of the desired methyl ester. Adapted from Hashimoto *et al*.⁸⁵

since measurement of the recovered fatty acid residue weight per unit weight tissue agrees well with total lipid content of the tissue as measured by other procedures such as solvent extraction and microwave drying of samples (procedure adapted from Elmore *et al.*,⁷⁵ by Murray, BE personal communication).

Diazomethane can easily convert free fatty acids to methyl esters in high yield in the presence of methanol, but cannot act on acylglycerides. It is therefore essential to hydrolyse or saponify lipid samples first.⁷⁶ Diazomethane, for all its advantages (short incubation times, mild reaction conditions, high methyl ester yield), has several preparative problems (toxic, explosive, potential carcinogen) and reportedly may give rise to artefacts associated with its use as indicated by some authors,⁷⁷ even though evidence appears to be lacking in the literature about this in relation to fatty acid methyl ester formation. This, perhaps, contributes to reluctance amongst laboratories to use it as a general procedure in the preparation of fatty acid methyl esters. There was an interesting report concerning unreliable estimates of serum non-esterified fatty acids with the use of diazomethane,⁷⁸ however, a subsequent report⁷⁹ was able to clarify that solvolysis of serum lipids took place to increase the free fatty acid content. In fact, numerous publications have used diazomethane quite satisfactorily for the methylation of their fatty acids.^{57,65,75,80–84} Trimethylsilyldiazomethane (TMS-DM) is an alternative, convenient and commercially available source of diazomethane. It is safer to handle and more stable with minimal side reactions. It can be obtained as a 2.0 M solution in diethyl ether or hexane. The esterification proceeds nearly instantaneously and quantitatively when an excess of TMS-DM is used in the presence of methanol. The reaction conditions are typically very mild (40 °C, 10 min) and the reaction can be easily monitored by the disappearance of its yellow colour and liberation of nitrogen gas (which drives the reaction in favour of methyl ester formation).

Excess reagent can be easily eliminated by evaporation under a stream of nitrogen gas and spillages rendered harmless by treatment with acetic acid.

It was demonstrated that the methyl esters prepared from a mixture of fatty acids by reaction with TMS-DM and methanol was quantitative and gave chromatograms which were completely identical to those of methyl esters prepared by the action of pure diazomethane.⁸⁵ Work carried out in the laboratory of one of the authors also demonstrates that saponification followed by use of TMS-DM as a methylating reagent gives identical fatty acid profiles to the general base-catalysed method represented by sodium methoxide in methanol when analysing identical lipid samples from a number of sources such as beef fat, muscle and menhaden fish oil (Fig 4). In addition, the usefulness of the technique described previously, allows for the analysis of a wide range of tissue and sample matrices without the need for extensive lipid extraction with the use of solvents (Fig 5). It is the opinion of these authors that TMS-DM promises a non-hazardous routine procedure for the preparation of methyl esters of fatty acids.

TMS-DM does not change the original isomer distribution nor alter the geometric configuration of conjugated double bonds; it also does not produce the methoxy artefacts associated with acid-catalysed methylations as reported by some researchers.^{57,59–61} However, in a recent study, Park *et al.*⁶⁰ found a small amount of artefacts attributable to TMS-DM (stated as new peaks generated from the sample by the procedure) and a group of impurities (stated as peaks that originated from reagents) at low retention times. They suggest that these are trimethylsilyl fatty acid derivatives and trimethylsilyl, respectively, and that they will interfere with short-chain fatty acids in GC. In our view, the comments relating to TMS-DM artefacts⁶⁰ need to be carefully considered, especially in view of the lack of detail in how the samples were processed. It must be stated that these authors also




Figure 4. Chromatogram excerpts (overlaid) illustrating the direct *transesterification* method using NaOCH₃ vs the TMS–diazomethane method for the methylation of fatty acids from lipid extracted from beef adipose tissue. Samples were methylated in duplicate and gave identical profiles. Insert is a magnified view of the C_{18:3} to CLA region. Column used was a Phenomenex BPX-70 (SGE), 120 m, 0.25 mm id, H₂ carrier gas, temperature programme for 42 min. Timebase given in minutes.




Figure 5. Chromatograms illustrating the range of different foodstuffs (sample matrices) that are easily processed using the saponification/TMS-DM procedure outlined in this review. Column used was a Phenomenex BPX-70 (SGE), 120 m, 0.25 mm id, H₂ carrier gas, temperature programme for 42 min. Timebase given in minutes.

reported artefacts due to the other methods employed in their study. In our experience, TMS-DM is a very satisfactory method for the methylation of free fatty acids with very little evidence of artefact formation (there are some contaminants found eluting early, well before C_{12:0}; however we find no evidence of artefacts eluting later than this fatty acid). It is crucial that methanol is present in large concentrations for the reaction to proceed completely and vigorously under mild conditions in a solvent system that adequately dissolves complex mixtures of fatty acids from tissue or lipid extracts (eg toluene—methanol at 1:2 v/v). We typically dissolve 30–40 mg total fatty acid in 1 ml of the above solvent with the addition of small amounts of TMS-DM (100–200 µl of a 2.0 M solution in hexane). With lower concentrations of methanol or its complete absence, artefacts are generated in the C_{18:0}–C_{23:0} region of our chromatograms. Presumably these artefacts represent partial or intermediate reaction products as well as trimethylsilyl derivatives.

SUGGESTED DERIVATIZATION

Gas chromatography is the primary method of choice for the analysis of the fatty acid content of lipid samples from many different sources because of its rapidity, high resolution and sensitivity.³⁹ Fatty acids must first be converted to non-polar, volatile derivatives such as methyl esters. However, care must be taken when deciding upon a suitable derivatization procedure, particularly if the analyst is also interested

in determining conjugated fatty acids such as CLA. No single procedure exists which simultaneously methylates all lipid components (triacylglycerides, free fatty acids, N-acyl lipids such as sphingolipids and glycosphingolipids, and very short chain fatty acids). The best method or combination of methods will depend almost entirely on the nature of the sample to be analysed. The papers by Shantha *et al*⁵⁴ and Kramer *et al*⁵⁷ have excellent summaries of the many varied methods available, particularly when referring to CLA analysis, and the reader is encouraged to refer to these publications.

Traditionally, when analysing lipid samples with a significant CLA content (milk and dairy products, ruminant meats) the recommendations clearly advise base-catalysed *transesterification*, often involving the use of sodium methoxide in methanol. However, a few reports also suggest that acid-catalysed methylations may be used if mild conditions are employed, often shortened times (5–20 min) at reduced temperatures (50 °C).^{45,59,60}

In this review, the authors have no problems in suggesting a slightly different approach which works well on both extracted lipid and oils, but especially so on different tissue matrices (eg adipose tissue, muscle tissue, fish, etc). However, the analysis of extracted lipid and pure oils by TMS-DM is not quite as efficient as methods based on sodium methoxide. Qualitatively, results are identical, all fatty acids are derivatized, but peak heights tend to be lower for samples processed by the saponification/TMS-DM

procedure by about 30%–40%. This difference may be a result of the difference in solubility of oils and lipids (or their respective solvents) in 5 M KOH 50% methanol vs 0.5 M sodium methoxide (NaOCH₃). In support of this is the observation that different solvents effect saponification—non-polar solvents such as diethylether and petroleum spirit reduce the efficiency of saponification whereas toluene–methanol solutions give the highest efficiency. In every case the smallest volume of added solvent ensures the highest efficiency of saponification. The saponification/TMS-DM procedure, however, has a large advantage over sodium methoxide in that it is not as sensitive to water in the samples. Hence, it is not necessary to lyophilize tissue and complex matrices prior to derivatization. Up to 1 g of tissue or 50 mg of oil or lipid extract can be successfully saponified using strong alkali in aqueous methanol. Following a simple set of clean up steps, the extracted fatty acids are dissolved in 1 ml of toluene–methanol (1:2 v/v) and converted to their methyl esters using 200–400 μmol TMS-diazomethane. There is no need to quench the reaction with acetic acid and after dissolving in n-hexane with 50 ppm of butylated hydroxytoluene (BHT) they are ready for injection onto the GC. This procedure will enable the analysis of all the esterified fatty acids as well as the free fatty acids (C_{11:0} to C_{22:6n-3}, branched chain fatty acids, cyclopropane fatty acids, furans, conjugated fatty acids, long chain PUFA, etc) in a sample, but it is important to realize that losses would be incurred for the volatile very small chain fatty acids and N-acyl lipids.

For very small chain fatty acids (as present in milk and butter fats) it would be wise to also produce the fatty acid isopropyl ester (FAIPE) analogues by the methods using isopropanol and H₂SO₄.^{86,87} For analysis of N-acyl lipids the acid-catalysed methods (HCl, BF₃, acetyl chloride and H₂SO₄), eg HCl in anhydrous methanol at 80 °C for 1 h,⁸⁸ are appropriate, but at the expense of correct CLA isomer determinations. For the determination of fatty acids in pure oils then the sodium methoxide method is more than appropriate.^{54,57,68}

ACKNOWLEDGEMENTS

N Aldai acknowledges a predoctoral fellowship awarded by the Spanish Ministry of Science and Technology (INIA). B Murray acknowledges support by the FIRM Programme administered by the Irish Department of Agriculture and Food.

REFERENCES

- Department of Health. Nutritional aspects of cardiovascular disease, in *Reports on Health and Social Subjects*, no 46. Her Majesty's Stationery Office, London (1994).
- Williams CM, Dietary fatty acids and human health. *Ann Zootech* 49:165–180 (2000).
- Enser M, The role of fats in human nutrition, in *Oils and Fats. Animal Carcass Fats*, Ed by Rossell B. Leatherhead Publishing, Leatherhead, Surrey, Vol 2, pp 77–122 (2001).
- Parodi PW, Conjugated octadecadienoic acids of milk fat. *J Dairy Sci* 60:1550–1553 (1977).
- Chin SF, Storkson JM, Ha YL and Pariza MW, Dietary sources of conjugated dienoic isomers of linoleic acid, a newly recognized class of anticarcinogens. *J Food Comp Anal* 5:185–197 (1992).
- Fritsche J, Rickert R and Steinhart H, Formation, contents, and estimation of daily intake of conjugated linoleic acid isomers and trans-fatty acids in foods, in *Advances in Conjugated Linoleic Acid Research*, Ed by Yurawecz MP, Mossoba MM, Kramer JKG, Pariza MW and Nelson GJ, Vol 1. AOCS-Press, Champaign, IL, pp 378–396 (1999).
- Cahoon EB, Carlson TJ, Ripp KG, Schweiger BJ, Cook GA, Hall SE and Kinney AJ. Biosynthetic origin of conjugated double bonds: Production of fatty acid components of high-value drying oils in transgenic soybean embryos. *Proc Natl Acad Sci USA* 96:12 935–12 940 (1999).
- Cahoon EB, Ripp KG, Hall SE and Kinney AJ. Formation of conjugated Δ8, Δ10-double bonds by Δ12-oleic-acid desaturase-related enzymes. *J Biol Chem* 276:2637–2643 (2001).
- Hamberg M, Metabolism of 6,9,12-octadecatrienoic acid in the red alga *Lithothamnion corallioides*: mechanism of formation of a conjugated tetraene fatty acid. *Biochem Biophys Res Commun* 188:1220–1227 (1992).
- Sehat N, Kramer JKG, Mossoba MM, Yurawecz MP, Roach JAG, Eulitz K, Morehouse KM and Ku Y, Identification of conjugated linoleic acid isomers in cheese by gas chromatography, silver ion high performance liquid chromatography and mass spectral reconstructed ion profiles. Comparison of chromatographic elution sequences. *Lipids* 33:963–971 (1998).
- Ip C, Chin SF, Scimeca JA and Pariza MW, Mammary cancer prevention by conjugated dienoic derivative of linoleic acid. *Cancer Res* 51:6118–6124 (1991).
- Ip C, Banni S, Angioni E, Carta G, McGinley J, Thompson HJ, Barbano D and Bauman D, Conjugated linoleic acid-enriched butter fat alters mammary gland morphogenesis and reduces cancer risk in rats. *J Nutr* 129:2135–2142 (1999).
- Houseknecht KL, Vanden Heuvel JP, Moya-Camarena SY, Portocarrero CP, Peck LW, Nickel KP and Belury MA, Dietary conjugated linoleic acid normalises impaired glucose tolerance in the Zucker diabetic fatty *falfa* rat. *Biochem Biophys Res Commun* 244:678–682 (1998).
- Li Y and Watkins BA, Conjugated linoleic acids alter bone fatty acid composition and reduce ex vivo prostaglandin E2 biosynthesis in rats fed *n-6* or *n-3* fatty acids. *Lipids* 33:417–425 (1998).
- Ha YL, Storkson J and Pariza MW, Inhibition of benz[α]pyrene-induced mouse forestomach neoplasia by conjugated dienoic derivatives of linoleic acid. *Cancer Res* 50:1097–1101 (1990).
- Nugteren DH, Inhibition of prostaglandin biosynthesis by 8-*cis*, 12-*trans*, 14-*cis*-eicosatrienoic acid and 5-*cis*, 8-*cis*, 12-*trans*, 14-*cis*-eicosatetraenoic acid. *Biochim Biophys Acta* 210:171–176 (1970).
- Blankson H, Stakkestad JA, Fagertun H, Thom E, Wadstein J and Gudmundsen O. Conjugated linoleic acid reduces body fat mass in overweight and obese humans. *J Nutr* 130:2943–2948 (2000).
- Zambell KL, Keim NL, Van Loan MD, Gale B, Benito P, Kelley DS and Nelson GJ. Conjugated linoleic acid supplementation in humans: effects on body composition and energy expenditure. *Lipids* 35:777–782 (2000).
- Fogerty AC, Ford GL and Svoronos D, Octadeca-9, 11-dienoic acid in foodstuffs and in the lipids of human blood and breast milk. *Nutr Rep Int* 38:937–944 (1988).
- EU Healthy Beef Project. A European Commission Research Project (QLRT-2000-31423). Executive Summary; Annual Report Year 2; www.healthybeef.iger.bbsrc.ac.uk [accessed 20 January 2005].
- Kepler CR and Tove SB, Biohydrogenation of unsaturated fatty acids. *J Biol Chem* 242:5686–5692 (1967).

- 22 Ogawa J, Matsumura K, Kishino S, Omura Y and Shimizu S, Conjugated linoleic acid accumulation via 10-Hydroxy-12-octadecenoic acid during microaerobic transformation of linoleic acid by *Lactobacillus acidophilus*. *Appl Environ Microbiol* **67**:1246–1252 (2001).
- 23 Alonso L, Cuesta EP and Gilliland SE, Production of free conjugated linoleic acid by *Lactobacillus acidophilus* and *Lactobacillus casei* of human intestinal Origin. *J Dairy Sci* **86**:1941–1946 (2003).
- 24 Coakley MR, Ross P, Nordgren M, Fitzgerald G, Devery R and Stanton C. Conjugated linoleic acid biosynthesis by human-derived Bifidobacterium species. *J Appl Microbiol* **94**:138–145 (2003).
- 25 Jiang J, Bjorck L and Fonden R. Production of conjugated linoleic acid by dairy starter cultures. *J Appl Microbiol* **85**:95–102 (1998).
- 26 Harfoot CG and Hazlewood GP, Lipid metabolism in the rumen, in *The Rumen Microbial Ecosystem*, 2nd edn, Ed by Hobson PN and Stewart CS. Blackie Academic & Professional, New York, pp. 382 (1997).
- 27 Demeyer D and Doreau M, Targets and procedures for altering ruminant meat and milk lipids. *Proc Nutr Soc* **58**:593–607 (1999).
- 28 Khanal RC and Dhiman TR. Biosynthesis of conjugated linoleic acid (CLA): a review. *Pakistan J Nutr* **3**:72–81 (2004).
- 29 Griinari JM and Bauman DE, Biosynthesis of conjugated linoleic acid and its incorporation into meat and milk in ruminants, in *Conjugated Linoleic Acid Research*, Ed by Yurawecz MP, Mossoba MM, Kramer JKG, Pariza MW and Nelson G, Vol 1. AOCS Press, Champaign, IL, pp 180–200 (1999).
- 30 Palmquist DL and Santora JE, Endogenous synthesis of rumenic acid in rodents and humans, in *Advances in Conjugated Linoleic Acid Research*, Ed by Yurawecz MP, Mossoba MM, Kramer JKG, Pariza MW and Nelson GJ, Vol 1. AOCS Press, Champaign, IL, pp 201–208 (1999).
- 31 Griinari JM, Corl BA, Lacy SH, Chouinard PY, Nurmela KVV and Bauman DE, Conjugated linoleic acid is synthesised endogenously in lactating cows by Δ^9 -desaturase. *J Nutr* **130**:2285–2291 (2000).
- 32 Corl BA, Baumgard LH, Dwyer DA, Griinari JM, Philips BS and Bauman DE, The role of Delta(9)-desaturase in the production of *cis*-9, *trans*-11 CLA. *J Nutr Biochem* **12**:622–630 (2001).
- 33 Parodi PW, Conjugated linoleic acid and other anticarcinogenic agents of bovine milk fat. *J Dairy Sci* **82**:1339–1349 (1999).
- 34 Adlof AB, Duval D and Emken EA, Biosynthesis of conjugated linoleic acid in humans. *Lipids* **35**:131–135 (2000).
- 35 Turpeinen AM, Mutanen M, Aro A, Salminen I, Basu S, Palmquist DL and Griinari JM, Bioconversion of vaccenic acid to conjugated linoleic acid in humans. *Am J Clin Nutr* **76**:504–510 (2002).
- 36 Child P, Aloe M and Mee D, Separation and quantification of fatty acids, sterols and bile acids in feces by gas chromatography as the butyl ester-acetate derivatives. *J Chromatogr* **415**:13–26 (1987).
- 37 Rosenfeld JM, Application of analytical derivatizations to the quantitative and qualitative determination of fatty acids. *Anal Chim Acta* **465**:93–100 (2002).
- 38 Werner AS, Luedecke LO and Shultz TD, Determination of conjugated linoleic acid in three Cheddar-type cheeses: effects of cheese cultures, processing and aging. *J Agric Food Chem* **40**:1817–1821 (1992).
- 39 Horning EC, Ahrens EH Jr, Lipsky SR, Mattson FH, Mead JF, Turner DA and Goldwater WH, Quantitative analysis of fatty acids by gas-liquid chromatography. *J Lipid Res* **5**:20–27 (1964).
- 40 Fritsche J and Steinhart H, Amounts of conjugated linoleic acid (CLA) in German foods and evaluation of daily intake. *Z Lebensm Unters Forsch A* **206**:77–82 (1998).
- 41 Precht D and Molckentin J, Analysis and seasonal variation of conjugated linoleic acid and further *cis*-/*trans*-isomers of C18:1 and C18:2 in bovine milk fat. *Kieler Milchwirtschaftl Forschungsberichte* **51**:63–78 (1999).
- 42 Kramer JKG and Zhou J, Conjugated linoleic acid and octadecenoic acids: Extraction and isolation of lipids. *Eur J Lipid Sci Technol* **103**:594–632 (2001).
- 43 Western RJ, Lau SS, Marriott PJ and Nichols PD, Positional and geometric isomer separation of FAME by comprehensive 2-D GC. *Lipids* **37**:715–724 (2002).
- 44 Shantha NC and Napolitano GE, Gas chromatography of fatty acids. *J Chromatogr* **624**:37–51 (1992).
- 45 Park SJ, Park CW, Kim SJ, Kim JK, Kim YR, Park KA, Kim JO and Ha YL, Methylation methods for the quantitative analysis of conjugated linoleic acid (CLA) isomers in various lipid samples. *J Agric Food Chem* **50**:989–996 (2002).
- 46 Morrison WR and Smith LM, Preparation of fatty acid methyl esters and dimethylacetals from lipids with boron fluoride-methanol. *J Lipid Res* **5**:600–608 (1964).
- 47 Metcalfe LD and Schmith AA, The rapid preparation of fatty acid esters for gas chromatographic analysis. *Anal Chem* **33**:363–364 (1961).
- 48 Metcalfe LD, Schmitz AA and Pelka JR, Rapid preparation of fatty acid esters from lipids for gas chromatographic analysis. *Anal Chem* **38**:514–515 (1966).
- 49 Fulk WK and Shorb MS, Production of an artifact during methanolysis of lipids by boron-trifluoride-methanol. *J Lipid Res* **11**:276–277 (1970).
- 50 Dawidowicz EA and Thompson TE, Artifacts produced by boron trifluoride methanolysis of a synthetic lectin containing cyclopropane fatty acids (1-2-dihydrosterculoyl-3-sn-phosphatidylcholine). *J Lipid Res* **12**:636–637 (1971).
- 51 Kramer JKG and Hulan HW, Artifacts produced during acid-catalysed methanolysis of sterol esters. *J Lipid Res* **17**:674–676 (1976).
- 52 Christie WW, *Gas Chromatography and Lipids*. The Oily Press, Ayr (1989).
- 53 Hammond EW, *Chromatography for the Analysis of Lipids*. CRC Press, Boca Raton, FL (1993).
- 54 Shantha NC, Decker EA and Hennig B, Comparison of methylation methods for the quantitation of conjugated linoleic acid isomers. *J AOAC Int* **76**:644–649 (1993).
- 55 Christie WW, Why I dislike boron trifluoride-methanol. *Lipid Technol* **6**:66–68 (1994).
- 56 Lough AK, The production of methoxy-substituted fatty acids as artefacts during the esterification of unsaturated fatty acids with methanol containing boron trifluoride. *Biochem J* **90**:4c–5c (1964).
- 57 Kramer JKG, Fellner V, Dugan MER, Sauer FD, Mossoba MM and Yurawecz MP, Evaluating acid and base catalysts in the methylation of milk and rumen fatty acids with special emphasis on conjugated dienes and total *trans* fatty acids. *Lipids* **32**:1219–1228 (1997).
- 58 Ichihara K, Shibahara A, Yamamoto K and Nakayama T, An improved method for rapid analysis of the fatty acids of glycerolipids. *Lipids* **31**:535–539 (1996).
- 59 Yamasaki M, Kishihara K, Ikeda I, Sugano M and Yamada K, A recommended esterification method for gas chromatographic measurement of conjugated linoleic acid. *J Am Oil Chem Soc* **76**:933–938 (1999).
- 60 Park Y, Albright KJ, Cai ZY and Pariza MW, Comparison of methylation procedures for conjugated linoleic acid and artifact formation by commercial (trimethylsilyl) diazomethane. *J Agric Food Chem* **49**:1158–1164 (2001).
- 61 Murrieta CM, Hess BW and Rule DC, Comparison of acidic and alkaline catalysts for preparation of fatty acid methyl esters from ovine muscle with emphasis on conjugated linoleic acid. *Meat Sci* **65**:523–529 (2003).
- 62 Yurawecz MP, Hood JK, Roach JAG, Mossoba MM, Daniels DH, Ku Y, Pariza MW and Chin SF, Conversion of allylic hydroxy oleate to conjugated linoleic acid and methoxy oleate by acid-catalysed methylation procedures. *J Am Oil Chem Soc* **71**:1149–1155 (1994).

- 63 Park Y and Pariza MW, Evidence that commercial calf and horse serum can contain substantial amounts of *trans*-10, *cis*-12 conjugated linoleic acid. *Lipids* **33**:817–819 (1998).
- 64 Hitchcock C and Hammond EW, in *Developments in Food Analysis*, 2nd edn, Ed by King RD. Applied Science, London (1980).
- 65 Yurawecz MP, Sehat N, Mossoba MM, Roach JAG, Kramer JKG and Ku Y, Variations in isomer distribution in commercially available conjugated linoleic acid. *Fett/Lipid* **101**:277–282 (1999).
- 66 *Official Methods and Recommended Practices of the American Oil Chemists' Society*, 4th edn, Ed by Firestone D. American Oil Chemists' Society, Champaign, IL, Official Method Ce 1b-89 (1989).
- 67 *Official Methods of Analysis of AOAC International*, 16th edn, Ed by Cunniff P, Vol 2. AOAC International, Arlington, VA, Official Method 969, p 33 (1995).
- 68 Christie WW, A simple procedure for rapid transmethylation of glycerolipids and cholesterol esters. *J Lipid Res* **23**:1072–1075 (1982).
- 69 Bannon CD, Breen GJ, Craske JD, Hai NT, Harper NL and O'Rourke KL, Analysis of fatty acid methyl esters with high accuracy and reliability. III. Literature review of and investigations into the development of rapid procedures for the methoxide-catalysed methanolysis of fats and oils. *J Chromatogr* **247**:71–89 (1982).
- 70 Luddy FE, Barfold RA and Riemenschneider RW, Direct conversion of lipid components to their fatty acid methyl esters. *J Am Oil Chem Soc* **37**:447–451 (1960).
- 71 Fritsche J, Fritsche S, Solomon MB, Mossoba MM, Yurawecz MP, Morehouse K and Ku Y, Quantitative determination of conjugated linoleic acid isomers in beef fat. *Eur J Lipid Sci Technol* **102**:667–672 (2000).
- 72 Schuchardt U and Lopes OC, Tetramethylguanidine-catalysed transesterification of fats and oils. A new method for rapid determination of their composition. *J Am Oil Chem Soc* **65**:1940–1941 (1988).
- 73 Misir R, Laarveld B and Blair R, Evaluation of a rapid method for preparation of fatty acid methyl esters for analysis by gas liquid chromatography. *J Chromatogr* **331**:141 (1985).
- 74 Fourie PC and Basson DS, Application of a rapid transesterification method for identification of individual fatty acids by gas chromatography on three different nut oils. *J Am Oil Chem Soc* **67**:18–20 (1990).
- 75 Elmore JS, Mottram DS, Enser M and Wood JD, Effect of the polyunsaturated fatty acid composition of beef muscle on the profile of aroma volatiles. *J Agric Food Chem* **47**:1619–1625 (1990).
- 76 Christie WW, *Lipid analysis*, 2nd edn. Pergamon Press, Oxford (1982).
- 77 Little JL, Derivatization of carboxylic acids with diazomethane and trimethylsilyldiazomethane: convenient methods and artefacts; <http://users.chartertn.net/slitttle/files/diazoalkanes.pdf> [accessed 20 January 2005].
- 78 Ingalls ST, Xu Y and Hoppel CL, Determination of plasma non-esterified fatty acids and triglyceride fatty acids by gas chromatography of their methyl esters after isolation by column chromatography on silica gel. *J Chromatogr B Biomed Applic* **666**:1–12 (1995).
- 79 Mueller H, Diazomethane as a highly selective fatty acid methylating reagent for use in gas chromatographic analysis. *J Chromatogr B* **679**:208–209 (1996).
- 80 Yurawecz MP, Hood JK, Mossoba MM, Roach JAG and Ku Y, Furan fatty acids determined as oxidation products of conjugated octadecadienoic acid. *Lipids* **30**:595–598 (1995).
- 81 Kramer JKG, Sehat N, Dugan MER, Mossoba MM, Yurawecz MP, Roach JAG, Eulitz K, Aalhus JL, Schaefer AL and Ku Y, Distributions of conjugated linoleic acid (CLA) isomers in tissue lipid classes of pigs fed a commercial CLA mixture determined by gas chromatography and silver ion-high-performance liquid chromatography. *Lipids* **33**:549–558 (1998).
- 82 Sehat N, Yurawecz MP, Roach JAG, Mossoba MM, Kramer JKG and Ku Y, Silver-ion high performance liquid chromatographic separation and identification of conjugated linoleic acid isomers. *Lipids* **33**:217–221 (1998).
- 83 Scollan ND, Choi NJ, Kurt E, Fisher AV, Enser M and Wood JD, Manipulating the fatty acid composition of muscle and adipose tissue in beef cattle. *Br J Nutr* **85**:115–124 (2001).
- 84 Adlof R, Copes LC and Walter EL, Changes in conjugated linoleic acid composition within samples obtained from a single source. *Lipids* **36**:315–317 (2001).
- 85 Hashimoto N, Aoyama T and Shioiri T, New methods and reagents in organic synthesis 14. A simple efficient preparation of methyl esters with trimethylsilyldiazomethane (TMSCHN₂) and its application to gas chromatographic analysis of fatty acids. *Chem Pharm Bull* **29**:1475–1478 (1981).
- 86 Wolff RL and Fabien RJ, Utilisation de l'isopropanol pour l'extraction de la matière grasse de produits laitiers et pour l'estérification subséquente des acides gras. *Le Lait* **69**:33–46 (1989).
- 87 Wolff RL and Sébédio JL, Geometrical isomers of linolenic acid in low-calorie spreads marketed in France. *J Am Oil Chem Soc* **68**:719–725 (1991).
- 88 Stoffel W, Chu F and Ahrens EH, Analysis of long chain fatty acids by gas-liquid chromatography. Micromethod for preparation of methyl esters. *Anal Chem* **31**:307–308 (1959).