

DETECTION OF  $1.6 \times 10^{10} M_{\odot}$  OF MOLECULAR GAS IN THE HOST GALAXY  
OF THE  $z = 5.77$  SDSS QUASAR J0927+2001

C. L. CARILLI,<sup>1</sup> R. NERI,<sup>2</sup> R. WANG,<sup>1,3</sup> P. COX,<sup>2</sup> F. BERTOLDI,<sup>4</sup> F. WALTER,<sup>5</sup> X. FAN,<sup>6</sup> K. MENTEN,<sup>7</sup> J. WAGG,<sup>1</sup> R. MAIOLINO,<sup>8</sup>  
A. OMONT,<sup>9</sup> MICHAEL A. STRAUSS,<sup>10</sup> D. RIECHERS,<sup>5</sup> K. Y. LO,<sup>11</sup> A. BOLATTO,<sup>12</sup> AND N. SCOVILLE<sup>13</sup>

Received 2007 June 24; accepted 2007 July 17; published 2007 August 14

ABSTRACT

We have detected emission by the CO(5–4) and (6–5) rotational transitions at  $z = 5.7722 \pm 0.0006$  from the host galaxy of the SDSS quasar J0927+2001 using the Plateau de Bure Interferometer. The peak line flux density for the CO(5–4) line is  $0.72 \pm 0.09$  mJy, with a line FWHM =  $610 \pm 110$  km s<sup>-1</sup>. The implied molecular gas mass is  $(1.6 \pm 0.3) \times 10^{10} M_{\odot}$ . We also detect the 90 GHz continuum at  $0.12 \pm 0.03$  mJy, consistent with a 47 K dust spectrum extrapolated from higher frequencies. J0927+2001 is the second example of a huge molecular gas reservoir within the host galaxy of a quasar within 1 Gyr of the big bang. Observations of J0927+2001 are consistent with a massive starburst coeval with a bright quasar phase in the galaxy, suggesting the rapid formation of both a supermassive black hole through accretion, and the stellar host spheroid, at a time close to the end of cosmic reionization.

*Subject headings:* galaxies: formation — galaxies: high-redshift — quasars: general — radio lines: galaxies — submillimeter

*Online material:* color figure

1. INTRODUCTION

Understanding the relationship between quasars and their host galaxies has become imperative since the discovery of the bulge mass–black hole mass correlation in nearby galaxies, a result which implies a fundamental relationship between the formation of supermassive black holes (SMBHs) and spheroidal galaxies (Ferrarese & Merrit 2000; Magorrian et al. 1998; Gebhardt et al. 2000). In our extensive study of the dust and gas content, and star formation activity, of the host galaxies of  $z > 2$ , luminous ( $M_{1450AA} < -26.1$ ), optically selected high-redshift quasars (Omont et al. 2003; Carilli et al. 2002; Bertoldi et al. 2003a; Petric et al. 2003; Beelen et al. 2006; Wang et al. 2007), we have found that roughly 30% of these sources are also hyperluminous infrared galaxies ( $L_{FIR} \sim 10^{13} L_{\odot}$ ). This FIR emission corresponds to thermal emission from warm dust. Searches for CO emission from these sources typically yield molecular gas masses  $\geq 10^{10} M_{\odot}$  (Solomon & Vanden Bout 2005). Such molecular gas represents the fuel for star formation and can serve as a dynamical tracer in a forming galaxy.

An important finding of our program is that the 30% fraction

of FIR luminous quasars continues to the highest redshifts,  $z \sim 6$ , approaching the end of cosmic reionization, and hence pushing toward the very first galaxies and SMBHs (e.g., Fan et al. 2006a, 2006b). A particularly enlightening example is the highest redshift quasar known, J1148+5251 at  $z = 6.419$ . The host galaxy has been detected in thermal dust, non-thermal radio continuum, and CO line emission (Walter et al. 2003; Bertoldi et al. 2003b; Carilli et al. 2004), with an FIR luminosity (integrated between 42.5 and 122  $\mu\text{m}$ ) of  $1.3 \times 10^{13} L_{\odot}$  (Beelen et al. 2006), a molecular gas mass of  $\sim 2 \times 10^{10} M_{\odot}$ , and a dust mass of  $7 \times 10^8 M_{\odot}$ . The dynamical mass to a radius of 2.5 kpc in the host galaxy, as estimated from high-resolution VLA imaging of the molecular gas distribution, is comparable to the gas mass, but much less than the mass predicted by the black hole mass–bulge mass relation measured at low redshift (Walter et al. 2004). We have also detected the dominant cooling line of interstellar gas, [C II] 158  $\mu\text{m}$ , from J1148+5251 (Maiolino et al. 2005). These observations demonstrate that large reservoirs of dust and metal enriched atomic and molecular gas can exist in massive galaxies within 1 Gyr of the big bang. The current observations suggest active star formation in the host galaxy, with a massive star formation rate of order  $10^3 M_{\odot} \text{ yr}^{-1}$ , although the question remains as to the contribution to dust heating by the AGN (Wang et al. 2007; Li et al. 2007b).

The source SDSS J092721.82+200123.7 (hereafter J0927+2001) is similar to J1148+5251 in many ways. J0927+2001 is a quasar selected from the Sloan Digital Sky Survey (York et al. 2000), with an optical redshift of  $z = 5.77 \pm 0.03$  (corresponding to  $t_{\text{univ}} = 1$  Gyr, assuming a standard concordance cosmology), based on fitting to the (relatively weak) Ly $\alpha$  + N v lines, and with  $L_{\text{bol}} = 7 \times 10^{13}$  erg s<sup>-1</sup> (Fan et al. 2006c; Wang et al. 2007). The black hole mass estimated from both the Eddington limit and UV line widths is  $\sim 10^9 M_{\odot}$  (Jiang et al. 2007).

J0927+2001 has been detected in 250 GHz continuum emission at the IRAM 30 m telescope, with a flux density of  $5.0 \pm 0.8$  mJy (R. Wang et al. 2007, in preparation), and at 350 GHz at the Caltech Submillimeter Observatory, with a flux

<sup>1</sup> National Radio Astronomy Observatory, P.O. Box O, Socorro, NM 87801; ccarilli@nrao.edu.

<sup>2</sup> Institut de Radio Astronomie Millimétrique (IRAM), 300 rue de la Piscine, Domaine Universitaire de Grenoble, 38406 St. Martin d’Hères, France.

<sup>3</sup> Astronomy Department, Peking University, Beijing 100871, China.

<sup>4</sup> Argelander-Institut für Radioastronomie, Universität Bonn, auf dem Hügel 71, Bonn 53121, Germany.

<sup>5</sup> Max-Planck-Institut für Astronomie, Königstuhl 17, Heidelberg, Germany.

<sup>6</sup> Steward Observatory, University of Arizona, Tucson, AZ 85721.

<sup>7</sup> Max-Planck-Institut für Radioastronomie, Auf dem Hügel 69, Bonn 53121, Germany.

<sup>8</sup> Osservatorio Astrofisico di Arcetri, Largo E. Fermi 5, 50125 Firenze, Italy.

<sup>9</sup> Institut d’Astrophysique de Paris, CNRS; and Université Pierre et Marie Curie, Paris, France.

<sup>10</sup> Princeton University Observatory, Princeton, NJ 08544.

<sup>11</sup> National Radio Astronomy Observatory, 520 Edgemont Road, Charlottesville, VA 22903.

<sup>12</sup> Department of Astronomy, 601 Campbell Hall, University of California, Berkeley, CA 94720.

<sup>13</sup> Robinson Laboratory, California Institute of Technology, Pasadena, CA 91125.




FIG. 1.—Spectra of the CO(5–4) (*top*), (6–5) (*middle*), and the sum of the (5–4) and (6–5) (*bottom*) emission from SDSS J0927+2001 at  $z = 5.77$ . Also shown are Gaussian fits to the lines with parameters as given in Table 1. All spectra are smoothed to  $70 \text{ km s}^{-1} \text{ channel}^{-1}$ , and each independent channel is displayed. The rms per channel varies across the band, although typical values in the combined (5–4) and (6–5) spectrum are  $0.3 \text{ mJy beam}^{-1}$ . [See the electronic edition of the *Journal* for a color version of this figure.]

density of  $18 \pm 5 \text{ mJy}$  (R. Wang et al. 2007, in preparation). These data, in combination with lower and higher frequency data, reveal a clear FIR excess over a standard quasar IR SED. This FIR excess corresponds to thermal emission from warm dust with a temperature  $\sim 47 \text{ K}$ . The dust mass is  $6.9 \times 10^8 M_{\odot}$ , and the FIR luminosity is  $1.2 \times 10^{13} L_{\odot}$ . A weak ( $45 \pm 14 \mu\text{Jy}$ ) radio continuum counterpart has been detected at 1.4 GHz, and the rest-frame radio through FIR SED is within the range defined by star-forming galaxies (Wang et al. 2007). The observations of J0927+2001 are consistent with a massive (stars  $> 5 M_{\odot}$ ) star formation rate of  $\sim 700 M_{\odot} \text{ yr}^{-1}$  (Kennicutt 1998), or a total star formation rate (stars  $> 0.1 M_{\odot}$ ) a factor 5.6 higher, assuming a Salpeter initial mass function, and assuming the warm (47 K) dust is heated by star formation.

In this Letter we present the discovery of CO(5–4) and (6–5) emission from J0927+2001. These observations reveal the necessary fuel for star formation and enable a number of key observations in the study of the earliest generation of massive galaxies and SMBHs.

## 2. OBSERVATIONS AND RESULTS

Our observations have been made with the improved Plateau de Bure Interferometer equipped with the new generation of receivers. With their increased sensitivity (detection of sub-

mJy lines in a single track) and increased bandwidth (2 of the 4 GHz with the current correlator), these new receivers allow for deep searches for faint redshifted molecular emission lines over bandwidths adequate to cover the typical uncertainties of optical redshifts of high-redshift quasars.

We searched for CO(5–4) emission from J0927+2001 with the Plateau de Bure Interferometer in 2007 April and May. The first track (8 hr) used two polarizations, and two frequency settings which covered a total bandwidth of 1.8 GHz ( $= 5700 \text{ km s}^{-1}$  or  $\Delta z = 0.13$ ) centered at  $z = 5.79$  [or 84.87 GHz for redshifted CO(5–4)]. After a possible line was identified in the first observations at  $z \sim 5.77$ , a second 8 hr track was observed centered on the line at 85.12 GHz, covering a total bandwidth of 1 GHz using dual polarization. This second track confirmed the (5–4) line with high significance. A third 8 hr track was then observed centered on the frequency of the CO(6–5) line redshifted to 102.10 GHz, which was also clearly detected. The spectral resolution for all observations was  $2.5 \text{ MHz channel}^{-1}$ . The phase stability, as measured on the phase calibrator (0851+202), was easily adequate for coherent integration in the D configuration, for which the interferometric synthesized beam had a FWHM  $\sim 5''$ . Absolute gain calibration was performed on the star MWC 349.

Figure 1 shows the spectra of the CO(5–4) and (6–5) emission lines from J0927+2001 at a spectral resolution of  $70 \text{ km s}^{-1} \text{ channel}^{-1}$ . The rms per channel varies slightly across the spectra, but typical values for the (5–4) spectrum are  $0.37 \text{ mJy beam}^{-1}$  for (5–4), and  $0.52 \text{ mJy beam}^{-1}$  for (6–5).

Both CO emission lines are clearly detected, as well as the underlying continuum emission at both frequencies. Gaussian fitting to the lines yield the parameters listed in Table 1, including the continuum level, the line peak flux density and FWHM, the velocity-integrated line flux, and the line luminosity,  $L'(\text{CO}) = 3.25 \times 10^{13} (1+z) I \Delta v D_A^2 \nu_{\text{obs}}^{-2} \text{ K km s}^{-1} \text{ pc}^2$ , where  $I \Delta v$  is the velocity-integrated flux density in  $\text{Jy km s}^{-1}$ ,  $D_A$  is the angular diameter distance in Gpc ( $D_A = 1.23 \text{ Gpc}$  at  $z = 5.77$ ), and  $\nu_{\text{obs}}$  is the observing frequency in GHz (Solomon et al. 1997). A line width of  $600 \pm 70 \text{ km s}^{-1}$  and an LSR line centroid of  $z = 5.7722 \pm 0.0006$  are derived from the spectrum obtained by merging the CO (5–4) and (6–5) data.

The measured continuum level for the combined data is  $0.12 \pm 0.03 \text{ mJy}$ . Combined with the continuum observations at 250 and 350 GHz, this yields a dust temperature of 47 K, assuming a modified graybody of index  $\beta = 1.6$  (see analysis in R. Wang et al. 2007, in preparation).

Figure 2 shows the image of the velocity-integrated line emission from the weighted sum of the (5–4) and (6–5) lines. The peak of the CO emission is at  $09^{\text{h}}27^{\text{m}}21.79^{\text{s}}, +20^{\circ}01'23.5''$  (J2000.0), or within  $0.2''$  of the optical position. The source appears unresolved at the present resolution of FWHM  $\sim 5''$ .

## 3. ANALYSIS

The ratio of the strength of the two CO transitions is  $L'_{6-5}/L'_{5-4} = 1.10 \pm 0.27$ . This ratio is consistent with constant

TABLE 1  
GAUSSIAN FITTING TO THE CO LINE EMISSION FROM J0927+2001

Transition	Continuum (mJy)	Line Peak (mJy)	FWHM (km s <sup>-1</sup> )	$I \Delta v$ (Jy km s <sup>-1</sup> )	$L'(\text{CO})$ (K km s <sup>-1</sup> pc <sup>2</sup> )
(5–4) .....	$0.12 \pm 0.03$	$0.72 \pm 0.09$	$610 \pm 110$	$0.44 \pm 0.07$	$(2.0 \pm 0.3) \times 10^{10}$
(6–5) .....	$0.13 \pm 0.09$	$1.25 \pm 0.25$	$550 \pm 150$	$0.69 \pm 0.13$	$(2.2 \pm 0.5) \times 10^{10}$




FIG. 2.—Image of the sum of the velocity-integrated CO(5–4) and CO(6–5) line emission from J0927+2001 at  $z = 5.77$ . Contour levels are  $0.14 \text{ Jy km s}^{-1} \text{ beam}^{-1}$ . The rms on the image is  $0.07 \text{ Jy km s}^{-1} \text{ beam}^{-1}$ . The cross indicates the position of the optical QSO.

brightness temperature (i.e., ratio = 1) up to CO(6–5). The high CO excitation in J0927+2001 is comparable to that seen in the host galaxy of J1148+5251 (Bertoldi et al. 2003b) and other high-redshift quasars (Solomon & Vanden Bout 2005; Carilli et al. 2002; A. Weiss et al. 2007, in preparation), as well as in the nuclei of nearby nuclear starburst galaxies, such as NGC 253 and Arp 220 (Greve et al. 2007; Bradford et al. 2003), implying warm ( $>50 \text{ K}$ ), dense ( $>10^4 \text{ cm}^{-3}$ ) gas. However, given the close spacing of the two transitions, observations of a lower order transition (1–0 or 2–1) are needed to better constrain the gas excitation conditions.

We derive the total molecular gas mass (dominated by  $\text{H}_2$ ) from the CO luminosity using the conversion factor for CO luminosity to total molecular gas mass appropriate for ultraluminous infrared galaxies (ULIRGs, or galaxies with  $L_{\text{FIR}} > 10^{12} L_{\odot} \text{ yr}^{-1}$ ; Downes & Solomon 1998), or  $X = 0.8 M_{\odot} (\text{K km s}^{-1} \text{ pc}^2)^{-1}$ , and assuming constant brightness temperature from CO(5–4) down to CO(1–0) [i.e., constant  $L(\text{CO})$ ]. The implied gas mass is  $M_{\text{H}_2} = 0.8 \times L_{\text{CO}(1-0)} = (1.6 \pm 0.3) \times 10^{10} M_{\odot}$ . Note that the CO luminosity to gas mass conversion factor is a factor 5 smaller for ULIRGs than it is for normal galaxies, such as the Milky Way. For a detailed discussion of the conversion of CO luminosity to gas mass in high-redshift, FIR-luminous galaxies, see Solomon & Vanden Bout (2005).

The ratio of FIR luminosity to CO luminosity has been used as a metric for the relative star formation efficiency in galaxies (i.e., proportional to the star formation rate per unit gas mass; Gao & Solomon 2004). For J0927+2001, we find  $L_{\text{FIR}}/L_{\text{CO}} = 650 L_{\odot} (\text{K km s}^{-1} \text{ pc}^2)^{-1}$ . In their review of molecular line emission from galaxies at  $z > 1$ , Solomon & Vanden Bout (2005) find that the values of  $L_{\text{FIR}}/L_{\text{CO}}$  range between roughly 100 and 1000, with a mean value of  $350 L_{\odot} (\text{K km s}^{-1} \text{ pc}^2)^{-1}$ , for galaxies with typical FIR luminosities between  $10^{12}$  and  $10^{13} L_{\odot}$ . For comparison, lower luminosity star-forming galaxies ( $L_{\text{FIR}} < 10^{11} L_{\odot}$ ) have  $L_{\text{FIR}}/L_{\text{CO}}$  ratios an order of magnitude smaller (Kennicutt 1998). J0927+2001 follows this general trend for increasing star formation efficiency with increasing FIR luminosity. These results suggest a relatively short

gas depletion timescale  $\equiv (\text{gas mass})/(\text{star formation rate}) \sim 10^7 \text{ yr}$ , implying a brief but very intense starburst. This timescale is comparable to the typical lifetime of luminous, high-redshift quasars derived from the clustering of quasars in the SDSS by Shen et al. (2007). Of course, these arguments assume the FIR excess is due to dust heated by star formation, and not the AGN.

Although the gas mass in J0927+2001 is comparable to that in J1148+5251, the line width of  $600 \text{ km s}^{-1}$  is almost a factor 2 broader than the CO emission from J1148+5251, and is at the high end of the CO line width distribution observed for high-redshift quasar host galaxies (Carilli & Wang 2006; Greve et al. 2005). A simple explanation for a broader line would be that the gas disk is more inclined to the sky plane in J0927+2001. High-resolution imaging of the CO distribution is required to determine the extent and dynamics of the molecular gas.

#### 4. DISCUSSION

J0927+2001 is the second example of an extreme molecular gas mass in a galaxy within 1 Gyr of the big bang. Like J1148+5251 at  $z = 6.42$ , J0927+2001 shows many characteristics of a coeval massive starburst in the host galaxy of the quasar (§ 1; R. Wang et al. 2007, in preparation). The observations presented herein reveal the requisite molecular gas reservoir to fuel the star formation. As in the case of J1148+5251, Walter et al. (2004) point out that the formation of the heavy elements, and in particular, the ISM processing required to form cooler molecular gas and dust, implies that star formation must have commenced very early in the host galaxy, at least a few hundred million years prior, or  $z > 8$ .

These results lead to the question, how are such massive galaxies and SMBHs formed within 1 Gyr of the big bang? Li et al. (2007a, 2007b) have addressed this question through multiscale cosmological simulations, including prescriptions for the complex processes of star formation and AGN feedback. They find that early galaxy and SMBH formation is possible in rare (comoving density  $\sim 1 \text{ Gpc}^{-3}$ ), high-density peaks (halo mass  $\sim 8 \times 10^{12} M_{\odot}$  at  $z \sim 6$ ) in the cosmic density field, through a series of gas-rich, massive mergers starting at  $z \sim 14$ . SMBH formation occurs through a combination of Eddington-limited accretion in each progenitor galaxy, plus rapid black hole mergers during galaxy interactions. The stellar spheroids are formed in merger-driven, extreme starbursts, with star formation rates exceeding  $10^3 M_{\odot} \text{ yr}^{-1}$  for short periods ( $\sim 10^7 \text{ yr}$ ). The ISM atomic abundances rapidly approach solar in the inner few kpc. Li et al. (2007a) suggest that such systems evolve into massive galaxies at the centers of the densest cluster environments seen today ( $\sim 10^{15} M_{\odot}$ ).

Li et al. (2007b) hypothesize that systems such as J1148+5251 and J0927+2001 may be in a late starburst phase, where the AGN has recently emerged from its dusty shroud and now potentially dominates the dust heating, even for the warm-dust component. The question of the relative contribution of star formation and the AGN to the heating of the warm dust that produces the observed (rest frame) FIR excesses will be discussed in detail in R. Wang et al. (2007, in preparation), and can be addressed through the observations suggested below.

One outstanding issue the Li et al. models do not address is the early formation of dust. Such early dust formation remains a puzzle, since the standard ISM dust formation mechanism, i.e., in the cool winds from evolved low-mass (AGB) stars,

may require timescales longer than the age of the universe at  $z \sim 6$ . One possible solution is dust formation associated with massive star formation (Stratta et al. 2007; Maiolino et al. 2004; Venkatesan et al. 2006; Dwek et al. 2007).

Such early coeval formation of galaxies and SMBHs has interesting implications for the interpretation of the cosmic Stromgren spheres around high-redshift quasars, as inferred from the Gunn-Peterson absorption spectra (Fan et al. 2006c). Masseli et al. (2007) and Lidz et al. (2007) point out that over the lifetime of the system, the integrated star formation and AGN emission may contribute roughly equally to the reionization of the regions immediately surrounding the quasar host galaxy.

The detection of the gas reservoir required to fuel star formation in the J0927+2001 quasar host galaxy is a key step in studying the formation of massive galaxies at very high redshift, but clearly further detailed observations are required to probe the host galaxy and SMBH formation in this system. Critical diagnostic observations include imaging of the CO dynamics to constrain the galaxy gravitational mass, and hence the black hole mass–bulge mass relation at high  $z$  (Shields et al. 2006; Walter et al. 2004), multiwavelength imaging of the dust to help constrain dust heating mechanisms, studies of lower order CO lines to determine gas excitation and total gas mass (Bertoldi et al. 2003b), detection of higher density gas

tracers, such as HCN, to determine the total mass of gas directly associated with star-forming clouds (Gao et al. 2007; Krumholz & Thompson 2007), and detection of the interstellar gas cooling lines, such as [C II] (Maiolino et al. 2005; Iono et al. 2006). The CO detection presented herein provides an accurate redshift for the host galaxy as well as direct evidence for a highly developed ISM, and hence is the first crucial step for future studies. For extreme luminosity objects such as 0927+2001, many of these studies can be performed with long integration times on current instruments, such as the Plateau de Bure Interferometer and Very Large Array. It will take the Atacama Large Millimeter Array, with its more than an order of magnitude increase in sensitivity relative to current millimeter interferometers, to perform similar studies on less extreme systems, such as the Ly $\alpha$  galaxies at  $z \sim 6$ .

The Plateau de Bure Interferometer is a facility of IRAM, supported by INSU/CNRS (France), MPG (Germany), and IGN (Spain). C. C., R. W., and J. W. acknowledge support from the Max Planck Society and the Alexander von Humboldt Foundation through the Max-Planck-Forschungspreise 2005. The National Radio Astronomy Observatory is a facility of the National Science Foundation, operated by Associated Universities, Inc. D. R. acknowledges support from the Deutsche Forschungsgemeinschaft (DFG) Priority Program 1177.

#### REFERENCES

- Beelen, A., et al. 2006, *ApJ*, 642, 694  
 Bertoldi, F., et al. 2003a, *A&A*, 406, L55  
 ———. 2003b, *A&A*, 409, L47  
 Bradford, C. M., et al. 2003, *ApJ*, 586, 891  
 Carilli, C. L., & Wang, R. 2006, *AJ*, 131, 2763  
 Carilli, C. L., et al. 2002, *AJ*, 123, 1838  
 ———. 2004, *AJ*, 128, 997  
 Downes, D., & Solomon, P. 1998, *ApJ*, 507, 615  
 Dwek, E., Galliano, F., & Jones, A. 2007, *ApJ*, 662, 927  
 Fan, X., Carilli, C. L., & Keating, B. 2006a, *ARA&A*, 44, 415  
 Fan, X., et al. 2006b, *AJ*, 131, 1203  
 ———. 2006c, *AJ*, 132, 117  
 Ferrarese, L., & Merritt, D. 2000, *ApJ*, 539, L9  
 Gao, Y., Carilli, C. L., Solomon, P., & Vanden Bout, P. 2007, *ApJ*, 660, L93  
 Gao, Y., & Solomon, P. 2004, *ApJ*, 606, 271  
 Gebhardt, K., Bender, R., & Bower, G. 2000, *ApJ*, 539, L13  
 Greve, T., Papadopoulos, P., Gao, Y., & Radford, S. 2007, *ApJ*, in press  
 Greve, T., et al. 2005, *MNRAS*, 359, 1165  
 Iono, D., et al. 2006, *ApJ*, 645, L97  
 Jiang, L., et al. 2007, *AJ*, submitted  
 Kennicutt, R., Jr. 1998, *ARA&A*, 36, 189  
 Krumholz, M. R., & Thompson, T. A. 2007, *ApJ*, submitted (arXiv:0704.0792)  
 Li, Y., et al. 2007a, *ApJ*, 665, 187  
 ———. 2007b, *ApJ*, submitted (arXiv:0706.3706)  
 Lidz, A., McQuinn, M., Zaldarriaga, M., Hernquist, L., & Dutta, S. 2007, *ApJ*, submitted (astro-ph/0703667)  
 Magorrian, J., et al. 1998, *AJ*, 115, 2285  
 Maiolino, R., et al. 2004, *Nature*, 431, 533  
 ———. 2005, *A&A*, 440, L51  
 Masseli, A., Gallerani, S., Ferrara, A., & Choudhury, T. 2007, *MNRAS*, 376, L34  
 Omont, A., et al. 2003, *A&A*, 398, 857  
 Petric, A. O., et al. 2003, *AJ*, 126, 15  
 Shen, Y., et al. 2007, *AJ*, 133, 2222  
 Shields, G. A., Menezes, K., Massart, C., & Vanden Bout, P. 2006, *ApJ*, 641, 683  
 Solomon, P., Radford, S., Downes, D., & Barrett, J. 1997, *ApJ*, 478, 144  
 Solomon, P., & Vanden Bout, P. 2005, *ARA&A*, 43, 677  
 Stratta, G., Maiolino, R., Fiore, F., & D'Elia, V. 2007, *ApJ*, 661, L9  
 Venkatesan, A., Nath, B. B., & Shull, J. M. 2006, *ApJ*, 640, 31  
 Walter, F., et al. 2003, *Nature*, 424, 406  
 ———. 2004, *ApJ*, 615, L17  
 Wang, R., et al. 2007, *AJ*, 134, 617  
 York, D. G., et al. 2000, *AJ*, 120, 1579