



Detection of Anti-*Neospora caninum* Antibodies in Cattle in Adana Province of Turkey

Funda EŞKİ¹ Armağan Erdem ÜTÜK²

¹ Çukurova University, Faculty of Ceyhan Veterinary Medicine, Department of Obstetrics & Gynecology, Adana, Turkey

² Çukurova University, Faculty of Ceyhan Veterinary Medicine, Department of Parasitology, Adana, Turkey

Received: 12.04.2018

Accepted: 08.06.2018

ABSTRACT

Neospora caninum is an important aborting agent in cattle. In addition to abortions, foetal deaths, resorption, mummification, autolysis, stillbirth, symptomatic or persistently infected calf births, decreased milk production, repeat breeder and premature culling occur in infected herds, which leads to economic losses. The aim of this study was to determine the prevalence of anti-*N. caninum* antibodies in cattle in Adana province of Turkey. For this purpose, 225 blood samples were collected from cattle in different breed, age and sex in 15 counties of Adana. Obtained sera were examined with c-ELISA test. At the end of the study, anti-*N. caninum* antibodies were detected in 24 out of 225 animals (10.7%). While the counties were grouped based on their altitudes (0-99 m, 100-500 m and ≥ 501 m), the cattle were grouped in respect of breeds (Holstein, crossbreed, others), ages (≤ 4 and ≥ 5) and sex (male and female). Correlation between seropositivity and variants (altitude, breed, age and sex) was investigated with chi-square (X^2) test. There was no statistical correlation between seropositivity and altitude, breed and sex ($P > 0.05$). Seropositivity rates were 15.8% at age of ≥ 5 and 6.9% at age of ≤ 4 in cattle. The correlation between age and seropositivity was statistically significant ($P < 0.05$). In this study, we evaluate the exposure rates of cattle to *N. caninum* as well as the correlation between seropositivity and different variants in Adana province of Turkey.

Keywords: *Neospora caninum*, Cattle, c-ELISA, Adana, Turkey

ÖZ

Türkiye'nin Adana Yöresi Sığırlarında Anti-*Neospora caninum* Antikorlarının Araştırılması

Neospora caninum sığırların önemli protozoal abort etkenlerinden biridir. Sığırlarda *Neospora* enfeksiyonlarında abortlara ilave olarak fetal ölüm ve rezorbsiyon, mumifikasyon, otoliz, ölü doğum, canlı ancak klinik semptom gösteren ya da persiste enfekte yavru doğumları şekillenir. Enfeksiyonun görüldüğü sürülerde ayrıca süt veriminde düşme, döl tutamama ve erken damızlıktan ayırma gibi ekonomik kayıplar da ortaya çıkabilir. Bu çalışmanın amacı Adana yöresi sığırlarında anti-*N. caninum* antikorlarının yaygınlığını belirlemektir. Bu amaçla Adana'nın 15 ilçesinden farklı ırk, yaş ve cinsiyette toplam 225 sığırdan kan alındı ve elde edilen serumlar c-ELISA testi ile incelendi. Çalışma sonucunda prevalans %10.7 (24/225) olarak belirlendi. İlçeler rakım özelliklerine göre 0-99 m, 100-500 m ve ≥ 501 m olarak, sığırlar ırk özelliklerine göre Holstein, melez ve diğerleri, yaşlarına göre de ≤ 4 ve ≥ 5 olarak gruplandırıldı. Rakım, ırk, yaş ve cinsiyet değişkenleri ile seropozitiflik arasındaki ilişki ki-kare (X^2) testi ile analiz edildi. Rakım, ırk ve cinsiyet değişkenleri ile seropozitiflik arasında istatistik açıdan bir fark bulunamadı ($P > 0.05$). 5 yaş ve üstü sığırlarda seropozitiflik oranı %15.8 iken, bu oranın 4 yaş ve altındaki sığırlarda %6.9'a düştüğü ve bu değerlerin istatistik açıdan önemli olduğu ($P < 0.05$) tespit edildi. Bu çalışma ile Adana yöresi sığırlarının etkene maruziyet durumları ve farklı değişkenler ile hastalığın yaygınlığı arasındaki ilişki değerlendirilmiştir.

Anahtar Kelimeler: *Neospora caninum*, Sığır, c-ELISA, Adana, Türkiye

INTRODUCTION

Abort in cattle is described as the loss of foetus between 42 and 260 days of gestation (Al-Samarai et al. 2012). Mechanic-traumatic factors, nutritional deficiencies, hormonal disorders and infections (bacteria, viruses and parasites) play important role in abortion etiology (Tulu et al. 2018). *Neospora caninum*, an obligate intracellular

protozoon in the phylum of Apicomplexa, is one of the most important infectious agents in cattle abortion (Schares et al. 1998; Anderson et al. 2000; Haddad et al. 2005). In the domestic cycle of the disease, cattle, sheep, goat, buffalo and horses are intermediate hosts, while dogs are final hosts of *N. caninum* (Dubey 2003; Dubey and Schares 2011).

The disease is transmitted horizontally and vertically between animals (Dubey 2007). Cattle are infected by contaminated food and water with sporulated oocysts in horizontal transmission, and carry the infection to their offsprings by endogenous and exogenous transplacental ways (Anderson et al. 2000). Endogenous transplacental transmission causes persistently infected individuals and expansion of the disease in cattle herds, while exogenous transplacental transmission causes abortion storms (Anderson et al. 2000; Dubey and Schares 2006).

In addition to abortions, foetal deaths, resorption, mummification, autolysis, stillbirth, symptomatic or persistently infected calf births, decreased milk production, repeat breeder and premature culling may occur in infected herds, which causes economic losses (Dubey 2003; Simsek et al. 2008; Piskin and Utuk 2009).

While the disease is a suspected cause of abortion in cattle herds, the final diagnosis is done through the examination of aborted foetus with histopathologic and molecular techniques (PCR, DNA sequencing). The prevalence of the disease in final and intermediate hosts can also be determined through serologic techniques (IFAT, ELISA) (Schares et al. 1999; Dubey 2003; Haddad et al. 2005; Guido et al. 2016).

Our hypothesis was that this study gives information about the seroprevalence of *N. caninum* in the sixth biggest and specific province of Turkey and one of the most detailed study about Neosporosis in Turkey and also reviewed the studies about Neosporosis in Turkey and worldwide.

The aim of this study was to determine the prevalence of anti-*N. caninum* antibodies as well as the correlation between seropositivity and different variants (altitude, breed, age and sex) in cattle in Adana, the sixth largest province of Turkey.

MATERIALS and METHODS

Adana is located to the south of the Taurus Mountains and on both sides of the Seyhan River. The city has major transportation routes connecting Europe to Asia. The altitude in Adana changes between 10 to 1453 m from one county to another. In this province, the Mediterranean climate is found in the lowlands, while the mountainous areas have a continental climate.

A total of 225 blood samples were collected from cattle in different breed, age and sex from 15 counties of Adana. Obtained sera were stored at -20°C until used. While the counties were grouped based on their altitudes (0-99 m, 100-500 m and ≥501 m) (Figure 1) the cattle was grouped in respect of breeds (Holstein, cross-breed, others), ages (≤4 and ≥5) and sex (male and female) (Table 1) Correlation between seropositivity and variants (altitude, breed, age and sex) were investigated with chi-square (X^2) test.




Figure 1. Location of Adana, study areas and their altitudes (Anonymous 2017).

Anti-*N. caninum* antibodies were detected with commercially available c-ELISA test kit (VMRD, USA) according to the manufacturer's recommendation. Percent inhibition values with ≥30 were accepted as positive while <30 accepted as negative.

This study was approved by the Ethics Committee of Adana Veterinary Control Institute (20.09.2016/2833).

RESULTS

At the end of the study, anti-*N. caninum* antibodies were detected in 24 out of 225 animals (10.7%). There was no statistical correlation between seropositivity and altitude, breed and sex ($P>0.05$). Seropositivity rates were 15.8% at age of ≥5 and 6.9% at age of ≤4 in cattle. The correlation between age and seropositivity was statistically significant ($P<0.05$) (Table 1).

Table 1. Number of cattle, seropositivity rates and correlation among different variants

Epidemiologic Data	No. Tested (n)	Seropositivity		P
		n (%)		
Altitude (m)	0-99	79	9 (11.4)	0.858
	100-500	69	8 (11.6)	
	≥501	77	7 (9.1)	
Breed	Holstein	170	20 (11.8)	0.576
	Cross-Breed	34	3 (8.8)	
	Other	21	1 (4.8)	
Age	≤4	130	9 (6.9)	0.033
	≥5	95	15 (15.8)	
Sex	Female	206	22 (10.7)	0.213
	Male	19	2 (10.5)	

Table 2. Serologic studies about *Neospora caninum* in cattle from different parts of the world

Continent	Country	Test	No. tested	No. positive	% positive	Reference	Mean Prevalence (%)	World (%)
Africa	Nigeria	ELISA	174	6	3.4	(Ayinmode et al. 2017)	14.78 (598/4046)	
	Morocco	iELISA	176	15	8.52	(Lucchese et al. 2016)		
	Egypt	iELISA	301	57	18.9	(Fereig et al. 2016)		
	Ethiopia	cELISA	2334	335	13.3	(Asmare et al. 2013)		
	Algerian	IFAT	799	157	19.64	(Ghalmi et al. 2012)		
	Sudan	cELISA	262	28	10.7	(Ibrahim et al. 2012)		
Asia	Iran	ELISA	1500	395	26.33	(Hosseinejad et al. 2017)	16.05 (1732/10787)	
	Iran	ELISA	492	63	12.80	(Gharekhanjani et al. 2014)		
	Iran	ELISA	266	28	10.5	(Nematollahi et al. 2011)		
	Iran	ELISA	237	76	32	(Youssefi et al. 2009)		
	Iran	ELISA	285	36	12.6	(Fard et al. 2008)		
	China	iELISA	510	210	41.2	(Qian et al. 2017)		
	China	ELISA	370	70	18.9	(Xia et al. 2011)		
	China	ELISA	540	72	13.3	(Wang et al. 2010)		
	Indonesia	ELISA	991	165	16.6	(Ichikawa-Seki et al. 2016)		
	Thailand	cELISA	445	52	11.7	(Jittapalapong et al. 2008)		
	Thailand	cELISA	549	30	5.5	(Kyaw et al. 2004)		
	Thailand	IFAT	904	54	6	(Suteeraparp et al. 1999)		
	Japan	IFAT	Dairy 145	29	20	(Koiwai et al. 2005)		
	Japan	IFAT	Beef 65	1	1.5	(Koiwai et al. 2005)		
	Pakistan	cELISA	641	277	43	(Nazir et al. 2013)		
	India	cELISA	427	35	8.2	(Meenakshi et al. 2007)		
	Europe	Slovak	iELISA	490	118	24.1		
Czech Republic		cELISA	546	3	0.5	(Bártová et al. 2015)		
Czech Republic		ELISA/IFAT	407	13	3.19	(Václavek et al. 2003)		
Serbia		cELISA	356	54	15.4	(Kuruca et al. 2013)		
Western Romania		cELISA	376	104	27.7	(Imre et al. 2012)		
Southern Romania		iELISA	258	104	40.31	(Mitrea et al. 2012)		
Romania		ELISA	901	312	34.6	(Gavrea et al. 2011)		
Italy		ELISA	3749	1114	29.7	(Gennero et al. 2007)		
Italy		ELISA	1140	126	11	(Otranto et al. 2003)		
France		ELISA	1924	107	5.6	(Ould-Amrouche et al. 1999)		
Germany		ELISA	Beef herd 106	43	41	(Bartels et al. 2006)		
Germany		ELISA	Dairy herd 100	50	50	(Bartels et al. 2006)		
Netherlands		ELISA	Beef herd 82	58	71	(Bartels et al. 2006)		
Netherlands		ELISA	Dairy herd 108	86	80	(Bartels et al. 2006)		
Spain		ELISA	Beef herd 372	171	46	(Bartels et al. 2006)		
Spain		ELISA	Dairy herd 291	184	63	(Bartels et al. 2006)		
Sweden	ELISA	Dairy herd 112	34	30	(Bartels et al. 2006)			
South America	Brazil	IFAT	75	5	6.7	(Cerqueira-Cézar et al. 2017)	53.76 (4797/18922)	
	Brazil	IFAT	575	559	97.2	(Cerqueira-Cézar et al. 2017)		
	Bahia	IFAT	447	63	14	(Gondim et al. 1999)		
	Goiás	IFAT	456	135	29.6	(Melo et al. 2006)		
	Mato Grosso do Sul	IFAT	2448	449	14.9	(Oshiro et al. 2007)		
	Minas Gerais	IFAT	575	559	97.2	(Guedes et al. 2008)		
	Minas Gerais	IFAT	1204	260	21.6	(Bruhn et al. 2013)		
	Para	IFAT	500	260	52	(Silva et al. 2017)		
	Parana	ELISA	1263	423	33	(Locatelli-Dittrich et al. 2008)		
	Pernambuco	IFAT	469	163	31.7	(Silva et al. 2008)		
	Rio Grande do Sul	IFAT	1549	276	17.8	(Corbellini et al. 2006)		
	Rondonia	IFAT	1011	114	11.2	(Aguiar et al. 2006)		
	Santa Catarina	IFAT	1518	466	30.6	(Fávero et al. 2017)		
	Sao Paulo	IFAT	1027	107	10.4	(Cardosa et al. 2012)		
	Tocantins	IFAT	192	48	25.0	(Martins et al. 2011)		
	Argentina	IFAT	290	59	20.3	(Moore et al. 2003)		
Paraguay	ELISA	879	262	29.8	(Osawa et al. 2002)			
Uruguay	ELISA	4444	589	13.25	(Bañales et al. 2006)			
North America	Mexico	ELISA	596	69	11.6	(García-Vázquez et al. 2009)	35.53 (1416/3985)	
	Mexico	ELISA	187	110	59	(García-Vázquez et al. 2002)		
	Costa Rica	iELISA	2743	1185	43.3	(Romero et al. 2005)		
	Venezuela	ELISA	459	52	11.3	(Lista-Alves et al. 2006)		

Table 3. Serologic studies about *Neospora caninum* in cattle from different parts of Turkey

Region	Province	Test	No. tested	No. positive	% positive	Reference	Mean Prevalence (%)	World (%)
Mediterranean	Burdur	cELISA	400	21	5.3	(Adanir et al. 2015)	5.3 (21/400)	
	Marmara							
	Sakarya	cELISA	92	10	9.2	(Öncel et al. 2003)	11.6 (52/470)	
	Tekirdağ	ELISA	124	7	5.6	(Biyikoğlu et al. 2003)		
	Kırklareli	ELISA	150	15	10	(Biyikoğlu et al. 2003)		
	Bursa	cELISA	104	20	19.23	(Eşki et al. 2016)		
Aegean	Afyonkarahisar	cELISA	485	102	21.03	(Çelik et al. 2013)	21.03 (102/485)	
Central Anatolia	Kayseri	cELISA	186	13	7	(İça et al. 2006)		
	Ankara	ELISA	453	46	10.15	(Vural et al. 2006)		
	Eskişehir	ELISA	387	21	5.43	(Vural et al. 2006)		
	Çankırı	ELISA	418	29	6.93	(Vural et al. 2006)		
	Kırıkkale	ELISA	434	142	32.72	(Vural et al. 2006)		
	Kırşehir	ELISA	409	80	19.55	(Vural et al. 2006)	15.89 (654/4114)	
	Yozgat	ELISA	369	75	20.32	(Vural et al. 2006)		
	Nevşehir	ELISA	392	20	5.10	(Vural et al. 2006)		13.06 (1023/7830)
	Kayseri	ELISA	425	46	10.82	(Vural et al. 2006)		
	Aksaray	ELISA	261	91	34.9	(Öcal et al. 2014)		
	Niğde	cELISA	264	70	26.51	(Karatepe and Karatepe 2016)		
	Kırşehir	cELISA	116	21	18.1	(Yıldız et al. 2017)		
	Eastern Anatolia	Bingöl	cELISA	149	7	4.69	(Aktaş et al. 2005)	
Elazığ		cELISA	120	18	15	(Aktaş et al. 2005)		
Malatya		cELISA	100	4	4	(Aktaş et al. 2005)		
Muş		cELISA	144	7	4.86	(Aktaş et al. 2005)	6.55 (120/1832)	
Kars		ELISA	301	6	2	(Akca et al. 2005)		
Elazığ		cELISA	183	15	8.19	(Şimşek et al. 2008)		
Van		cELISA	450	22	4.88	(Alan et al. 2011)		
Erzurum		cELISA	385	41	10.65	(Balkaya et al. 2012)		
Southeastern Anatolia	Şanlıurfa	cELISA	305	23	7.5	(Sevgili et al. 2005)	7.5 (23/305)	
Black Sea	Samsun	cELISA	224	51	22.7	(Kaya S et al. 2011)	22.7 (51/24)	

DISCUSSION and CONCLUSION

Different studies have been conducted to evaluate the correlation between the seroprevalence of *N.caninum* and several variables such as breed, age, sex, altitude, etc., but have yielded different results. In some studies, a statistically significant correlation is found between the seroprevalence of *N.caninum* and the variants like breed, age, sex and climatic conditions (Guimarães et al. 2004; Jittapalpong et al. 2008; Asmare et al. 2013; Nazir et al. 2013; Macedo et al. 2017; Yıldız et al. 2017), while others indicate the opposite (Aktaş et al. 2005; Simsek et al. 2008; İbrahim et al. 2012; Imre et al. 2012; Fereig et al. 2016; Gharekhani et al. 2014; Adanir et al. 2015). In this study, no correlation was found between the seropositivity and the variants like breed, sex, and altitude ($P>0.05$). However, the rate of seropositivity in cattle aged ≥ 5 was 15.8%, while it dropped to 6.9% in cattle aged ≤ 4 , and these values were found to be statistically significant ($P<0.05$). The reason for this is considered to be the host-parasite interaction in cattle, increasing with age.

When the seroprevalence values in different studies from around the world are considered, it can be observed that there exists a dramatic fluctuation in respect of the serological methods used, the differences in cut-off values even when the same method is employed, sampling methods as well as sample size (Wapenaar et al. 2007; Guido et al. 2016; Cerqueira-Cézar et al. 2017).

For instance, in a study conducted in 2003 in the Minas Gerais region of Brazil (Ragozo et al. 2003), the rate of positivity was found to be 11.1% (4/36) at 1/25 titer by IFA test, while another study in 2008 reported the rate of

positivity in the same region as 97.2% (559/575) at 1/200 titer with the use of same test (Guedes et al. 2008). Another study in Brazil in 2004 at Mato Grosso do Sul region (Andreotti et al. 2004) reported the seropositivity as 7.7% (7/91) with ELISA, while in another study in 2010 in the same region (Andreotti et al. 2010), the seropositivity was found to be 62.5% (687/1098) at 1/50 titre by IFA. When the studies in Turkey are concerned, a study in 2013 in Afyon province reported the rate of seropositivity as 21.03% (102/485) tested by c-ELISA (Çelik et al. 2013), while, in another study in 2016 in the same province, the rate of seropositivity was detected to be 3.8% (3/80) by in house ELISA (Zhou et al. 2016).

In the studies performed to find the seroprevalence of *N.caninum* in a specific region, it is significant to ensure the appropriate sample size and its representation of the region, to use same testing methods and cut-off values in regional and nation-wide studies as well as to consider such factors as species, breed, age, sex and various geographical parameters to reach accurate epidemiological data and for the control of the disease. In this study, a total of 225 cattle from 15 counties in Adana, with various climatic and geographic conditions, were grouped in respect of breed, age and sex; whilst the counties were grouped based on altitude. At the end of the study, through face-to-face interviews, the cattle owners were informed of the disease and its effect on their herds as well as the ways for its prevention and control.

The main purpose of cattle breeding is to obtain commercially valuable products such as meat, milk and leather from animals, and to get regularly one calf every year to ensure the continuity of the herds. Increasing feed and treatment costs affect farming negatively; and

sometimes, income from meat and milk barely covers the basic expenses. In such cases, the largest gain of the business is the offspring born healthy (Alpan 1994). In this context, abortions have become one of the most significant economic problems of the breeder (Juyal et al. 2011).

Nowadays, *N.caninum* is among the most important aborting agents (Hosseinnejadet al. 2017). Serologic studies (c-ELISA, i-ELISA, IFAT) indicate that the prevalence of the disease is 11.3-59% in North America, 6.7-97.2% in South America, 1.5-43% in Asia, 0.5-80% in Europe and 3.4-19.64% in Africa (Table 2). Seroprevalence of *N.caninum* Turkey, on the other hand, was determined as 2-34.9% with ELISA test (Table 3).

According to the literature we examined, the mean seroprevalence of *N.caninum* is 13.06% (1023/7830) in Turkey (Table 3), 35.53% (1416/3985) in North America, 53.76% (4797/18922) in South America, 16.05% (1732/10787) in Asia, 23.69% (2681/11318) in Europe and 14.78% (598/4046) in Africa. The overall assessment of the mean seroprevalence in the world can be determined as 22.88% (11224/49058) (Table 2). In this study, we saw that the mean seroprevalence of Turkey, as a bridge between Asia, Europe and Africa, is lower than the mean seroprevalence of other continents and the world. At the end of the study, anti-*N.caninum* antibodies were detected in 10.7% of cattle in Adana province, and this rate is lower than the mean seroprevalence of Turkey and the world.

ACKNOWLEDGEMENT

The authors extended thanks to Scientific Research Coordination Unit of Cukurova University as a Project (TSA-2017-9127) and a part of this study was orally presented in International Mediterranean Science and Engineering Congress (IMSEC 2017).

REFERENCES

- Adanir R, Çetin Y, Kocamüftüoğlu M, Köse O (2015). Seroprevalence of *Neospora caninum* in cows in Burdur Region. Investigation of its relationship with abortions and infertility. Proc XVII International Congress on Animal Hygiene", Košice, Slovakia, 7-11 June, 2015, pp.223-224.
- Aguiar DM, Cavalcante GT, Rodrigues AAR, Labruna MB, Camargo LMA, Camargo EP, Gennari SM (2006). Prevalence of anti-*Neospora caninum* antibodies in cattle and dogs from Western Amazon, Brazil, in association with some possible risk factors. *Vet Parasitol*, 142, 71-77.
- Akca A, Gokce HI, Guy CS, McGarry JW, DJ Williams (2005). Prevalence of antibodies to *Neospora caninum* in local and imported cattle breeds in the Kars Province of Turkey. *Res Vet Sci*, 78, 123-126.
- Aktaş M, Ecmel ÇŞ, Altay K, Şimşek S, Ütük AE, Köroğlu E, Dumanlı N (2005). Doğu Anadolu Bölgesi'nin bazı illerinde bulunan sığırlarda *Neospora caninum*'un araştırılması. *Türkiye Parazitoloj Derg*, 29, 22-25.
- Alan M, Çetin Y, Şendağ S, Akkan HA, Karaca M (2011). Seroprevalence of antibodies against *Neospora caninum* in cows in Van Province. *Kafkas Univ Vet Fak*, 17, 767-771.
- Alpan O (1994). Sığır yetiştiriciliği ve besiciliği. 4. Basım, Şahin Matbaası, Ankara.
- Al-Samarai FR, Abdulrahman YK, Ibrahim WI, Al-Nedawi AM (2012). Effect of some environmental factors on abortion and offspring sex ratio in holstein cows in Iraq. *I J S N*, 3, 361-365.
- Anderson ML, Andrianarivo AG, Conrad PA (2000). Neosporosis in cattle. *Anim Reprad Sci*, 60-61, 417-431.
- Andreotti R, Barros JC, Pereira AR, Oshiro LM, Cunha RC, Figueiredo LFN (2010). Association between seropositivity for *Neospora caninum* and reproductive performance of beef heifers in the Pantanal of Mato Grosso do Sul, Brazil. *Rev Bras Parasitol Vet*, 19, 119-123.
- Andreotti R, Pinckney RD, Pires PP, Silva EAE (2004). Evidence of *Neospora caninum* in beef cattle and dogs in the state of Mato Grosso do Sul, center-western region, Brazil. *Rev Bras Parasitol Vet*, 13, 129-131.
- Anonymous (2017). Map of Adana. Available at http://cografyaharita.com/haritalarim/4l_adana_ili_haritasi.png (accessed December 21 2017, 12.50 pm).
- Asmarek K, Regassa F, Robertson LJ, Skjerve E (2013). Seroprevalence of *Neospora caninum* and associated risk factors in intensive or semi-intensively managed dairy and breeding cattle of Ethiopia. *Vet Parasitol*, 193, 85-94.
- Ayinmode A, Akinseye V, Schares G, Cadmus S (2017). Serological survey of toxoplasmosis, neosporosis and brucellosis among cattle herds in Oyo State, South-Western Nigeria. *Afr J Infect Dis*, 11, 95-101.
- Balkaya I, Bastem Z, Avcioglu H, Onalan SK (2012). Seroprevalence of *Neospora caninum* Antibodies in Cattle in Eastern Turkey. *Isr J Vet Med*, 67, 109-112.
- Bañales P, Fernandez L, Repiso MV, Gil A, Dargatz DA, Osawa T (2006). A nationwide survey on seroprevalence of *Neospora caninum* infection in beef cattle in Uruguay. *Vet Parasitol*, 139, 15-20.
- Bartels CJM, Arnaiz-Seco JI, Ruiz-Santa-Quitera A, Björkman C, Frössling J, von Blumröder D, Conraths FJ, Schares G, van Maanen C, Wouda W, Ortega-Mora LM (2006). Supranational comparison of *Neospora caninum* seroprevalences in cattle in Germany, The Netherlands, Spain and Sweden. *Vet Parasitol*, 137, 17-27.
- Bártová E, Sedlák K, Budíková M (2015). A study of *Neospora caninum* and *Toxoplasma gondii* antibody seroprevalence in healthy cattle in the Czech Republic. *Ann Agric Environ Med*, 22, 32-34.
- Bıyıkoğlu G, Öncel T, Bağcı Ö (2003). Trakya sığırlarında *Neospora caninum*'un seroprevalansı. "13. Ulusal Parazitoloji Kongresi", 8-12 Eylül, Konya, Türkiye.
- Bruhn FRP, Daher DO, Lopes E, Barbieri JM, Rocha CMBM, Guimarães AM (2013). Factors associated with seroprevalence of *Neospora caninum* in dairy cattle in southeastern Brazil. *Trop Anim Health Prod*, 45, 1093-1098.
- Cardoso JMS, Amaku M, Araújo AJUS, Gennari SM (2012). A longitudinal study of *Neospora caninum* infection on three dairy farms in Brazil. *Vet Parasitol*, 187, 553-557.
- Çelik HA, Kozan E, Eser M, Yılmaz O, Birdane MK, Sarımehtemetoğlu HO (2013). A research on seroprevalence of *Neospora caninum* in cattle. *Ankara Üniv Vet Fak Derg*, 60, 99-102.
- Cerqueira-Cézar CK, Calero-Bernal R, Dubey JP, Gennari SM (2017). All about neosporosis in Brazil. *Rev Bras Parasitol Vet*, 26, 253-279.
- Corbellini LG, Smith DR, Pescador CA, Schmitz M, Correa A, Steffen DJ, Driemeier D (2006). Herd-level risk factors for *Neospora caninum* seroprevalence in dairy farms in southern Brazil. *Prev Vet Med*, 74, 130-141.
- Dubey JP (2003). Review of *Neospora caninum* and neosporosis in animals. *Korean J Parasitol*, 41, 1-16.
- Dubey JP, Schares G (2006). Diagnosis of bovine neosporosis. *Vet Parasitol*, 140, 1-34.
- Dubey JP, Schares G (2011). Neosporosis in animals - The last five years. *Vet Parasitol*, 180, 90-108.
- Dubey JP, Schares G, Ortega-Mora LM (2007). Epidemiology and control of Neosporosis and *Neospora caninum*. *Clin Microbiol Rev*, 20, 323-367.
- Eşki F, Önat K, Günaydın E, Pekmaya S, Çetin N, Ütük AE (2016). Detection of *Neospora caninum*, *Toxoplasma gondii*, *Chlamydomphila abortus* and *Coxiella burnetti* antibodies in aborted Holstein Cows. "1st International Mediterranean Science and Engineering Congress", 26-28 October, Adana, Turkey, p. 4550.
- Fard SRN, Khalili M, Aminzadeh A (2008). Prevalence of antibodies to *Neospora caninum* in cattle in Kerman province, South East Iran. *Vet Arhiv*, 78, 253-259.
- Fávero JF, Silva AS, Campigotto G, Machado G, Barros LD, Garcia JL, Vogel FF, Mendes RE, Stefani LM (2017). Risk factors for *Neospora caninum* infection in dairy cattle and their possible cause-effect relation for disease. *Microb Pathog*, 110, 202-207.
- Fereig RM, AbouLaila MR, Mohamed SGA, Mahmoud HYAH, Ali AO, Ali AF, Hilali M, Zaide A, Mohamed AEA, Nishikawa Y (2016). Serological detection and epidemiology of *Neospora caninum* and *Cryptosporidium parvum* antibodies in cattle in southern Egypt. *Acta Tropica*, 162, 206-211.
- García-Vázquez Z, Cruz-Vázquez C, Medina-Espinoza L, García-Tapia D, Chavarria-Martinez B (2002). Serological survey of *Neospora caninum* infection in dairy cattle herds in Aguascalientes, Mexico. *Vet Parasitol*, 106, 115-120.

- García-Vazquez Z, Rosario-Cruz R, Mejía-Estrada F, Rodríguez-Vivas I, Romero-Salas D, Fernandez-Ruvalcaba M, Cruz-Vazquez C (2009). Seroprevalence of *Neospora caninum* antibodies in beef cattle in three southern states of Mexico. *Trop Anim Health Prod*, 41, 749-753.
- Gavrea RR, Iovu A, Losson B, Cozma V (2011). Seroprevalence of *Neospora caninum* in dairy cattle from north-west and centre of Romania. *Parasite*, 18, 349-351.
- Gennero MS, Bergagna S, Pasino M, Barbaro A, Romano A, Trisciuglio A, Ezio F (2007). *Neospora caninum* serological survey in cattle from the piedmont region (NORTHWESTERN ITALY). *Rev Epid San Anim*, 51, 65-68.
- Ghalmi F, China B, Ghalmi A, Hammitouche D, Losson B (2012). Study of the risk factors associated with *Neospora caninum* seroprevalence in Algerian cattle populations. *Res Vet Sci*, 93, 655-661.
- Gharekhani J, Haddadzadeh H, Bahonar A (2014). Prevalence of immunoglobulin G (IgG) antibody to *Neospora caninum* in dairy cattle of Hamedan province, west of Iran. *Vet Res Forum*, 5, 149-152.
- Gondim LFP, Sartor IF, Monteiro LA, Haritani M (1999). *Neospora caninum* infection in an aborted bovine foetus in Brazil. *New Zeal Vet J*, 47, 35.
- Guedes MHP, Guimarães AM, Rocha CMBM, Hirsch C (2008). Frequência de anticorpos anti-*Neospora caninum* em vacas e fetos provenientes de municípios do sul de Minas Gerais. *Rev Bras Parasitol Vet*, 17, 189-194.
- Guido S, Katzer F, Nanjiani I, Milne E, Innes EA (2016). Serology-based diagnostics for the control of bovine neosporosis. *Trends Parasitol*, 32, 131-143.
- Guimarães JS, Souza SLP, Bergamaschi DP, Gennari SM (2004). Prevalence of *Neospora caninum* antibodies and factors associated with their presence in dairy cattle of the north of Paraná state, Brazil. *Vet Parasitol*, 124, 1-8.
- Haddad JPA, Dohoo IR, VanLeewen JA (2005). A review of *Neospora caninum* in dairy and beef cattle - a Canadian perspective. *Can Vet J*, 46, 230-243.
- Hosseininejad M, Mahzounieh M, Esfandabadi NS (2017). *Neospora caninum* suspects as one of the most important causes of abortion in large dairy farms in Isfahan, Iran. *Iran J Parasitol*, 12, 408-412.
- Ibrahim AME, Elfahal AM, Hussein ARME (2012). First report of *Neospora caninum* infection in cattle in Sudan. *Trop Anim Health Prod*, 44, 769-772.
- İça A, Yildirim A, Düzlü Ö, İnci A (2006). Seroprevalence of *Neospora caninum* in cattle in the region of Kayseri. *Türkiye Parazitoloj Derg*, 30, 92-94.
- Ichikawa-Seki M, Guswanto A, Allamanda P, Mariamah ES, Wibowo PE, Nishikawa Y (2016). Seroprevalence of antibody to NcSAG1 antigen of *Neospora caninum* in cattle from Western Java, Indonesia. *J Vet Med Sci*, 78, 121-123.
- Imre K, Morariu S, Ilie MS, Imre M, Ferrari N, Genchi C, Darabus G (2012). Serological survey of *Neospora caninum* infection in cattle herds from western Romania. *J Parasitol*, 98, 683-685.
- Jittapalapong S, Sangwaranond A, Inpankaew T, Phasuk C, Pinyopanuwat N, Chimnoi W, Kengradomkij C, Saengow S, Pumphom P, Arunwipat P, Anakewit T, Robertson ID (2008). Seroprevalence of *Neospora caninum* infections of dairy cows in the North-east of Thailand. *Kasetsart Journal (Nat Sci)*, 42, 61-66.
- Juyal PD, Bal MS, Singla LD (2011). Economic impact, diagnostic investigations and management of protozoal abortions in farm animals. *All India SMVS Dairy Business Directory* 11, 39-46.
- Karatepe B, Karatepe M (2016). Seroprevalence of *Neospora caninum* in cattle in Nigde Province, Turkey. *Isr J Vet Med*, 71, 39-42.
- Kaya S, Kurt M, Mustafa A, Cenk SB, Ali TG, Şinasi U (2011). Samsun yöresinde brucellosis yönünden negatif olan sığırlarda *Neospora caninum* seropozitifliği. "17th National Parasitology Congress andCaucasian and Middle East Symposium on Parasitic Diseases"4-10 September, Kars, Turkey.
- Koiwai M, Hamaoka T, Haritani M, Shimizu S, Tsutsui T, Eto M, Yamane I (2005). Seroprevalence of *Neospora caninum* in dairy and beef cattle with reproductive disorders in Japan. *Vet Parasitol*, 130, 15-18.
- Koiwai M, Hamaoka T, Haritani M, Shimizu S, Zeniya Y, Eto M, Yokoyama R, Tsutsui T, Kimura K, Yamane I (2006). Nationwide seroprevalence of *Neospora caninum* among dairy cattle in Japan. *Vet Parasitol*, 135, 175-179.
- Kuruca L, Spasojević-Kosić L, Simin S, Savović M, Laus S, Lalosević V (2013). *Neospora caninum* antibodies in dairy cows and domestic dogs from Vojvodina, Serbia. *Parasite*, 20, 40.
- Kyaw T, Virakul P, Muangyai M, Suwimonteerabutr J (2004). *Neospora caninum* seroprevalence in dairy cattle in central Thailand. *Vet Parasitol*, 121, 255-263.
- Lista-Alves D, Palomares-Naveda R, Garcia F, Obando C, Arrieta D, Hoet AE (2006). Serological evidence of *Neospora caninum* in dual-purpose cattle herds in Venezuela. *Vet Parasitol*, 136, 347-349.
- Locatelli-Dittrich R, Machado PC, Fridlund-Plugge N, Richartz RRTB, Montiani-Ferreira F, Patrício LFL, Patrício MAC, Joineau MG, Pieppe M (2008). Determinação e correlação de anticorpos anti-*Neospora caninum* em bovinos e cães do Paraná, Brasil. Determination and correlation of anti-*Neospora caninum* antibodies in cattle and dogs from Paraná, Brazil]. *Rev Bras Parasitol Vet*, 17, 191-196.
- Luchese L, Benkirane A, Hakimi I, Idrissi AE, Natale A (2016). Seroprevalence study of the main causes of abortion in dairy cattle in Morocco. *Vet Ital*, 52, 13-19.
- Macedo CAB, Macedo MFSB, Miura AC, Taroda A, ST Cardim, Innes AE, Katzer F, Cantón G, Chianini F, Headley SA, Garcia JL (2017). Occurrence of abortions induced by *Neospora caninum* in dairy cattle from Santa Catarina, southern Brazil. *Rev Bras Parasitol Vet*, 26, 292-298.
- Martins NÉX, Freschi CR, Baptista F, Machado RZ, Freitas FLC, Almeida KS (2011). Ocorrência de anticorpos anti-*Neospora caninum* em vacas lactantes do município de Araguaína, estado do Tocantins, Brasil. *Rev Patol Trop*, 40, 231-238.
- Meenakshi SKS, Ball MS, Kumar H, Sharma S, Sidhu PK, Sreekumar C, JP Dubey (2007). Seroprevalence of *Neospora caninum* antibodies in cattle and water buffaloes in India. *J Parasitol*, 93, 1374-1377.
- Melo DPG, Silva AC, Ortega-Mora LM, Bastos SA, Boaventura CM (2006). Prevalência de anticorpos anti-*Neospora caninum* em bovinos das microrregiões de Goiânia e Anápolis, Goiás, Brasil. [Prevalence of antibodies anti-*Neospora caninum* in bovines from Anápolis and Goiânia micro regions, Goiás, Brazil]. *Rev Bras Parasitol Vet*, 15, 105-109.
- Mitreă IL, Enăchescu V, Radulescu R, Ionita M (2012). Seroprevalence of *Neospora caninum* infection on dairy cattle in farms from southern Romania. *J Parasitol*, 98, 69-72.
- Moore D, Reichel M, Spath E, Campero C (2013). *Neospora caninum* causes severe economic losses in cattle in the humid pampa region of Argentina. *Trop Anim Health Prod*, 45, 1237-1241.
- Nazir MM, Maqbool A, Khan MS, Sajjid A, Lindsay DS (2013). Effects of age and breed on the prevalence of *Neospora caninum* in commercial dairy cattle from Pakistan. *J Parasitol*, 99, 368-370.
- Nematollahi A, Jaafari R, Moghaddam GH (2011). Seroprevalence of *Neospora caninum* infection in dairy cattle in Tabriz, Northwest Iran. *Iran J Parasitol*, 6, 95-98.
- Öcal N, Atmaca HT, Albay MK, Deniz A, Kalender H, Yıldız K, Kul O (2014). A new approach to *Neospora caninum* infection epidemiology. neosporosis in integrated and rural dairy farms in Turkey. *Turk. J Vet Anim Sci*, 38, 161-168.
- Öncel T, Biyikoğlu G (2003). *Neosporosis caninum* in dairy cattle in Sakarya, Turkey. *Uludağ Univ Vet Fak Derg*, 22, 87-89.
- Osawa T, Wastling J, Acosta L, Ortellado C, Ibarra J, Innes EA (2002). Seroprevalence of *Neospora caninum* infection in dairy and beef cattle in Paraguay. *Vet Parasitol*, 110, 17-23.
- Oshiro LM, Matos MFC, Oliveira JM, Monteiro LARC, Andreotti R (2007). Prevalence of anti-*Neospora caninum* antibodies in cattle from the state of Mato Grosso do Sul, Brazil. *Rev Bras Parasitol Vet*, 16, 133-138.
- Otranto D, Iazari AL, Testini G, Traversa D, di Regalbano AF, Badan M, Capelli G (2003). Seroprevalence and associated risk factors of neosporosis in beef and dairy cattle in Italy. *Vet Parasitol*, 118, 7-18.
- Ould-Amrouche A, Klein F, Osdoit C, Mohamed HO, Touratier A, Sanaa M, Mialot JP (1999). Estimation of *Neospora caninum* seroprevalence in dairy cattle from Normandy, France. *Vet Res*, 30, 531-538.
- Piskin FC, Utuk AE (2009). Seroprevalence of *Neospora caninum* in cows with stillbirth and abortion. *J Etlik Vet Microbiol*, 20, 23-26.
- Qian W, Wang T, Yan W, Zhang M, Han L, Xue R, Song S, Lv C (2017). Seroprevalence and first multilocus microsatellite genotyping of *Neospora caninum* in dairy cattle in Henan, central China. *Vet Parasitol*, 244, 81-84.
- Ragozo AMAR, Paula VSO, Souza SLP, Bergamaschi DP, Gennari SM (2003). Ocorrência de anticorpos anti-*Neospora caninum* em soros bovinos procedentes de seis estados brasileiros. *Rev Bras Parasitol Vet*, 12, 33-37.
- Romero JJ, Van Breda S, Vargas B, Dolz G, Frankena K (2005). Effect of neosporosis on productive and reproductive performance of dairy cattle in Costa Rica. *Theriogenology* 64, 1928-1939.
- Schares G, Conraths FJ, Reichel MP (1999). Bovine neosporosis. comparison of serological methods using outbreak sera from a dairy herd in New Zealand. *Int J Parasitol*, 29, 1659-1667.

- Schares G, Peters M, Wurm R, Bärwald A, Conraths FJ (1998).** The efficiency of vertical transmission of *Neospora caninum* in dairy cattle analysed by serological techniques. *Vet Parasitol*, 80, 87-98.
- Sevgili M, Altaş MG, Keskin O (2005).** Seroprevalence of *Neospora caninum* in Cattle in the Province of Şanlıurfa. *Turk J Vet Anim Sci*, 29, 127-130.
- Silva JB, Nicolino RR, Fagundes GM, Bomjardim HA, Reis ASB, Lima DHS, Oliveira CMC, Barbosa JD, Fonseca AH (2017).** Serological survey of *Neospora caninum* and *Toxoplasma gondii* in cattle (*Bos indicus*) and water buffaloes (*Bubalus bubalis*) in ten provinces of Brazil. *Comp Immunol Microbiol Infect Dis*, 52, 30-35.
- Silva MIS, Almeida MÃO, Mota RA, Pinheiro-Junior JW, Rabelo SSA (2008).** Fatores de riscos associados à infecção por *Neospora caninum* em matrizes bovinas leiteiras em Pernambuco. [Risk factors associated to *Neospora caninum* infection in dairy cows in Pernambuco]. *Ci Anim Bras*, 9, 455-461.
- Simsek S, Utuk AE, Koroglu E, Dumanlı N, Risvanli A (2008).** Seroprevalance of *Neospora caninum* in repeat breeder dairy cows in Turkey. *Arch Tierz Dummerstorf*, 51, 143-148.
- Špilovská S, Reiterová K, Antolová D (2015).** *Neospora caninum*-associated abortions in Slovak dairy farm. *Iran J Parasitol*, 10, 96-101.
- Suteeraparp P, Pholpark S, Pholpark M, Charoenchai A, Chompoochan T, Yamane I, Kashiwazaki Y (1999).** Seroprevalence of antibodies to *Neospora caninum* and associated abortion in dairy cattle from central Thailand. *Vet Parasitol*, 86, 49-57.
- Tulu D, Deresa B, Begna F, Gojam A (2018).** Review of common causes of abortion in dairy cattle in Ethiopia. *J Vet Med Anim Health*, 10, 1-13.
- Václavěk P, Koudela B, Modrý D, Sedlák K (2003).** Seroprevalence of *Neospora caninum* in aborting dairy cattle in the Czech Republic. *Vet Parasitol*, 115, 239-245.
- Vural G, Aksoy E, Bozkir M, Kuçukayan U, Erturk A (2006).** Seroprevalence of *Neospora caninum* in dairy cattle herds in Central Anatolia, Turkey. *Vet Arhiv*, 76, 343-349.
- Wang C, Wang Y, Zou X, Zhai Y, Gao J, Hou M, Zhu XQ (2010).** Seroprevalence of *Neospora caninum* infection in dairy cattle in Northeastern China. *J Parasitol*, 96, 451-452.
- Wapenaar W, Barkema HW, VanLeeuwen JA, McClure JT, O'Handley RM, Kwok OCH, Thulliez P, Dubey JP, Jenkins MC (2007).** Comparison of serological methods for the diagnosis of *Neospora caninum* infection in cattle. *Vet Parasitol*, 143, 166-173.
- Xia HY, Zhou DH, Jia K, Zeng XB, Zhang DW, She LX, Lin RQ, Yuan ZG, Li SJ, Zhu XQ (2011).** Seroprevalence of *Neospora caninum* infection in dairy cattle of Southern China. *J Parasitol*, 97, 172-173.
- Yıldız K, Gökpınar S, Sürsal N, Değirmenci R (2017).** Kırşehir İli Çiçekdağı İlçesi'nde Yetiştirilen Süt İneklerinde *Neospora caninum*'ün Seroprevalansı. *Türkiye Parazitoloj Derg*, 41, 135-138.
- Youssefi MR, Arabkhazaeli F, Hassan ATM (2009).** Seroprevalence of *Neospora caninum* Infection in Rural and Industrial Cattle in Northern Iran. *Iran J Parasitol*, 4, 15-18.
- Zhou M, Cao S, Sevinc F, Sevinc M, Ceylan O, Liu M, Wang G, Mounouni PFA, Jirapattharasate C, Suzuki H, Nishikawa Y, Xuan X (2016).** Enzyme-linked immunosorbent assays using recombinant TgSAG2 and NcSAG1 to detect *Toxoplasma gondii* and *Neospora caninum*-specific antibodies in domestic animals in Turkey. *J Vet Med Sci*, 78, 1877-1881.