

Detection of *Campylobacter jejuni* in rectal swab samples from *Rousettus amplexicaudatus* in the Philippines

Yuki HATTA¹), Tsutomu OMATSU¹)*, Shinobu TSUCHIAKA¹), Yukie KATAYAMA¹), Satoshi TANIGUCHI²), Joseph S MASANGKAY³), Roberto PUENTESPINA Jr⁴), Eduardo ERES⁵), Edison COSICO⁵), Yumi UNE⁶), Yasuhiro YOSHIKAWA⁷), Ken MAEDA⁸), Shigeru KYUWA⁹)* and Tetsuya MIZUTANI¹)

¹)Research and Education Center for Prevention of Global Infectious Diseases of Animals, Tokyo University of Agriculture and Technology, 3-5-8 Saiwai-cho, Fuchu-shi, Tokyo 183-8509, Japan

²)Special Pathogens Laboratory, Department of Virology I, National Institute of Infectious Diseases, 4-7-1 Gakuen, Musashimurayama, Tokyo 208-0011, Japan

³)College of Veterinary Medicine, University of the Philippines Los Baños, 4031 Los Baños, Laguna, Philippines

⁴)University of the Philippines Mindanao, Bago-Oshiro, Davao, Philippines

⁵)Museum of Natural History, University of the Philippines Los Baños, 4031 Los Baños, Laguna, Philippines

⁶)Laboratory of Veterinary Pathology, School of Veterinary Medicine, Azabu University, 1-17-1 Fuchinobe, Sagami-hara, Kanagawa 252-5201, Japan

⁷)Department of Animal Risk Management, Chiba Institute of Science, Choshi, Chiba 288-0025, Japan

⁸)Laboratory of Veterinary Microbiology, Joint Faculty of Veterinary Medicine, Yamaguchi University, 1677-1 Yoshida, Yamaguchi 753-8515, Japan

⁹)Department of Biomedical Science, Graduate School of Agricultural and Life Sciences, The University of Tokyo, 1-1-1 Yayoi, Bunkyo-ku, Tokyo 113-8657, Japan

(Received 30 October 2015/Accepted 14 April 2016/Published online in J-STAGE 25 April 2016)

ABSTRACT. Bats are the second diversity species of mammals and widely distributed in the world. They are thought to be reservoir and vectors of zoonotic pathogens. However, there is scarce report of the evidence of pathogenic bacteria kept in bats. The precise knowledge of the pathogenic bacteria in bat microbiota is important for zoonosis control. Thus, metagenomic analysis targeting the V3-V4 region of the 16S rRNA of the rectal microbiota in *Rousettus amplexicaudatus* was performed using high throughput sequencing. The results revealed that 103 genera of bacteria including *Campylobacter* were detected. *Campylobacter* was second predominant genus, and *Campylobacter coli* and *Campylobacter jejuni* were identified in microbiome of *R. amplexicaudatus*. *Campylobacteriosis* is one of the serious bacterial diarrhea in human, and the most often implicated species as the causative agent of campylobacteriosis is *C. jejuni*. Therefore, we investigated the prevalence of *C. jejuni* in 91 wild bats with PCR. As a result of PCR assay targeted on 16S-23S intergenic spacer, partial genome of *C. jejuni* was detected only in five *R. amplexicaudatus*. This is the first report that *C. jejuni* was detected in bat rectal swab samples. *C. jejuni* is the most common cause of campylobacteriosis in humans, transmitted through water and contact with livestock animals. This result indicated that *R. amplexicaudatus* may be a carrier of *C. jejuni*.

KEY WORDS: 16S rRNA, bat, *Campylobacter*, metagenome, microbiota

doi: 10.1292/jvms.15-0621; *J. Vet. Med. Sci.* 78(8): 1347-1350, 2016

Bats are the second diversity species of mammals in the world, consisted of approximately 1,100 species [25]. Bats have a variety of food choices, including fruits, insects, small vertebrate and blood. Furthermore, bats are known as reservoir hosts and vectors of zoonotic pathogens. Previous and ongoing researches in bats mainly focus on viruses,

including Rabies virus [9], Nipah virus [5, 6] and Hendra virus [18] as well as European and Australian bat lyssaviruses [3]. In contrast, the knowledge of pathogenic bacteria in bats is scarce. Only a few bacteria pathogenic to humans, such as *Salmonella* spp. [23] and *Clostridium* spp. [14], were isolated from bats. Therefore, it is important to characterize rectal microbiota and identify pathogenic agents in bats from the viewpoint of public health.

Gut microbiota has been investigated and characterized by a variety of methods. These methods include culture-based analysis, DNA sequencing using the Sanger method and fluorescence *in situ* hybridization targeting the 16S rRNA gene [10]. Most of the previous approaches are limited in scope. Traditional DNA-based approaches have targeted phylogenetic genes, which provide limited information, especially when a large amount of microbial groups must be classified.

The number of metagenomic studies has increased in recent years, because of the availability of high throughput sequencing technologies. High throughput sequencing al-

*CORRESPONDENCE TO: OMATSU, T., Research and Education Center for Prevention of Global Infectious Diseases of Animals, Tokyo University of Agriculture and Technology, Saiwai-cho, Fuchu-shi, Tokyo 183-8509, Japan. e-mail: tomatsu@cc.tuat.ac.jp
KYUWA, S., Department of Biomedical Science, Graduate School of Agricultural and Life Sciences, The University of Tokyo, 1-1-1 Yayoi, Bunkyo-ku, Tokyo 113-8657, Japan.
e-mail: akyuwa@mail.ecc.u-tokyo.ac.jp

©2016 The Japanese Society of Veterinary Science

This is an open-access article distributed under the terms of the Creative Commons Attribution Non-Commercial No Derivatives (by-nc-nd) License <<http://creativecommons.org/licenses/by-nc-nd/4.0/>>.

lows the production of millions of short sequence reads in a single run and do not require additional cloning steps like the traditional Sanger sequencing [17]. The enormous amount of data collected by high throughput sequencing indicates that a variety of microbiota has not been analyzed previously [16]. High throughput sequencing technologies have proven usefulness for studying the diversity and dynamics of bacterial species, even in complex systems like the gut [12].

Despite of the importance from the viewpoint of public health, little information on bat microbiota is available. In this study, we performed metagenomic analysis of the 16S rRNA of rectal microbiota in wild bats using high throughput sequencing to detect pathogenic bacteria.

A total of 88 megachiropteras, comprised of 67 *Rousetus amplexicaudatus*, 14 *Eonycteris spelaea*, 5 *Cynopterus brachyotis* and 2 *Macroglossus minimus*, and 3 microchiropteras comprised of 3 *Hipposideros diadema* were collected from five sites located in the Davao region of the Philippines: Dadatan (6°56'N; 125°40'E), Guilon (7°03'N; 125°44'E), Lavigan (6°16'N; 126°11'E), the forest of Davao (7°11'N; 125°25'E) and Sion (6°56'N; 125°45'E). Bats were captured with the permission of the regional executive director of the Department of Environment and Natural Resources (DENR).

Rectal swab samples were collected under anesthesia and suspended in 500 μ l of PBS. The samples were stored at -80°C until DNA extraction. To extract DNA, 250 μ l of the swab suspension were centrifuged at 10,000 \times g for 5 min, and supernatant was removed. DNA samples were extracted from the pellet using prepGEM Bacteria (ZyGEM, Hamilton, New Zealand) according to the manufacturer's protocol.

To identify the microbiome, DNA sample from *R. amplexicaudatus* captured at Lavigan was used for high throughput sequencing analysis. For high throughput sequencing of the 16S rRNA gene, PCR fragments of the V3-V4 region were amplified using the universal primers F341 (5'-CCTACGGGNGGCWGCAG-3') and R805 (5'-GACTACHVGGGTATCTAATCC-3') [11]. These primers were modified according to the high throughput sequencing protocol (Illumina, San Diego, CA, U.S.A.). For PCR, the reaction mixture (50 μ l) contained 25 μ l Premix EX Taq (TaKaRa-Bio, Otsu, Japan), 1 μ M each primer and 5 ng isolated DNA. The PCR conditions were as follows: 98°C for 3 min; 25 cycles of 98°C for 10 sec, 55°C for 30 sec, 72°C for 30 sec; and a final extension at 72°C for 10 min. The PCR products were purified using the MonoFas DNA purification kit (GL Science, Tokyo, Japan) and used as template in the second PCR round to attach dual indices and Illumina sequencing adapters. For the second round PCR, the reaction mixture (50 μ l) contained 25 μ l Premix EX Taq, 1 μ M each primer and 10 ng purified amplicons. The PCR conditions were as follows: 98°C for 3 min; 12 cycles of 98°C for 10 sec, 55°C for 30 sec, 72°C for 30 sec; and a final extension at 72°C for 10 min. The second round PCR products were purified as above and then were adjusted to 10–20 ng of DNA. Metagenomic sequencing was carried out on a MiSeq sequencer (Illumina) using the MiSeq Reagent kit v3 (600 cycles) with 300 paired-end reads, and data analysis was performed using the MiSeq Reporter software with the Greengenes database.

Table 1. Family level classification of microbiome in *R. amplexicaudatus*

Family ^{a)}	Reads (%)
<i>Clostridiaceae</i>	162,921 (69.3)
<i>Campylobacteraceae</i>	56,852 (24.2)
<i>Enterobacteriaceae</i>	11,898 (5.1)
<i>Bacillaceae</i>	589 (0.3)
<i>Streptococcaceae</i>	467 (0.2)
<i>Enterococcaceae</i>	350 (0.1)
Others	602 (0.3)
Unclassified	850 (0.4)
Total	234,529

a) Family obtained over 0.1% of total reads was shown.

The metagenomic reads are stored at the National Center for Biotechnology Information Sequence Read Archive under BioProject PRJDB4733 (Accession nos. experiment: DRX053603, run: DRR059024).

A total of 469,058 reads were obtained. Of those, 619 reads contained an ambiguous base (N). In the diversity of microbiota, the predominant phyla were Firmicutes. Family level classification resulted in the identification of 66 families. The predominant families were *Clostridiaceae* (69.5%), *Campylobacteraceae* (24.2%) and *Enterobacteriaceae* (5.1%) (Table 1). In Genus level classification, 103 genera were identified. The predominant genera were *Clostridium* (65.7%) and *Campylobacter* (24.2%) (Table 2). Species level classification resulted in the identification of 170 species. The predominant species were *Clostridium butyricum* (21.0%) and *Campylobacter coli* (19.3%) (Table 3). In a previous study, 71 genera were identified on metagenomics analysis of the fecal bacterial flora of *Myotis daubentonii*, and genera *Leuconostoc*, *Betaproteobacteria* and *Enterobacter* dominated [27]. Common genera between *R. amplexicaudatus* and *M. daubentonii* were 21, and dominated genera were different between them. These differences might be due to the difference of food habit. *R. amplexicaudatus* is frugivorous, and *M. daubentonii* is insectivorous.

Campylobacter was second predominant genus, and *C. coli* and *C. jejuni* were identified in microbiome of *R. amplexicaudatus*. Campylobacteriosis is one of the most important infectious disease in children and has increased over the world in the last decade [2, 13]. Campylobacteriosis is attributed to contamination of poultry product [1, 20], or drinking and environmental water [19, 24]. In *Campylobacter* spp., *C. jejuni* is the most common cause of acute bacterial diarrhea in humans [8]. Genus *Campylobacter* dominated secondly in the flora of *R. amplexicaudatus*, and to further evaluate the risk of spreading *Campylobacter* spp. by wild bats, we investigated the prevalence of *C. jejuni* in these 91 bats. To determine the prevalence of *C. jejuni* in bats, conventional PCR assay was conducted on all the fecal DNAs. The intergenic spacer 16S–23S (ITS) of *C. jejuni* was amplified by PCR using the primers CampyForw (5'-CTGATAAGGGTGAGGTCACAAGT-3') and CampyRev (5'-CTTGCTTGTGACTCTTAACAATG-3') [7]. The

Table 2. Genus level classification of microbiome in *R. amplexicaudatus*

Genus ^{a)}	Reads (%)
<i>Clostridium</i>	154,182 (65.7)
<i>Campylobacter</i>	56,846 (24.2)
<i>Escherichia</i>	9,720 (4.1)
<i>Sarcina</i>	7,495 (3.2)
<i>Serratia</i>	1,526 (0.7)
<i>Bacillus</i>	588 (0.3)
<i>Streptococcus</i>	379 (0.2)
<i>Enterococcus</i>	328 (0.1)
<i>Alkaliphilus</i>	273 (0.1)
<i>Gemella</i>	149 (0.1)
<i>Chlamydomphila</i>	128 (0.1)
Others	612 (0.3)
Unclassified	2,303 (1.0)
Total	235,429

a) Genus obtained over 0.1% of total reads was shown.

reaction mixture (50 μ l) contained 25 μ l Premix EX Taq (TaKaRa-Bio), 1 μ M each primer and 5 ng isolated DNA. The PCR conditions were as follows: 98°C for 3 min; 35 cycles of 98°C for 10 sec, 60°C for 30 sec, 72°C for 30 sec; and a final extension at 72°C for 10 min. PCR products were purified as described above, and then, the products were sequenced using the BigDye Terminator v3.1 Cycle Sequence kit (Applied Biosystems, Foster City, CA, U.S.A.) on a 3130 Genetic Analyzer (Applied Biosystems) and analyzed by Blast (<http://blast.ncbi.nlm.nih.gov/Blast.cgi>) for validation. These results confirm the accurate identification of *C. jejuni*. From 91 samples, *C. jejuni* was detected only in *R. amplexicaudatus* with a prevalence of 7.5% (5/67), and these five bats were captured at two sites, three at Lavigan and two at Sion. Another 43 bats (17 *R. amplexicaudatus* and 3 *H. diadema* at Lavigan, 22 *R. amplexicaudatus* and 1 *E. spelaea* at Sion) were negative at both sites.

The genus *Campylobacter* currently consists of 25 species, some of which are important human pathogens. *Campylobacter enteritis* causes acute bacterial diarrhea in humans, mainly in the developed world. The most often implicated species as the causative agent of campylobacteriosis is *C. jejuni*, followed by *C. coli* [26], *C. upsaliensis* [15] and *C. lari* [22]. Palmer and colleagues speculated that human campylobacteriosis may be the result of water contaminated by bird or bat feces [21], and enteric bacteria were isolated or detected with high throughput sequencing from bats [1, 27]. In these studies, only microchiroptera was surveyed, and *Campylobacter* spp. were not isolated or detected from their fecal or enteric swab samples. In this study, however, we detected *C. jejuni* in rectal swab samples from megachiroptera, *R. amplexicaudatus*, for the first time. *C. jejuni*, especially, could enter a viable, but non-culturable (VBNC) state under stressed condition, such as nutrient starvation, osmotic shock and fluctuations in temperature and pH and is thought to be a possible cause of water borne infections. *C. jejuni* in VBNC form was able to remain in this form for several months and then recover their culturability in the mouse intestine [4].

Table 3. Species level classification of microbiome in *R. amplexicaudatus*

Species ^{a)}	Reads (%)
<i>Clostridium butyricum</i>	49,319 (21.0)
<i>Campylobacter coli</i>	45,167 (19.3)
<i>Clostridium saccharobutylicum</i>	11,320 (4.8)
<i>Clostridium saccharoperbutylacetonicum</i>	10,219 (4.4)
<i>Escherichia albertii</i>	9,396 (4.0)
<i>Clostridium cavendishii</i>	9,249 (3.9)
<i>Sarcina maxima</i>	7,411 (3.2)
<i>Campylobacter helveticus</i>	3,970 (1.7)
<i>Clostridium chartatabidum</i>	3,011 (1.3)
<i>Campylobacter peloridis</i>	2,886 (1.2)
<i>Clostridium cadaveris</i>	2,353 (1.0)
<i>Clostridium histolyticum</i>	1,746 (0.7)
<i>Serratia entomophila</i>	1,524 (0.6)
<i>Clostridium vincentii</i>	999 (0.4)
<i>Clostridium sardiniense</i>	759 (0.3)
<i>Clostridium baratii</i>	754 (0.3)
<i>Bacillus butanolivorans</i>	478 (0.2)
<i>Campylobacter insulaenigrae</i>	278 (0.1)
<i>Campylobacter subantarcticus</i>	272 (0.1)
<i>Campylobacter volucris</i>	262 (0.1)
<i>Clostridium taeniosporum</i>	261 (0.1)
<i>Clostridium bovipellis</i>	242 (0.1)
<i>Campylobacter jejuni</i>	237 (0.1)
<i>Clostridium nitrophenolicum</i>	234 (0.1)
<i>Clostridium sulfidigenes</i>	219 (0.1)
<i>Enterococcus lactis</i>	214 (0.1)
<i>Clostridium chromoreductans</i>	211 (0.1)
<i>Clostridium fallax</i>	141 (0.1)
Others	1,222 (0.5)
Unclassified	70,175 (29.9)
Total	234,529

a) Species obtained over 0.1% of total reads was shown.

Therefore, *C. jejuni* might also be transmitted from wild bats to humans via water contaminated by bat feces.

In conclusion, the microbiota of bat was characterized using high throughput sequencing. Furthermore, we showed *R. amplexicaudatus* is potential carrier of *C. jejuni* in the Philippines. Bats can acquire infectious agents from their diet and other environmental sources and subsequently may transfer those agents to humans and livestock. Therefore, it is necessary to detect other bacterial pathogen to human from bats continuously.

ACKNOWLEDGMENT. This work was supported by a grant from the Ministry of Education, Science, Sports and Culture, Japan (JSPS KAKENHI Grant Number 25304043).

REFERENCES

- Adesiyun, A. A., Stewart-Johnson, A. and Thompson, N. N. 2009. Isolation of enteric pathogens from bats in Trinidad. *J. Wildl. Dis.* **45**: 952–961. [Medline] [CrossRef]
- Allos, B. M. 2001. *Campylobacter jejuni* Infections: update on emerging issues and trends. *Clin. Infect. Dis.* **32**: 1201–1206.

- [Medline] [CrossRef]
3. Badrane, H., Bahloul, C., Perrin, P. and Tordo, N. 2001. Evidence of two Lyssavirus phylogroups with distinct pathogenicity and immunogenicity. *J. Virol.* **75**: 3268–3276. [Medline] [CrossRef]
 4. Baffone, W., Casaroli, A., Citterio, B., Pierfelici, L., Campana, R., Vittoria, E., Guaglianone, E. and Donelli, G. 2006. *Campylobacter jejuni* loss of culturability in aqueous microcosms and ability to resuscitate in a mouse model. *Int. J. Food Microbiol.* **107**: 83–91. [Medline] [CrossRef]
 5. Chua, K. B., Koh, C. L., Hooi, P. S., Wee, K. F., Khong, J. H., Chua, B. H., Chan, Y. P., Lim, M. E. and Lam, S. K. 2002. Isolation of Nipah virus from Malaysian Island flying-foxes. *Microbes Infect.* **4**: 145–151. [Medline] [CrossRef]
 6. Chua, K. B., Bellini, W. J., Rota, P. A., Harcourt, B. H., Tamin, A., Lam, S. K., Ksiazek, T. G., Rollin, P. E., Zaki, S. R., Shieh, W., Goldsmith, C. S., Gubler, D. J., Roehrig, J. T., Eaton, B., Gould, A. R., Olson, J., Field, H., Daniels, P., Ling, A. E., Peters, C. J., Anderson, L. J. and Mahy, B. W. 2000. Nipah virus: a recently emergent deadly paramyxovirus. *Science* **288**: 1432–1435. [Medline] [CrossRef]
 7. Fontanot, M., Iacumin, L., Cecchini, F., Comi, G. and Manzano, M. 2014. Rapid detection and differentiation of important *Campylobacter* spp. in poultry samples by dot blot and PCR. *Food Microbiol.* **43**: 28–34. [Medline] [CrossRef]
 8. Friedman, C. R., Neimann, J., Wegener, H. C. and Tauxe, R. V. 2000. Epidemiology of *Campylobacter jejuni* infections in the United States and other industrialized nations. pp. 121–138. In: *Campylobacter*, 2nd ed., (Nachamkin, I. and Blaser, M.J. eds.), ASM International, Washington.
 9. Gibbons, R. V. 2002. Cryptogenic rabies, bats, and the question of aerosol transmission. *Ann. Emerg. Med.* **39**: 528–536. [Medline] [CrossRef]
 10. Gomez-Alvarez, V., Revetta, R. P. and Santo Domingo, J. W. 2012. Metagenomic analyses of drinking water receiving different disinfection treatments. *Appl. Environ. Microbiol.* **78**: 6095–6102. [Medline] [CrossRef]
 11. Herlemann, D. P., Labrenz, M., Jürgens, K., Bertilsson, S., Waniek, J. J. and Andersson, A. F. 2011. Transitions in bacterial communities along the 2000 km salinity gradient of the Baltic Sea. *ISME J.* **5**: 1571–1579. [Medline] [CrossRef]
 12. Human Microbiome Project Consortium 2012. Structure, function and diversity of the healthy human microbiome. *Nature* **486**: 207–214. [Medline] [CrossRef]
 13. Kaakoush, N. O., Castaño-Rodríguez, N., Mitchell, H. M. and Man, S. M. 2015. Global Epidemiology of *Campylobacter* Infection. *Clin. Microbiol. Rev.* **28**: 687–720. [Medline] [CrossRef]
 14. Klite, P. D. 1965. Intestinal bacterial flora and transit time of three neotropical bat species. *J. Bacteriol.* **90**: 375–379. [Medline]
 15. Labarca, J. A., Sturgeon, J., Borenstein, L., Salem, N., Harvey, S. M., Lehnkering, E., Reporter, R. and Mascola, L. 2002. *Campylobacter upsaliensis*: Another pathogen for consideration in the United States. *Clin. Infect. Dis.* **34**: E59–E60. [Medline] [CrossRef]
 16. Lane, D. J., Pace, B., Olsen, G. J., Stahl, D. A., Sogin, M. L. and Pace, N. R. 1985. Rapid determination of 16S ribosomal RNA sequences for phylogenetic analyses. *Proc. Natl. Acad. Sci. U.S.A.* **82**: 6955–6959. [Medline] [CrossRef]
 17. Liu, G. E. 2009. Applications and case studies of the next-generation sequencing technologies in food, nutrition and agriculture. *Recent Pat. Food Nutr. Agric.* **1**: 75–79. [Medline] [CrossRef]
 18. Murray, K., Rogers, R., Selvey, L., Selleck, P., Hyatt, A., Gould, A., Gleeson, L., Hooper, P. and Westbury, H. 1995. A novel morbillivirus pneumonia of horses and its transmission to humans. *Emerg. Infect. Dis.* **1**: 31–33. [Medline] [CrossRef]
 19. Moore, J., Caldwell, P. and Millar, B. 2001. Molecular detection of *Campylobacter* spp. in drinking, recreational and environmental water supplies. *Int. J. Hyg. Environ. Health* **204**: 185–189. [Medline] [CrossRef]
 20. Nesbit, E. G., Gibbs, P., Dreesen, D. W. and Lee, M. D. 2001. Epidemiologic features of *Campylobacter jejuni* isolated from poultry broiler houses and surrounding environments as determined by use of molecular strain typing. *Am. J. Vet. Res.* **62**: 190–194. [Medline] [CrossRef]
 21. Palmer, S. R., Gully, P. R., White, J. M., Pearson, A. D., Suckling, W. G., Jones, D. M., Rawes, J. C. and Penner, J. L. 1983. Water-borne outbreak of *campylobacter* gastroenteritis. *Lancet* **i**: 287–290. [Medline] [CrossRef]
 22. Prasad, K. N., Dixit, A. K. and Ayyagari, A. 2001. *Campylobacter* species associated with diarrhoea in patients from a tertiary care centre of north India. *Indian J. Med. Res.* **114**: 12–17. [Medline]
 23. Reyes, A. W., Rovira, H. G., Masangkay, J. S., Ramirez, T. J., Yoshikawa, Y. and Baticados, W. N. 2011. Polymerase chain reaction assay and conventional isolation of *Salmonella* spp. from Philippine bats. *Acta Sci. Vet.* **39**: 947.
 24. Schallenberg, M., Bremer, P. J., Henkel, S., Launhardt, A. and Burns, C. W. 2005. Survival of *Campylobacter jejuni* in water: effect of grazing by the freshwater crustacean *Daphnia carinata* (Cladocera). *Appl. Environ. Microbiol.* **71**: 5085–5088. [Medline] [CrossRef]
 25. Schipper, J., Chanson, J. S., Chiozza, F., Cox, N. A., Hoffmann, M., Katariya, V., Lamoreux, J., Rodrigues, A. S., Stuart, S. N., Temple, H. J., Baillie, J., Boitani, L., Lacher, T. E. Jr., Mittermeier, R. A., Smith, A. T., Absolon, D., Aguiar, J. M., Amori, G., Bakkour, N., Baldi, R., Berridge, R. J., Bielby, J., Black, P. A., Blanc, J. J., Brooks, T. M., Burton, J. A., Butynski, T. M., Catullo, G., Chapman, R., Cokeliss, Z., Collen, B., Conroy, J., Cooke, J. G., da Fonseca, G. A., Derocher, A. E., Dublin, H. T., Duckworth, J. W., Emmons, L., Emslie, R. H., Festa-Bianchet, M., Foster, M., Foster, S., Garshelis, D. L., Gates, C., Gimenez-Dixon, M., Gonzalez, S., Gonzalez-Maya, J. F., Good, T. C., Hammerson, G., Hammond, P. S., Happold, D., Happold, M., Hare, J., Harris, R. B., Hawkins, C. E., Haywood, M., Heaney, L. R., Hedges, S., Helgen, K. M., Hilton-Taylor, C., Hussain, S. A., Ishii, N., Jefferson, T. A., Jenkins, R. K., Johnston, C. H., Keith, M., Kingdon, J., Knox, D. H., Kovacs, K. M., Langhammer, P., Leus, K., Lewison, R., Lichtenstein, G., Lowry, L. F., Macavoy, Z., Mace, G. M., Mallon, D. P., Masi, M., McKnight, M. W., Medellín, R. A., Medici, P., Mills, G., Moehlan, P. D., Molur, S., Mora, A., Nowell, K., Oates, J. F., Olech, W., Oliver, W. R., Oprea, M., Patterson, B. D., Perrin, W. F., Polidoro, B. A., Pollock, C., Powel, A., Protas, Y., Racey, P., Ragle, J., Ramani, P., Rathbun, G., Reeves, R. R., Reilly, S. B., Reynolds, J. E. 3rd., Rondinini, C., Rosell-Ambal, R. G., Rulli, M., Rylands, A. B., Savini, S., Schank, C. J., Sechrest, W., Self-Sullivan, C., Shoemaker, A., Sillero-Zubiri, C., De Silva, N., Smith, D. E., Srinivasulu, C., Stephenson, P. J., van Strien, N., Talukdar, B. K., Taylor, B. L., Timmins, R., Tirira, D. G., Tognelli, M. F., Tsytulina, K., Veiga, L. M., Vié, J. C., Williamson, E. A., Wyatt, S. A., Xie, Y. and Young, B. E. 2008. The status of the world's land and marine mammals: diversity, threat, and knowledge. *Science* **322**: 225–230. [Medline] [CrossRef]
 26. Vandamme, P. 2000. Taxonomy of the family *Campylobacteraceae*. pp. 3–26. In: *Campylobacter*, 2nd ed. (Irving, N. and Martin, J. B. eds.), ASM Press, Washington.
 27. Veikkolainen, V., Vesterinen, E. J., Lilley, T. M. and Pulliainen, A. T. 2014. Bats as reservoir hosts of human bacterial pathogen, *Bartonella mayotimonensis*. *Emerg. Infect. Dis.* **20**: 960–967. [Medline] [CrossRef]