

Detection of QTLs for grain protein content in durum wheat

A. Blanco · R. Simeone · A. Gadaleta

Published online: 20 June 2006
© Springer-Verlag 2006


Theor Appl Genet (2006) 112:1195–1204.
DOI: 10.1007/s00122-006-0221-6

The authors regret that the legends accompanying Figs. 1 and 2 were not fully correct. The figures are reproduced here with the correct legends.

The online version of the original article can be found at <http://dx.doi.org/10.1007/s00122-006-0221-6>

A. Blanco (✉) · R. Simeone · A. Gadaleta
Department of Environmental and Agro-Forestry Biology
and Chemistry, University of Bari, Via Amendola, 165/A,
70126 Bari, Italy
e-mail: blanco@agr.uniba.it

Fig. 1 Frequency distributions of grain protein content values of three genotypes based on *Xcfa2164-2A* in **a** F3 progenies grown at Valenzano in 2001 and **b, c** F4 progenies grown at Valenzano and Gaudiano in 2002, respectively. 1 Marker allele from cv. Latino (low grain protein content); 2 marker allele from 3BIL-85 (high grain protein content)






Fig. 2 Linkage map of the 2A chromosome segment introgressed from var. *dicoccoides* in the backcross inbred line 3BIL-85 (*black region*). Map distances (cM) and microsatellite markers significantly associated to grain protein content are shown on the *left* and *right* sides. The centromere (*filled arrow*) and the introgressed segment were positioned according to previously published maps (Roder et al. 1998; Blanco et al. 2004)