



International Journal of Pharmacology

ISSN 1811-7775

science
alert

ansinet
Asian Network for Scientific Information

Determination of Antioxidant Properties of *Gypsophila bitlisensis* Bark.

¹Mesut Işık, ²Mustafa Korkmaz, ³Ercan Bursal, ^{4,5}İlhami Gülçin, ¹Ekrem Köksal and ¹Hatice Tohma

¹Department of Chemistry, Faculty of Science and Arts, Erzincan University, Turkey

²Department of Biology, Faculty of Science and Arts, Erzincan University, Turkey

³Department of Nursing, School of Health, Muş Alparslan University, Turkey

⁴Department of Chemistry, Faculty of Sciences, Atatürk University, Turkey

⁵Fetal Programming of Diseases Research Chair, Department of Zoology, College of Science, King Saud University, Riyadh, Saudi Arabia

ARTICLE INFO

Article History:

Received: January 06, 2015

Accepted: April 09, 2015

Corresponding Author:

Hatice Tohma,

Department of Chemistry,
Faculty of Science and Arts,
Erzincan University,
Turkey

Tel: +90 446 226 66 66 (40015)

ABSTRACT

Oxidative stress has been suggested to explain the mechanism of numerous diseases and antioxidants have been proposed to prevent cellular oxidative damage, therefore disease progression. Antioxidant capacity of many plants, fruits and vegetables have been studied by the researchers, however there was no data regarding the antioxidant properties of *Gypsophila bitlisensis*. The species is an endemic and natural plant of Turkey. We performed three assays to reveal antioxidant capacity of ethanol extract of *Gypsophila bitlisensis*: FRAP, DPPH and CUPRAC. Trolox and α -tocopherol have been used as standards. We also determined phenolic content of the extract. The results showed that *Gypsophila bitlisensis* does not have significant antioxidant capacity and phenolic content. On the basis of these findings, we concluded that *Gypsophila* does not have considerable antioxidant activity in these *in vitro* assays.

Key words: Antioxidant, *Gypsophila bitlisensis*, oxidative stress, FRAP, DPPH, CUPRAC

INTRODUCTION

Under normal physiological conditions, Reactive Oxygen Species (ROS) are produced. These species play important role in signalling, cellular redox cycling and activation of some enzymes. The excess ROS are removed from the cell, as they are also capable of oxidizing biomacromolecules. If ROS are not efficiently removed by antioxidants defences (enzymatic, non-enzymatic or dietary origin) oxidative stress occurs (Bursal and Gulcin, 2011). Chronic oxidative stress results in oxidative modification of proteins, lipids and nucleic acid which can initiate abnormal tissue functions and diseases. Consumption of foods containing natural antioxidants can be used to eliminate the damage caused by ROS and reduce oxidative stress related impairments. Therefore, assessing the antioxidant capacity of the food has become one of the interests in the research area (Bursal and Gulcin, 2011).

Generally, mitochondria and cytochrome 450 enzymes are the main producers of ROS. The factors contributing ROS formation could be endogenous such as pathophysiology, exercise, drug metabolites and ion influx, or exogenous such as drugs, toxicants, pathogens and dietary ingredients. The primary free radicals in the cells are superoxide ($O_2^{\bullet-}$) and Nitric Oxide (NO). The $O_2^{\bullet-}$ is generated as a consequences of incomplete reduction of oxygen in electron transport systems or by specific enzymatic reactions whereas NO is the product of some specific enzymes (the nitric oxide synthases). There are also nonradical oxygen species like H_2O_2 . This is particularly interesting since it is neutral, membrane permeable and can oxidise thiol of cysteine residues which alter the activity of target protein and has role in signalling pathways. The H_2O_2 can also give rise to OH^{\bullet} radical via Fenton reaction in the presence of metal ions. Hydroxyl radicals also form due to indirect effect of X- and gamma-rays through the radiolysis of water (Von Sonntag, 2006).

There are numerous studies showing the direct effect of ROS on disease progression. Perhaps the most accepted idea of involvement of ROS in human disease and toxicology comes from the detection and determination of lipid peroxidation (Valko *et al.*, 2004). For example malondialdehyde is a chain cleavage product of peroxy radicals and it was elevated in patients with liver cancer (Arslan *et al.*, 2014). Attack of peroxy radicals may result in DNA cleavage and protein backbone modification (Valko *et al.*, 2004). In addition, H₂O₂ has been linked to the formation and accumulation of amyloid beta in brain and retina cells which results in lens opacification (Melov *et al.*, 2005). The H₂O₂ has been also suggested to be generated in early stages of aggregate formation in Alzheimer diseases (Tabner *et al.*, 2005). On the other hand, hydroxyl radical is also harmful to living cell by causing damage in DNA. Strand breaks, complex DNA lesions, protein-DNA cross-links and protein and DNA adducts have been suggested to be results of hydroxyl radical induced damage (Cadet and Wagner, 2014).

Despite detrimental effect of ROS, cell has antioxidant defence system which removes ROS and therefore help preventing cancer, heart diseases and stroke (Valko *et al.*, 2004). It has been reported that consumption of vegetables is related to reduced risk of digestive tract neoplasms and fruits is related to reduced risk of cancers of the upper digestive tract, stomach and urinary tract (La Vecchia *et al.*, 2001). Moreover, dietary intake of antioxidants has been shown to have positive impact on immune system, cardiovascular diseases, cataracts, as well as brain dysfunction (Ames *et al.*, 1993).

Studies showing the beneficial effects of specific antioxidants are also accumulating. For example N-acetylcysteine, a thiol containing antioxidant, has been shown to protect rats from memory impairment after sepsis induction (Barichello *et al.*, 2007). Also a recent study has shown that early induction of vitamin E and lycopene suppressed allergenic airway disease in the later life in mice (Hansbro *et al.*, 2014). Consumption of food rich in vitamin E and C has been related to delayed development of the various forms of cataracts (Jacques *et al.*, 1994).

Numerous foods have been reported as source of natural antioxidants and inadequate consumption of these foods has been related to increased risk of cancer (Block *et al.*, 1992). Thus discovery of novel species is useful in terms of preventing diseases and delaying aging. *Gypsophila* L. (family: Caryophyllaceae), are annual, biennial or perennial suffrutescent herbs with linear-subulate leaves which mainly grown in Mediterranean zone (Yucekutlu and Bildaci, 2008). There are around 150 *Gypsophila* species which are commonly used in the industry and medicine to treat some diseases such as hepatitis, gastritis and bronchitis and also used as expectorant (Simeonova *et al.*, 2013; Yucekutlu and Bildaci, 2008). *Gypsophila* is the third biggest genus of Caryophyllaceae family in Turkey. The genus has 55 species in the country and 33 species of them are endemics. All members of the genera are known as Coven and developed roots are economically very important because of their

different amount of saponin contents. Extract produced from their roots are known as fire extinguisher, gold polishing, cleaner and softener of delicate fabrics and crispness giving to halva. The extracts are often used for making liqueur, herbal cheese, ice cream and different foods. Some species of the genus are Boron (B) hyperaccumulators (Korkmaz and Ozelik, 2011). *Gypsophila* has shown to have moderate antimicrobial activity and substantial effect on fungi (Shafagha and Shafaghatlonbar, 2011). The presence of triterpene saponins (Arslan *et al.*, 2013), sterols, flavonoids (Darmograi *et al.*, 1969), triterpens in different *Gypsophila* species (Krsteva *et al.*, 2014) were indicated. *Gypsophila* saponins have been also reported to have *in vitro* cytoprotective effect on galactosamine induced hepatotoxicity (Braut Boucher *et al.*, 1990). It was proposed that *Gypsophila* might have anticancer activity due to containing an active component called gypsogenin at high concentrations (Emirdag-Ozturk *et al.*, 2014). In addition, antioxidant effect of saponarin isolated from *Gypsophila trichotoma* was determined, however, there is no report on overall *in vitro* antioxidant properties of the genus *Gypsophila*. Therefore, it was considered useful to determine the antioxidant properties of *Gypsophila bitlisensis* in order to provide reference to the consumers.

MATERIALS AND METHODS

Chemicals: The 2,9-dimethyl-1,10-phenanthroline (neocuproine), the stable free radical 1,1-diphenyl-2-picrylhydrazyl (DPPH•), 3-(2-Pyridyl)-5,6-bis (4-phenyl-sulfonic acid)-1,2,4-triazine (Ferrozine®), 6-hydroxy-2, 5,7,8-tetramethylchroman-2-carboxylic acid (Trolox), α-tocopherol, CuCl₂, Folin-Ciocalteu reagent and gallic acid were obtained from Sigma (Sigma-Aldrich GmbH, Sternheim, Germany). All other chemicals used were analytical grade and obtained from Merck.

Sample preparation: This plant material was collected between Büyükyayla and Mezra villages in a wet habitat from Çayırılı-Erzincan, Turkey in 2011. The plant samples of the species were identified and some of them were deposited in Erzincan University Herbarium with the sample number of Korkmaz: 2881.

For Ethanol Extract of *Gypsophila* (EEG), about 25 g of *Gypsophila bitlisensis* was grounded to get fine powder and then mixed with 100 mL ethyl alcohol on a magnetic stirring for 1 h. Remaining residue was re-extracted until colourless extraction solvent was obtained (final volume of 600 mL). After extracting the sample through Whatman No. 1 paper, ethanol was removed using a rotary evaporator (RE 100 Bibby, Stone Staffordshire England) at 50°C to obtain dry extract. The extracts were stored at -20°C in dark plastic bottle until experimental studies.

Ferric cyanide (Fe³⁺) reducing antioxidant power assay (FRAP): The FRAP assay was performed as previously described (Bursal and Koksak, 2011). The method is based on the spectrophotometric measurement of complex formed when

ferric tripyridyl triazene (Fe^{3+} -TPTZ) complex was reduced to the ferrous (Fe^{2+}) ion at 700 nm. Briefly, different concentrations of the samples (10, 20 and 30 $\mu\text{g mL}^{-1}$) in 0.75 mL water was mixed with 1 mL of sodium phosphate buffer (0.2 M, pH 6.6) and 1 mL (1%) of potassium ferricyanide (1%). After incubating the mixture at 50°C for 20 min, trichloroacetic acid (10%) was added to mixture to acidify. Absorbance was measured by using distilled water as blank and for control. Higher absorbance indicates better reducing capacity of the sample.

Cupric ions reducing assay (CUPRAC): The protocol for determining the capacity of reducing power: The capacity of sample to reduce cupric ions (Cu^{2+}) was determined by CUPRAC assay as previously described (Bursal and Koksall, 2011). A volume of 0.25 mL CuCl_2 (0.01 M), 0.25 mL neocuproine (7.5 mM) in ethanol and 0.25 mL NH_4Ac (1 M) was mixed with standards and sample solutions at different concentrations (10, 20 and 30 $\mu\text{g mL}^{-1}$). Total volume was adjusted to 2 mL with distilled water and the mixture was kept in the room temperature for 30 min. The absorbance values were read at 450 nm against blank.

1, 1-diphenyl-2-picryl-hydrazil (DPPH) free radical scavenging activity: The DPPH free radical scavenging activities of *Gypsophila bitlisensis* was measured according to the method described by Blois (1958). In the method, the purple colour exhibited by the stable DPPH radical is bleached by free radical scavenging activity of the sample which could be monitored at an absorbance of 517 nm. Briefly, different concentrations (10, 20 and 30 $\mu\text{g mL}^{-1}$) of ethanol extract from *Gypsophila bitlisensis* were prepared and volume adjusted to 3 mL with ethanol. Alcoholic solution of DPPH (1 mL, 0.1 M) was added to sample, followed by incubation in the dark for 30 min. A reduction in the absorbance indicates DPPH• scavenging activity.

Determination of total phenolic content: Total phenolic analysis was performed using Folin-Ciocalteu reagent as previously described (Bursal and Koksall, 2011). The sample test (1 mg) was mixed with 23 mL water, then Folin-Ciocalteu phenol reagent (0.5 mL) and 3 min later, 2% Na_2CO_3 (1.5 mL) were added. The mixture was vortexed and kept in the room temperature for 30 min. Absorbance of sample was measured at 760 nm. Gallic acid was used as standard phenolic compound and calculation was made on the basis of a standard curve of gallic acid. The phenolic content of *Gypsophila bitlisensis* was calculated by using a standard curve and expressed as gallic acid equivalents per mg extract.

Statistical analysis: The results were expressed as the Mean±Standard Deviation of three separate experiments performed in triplicate. Data were analysed using one-way ANOVA followed by LSD in SPSS. Differences were considered statistically significant when $p < 0.05$.

RESULTS AND DISCUSSION

Oxidative stress has been suggested as a causative factor for progression of diseases as well as aging. There are number of studies which link biological oxidative damage to ROS formation. Therefore the assessments of antioxidant characteristics of substances which have scavenging properties of ROS, have been a growing interest.

Antioxidants can act as either stopping the free radicals formation in the first place or stopping radical chain reactions, thus preventing biomolecular oxidation (Karadag *et al.*, 2009). There are two working mechanisms of antioxidants, the one is hydrogen atom transfer and the other is electron transfer (Gulcin, 2012). The response of antioxidants to different radicals and oxidant might be different (Prior *et al.*, 2005). Due to the involvement of different mechanism, there is no universal single assay to measure antioxidant capacity (Celik *et al.*, 2010). Therefore different assays should be performed to evaluate antioxidant capacity.

In this study; we examined the antioxidant capacity of *Gypsophila bitlisensis* species, since its wide use in industry and medicine. Characterization of antioxidant properties of *Gypsophila bitlisensis* was determined and compared to standard antioxidants: Trolox and α -tocopherol. For those comparison three different assays were employed: Ferric cyanide (Fe^{3+}) reducing antioxidant power assay (FRAP), cupric ions reducing assay (CUPRAC) and DPPH• scavenging activity.

The FRAP assay has been widely used in the literature to determine the antioxidant capacity of the sample since it is fast, simple, robust and does not require specialized equipment. The ability of FRAP assay to measure antioxidant substance has been confirmed by the studies which showed the beneficial effect of the food on health and corresponding antioxidant capacity measured by FRAP assay. For example, green tea extract has been shown to prevent collagen induced arthritis in mice (Haqqi *et al.*, 1999) and FRAP assay has distinctly showed that green tea extracts increases total antioxidant capacity of heart, aorta and duodenum but not plasma in mice (Koutelidakis *et al.*, 2014). Additionally, grape seed extracts has been shown to prevented colorectal cancer in mice (Velmurugan *et al.*, 2010) and its antioxidant properties has been elucidated with FRAP method (Guo *et al.*, 2003).

There are other assays to measure antioxidant capacity of the samples such as ABTS radical scavenging activity, Oxygen Radical Absorbance Capacity (ORAC) or the Total Radical-Trapping Antioxidant Parameter (TRAP assay). Each of these assays has advantages over the other. A comparative study has shown that although both DPPH and ABTS assay showed positive correlation with antioxidant capacity of foods, ABTS result were higher for fruits, vegetables and beverage as compared to DPPH assay (Floegel *et al.*, 2011). In addition, antioxidant capacity of methanol extract of guava fruit was estimated as 31.1, 25.2, 26.1 and 21.3 determined by the ABTS, DPPH, FRAP and ORAC assay, respectively (Thaipong *et al.*, 2006). On the other hand, antioxidant




Fig. 1: Ferric cyanide (Fe³⁺) reducing antioxidant power of *Gypsophila bitlisensis* and standards by FRAP assay




Fig. 2: Reducing power of ethanol extracts of *Gypsophila bitlisensis* and standards by CUPRAC assay

capacity of blackberry fruit were 4.19, 6.82 and 4.92 as measured by ABTS, DPPH and FRAP (Ozgen *et al.*, 2006). These results indicated that antioxidant capacity may vary depending on the method, extraction solution, sample and pH used.

It has been shown that both ABTS and FRAP methods are reliable in terms of determining Total Antioxidant Capacity (TAC) of plant foods (Biskup *et al.*, 2013). Although, quercetin, rutin, caffeic acid and chlorogenic acid exhibited comparable results for ABTS and FRAP assays, ferulic acid and catechin indicated higher TAC when ABTS method was used (Nilsson *et al.*, 2005). The reason for higher activity might be the usage of water as a soluble in ABTS method which could increase the amount of water-soluble antioxidants in the sample (Nilsson *et al.*, 2005). It was also mentioned that FRAP method is not responsive to thiol-type antioxidant measurements (Koksal and Gulcin, 2008). In this study, *Gypsophila bitlisensis* has shown slight but significant antioxidant activity at 20 µg mL⁻¹ concentration and no further increase was observed at 30 µg mL⁻¹ (Fig. 1). Therefore, this result showed that *Gypsophila bitlisensis* does not consist of large amount of antioxidants, as it is not highly detectable by using FRAP assay.

In order to monitor the antioxidants that may not be detectable using FRAP assay, we performed CUPRAC assay. The principle of the method is based on the reduction of Cu²⁺-neocuproine complex to Cu⁺-neocuproine complex by antioxidants (Segundo *et al.*, 2015). It has been first proposed by Apak *et al.* (2005) and was shown to be applicable to different biological samples including human serum




Fig. 3: Radical scavenging activity of *Gypsophila bitlisensis* and standards by DPPH assay

food and plant materials (Bener *et al.*, 2010). The advantages of CUPRAC method are that it is capable of oxidizing thiol containing antioxidants, the reagents is more stable and accessible than other chromogenic reagents such as ABTS and DPPH and the reaction is carried at pH 7 which simulates physiological pH. There has been no report on evaluating the Cu reducing capacity of *Gypsophila bitlisensis* extract in the literature. This study, for the first time, showed that *Gypsophila bitlisensis* does not have a significant Cu reducing capacity as compared to trolox (Fig. 2).

The DPPH assay is also performed to monitor hydrogen atom and electron donating ability of *Gypsophila bitlisensis* extract. The mechanism is based on decolouring of purple DPPH by antioxidants which can be monitored by absorbance decrease at 517 nm. The method is rapid, easy and UV-spectrometer based. This method is powerful to show the antioxidant activity of sample as shown in previous studies. For example, using DPPH assay, researchers have shown antioxidant capacity of different plants such as fringe tree (Gulcin *et al.*, 2006), vegetables (cauliflower) (Koksal and Gulcin, 2008), fruits (guava) (Thaipong *et al.*, 2006) leaves (*Melissa officinalis*) (Koksal *et al.*, 2011). In this study, *Gypsophila bitlisensis* extract did not show a significant reduction in DPPH scavenging activity (Fig. 3).

The phenolic compounds such as lignin, flavonoid or tannin can be considered as a large group of secondary metabolites of plants (Blokhina and Fagerstedt, 2010). Polyphenol structures have ideal properties for antioxidant system and some polyphenols can show more antioxidant activity *in vitro* than vitamin E or C. Furthermore, different mechanisms are proposed (Rice-Evans *et al.*, 1997). The food phenolics exhibit important structural diversity and can be categorized as many compounds class. The total phenolic content in many plants have been determined such as sumac (Bursal and Koksal, 2011), orange (Rapisarda *et al.*, 1999), cherry stem (Bursal *et al.*, 2013) and cauliflower (Koksal and Gulcin, 2008). The method used for determination total phenolic content in *Gypsophila bitlisensis* was already explained in the study of Koksal *et al.* (2011). In this study, we have found that total phenolic amount of *Gypsophila bitlisensis* was 8.75 µg GAE.

Gypsophila saponin has been found to have blood cholesterol lowering effect in rats, due to the possible

interference with Fe metabolism (Southon *et al.*, 1988). Therefore it would be useful to determine Fe³⁺ reducing capacity of *Gypsophila bitlisensis* which may explain antioxidant characteristics of it, if has any.

In summary oxidative stress theory has been widely accepted to explain cellular damage in diseases and aging. Therefore discoveries of plant rich in antioxidant are important to treat and prevent the oxidation of cellular components. In this study we have shown that *Gypsophila bitlisensis* does not have considerable antioxidant capacity as measured by three different assays: The FRAP, CUPRAC and DPPH scavenging activity. Further studies should be performed to explore herbs and plants that can be used in medicine.

ACKNOWLEDGMENTS

The authors gratefully acknowledge Scientific Research Projects Board of Erzincan University for the financial supports (grant numbers are FEN-A-220114-0062 and 2011-BAP-10.01.05).

REFERENCES

- Ames, B.N., M.K. Shigenaga and T.M. Hagen, 1993. Oxidants, antioxidants and the degenerative diseases of aging. *Proc. Nat. Acad. Sci. USA.*, 90: 7915-7922.
- Apak, R., K. Guclu, M. Ozyurek, S.E. Karademir and M. Altun, 2005. Total antioxidant capacity assay of human serum using copper (II)-neocuproine as chromogenic oxidant: The CUPRAC method. *Free Radical Res.*, 39: 949-961.
- Arslan, I., A. Celik and M.F. Melzig, 2013. Nebulosides A-B, novel triterpene saponins from under-ground parts of *Gypsophila arrostii* Guss. var. *nebulosa*. *Bioorg. Med. Chem.*, 21: 1279-1283.
- Arslan, A., H. Demir, M.F. Ozbay and H. Arslan, 2014. Evaluation of lipid peroxidation and some antioxidant activities in patients with primary and metastatic liver cancer. *J. Cancer Therapy*, 5: 192-197.
- Barichello, T., R.A. Machado, L. Constantino, S.S. Valvassori and G.Z. Reus *et al.*, 2007. Antioxidant treatment prevented late memory impairment in an animal model of sepsis. *Crit. Care Med.*, 35: 2186-2190.
- Bener, M., M. Ozyurek, K. Guclu and R. Apak, 2010. Development of a low-cost optical sensor for cupric reducing antioxidant capacity measurement of food extracts. *Anal. Chem.*, 82: 4252-4258.
- Biskup, I., I. Golonka, A. Gamian and Z. Sroka, 2013. Aktywnosc przeciwutleniajaca wybranych fenoli oznaczona testem ABTS i FRAP [Antioxidant activity of selected phenols estimated by ABTS and FRAP methods]. *Postepy Higieny Medycyny Doswiadczalnej*, 67: 958-963, (In Polish).
- Block, G., B. Patterson and A. Subar, 1992. Fruit, vegetables and cancer prevention: A review of the epidemiological evidence. *Nutr. Cancer*, 18: 1-29.
- Blois, M.S., 1958. Antioxidant determinations by the use of a stable free radical. *Nature*, 181: 1199-1200.
- Blokhina, O. and K.V. Fagerstedt, 2010. Oxidative metabolism, ROS and NO under oxygen deprivation. *Plant Physiol. Biochem.*, 48: 359-373.
- Braut-Boucher, F., S. Achard-Ellouk, D. Pauthe-Dayde, M. Henry and H. Hoellinger, 1990. Cytoprotective effects of *Gypsophila saponins* towards isolated rat hepatocytes. *Food Addit. Contam.*, 7: S127-S130.
- Bursal, E. and E. Koksals, 2011. Evaluation of reducing power and radical scavenging activities of water and ethanol extracts from sumac (*Rhus coriaria* L.). *Food Res. Int.*, 44: 2217-2221.
- Bursal, E. and I. Gulcin, 2011. Polyphenol contents and *in vitro* antioxidant activities of lyophilised aqueous extract of kiwifruit (*Actinidia deliciosa*). *Food Res. Int.*, 44: 1482-1489.
- Bursal, E., E. Koksals, I. Gulcin, G. Bilsel and A.C. Goren, 2013. Antioxidant activity and polyphenol content of cherry stem (*Cerasus avium* L.) determined by LC-MS/MS. *Food Res. Int.*, 51: 66-74.
- Cadet, J. and J.R. Wagner, 2014. Oxidatively generated base damage to cellular DNA by hydroxyl radical and one-electron oxidants: Similarities and differences. *Arch. Biochem. Biophys.*, 557: 47-54.
- Celik, S.E., M. Ozyurek, K. Guclu and R. Apak, 2010. Solvent effects on the antioxidant capacity of lipophilic and hydrophilic antioxidants measured by CUPRAC, ABTS/persulphate and FRAP methods. *Talanta*, 81: 1300-1309.
- Darmograi, V.N., P.E. Krivenchuk and V.I. Litvinenko, 1969. [Flavonoids of (*Gypsophila paniculata* L.)]. *Farmatsiia*, 18: 30-32, (In Russian).
- Emirdag-Ozturk, S., T. Karayildirim, A. Capci-Karagoz, O. Alankus-Caliskan, A. Ozmen and E. Poyrazoglu-Coban, 2014. Synthesis, antimicrobial and cytotoxic activities and structure-activity relationships of gypsogenin derivatives against human cancer cells. *Eur. J. Med. Chem.*, 82: 565-673.
- Floegel, A., D.O. Kim, S.J. Chung, S.I. Koo and O.K. Chun, 2011. Comparison of ABTS/DPPH assays to measure antioxidant capacity in popular antioxidant-rich US foods. *J. Food Comp. Anal.*, 24: 1043-1048.
- Gulcin, I., 2012. Antioxidant activity of food constituents: An overview. *Arch. Toxicol.*, 86: 345-391.
- Gulcin, I., R. Elias, A. Gepdiremen and L. Boyer, 2006. Antioxidant activity of lignans from fringe tree (*Chionanthus virginicus* L.). *Eur. Food Res. Technol.*, 223: 759-767.
- Guo, C., J. Yang, J. Wei, Y. Li, J. Xu and Y. Jiang, 2003. Antioxidant activities of peel, pulp and seed fractions of common fruits as determined by FRAP assay. *Nutr. Res.*, 23: 1719-1726.
- Hansbro, P.M., J.C. Horvat, R.Y. Kim, J.R. Mayall and J.W. Pinkerton *et al.*, 2014. Antioxidant treatment suppresses the progression of early-life infection-induced severe asthma and pathology in later-life. *Am. J. Respir. Crit. Care Med.*, 189: A1291-A1291.

- Haqqi, T.M., D.D. Anthony, S. Gupta, N. Ahmad, M.S. Lee, G.K. Kumar and H. Mukhtar, 1999. Prevention of collagen-induced arthritis in mice by a polyphenolic fraction from green tea. Proc. Natl. Acad. Sci. USA., 96: 4524-4529.
- Jacques, P.F., L.T. Chylack Jr. and A. Taylor, 1994. Relationships between Natural Antioxidants and Cataract Formation. In: Natural Antioxidants in Human Health and Disease, Frei, B (Ed.). 1st Edn., Chapter 18, Academic Press, San Diego, USA., ISBN-13: 978-0122669750, pp: 515-533.
- Karadag, A., B. Ozcelik and S. Saner, 2009. Review of methods to determine antioxidant capacities. Food Anal. Methods, 2: 41-60.
- Koksal, E. and I. Gulcin, 2008. Antioxidant activity of cauliflower (*Brassica oleracea* L.). Turk. J. Agric. For., 32: 65-78.
- Koksal, E., E. Bursal, E. Dikici, F. Tozoglu and I. Gulcin, 2011. Antioxidant activity of *Melissa officinalis* leaves. J. Med. Plants Res., 5: 217-222.
- Korkmaz, M. and H. Ozcelik, 2011. Economic importance of *Gypsophila* L., *Ankyropetalum* Fenzl and *Saponaria* L. (Caryophyllaceae) taxa of Turkey. Afr. J. Biotechnol., 10: 9533-9541.
- Koutelidakis, A.E., D. Kizis, K. Argyri, A. Kyriakou, M. Komaitis and M. Kapsokefalou, 2014. The effect of iron and fat in a diet containing green tea extract (*Camellia sinensis*) on the antioxidant capacity of some organs and the mRNA expression of specific genes in mice. J. Med. Food, 17: 1232-1238.
- Krasteva, I., M. Yotova, D. Yosifov, N. Benbassat, K. Jenett-Siems and S. Konstantinov, 2014. Cytotoxicity of gypsogenic acid isolated from *Gypsophila trichotoma*. Pharmacogn. Mag., 10: 430-433.
- La Vecchia, C., A. Altieri and A. Tavani, 2001. Vegetables, fruit, antioxidants and cancer: A review of Italian studies. Eur. J. Nutr., 40: 261-267.
- Melov, S., N. Wolf, D. Strozyk, S.R. Doctrow and A.I. Bush, 2005. Mice transgenic for Alzheimer disease β -amyloid develop lens cataracts that are rescued by antioxidant treatment. Free Radical Biol. Med., 38: 258-261.
- Nilsson, J., D. Pillai, G. Onning, C. Persson, A. Nilsson and B. Akesson, 2005. Comparison of the 2,2'-azino-bis-3-ethylbenzothiazole-6-sulfonic acid (ABTS) and Ferric Reducing Anti-oxidant Power (FRAP) methods to assess the total antioxidant capacity in extracts of fruit and vegetables. Mol. Nutr. Food Res., 49: 239-246.
- Ozgen, M., R.N. Reese, A.Z. Tulio Jr., J.C. Scheerens and A.R. Miller, 2006. Modified 2,2'-azino-bis-3-ethylbenzothiazole-6-sulfonic acid (ABTS) method to measure antioxidant capacity of selected small fruits and comparison to Ferric Reducing Antioxidant Power (FRAP) and 2,2'-diphenyl-1-picrylhydrazyl (DPPH) methods. J. Agric. Food Chem., 54: 1151-1157.
- Prior, R.L., X. Wu and K. Schaich, 2005. Standardized methods for the determination of antioxidant capacity and phenolics in foods and dietary supplements. J. Agric. Food Chem., 53: 4290-4302.
- Rapisarda, P., A. Tomaino, R. lo Cascio, F. Bonina, A. de Pasquale and A. Saija, 1999. Antioxidant effectiveness as influenced by phenolic content of fresh orange juices. J. Agric. Food Chem., 47: 4718-4723.
- Rice-Evans, C.A., N. Miller and G. Paganga, 1997. Antioxidant properties of phenolic compounds. Trends Plant Sci., 2: 152-159.
- Segundo, M.A., I.V. Toth, L.M. Magalhaes and S. Reis, 2015. Automatic Flow Injection Analysis (FIA) Determination of Total Reducing Capacity in Serum and Urine Samples. In: Advanced Protocols in Oxidative Stress III, Armstrong, D. (Ed.). Chapter 20, Springer, New York, ISBN: 978-1-4939-1440-1, pp: 277-284.
- Shafagha, A. and M. Shafaghatlonbar, 2011. Antimicrobial activity and chemical constituents of the essential oils from flower, leaf and stem of *Gypsophila bicolor* from Iran. Nat. Prod. Commun., 6: 275-276.
- Simeonova, R., V. Vitcheva, M. Kondeva-Burdina, I. Krasteva, V. Manov and M. Mitcheva, 2013. Hepatoprotective and antioxidant effects of saponarin, isolated from *Gypsophila trichotoma* Wend. on paracetamol-induced liver damage in rats. BioMed Res. Int. 10.1155/2013/757126
- Southon, S., I.T. Johnson, J.M. Gee and K.R. Price, 1988. The effect of *Gypsophila saponins* in the diet on mineral status and plasma cholesterol concentration in the rat. Br. J. Nutr., 59: 49-55.
- Tabner, B.J., O.M. El-Agnaf, S. Turnbull, M.J. German and K.E. Paleologou *et al.*, 2005. Hydrogen peroxide is generated during the very early stages of aggregation of the amyloid peptides implicated in Alzheimer disease and familial British dementia. J. Biol. Chem., 280: 35789-35792.
- Thaipong, K., U. Boonprakob, K. Crosby, L. Cisneros-Zevallos and D.H. Byrne, 2006. Comparison of ABTS, DPPH, FRAP and ORAC assays for estimating antioxidant activity from guava fruit extracts. J. Food Comp. Anal., 19: 669-675.
- Valko, M., M. Izakovic, M. Mazur, C.J. Rhodes and J. Telser, 2004. Role of oxygen radicals in DNA damage and cancer incidence. Mol. Cell. Biochem., 266: 37-56.
- Velmurugan, B., R.P. Singh, N. Kaul, R. Agarwal and C. Agarwal, 2010. Dietary feeding of grape seed extract prevents intestinal tumorigenesis in APC^{min/+} mice. Neoplasia, 12: 95-102.
- Von Sonntag, C., 2006. Free-Radical-Induced DNA Damage and Its Repair: A Chemical Perspective. Springer, New York, ISBN: 9783540261209, Pages: 523.
- Yucekutlu, A.N. and I. Bildaci, 2008. Determination of plant saponins and some of *Gypsophila* Species: A review of the literature. Hacettepe J. Biol. Chem., 36: 129-135.