



Determination of Chitin Content in Insects: An Alternate Method Based on Calcofluor Staining

Bianca Santos Henriques¹, Eloi Souza Garcia^{1,2}, Patricia Azambuja¹ and Fernando Ariel Genta^{1,2*}

¹ Laboratory of Insect Physiology and Biochemistry, Oswaldo Cruz Institute – Oswaldo Cruz Foundation (IOC-FIOCRUZ), Rio de Janeiro, Brazil, ² National Institute of Science and Technology for Molecular Entomology (INCT-EM), Cidade Universitária, Rio de Janeiro, Brazil

OPEN ACCESS

Edited by:

Xanthe Vafopoulou,
York University, Canada

Reviewed by:

Sheila Ons,
National University of La Plata,
Argentina

Ana C. A. Melo,
Federal University of Rio de Janeiro,
Brazil

*Correspondence:

Fernando Ariel Genta
genta@ioc.fiocruz.br;
gentafernando@gmail.com

Specialty section:

This article was submitted to
Invertebrate Physiology,
a section of the journal
Frontiers in Physiology

Received: 29 November 2019

Accepted: 31 January 2020

Published: 18 February 2020

Citation:

Henriques BS, Garcia ES,
Azambuja P and Genta FA (2020)
Determination of Chitin Content
in Insects: An Alternate Method
Based on Calcofluor Staining.
Front. Physiol. 11:117.
doi: 10.3389/fphys.2020.00117

Chitin is an aminopolysaccharide present in yeast cells and arthropod cuticle and is one of the most abundant biopolymers. The conventional methods for the quantitation of chitin content in biological samples are based on its hydrolysis (acid or enzymatic), and the assessment of the byproduct, glucosamine. However, previously described methodologies are time-consuming, laborious, low throughput, and not applicable to insect samples in many cases. Here we describe a new approach to chitin content quantitation based on calcofluor fluorescent brightener staining of samples, followed by microplate fluorescence readings. Calcofluor is a specific chitin stain commonly used for topological localization of the polymer. The protocol was tested in three important disease vector species, namely *Lutzomyia longipalpis*, *Aedes aegypti*, and *Rhodnius prolixus*, and then compared to a classic colorimetric chitin assessment method. Results show that chitin content in the tested insects can vary largely in a range of 8–4600 micrograms of chitin per insect, depending on species, sex, and instar. Comparisons between measurements from the previous protocol and calcofluor method showed statistically significant differences in some samples. However, the difference might be due to interference in the classic method from non-chitin sources of glucosamine and reducing agents. Furthermore, chitinase hydrolysis reduces the total chitin mass estimated between 36 and 74%, consolidating the fluorescent measurements as actual stained chitin in the same extent that was observed with the standard protocol. Therefore, the calcofluor staining method revealed to be a fast and reliable technique for chitin quantitation in homogenized insect samples.

Keywords: chitin, polysaccharide, calcofluor, insect, quantitation, technique

INTRODUCTION

Chitin is a large hydrophobic homopolymer chain of β -(1-4)-linked *N*-acetyl-D-glucosamine synthesized by membrane-bound chitin synthase enzymes (EC 2.4.1.16) (Cohen, 2010). It is an aminopolysaccharide present in both fungal cell wall and arthropod exoskeleton, such as insects, and is responsible for providing rigidity and structural integrity (Cohen, 1993, 2010; Pillai et al., 2009).

The determination of chitin amounts has been previously described mainly for yeast cells. However, it has been done based on the assessment of respective glucosamine release through chitin acid hydrolysis (Paech and Tracey, 1956; Chen and Johnson, 1983) or deacetylation (Ride and Drysdale, 1972; Lehmann and White, 1975), which are laborious and time-consuming processes.

Chitin quantitation was also achieved through the enzymatic hydrolysis of chitin and chitosan, still configuring a demanding and challenging technique (Subramanyam and Rao, 1987). Other methods propose a direct quantification of chitin levels through flow cytometry in fungi cells stained with specific chitin dye, calcofluor white (Costa-de-Oliveira et al., 2013). Nevertheless, this approach is not applicable to insect samples.

Calcofluor white is a fluorescent brightener widely used in the topological localization of chitin, commonly used in yeast fluorescent microscopy. This fluorochrome has the ability to make hydrogen bonds oriented to the structural polysaccharide microfibrils (Herth and Schnepf, 1980; Herth, 1980).

Here we describe a fast and simple protocol to determine chitin content in homogenized insect samples stained with calcofluor, based on fluorometric microplate readings.

MATERIALS AND EQUIPMENT

Materials

Steel beads (11079112ss, BioSpec Products); calcofluor white fluorescent brightener (F3543, SIGMA); *Streptomyces griseus chitinase* (C6137, Sigma).

Equipment

Mini-Bead Beater (BioSpec Products); fluorimeter microplate reader (SpectraMax® Gemini XPS, Molecular Devices).

MATERIALS AND METHODS

Rhodnius prolixus

Rhodnius prolixus were reared in the Laboratory of Insect Biochemistry and Physiology (LABFISI – IOC/FIOCRUZ) maintained at 28°C and relative humidity between 60 and 70%. Insects were fed through artificial apparatus upon latex membrane with defibrinated rabbit blood (Garcia et al., 1984). Blood was provided by the Laboratory Animals Creation Center of Fiocruz (Cecal), respecting the guidelines established by the Ethics Committee on Animal Use (CEUA) established by Fiocruz (license number LW019/17). Samples were prepared using starving first instar nymphs collected 7 days after egg hatching, or starving male adults (at least 2 weeks from the last blood meal).

Lutzomyia longipalpis

Adult sandflies were obtained from a colony descendant from individuals initially captured in Jacobina, Brazil, maintained at LABFISI. Larvae were fed with a mix of rabbit feces, rabbit food, and soil throughout all larval stages, with the addition

of white soy protein (bran) and cereal flakes (Neston) (1:1) during third and fourth instars. After molting, adults were kept with 70% sucrose solution (w/v), and females were blood fed in anesthetized hamster (Ketamine, 200 mg/kg) (CEUA L-029/2016) (Moraes et al., 2014). Samples were prepared with adult males or females collected 3–5 days after emergence from pupae, and fed with sugar only.

Aedes aegypti

Eggs from the Rockefeller strain were obtained from the Laboratory of Physiology and Control for Arthropod Vectors (LAFICAVE – IOC/FIOCRUZ, License CEUA L-004/2018). The hatching of eggs was induced by the addition of distilled water in plastic cups. Larvae were reared until the adult stage in dechlorinated water containing 0.1 g of cat food (Whiskas®, Purina, Brazil) in LABFISI at 28°C and relative humidity between 60 and 70% with 12-h light and dark cycle (Souza et al., 2016). Samples were prepared with starving adult males or females collected 1–2 days after emergence.

Chitin Measurement

Preparation of Samples

Insects were homogenized using a Mini-Bead Beater (BioSpec Products) and 1.2 mm stainless steel beads (11079112ss, BioSpec Products). *R. prolixus* first instar nymphs were pooled in groups of 10 in 200 µL, and male adults were homogenized individually in 1000 µL distilled water, followed by samples dilution. *L. longipalpis* were grouped in pools of 20 insects in 200 µL distilled water and *A. aegypti* adults in pools of 20 insects in 200 µL distilled water.

Lehmann and White Method (1975)

Chitin content from samples was determined according to Lehmann and White (1975). Aliquots of 20 µL of *R. prolixus* samples, or 40 µL for *L. longipalpis* and *A. aegypti*, were centrifuged at 21,000 × g for 5 min at room temperature. Pellets were resuspended in 200 µL 3% (w/v) sodium dodecyl sulfate and incubated at 100°C for 15 min. After cooling, samples were centrifuged again, washed twice with 200 µL water and resuspended in 150 µL of 2.1 M KOH. Samples were heated to 130°C for 1 h then brought to room temperature for the addition of 400 µL 75% (v/v) ethanol and incubation on ice for 15 min. After that, 60 µL of the supernatant of 1 g suspension of Celite® 545 in 12.5 mL 75% ethanol (rested for 2 min) were added to the samples, following centrifugation and washing with 400 µL 40% ethanol solution.

Pellets containing insoluble chitosan were then resuspended in 125 µL water, plus 125 µL of 5% (w/v) NaNO₂ and 125 µL 5% (w/v) KHSO₄ and samples were incubated at room temperature for 15 min. The same was done with 125 µL water as a control. After centrifugation (1,500 × g, 2 min, 4°C), two aliquots were taken from each tube as replicates and combined each with 50 µL of ammonium sulfamate 12.5% (w/v) and 50 µL of fresh 12.5% (w/v) MBTH (3-methyl-benzo-2-thiazolone-hydrazone). Samples were incubated at 100°C for 3 min and left to cool, then 50 µL of 0.83% (w/v) FeCl₃·6H₂O, and samples were incubated at room temperature for 25 min. Aliquots of 200 µL were

transferred to a 96-well microplate and absorbance at 650 nm was measured using a spectrophotometer (SpectraMax® 190, Molecular Devices). When necessary, samples were diluted in water to reduce optical density to the range of the reader.

Calcofluor Method

Aliquots were taken from the different species samples and added to 100 μ L of a solution of calcofluor white fluorescent brightener (F3543, SIGMA), at a concentration of 1 mg/mL for *R. prolixus* nymphs and *L. longipalpis* adults or 50 mg/mL in dimethyl sulfoxide (DMSO) for *R. prolixus* adult males and *A. aegypti* adults. Higher concentration of calcofluor was used for the last samples due to an observed underestimation of the amounts of chitin (see section “Discussion”). Experiments required different volume input for each type of sample (Table 1).

Samples were incubated in the dark for 15 min for chitin binding to the stain, then centrifuged $21,000 \times g$ for 5 min. and the pellets were washed two times with 200 μ L distilled water. Pellets were resuspended in 200 μ L distilled water and transferred to a black 96-well microplate (Costar, 3915, Corning). Fluorescent intensity was determined using a microplate reader (SpectraMax® Gemini XPS, Molecular Devices) with excitation at λ_{ex} 355 nm and emission at λ_{em} 433 nm.

Chitin hydrolysis

Samples of *R. prolixus* first instar nymphs and adult males; adult males and females of *L. longipalpis*; and adult males and females of *A. aegypti* were prepared (see item “Preparation of Samples” for details). Aliquots of 70 μ L were separated and combined with 70 μ L of 1 mg/mL chitinase from *Streptomyces griseus* (C6137, Sigma) in 50 mM potassium phosphate, pH 6.0, 20 mM sodium azide. Control groups consisted of tubes containing samples and 70 μ L of the same buffer. Samples were incubated at 37°C for 18 h, then filled to 350 μ L with distilled water and aliquots were taken for chitin content determination following both protocols.

Preparation of Colloidal Chitin Suspension

Colloidal chitin suspension was prepared following previously described methodology (Hsu and Lockwood, 1975). Briefly, 2 g of commercial chitin from shrimp shells (C7170, SIGMA) were dissolved in 40 mL of concentrated hydrochloric acid (36.5–38%, Vetec, V000154) by magnetic stirring for 50 min. Chitin was then precipitated by adding it to 100 mL of distilled water. The colloidal suspension was centrifuged, and the pellet washed with distilled water until pH was above 3.5.

Determination of Colloidal Chitin Suspension Concentration

For the determination of the colloidal chitin suspension concentration, masses between 1 and 20 mg of chitin (C7170, SIGMA) and 400 μ L of distilled water were added to glass test tubes, followed by the addition of 400 μ L of phenol 5% (w/v). Then one milliliter of sulfuric acid (95–97%) was pipetted rapidly directly onto the liquid surface, in order to obtain proper mixing of the reagents. Tubes were incubated for 10 min, shaken and incubated for 10 more min at

30°C (Dubois et al., 1956). Volumes of 200 μ L of samples were transferred to a 96-well microplate and absorbance at 490 nm was measured using a spectrophotometer (SpectraMax® 190, Molecular Devices). Chitin weight was plotted against corresponding absorbance values resulting in a standard curve. The same was done using samples of colloidal chitin to confirm concentrations. Slopes from the two curves were used to estimate the suspension concentration.

Standard Curve and Chitin Content Determination

For the calculations of the amount of chitin recovered from samples, a standard curve was prepared using increasing volumes of colloidal chitin suspension at 4.24 μ g/ μ L, by plotting the weight of chitin added against corresponding absorbance or fluorescence. Chitin content was determined using linear regression prediction, taking into account dilution, sample volume input and volume per insect.

Statistical Analyses

All data obtained from chitin quantitation assays were analyzed by Two-way analysis of variance (ANOVA) by using the software GraphPad Prism version 6.04 for Windows (San Diego, CA, United States). All quantitation graphs denote means of at least ten, up to 40 samples, and respective standard errors of the means (SEM).

RESULTS

A colloidal chitin suspension was prepared according to Hsu and Lockwood (1975) to be used as a standard for the estimation of the unknown chitin content from samples. Authors infer that the protocol for the suspension preparation recovers at least 85% of the chitin input dissolved in Hydrochloric acid, therefore, the concentration of the suspension needed to be evaluated. We used the colorimetric method for the determination of sugars described by Dubois et al. (1956), applying the protocol to increasing masses of powdered chitin from shrimp shells (Figure 1A) or increasing volumes of the colloidal chitin suspension prepared (Figure 1B). Weights and volumes, respectively, were plotted against optical density obtained at 490 nm, and slopes used to estimate the concentration of the suspension at 4.24 μ g/ μ L.



For the calculation of the corresponding amount of chitin contained in the tested samples, it was necessary to build standard curves (Figure 2) applying both methods to increasing masses of chitin suspended in a colloidal suspension. Weights of chitin were plotted against the optic density, for the Lehmann and White (1975) methodology (Figure 2A), or against the fluorescent intensity (Figure 2B), for the calcofluor staining quantitation method. Colloidal chitin suspension contained 4.24 μ g/ μ L, and curves presented slopes of 0.02 and 809.16, respectively, both with $R^2 = 0.99$.

Linear regression was applied to estimate the unknown amount of chitin in the tested samples by fitting the obtained optical density or fluorescence to the respective standard curves.

TABLE 1 | Detailed methodology for the three insect species, *Rhodnius prolixus*, *Lutzomyia longipalpis*, and *Aedes aegypti*.

Insect Species	Stage	Sex	Number of insects	Total H ₂ O volume (μL)	Volume input		Insect equivalent input	
					Lehmann and White (μL)	Calcofluor quantitation method (μL)	Lehmann and White	Calcofluor quantitation method
<i>R. prolixus</i>	First instar	-	10	200	80	20	0.8	0.2
	adult	Male	1	1000	20	15	0.004	0.003
<i>L. longipalpis</i>	Adult	Male	40	200	50	25	2	1
		Female	20		100	50		
<i>A. aegypti</i>	Adult	Male	20	200	50	50	1	1
		Female			20	30	0.4	0.6

Data includes stage, sex, number of insects, total volume of distilled water for sample homogenization, volume input in quantitation assays after chitinase hydrolysis incubation and number of insects contained in volume input for chitin determination (considering sample dilution).



Results were then subjected to adjustments considering the number of insects homogenized, the total volume used, the volume input in the chitinase hydrolysis step, and volume input in each quantitation assay.

For the purpose of testing the application of the described method in homogenized insect samples, we used three insect species, that are important disease vectors and research models, namely *Lutzomyia longipalpis*, *Aedes aegypti*, and *Rhodnius prolixus*. Aliquots of homogenized samples were submitted to hydrolysis using a commercial chitinase enzyme from the bacterial species *Streptomyces griseus*. This step was included in order to verify if the enzyme activity would reduce the total amount of chitin calculated using both methods, and make sure what was being measured was, in fact, the desired polysaccharide.

The assessment of the chitin content of samples containing a pool of 40 males or 20 females of *L. longipalpis*, homogenized in 200 μ L of distilled water, resulted in similar amounts of chitin in both methodologies (Figure 3). Quantitation assays were carried out in both control samples and samples incubated in the presence of chitinase. Using the calcofluor staining method we obtained a result of 7.57 ± 0.38 micrograms of chitin per insect for control males and 2.64 ± 0.02 μ g/insect for the chitinase hydrolyzed samples, a reduction of 65% in the total mass of the polysaccharide ($\Delta = 4.93$ μ g/insect) (Figure 3A). Meanwhile, using the Lehmann and White method in the same group of insect the resulting measurements were 9.79 ± 0.30 μ g/insect for control and 6.99 ± 0.12 μ g/insect for the chitinase treated group (Figure 3A), meaning a reduction of 29% ($\Delta = 2.80$ μ g/insect).

In the female *L. longipalpis* control group, the quantitation with the fluorescent stain resulted in a total mass of 11.96 ± 0.54 μ g of chitin/insect, while the chitinase treated group resulted in 3.13 ± 0.02 μ g/insect (Figure 3B), accounting for a reduction of 74% ($\Delta = 8.83$ μ g/insect). For the Lehmann and White chitin assessment method, in the control group results showed a total of 13.50 ± 0.61 μ g/insect, and 7.99 ± 0.22 μ g of chitin per insect after hydrolysis (Figure 3B), reducing the polymer content calculated in this group of insects by 41% ($\Delta = 5.51$ μ g/insect).

For *Aedes aegypti*, adult males or females were gathered in pools of 20 insects and subsequently homogenized in a total volume of 200 μ L of distilled water. Samples were divided into aliquots for the control groups and treatment with chitinase (Figure 4). Using calcofluor fluorescent brightener for the total chitin mass assessment in control samples of male *A. aegypti* resulted in 24.6 ± 0.7 μ g of chitin/insect (Figure 4A). Chitinase treatment reduced the polysaccharide amount by 46%, resulting in 13.2 ± 0.6 μ g/insect ($\Delta = 11.3$ μ g/insect) (Figure 4A). The method described by Lehmann and White (1975) for the male *A. aegypti* control group resulted in a total of 29.9 ± 0.8 μ g of chitin per insect, while the chitinase treatment reduced it by 44%, with a total residue of 16.6 ± 0.4 μ g/insect.

Female *A. aegypti* chitin measured with calcofluor gave a mean of 47 ± 3 μ g/insect (Figure 4B). Treatment with chitinase from *S. griseus* reduced the polymer detection by 36%, corresponding to 30 μ g of chitin/insect (Figure 4B). Colorimetric quantitation by deacetylation (Lehmann and White, 1975) resulted in a calculated mass of 60 ± 2 μ g of

chitin/insect in the control group, while the chitinase treated group had a value corresponding to 35 ± 2 μ g/insect after treatment (Figure 4B). In this case, hydrolyzation accounted for a reduction of 41% in the estimated total mass of chitin ($\Delta = 25$ μ g/insect).

In the case of the hemipteran species *R. prolixus*, chitin quantitation in first instar nymphs was performed in homogenized pools of 10 insects in 200 μ L of distilled water. After incubation with chitinase, sample's chitin content was assessed comparatively using both fluorometric and colorimetric assays (Figure 5). Calcofluor staining resulted in an estimated total mass of chitin of 41.4 ± 3.3 μ g/insect in the control group, and 14.1 ± 0.8 μ g of chitin per insect after chitinase treatment (Figure 5A). Hydrolysis resulted in a reduction of 66% ($\Delta = 27.3$ μ g/insect). Comparatively, the previously described quantitation method estimated a total of 43.1 ± 1.5 μ g/insect for control samples, and 14.9 μ g/insect after chitinase hydrolyzation (Figure 5A), equally amounting to a 66% reduction in chitin concentration ($\Delta = 28.3$ μ g/insect).



For adults of the same species, males were homogenized individually in 1 mL of distilled water and processed similarly to previous groups. Chitin mass assessment using the calcofluor method estimated a mean of 4600 ± 700 μ g/insect, and 1900 ± 100 μ g of chitin/insect after sample hydrolysis with chitinase (Figure 5B). Using the Lehmann and White (1975) method, the quantitation of chitin in *R. prolixus* adult males totalized a mass of 6406 ± 416 μ g/insect in the control group, and 3682 ± 350 μ g/insect following the chitinase incubation step (Figure 5B). Polysaccharide hydrolysis resulted in a reduction of 59% ($\Delta = 2753$ μ g/insect), and 43% ($\Delta = 2724$ μ g/insect), respectively.

Alternatively, similar results were obtained by homogenizing bigger insects using a mortar and pestle. Tissues were covered with liquid nitrogen, homogenized while frozen and resuspended in distilled water. For smaller insects, homogenization using a pellet pestle cordless motor (Z359971 Sigma) also gave similar results (data not shown). All measurements are summarized in Table 2.

DISCUSSION

Chitin is a polysaccharide composed of an extended hydrogen-bonded chain of repeated β -(1-4)-linked *N*-acetyl-D-glucosamine. This biopolymer is one of the most abundant regarding biomass in nature, second only to cellulose, being synthesized by numerous living organisms, mainly arthropods, and fungi (Rinaudo, 2006; Cohen, 2010). Amongst the Arthropoda, insects rely on this bioproduct for fundamental physiological functions, and it is present in various extracellular structures, such as the integument (Cohen, 2010).

Insect cuticle is composed of layers of chitin separated by layers of protein, as X-ray analyses data suggested (Fraenkel and Rudall, 1946), possibly containing variations of the purely poly-*N*-acetyl-glucosamine, with about one glucosamine in six residues, and about 5% bound water even in thoroughly dried



samples (Rudall, 1963). Generally, protein is the predominant compound in insect cuticle, corresponding to a range of 40–75% (p/p) of crude protein on dry weight basis, varying depending on the species and life cycle stage (Finke, 2007). In the exuviae, protein varies between 44 and 61%, and chitin can account for a range of 30–42% (Kramer et al., 1995).

This biopolymer has become a new and attractive target for pest control, because of its differential distribution amongst living organisms, which would reduce possible unwanted health risks for vertebrates (Cohen, 1993). The insecticide class of Insect Growth Regulators was introduced in 1972 with the discovery of the effects of Diflubenzuron, the first prototype for the benzoylurea chitin synthesis inhibitor insecticides

(Verloop and Ferrell, 1977). Because of the potent insecticidal activities of the class, chitin quantitation emerges as an important diagnosis of the effects of these compounds in insect chitin physiology. Besides that, a rapid and fast chitin measurement technique would allow the high throughput exploration of resistance mechanisms linked to changes in chitin quantity, composition and structure in field samples. It is noteworthy to point out that resistance against chitin synthesis inhibitors was already established in mosquitoes (Araújo et al., 2019; Porretta et al., 2019) and other insects (Wang et al., 2019), and that changes in cuticle composition are the basis of an important resistance mechanism related to reduced penetration of toxic molecules (Balabanidou et al., 2018).




TABLE 2 | Chitin amounts per insect in three different species, as measured by with Lehmann and White and Calcofluor white staining techniques.

Insect Species	Lehmann and White microplate procedure				Calcofluor white staining			
	Before chitinase treatment (μ g)		After chitinase treatment (μ g)		Before chitinase treatment (μ g)	After chitinase treatment (μ g)		
	mean \pm SEM	mean \pm SEM	mean \pm SEM	Net change	mean \pm SEM	mean \pm SEM	Net change	
<i>R. prolixus</i>	First instar	–	43 \pm 2	14.9 \pm 0.9	28.3	41 \pm 3	14.1 \pm 0, 8	27.3
	Adult	Male	6406 \pm 416	3682 \pm 350	2724	4669 \pm 729	1916 \pm 198	2753
<i>L. longipalpis</i>	Adult	Male	9.8 \pm 0.3	7.0 \pm 0.1	2.80	7.6 \pm 0.4	2.64 \pm 0.02	4.93
		Female	13.5 \pm 0.6	8.0 \pm 0.2	5.51	12.0 \pm 0.5	3.13 \pm 0.07	8.83
<i>A. aegypti</i>	Adult	Male	29.9 \pm 0.8	16.6 \pm 0.4	13.2	24.6 \pm 0.7	13.2 \pm 0.6	11.3
		Female	60 \pm 2	35 \pm 2	25	47 \pm 3	30 \pm 2	17

Measurements were performed in crude homogenates, before and after treatment with commercial chitinase from *Streptomyces griseus*. Values are means with standard error of the mean (SEM), μ g per insect, from at least 10 biological replicates. See section "Materials and Methods" for details.

Chitin quantitation has been proposed through different approaches, mainly focusing on the application of the methods in fungal cell wall analyses. Protocols described chitin acid hydrolysis using hydrochloric acid (Paech and Tracey, 1956; Chen and Johnson, 1983) or deacetylation through high-temperature incubation with saturated potassium hydroxide (Ride and Drysdale, 1972; Lehmann and White, 1975) and measured the chitin content colorimetrically from samples as the equivalent in glucosamine released, using glucosamine standard curves. Alternative methods emerged, proposing the direct assessment of the polysaccharide in question using the products of chitinase hydrolyzates (Subramanyam and Rao, 1987). However, still correlating the total amount of chitin to the *N*-acetyl-glucosamine released by the hydrolysis catalyzed by the enzyme. Moreover, despite being applicable to insect samples, these protocols are time-consuming, extremely laborious, and low-throughput, which reduces the number of samples that can be processed at the same time.

Direct chitin content evaluation has also been achieved in an analytical chemistry approach, such as the use of micro

distillation (Kan et al., 1964), gas chromatography (Sweeley et al., 1963), or colloidal titration (Tôei and Kohara, 1976). Alternatively, previous works involving the use of insects as sustainable nutritional sources calculate the chitin concentration as the corrected measurement for the acid detergent fiber (ADF) fraction minus the percentage that corresponds to amino acids (Finke, 2002). This method of analysis can overestimate the quantitation, because the amino acids methionine, cysteine, and tryptophan were not included in the ADF fraction measurement (Finke, 2007).

The use of the chitin specific stain, calcofluor fluorescent brightener, has been previously proposed while associated with the use of flow cytometry to quantify the amount of chitin present in the fungal cell wall (Costa-de-Oliveira et al., 2013). Authors quantified the fluorescent intensity of stained cells in some yeast strain isolates, and results are showed as a mean staining index. Furthermore, Costa-de-Oliveira et al. (2013) infer that measurements are in agreement with previous works (Bulawa et al., 1995; Plaine et al., 2008; Walker et al., 2008),

but the arbitrary units used in the data seem to be difficult to extrapolate.

With the known use of calcofluor for topological localization of chitin in fungal cells and the known affinity of the stain to its target, we were able to quantify successively small amounts of chitin in homogenized samples of the proposed insect species. Our calcofluor staining method results in calculated measurements for insects individually, through microplate fluorescent intensity readings. An adapted protocol of Lehmann and White (1975) for microplate assays was chosen as a comparative control method. Several previous works involving chitin assessment in insect samples have used the Lehmann and White technique (Lehmann and White, 1975; Zhang and Zhu, 2006; Rezende et al., 2008; Farnesi et al., 2012; Chen et al., 2013), and it has been similarly applied to other arthropods (Hackman and Goldberg, 1981). However, to make the measurements obtained with both techniques comparable to each other, we decided to use colloidal chitin instead of glucosamine for the calibration curves.

Our results show that *L. longipalpis* females have more chitin content than males, with a composition of about 12 μg of chitin per insect, while the individual male body is composed of about 8 μg of chitin. The same tendency is seen in *A. aegypti*, with females having about 47 μg chitin/insect, and males around 25 μg of chitin per insect. The females tend to be larger than males in both cases and may have eggs in the ovaries (Farnesi et al., 2015; Souza et al., 2019), which would account for the differences seen in measurements. *Rhodnius prolixus* first instar nymphs have 41 μg of chitin/insect, and adult males increase this total to a mean of 4600 μg /insect, an enhancement of 1000 fold in the concentration of the polymer, which is consistent with the growth of the insects.

For *L. longipalpis* and *A. aegypti* females the measured chitin content corresponds to about 3 and 4% of the insects' total wet weight, respectively, and for *R. prolixus*, it corresponds to 10% for first instar nymphs' wet weight and 8% for adult males. The percentage of the insect weight that corresponds to chitin is similar to what is seen in other insect species. Previous works described that amongst the studied species the chitin content could range from 0,27%, as is the case of the bee brood, to 5% in mealworm beetles (Finke, 2007, 2013).

Comparisons between the two methods resulted in similar amounts of chitin in all tested cases, although some statistically significant differences were observed. Differences may be due to interference in the control method caused by the presence of glucosamine from non-chitin sources. Subramanyam and Rao (1987) verified that a significant part of the measured chitin accounts for hexosamine contaminants, even in commercialized chitin.

Chitinase hydrolysis was used as a proof of concept, showing that the fluorescence obtained from calcofluor stained samples related to the amount of chitin present. As expected, in all species, incubation with the enzyme resulted in a significant reduction of the total amount of chitin estimated by the calculations. Residual amounts of staining were observed in all

hydrolyzate samples, possibly because of the unbalanced ratio of enzyme to the substrate (≥ 0.0002 units per microliter of the sample) or the incubation length (18 h). Chitinase preparations from cockroaches may take up to 6 days to digest about 10% of the substrate (Waterhouse et al., 1961). Nevertheless, when saturated to ten units of enzyme per nmol of chitin, chitinase is able to completely solubilize the chitin present in fungal cell walls in 10 h (Subramanyam and Rao, 1987). In *L. longipalpis*, the signal reduction after hydrolysis showed to be less effective in the colorimetric method when compared to calcofluor staining. This data probably suggests that the chitinase activity in this particular sample results in the noteworthy production of free glucosamine, which has been recorded to represent up to about 10% of chitin digests (Waterhouse et al., 1961). Besides that, the treatment with chitinase seems to be less effective in *A. aegypti* samples with respect to the other species analyzed (Table 2).

Interestingly, the calcofluor fluorescent brightener at a concentration of 1 mg/mL was efficient in the quantitation of smaller insects, namely both males and females *L. longipalpis*, and *R. prolixus* first instar nymphs. However, in *R. prolixus* adults, and *A. aegypti* males and females, we observed underestimations in chitin quantitation (data not shown). This may be related to the higher amount of chitin present in the insect's body (*R. prolixus*) or to different accessibility of the chitin microfibrils to the stain (*A. aegypti*). Dilution of the samples resulted in low reproducibility, probably because of the difficulty of getting homogeneous aliquots from a diluted suspension. Therefore, the stain concentration was enhanced to 50 mg/mL, and for that, the solvent of the stock solution was changed to DMSO. In this way, the sample and calcofluor concentration need to be taken into account, since chitin saturation may result in a low polymer/stain ratio, and, consequently, ineffective quantitation due to underestimation. Thus the volume input was represented as an insect equivalent (Table 1), meaning the number of insects per sample that are necessary for a correct quantitation.

Doubling sample concentrations returns doubled fluorescent intensity signal, but does not change final estimation in chitin concentration per insect (data not shown). It is important to notice that the high background observed after chitinase treatment was similar in both techniques tested, meaning that calcofluor might be used for relative comparison between insects groups in the same way as the Lehmann and White technique. Interestingly, the net change observed after chitinase treatment was very similar for both techniques (Table 2), suggesting that they are detecting the same chemical entity. In the future, it would be important to characterize the source of this residual background, as it would correspond to undigested chitin or other glucosamine containing molecules, as glycoproteins or glycolipids.

The chitin determination protocol described herein constitute a simple, fast, and considerably less laborious method in comparison with previous quantitation methods. However, the necessity of a more sophisticated equipment, as a fluorescence microplate reader, may be a restriction for its application in basic-equipped laboratories. Furthermore, analyses demonstrate

that the assay is consistent and reliable, suggesting a possible incorporation in screening for insecticide resistance mechanisms against insect growth regulators (IGRs).

DATA AVAILABILITY STATEMENT

All datasets generated for this study are included in the article/supplementary material.

AUTHOR CONTRIBUTIONS

FG and BH conceived and designed the study. BH performed the experiments and data analysis. PA and EG gave important intellectual contributions.

REFERENCES

- Araújo, A. P., de Paiva, M. H. S., Cabral, A. M., Cavalcanti, A. E. H. D., Pessoa, L. F. F., Diniz, D. F. A., et al. (2019). Screening *Aedes aegypti* (Diptera: Culicidae) populations from Pernambuco, Brazil for resistance to temephos, Diflubenzuron, and Cypermethrin and characterization of potential resistance mechanisms. *J. Insect Sci.* 19, 1–15. doi: 10.1093/jisesa/iez054
- Balabanidou, V., Grigoraki, L., and Vontas, J. (2018). Insect cuticle: a critical determinant of insecticide resistance. *Curr. Opin. Insect. Sci.* 27, 68–74. doi: 10.1016/j.cois.2018.03.001
- Bulawa, C. E., Miller, D. W., Henry, L. K., and Becker, J. M. (1995). Attenuated virulence of chitin-deficient mutants of *Candida albicans*. *Proc. Natl. Acad. Sci. U.S.A.* 92, 10570–10574. doi: 10.1073/pnas.92.23.10570
- Chen, G. C., and Johnson, B. R. (1983). Improved colorimetric determination of cell wall chitin in wood decay fungi. *Appl. Environ. Microbiol.* 46, 13–16. doi: 10.1128/aem.46.1.13-16.1983
- Chen, L., Yang, W.-J., Cong, L., Xu, K.-K., and Wang, J.-J. (2013). Molecular cloning, characterization and mRNA expression of a chitin synthase 2 gene from the oriental fruit fly, *Bactrocera dorsalis* (Diptera: Tephritidae). *Int. J. Mol. Sci.* 14, 17055–17072. doi: 10.3390/ijms140817055
- Cohen, E. (1993). Chitin synthesis and degradation as targets for pesticide action. *Arch. Insect Biochem. Physiol.* 22, 245–261. doi: 10.1002/arch.940220118
- Cohen, E. (2010). *Chitin Biochemistry: Synthesis, Hydrolysis and Inhibition. Advances in Insect Physiology*, 1st Edn, Vol. 38. Amsterdam: Elsevier Ltd.
- Costa-de-Oliveira, S., Silva, A. P., Miranda, I. M., Salvador, A., Azevedo, M. M., Munro, C. A., et al. (2013). Determination of chitin content in fungal cell wall: an alternative flow cytometric method. *Cytometry A* 83, 324–328. doi: 10.1002/cyto.a.22250
- Dubois, M., Gilles, K. A., Hamilton, J. K., Rebers, P. A., and Smith, F. (1956). Colorimetric method for determination of sugars and related substances. *Anal. Chem.* 28, 350–356. doi: 10.1021/ac60111a017
- Farnesi, L. C., Brito, J. M., Linss, J. G., Pelajo-Machado, M., Valle, D., and Rezende, G. L. (2012). Physiological and morphological aspects of *Aedes aegypti* developing larvae: effects of the chitin synthesis inhibitor novaluron. *PLoS One* 7:e30363. doi: 10.1371/journal.pone.0030363
- Farnesi, L. C., Menna-Barreto, R. F. S., Martins, A. J., Valle, D., and Rezende, G. L. (2015). Physical features and chitin content of eggs from the mosquito vectors *Aedes aegypti*, *Anopheles aquasalis* and *Culex quinquefasciatus*: connection with distinct levels of resistance to desiccation. *J. Insect. Physiol.* 83, 43–52. doi: 10.1016/j.jinsphys.2015.10.006
- Finke, M. D. (2002). Complete nutrient composition of commercially raised invertebrates used as food for insectivores. *Zoo Biol.* 21, 269–285. doi: 10.1002/zoo.10031
- Finke, M. D. (2007). Estimate of chitin in raw whole insects. *Zoo Biol.* 26, 105–115. doi: 10.1002/zoo
- Finke, M. D. (2013). Complete nutrient content of four species of feeder insects. *Zoo Biol.* 32, 27–36. doi: 10.1002/zoo.21012

FUNDING

This research was supported by the Brazilian research agencies FAPERJ, CNPq, CAPES, and FIOCRUZ (PAPES, PAEF, and PROEP Programs). BH received M.S. and Ph.D. grants, respectively from CNPq (Parasitology Postgraduation Program) and CAPES/FAPERJ (Cellular and Molecular Biology Postgraduation Program).

ACKNOWLEDGMENTS

The authors thank Zildo Alves da Cruz (IOC-FIOCRUZ), José Carlos Nascimento (IOC-FIOCRUZ), and Samara Graciane da Costa (IOC-FIOCRUZ) for technical assistance in the care of the insect facilities.

- Fraenkel, G., and Rudall, K. M. (1946). The structure of insect cuticles. *Proc. R. Soc. B* 134, 111–143.
- García, E. S., Azambuja, P., and Contreras, V. (1984). “Large-scale rearing of *Rhodnius prolixus* and production of metacyclic trypomastigotes of *Trypanosoma cruzi*,” in *Genes and Antigens of Parasites, a Laboratory Manual*, 2nd Edn, ed. C. M. Morel, (Rio de Janeiro: Fundação Oswaldo Cruz), 44–47.
- Hackman, R. H., and Goldberg, M. (1981). A method for determinations of microgram amounts of chitin in arthropod cuticles. *Anal. Biochem.* 110, 277–280. doi: 10.1016/0003-2697(81)90192-5
- Herth, W. (1980). Calcofluor white and congo red inhibit chitin microfibril assembly of poteriocochromonas: evidence for a gap between polymerization and microfibril formation. *J. Cell Biol.* 87, 442–450. doi: 10.1083/jcb.87.2.442
- Herth, W., and Schnepf, E. (1980). The fluorochrome, calcofluor white, binds oriented to structural polysaccharide fibrils. *Protoplasma* 105, 129–133. doi: 10.1007/BF01279855
- Hsu, S. C., and Lockwood, J. L. (1975). Powdered chitin agar as a selective medium for enumeration of actinomycetes in water and soil. *Appl. Microbiol.* 29, 422–426. doi: 10.1128/aem.29.3.422-426.1975
- Kan, M., Suzuki, F., and Kashiwagi, H. (1964). Microdetermination of formyl group in organic compounds in the presence of acetyl group and vice versa. *Microchem. J.* 8, 42–50. doi: 10.1016/0026-265X(64)90010-4
- Kramer, K. J., Hopkins, T. L., and Schaefer, J. (1995). Applications of solids NMR to the analysis of insect sclerotized structures. *Insect Biochem. Mol. Biol.* 25, 1067–1080. doi: 10.1016/0965-1748(95)00053-4
- Lehmann, P. F., and White, L. O. (1975). Chitin assay used to demonstrate renal localization and cortisone enhanced growth of *Aspergillus fumigatus* mycelium in mice. *Infect. Immun.* 12, 987–992. doi: 10.1128/iai.12.5.987-992.1975
- Moraes, C. S., Diaz-Albiter, H. M., do Faria, M. V., Sant’Anna, M. R. V., Dillon, R. J., and Genta, F. A. (2014). Expression pattern of glycoside hydrolase genes in *Lutzomyia longipalpis* reveals key enzymes involved in larval digestion. *Front. Physiol.* 5:276. doi: 10.3389/fphys.2014.00276
- Paech, K., and Tracey, M. V. (1956). *Modern Methods of Plant Analysis*. Cham: Springer.
- Pillai, C. K. S., Paul, W., and Sharma, C. P. (2009). Chitin and chitosan polymers: chemistry, solubility and fiber formation. *Prog. Polym. Sci.* 34, 641–678. doi: 10.1016/j.progpolymsci.2009.04.001
- Plaine, A., Walker, L., Da Costa, G., Mora-Montes, H. M., McKinnon, A., Gow, N. A. R., et al. (2008). Functional analysis of *Candida albicans* GPI-anchored proteins: roles in cell wall integrity and caspofungin sensitivity. *Fungal Genet. Biol.* 45, 1404–1414. doi: 10.1016/j.fgb.2008.08.003
- Porretta, D., Fotakis, E. A., Mastrantonio, V., Chaskopoulou, A., Michaelakis, A., Koulos, I., et al. (2019). Focal distribution of di fl ubenzuron resistance mutations in *Culex pipiens* mosquitoes from Northern Italy. *Acta Trop.* 193, 106–112. doi: 10.1016/j.actatropica.2019.02.024
- Rezende, G. L., Martins, A. J., Gentile, C., Farnesi, L. C., Pelajo-Machado, M., Peixoto, A. A., et al. (2008). Embryonic desiccation resistance in *Aedes aegypti*:

- presumptive role of the chitinized serosal cuticle. *BMC Dev. Biol.* 8:82. doi: 10.1186/1471-213X-8-82
- Ride, J., and Drysdale, R. B. (1972). A rapid method for the chemical estimation of filamentous fungi in plant tissue. *Physiol. Plant Pathol.* 2, 7–15. doi: 10.1016/0048-4059(72)90043-4
- Rinaudo, M. (2006). Chitin and chitosan: properties and applications. *Prog. Polym. Sci.* 31, 603–632. doi: 10.1016/j.propolymsci.2006.06.001
- Rudall, K. M. (1963). The chitin/protein complexes of insect Cuticles. *Adv. Insect Physiol.* 1, 257–313. doi: 10.1016/S0065-2806(08)60177-0
- Souza, R. S., Diaz-Albiter, H. M., Dillon, V. M., Dillon, R. J., and Genta, F. A. (2016). Digestion of yeasts and beta-1,3-glucanases in mosquito larvae: physiological and biochemical considerations. *PLoS One* 11:e0151403. doi: 10.1371/journal.pone.0151403
- Souza, R. S., Virginio, F., Riback, T. I. S., Suesdek, L., Barufi, J. B., and Genta, F. A. (2019). Microorganism-based larval diets affect mosquito development, size and nutritional reserves in the yellow fever mosquito *Aedes aegypti* (Diptera: Culicidae). *Front. Physiol.* 10:152. doi: 10.3389/fphys.2019.00152
- Subramanyam, C., and Rao, S. L. N. (1987). An enzymic method for the determination. *J. Biosci.* 12, 125–129.
- Sweeley, C. C., Bentley, R., Makita, M., and Wells, W. W. (1963). Gas-liquid chromatography of trimethylsilyl derivatives of sugars and related substances. *J. Am. Chem. Soc.* 85, 2497–2507. doi: 10.1021/ja00899a032
- Tôei, K., and Kohara, T. (1976). Naturally occurring or synthesized polyelectrolytes, whether positive or Reagents. *Anal. Chim. Acta* 83, 59–65.
- Verloop, A., and Ferrell, C. D. (1977). *Benzoylphenyl Ureas a New Group of Larvicides Interfering with Chitin Deposition*. Washington, DC: American Chemical Society, 237–270.
- Walker, L. A., Munro, C. A., De Bruijn, I., Lenardon, M. D., McKinnon, A., and Gow, N. A. R. (2008). Stimulation of chitin synthesis rescues *Candida albicans* from echinocandins. *PLoS Pathog.* 4:e1000040. doi: 10.1371/journal.ppat.1000040
- Wang, Y., Li, Y.-Z., Li, G.-Q., Wan, P.-J., and Li, C. (2019). Identification of cuticular protein genes in the colorado potato beetle *Leptinotarsa decemlineata* (Coleoptera: Chrysomelidae). *J. Econ. Entomol.* 112, 912–923. doi: 10.1093/jeetoy396
- Waterhouse, D. F., Hackman, R. H., and McKellar, J. W. (1961). An investigation of chitinase activity in cockroach and termite extracts. *J. Insect Physiol.* 6, 96–112. doi: 10.1016/0022-1910(61)90032-4
- Zhang, J., and Zhu, K. Y. (2006). Characterization of a chitin synthase cDNA and its increased mRNA level associated with decreased chitin synthesis in *Anopheles quadrimaculatus* exposed to diflubenzuron. *Insect Biochem. Mol. Biol.* 36, 712–725. doi: 10.1016/j.ibmb.2006.06.002

Conflict of Interest: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2020 Henriques, Garcia, Azambuja and Genta. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.