

# DETERMINATION OF THE STRUCTURE OF ALGEBRAIC CURVATURE TENSORS BY MEANS OF YOUNG SYMMETRIZERS

BERND FIEDLER

ABSTRACT. For a positive definite fundamental tensor all known examples of Osserman algebraic curvature tensors have a typical structure. They can be produced from a metric tensor and a finite set of skew-symmetric matrices which fulfil Clifford commutation relations. We show by means of Young symmetrizers and a theorem of S. A. Fulling, R. C. King, B. G. Wybourne and C. J. Cummins that every algebraic curvature tensor has a structure which is very similar to that of the above Osserman curvature tensors. We verify our results by means of the Littlewood–Richardson rule and plethysms. For certain symbolic calculations we used the Mathematica packages `MathTensor`, `Ricci` and `PERMS`.

## 1. INTRODUCTION

Let  $\mathcal{T}_r V$  be the vector space of the  $r$ -times covariant tensors  $T$  over a finite-dimensional  $\mathbb{K}$ -vector space  $V$ ,  $\mathbb{K} = \mathbb{R}$  or  $\mathbb{K} = \mathbb{C}$ .

**Definition 1.1.** A tensor  $T \in \mathcal{T}_4 V$  is called an *algebraic curvature tensor* iff  $T$  has the *index commutation symmetry*

$$(1.1) \quad \forall u, x, y, z \in V : \quad T(u, x, y, z) = -T(u, x, z, y) = T(y, z, u, x)$$

and fulfills the *first Bianchi identity*

$$(1.2) \quad \forall u, x, y, z \in V : \quad T(u, x, y, z) + T(u, y, z, x) + T(u, z, x, y) = 0.$$

If we consider the coordinates  $T_{ijkl}$  of a tensor  $T \in \mathcal{T}_r V$  then  $T$  is an algebraic curvature tensor iff its coordinates satisfy

$$(1.3) \quad T_{ijkl} = -T_{jikl} = -T_{ijlk} = T_{klij}$$

and

$$(1.4) \quad T_{ijkl} + T_{iklj} + T_{iljk} = 0.$$

with respect to every basis of  $V$ .

We assume that  $V$  possesses a fundamental tensor  $g \in \mathcal{T}_2 V$  (of arbitrary signature) which can be used for raising and lowering of tensor indices.

**Definition 1.2.** Let  $T \in \mathcal{T}_4 V$  be an algebraic curvature tensor and  $x \in V$  be a vector with  $|g(x, x)| = 1$ . The *Jacobi operator*  $J_T(x)$  of  $T$  and  $x$  is the linear operator  $J_T(x) : V \rightarrow V$ ,  $J_T(x) : y \mapsto J_T(x)y$  that is defined by

$$(1.5) \quad \forall w \in V : \quad g(J_T(x)y, w) = T(y, x, x, w).$$

Furthermore we use the following operators  $\gamma$  and  $\alpha$ :

**Definition 1.3.** (1) Let  $S \in \mathcal{T}_2V$  be a symmetric tensor of order 2, i.e. the coordinates of  $S$  satisfy  $S_{ij} = S_{ji}$ . We define a tensor  $\gamma(S) \in \mathcal{T}_4V$  by

$$(1.6) \quad \gamma(S)_{ijkl} := \frac{1}{3} (S_{il}S_{jk} - S_{ik}S_{jl}) .$$

(2) Let  $A \in \mathcal{T}_2V$  be a skew-symmetric tensor of order 2, i.e. the coordinates of  $A$  satisfy  $A_{ji} = -A_{ij}$ . We define a tensor  $\alpha(A) \in \mathcal{T}_4V$  by

$$(1.7) \quad \alpha(A)_{ijkl} := \frac{1}{3} (2A_{ij}A_{kl} + A_{ik}A_{jl} - A_{il}A_{jk}) .$$

Tensors of type  $\gamma(S)$  and  $\alpha(A)$  occur in formulas of so-called *Osserman tensors* (see [24]).

**Definition 1.4.** An algebraic curvature tensor  $T$  is called *spacelike Osserman* (respectively *timelike Osserman*) if the eigenvalues of  $J_T(x)$  are constant on  $S^+(V) := \{x \in V \mid g(x, x) = +1\}$  (respectively  $S^-(V) := \{x \in V \mid g(x, x) = -1\}$ ).

Since the fundamental tensor  $g$  connects an algebraic curvature tensor  $T$  with its Jacobi operator  $J_T(x)$  in (1.5), the Osserman property depends not only on  $T$  but also on  $g$ . It is known that spacelike Osserman and timelike Osserman are equivalent notions so one simply says *Osserman*.

If  $R$  is the Riemann tensor of a Riemannian manifold  $(M, g)$  which is locally a rank one symmetric space or flat, then the eigenvalues of  $J_R(x)$  are constant on the unit sphere. Osserman [37] wondered if the converse held. This question is known as the *Osserman conjecture*.

The correctness of the Osserman conjecture has been established for Riemannian manifolds  $(M, g)$  in all dimensions  $\neq 8, 16$  (see [7, 36]) and for Lorentzian manifolds  $(M, g)$  in all dimensions (see [1, 20]). The situation is much more complicated in the case of a pseudo-Riemannian metric with signature  $(p, q)$ ,  $p, q \geq 2$ . It is known that there exist pseudo-Riemannian Osserman manifolds which are not locally homogeneous [2, 6, 22]. Furthermore there are counter examples to the conjecture if the manifold  $(M, g)$  has signature  $(2, 2)$  and if the Jacobi operator is not diagonalizable [2, 3]. (See also the examples in [21, 38]). Thus the question Osserman raised has a negative answer in the higher signature setting. More extensive bibliographies in the subject can be found in the books [26] by P. B. Gilkey and [23] by E. García-Río, D. N. Kupeli and R. Vázquez-Lorenzo.

Examples of Osserman algebraic curvature tensors can be constructed in the following way. Let  $\{C_i\}_{i=1}^r$  be a finite set of real, skew-symmetric  $(\dim V \times \dim V)$ -matrices that satisfy the Clifford commutation relations

$$(1.8) \quad C_i \cdot C_j + C_j \cdot C_i = -2\delta_{ij} .$$

We define

$$(1.9) \quad T_C(x, y)z := g(y, C \cdot z) C \cdot x - g(x, C \cdot z) C \cdot y - 2g(x, C \cdot y) C \cdot z ,$$

$$(1.10) \quad T_0(x, y)z := g(y, z)x - g(x, z)y, \quad x, y, z \in V ,$$

$$(1.11) \quad T := \lambda_0 T_0 + \lambda_1 T_{C_1} + \dots + \lambda_r T_{C_r} .$$

Now if  $g$  is positive definite, then  $T$  is an algebraic curvature tensor that is *Osserman*. (See [25, 27, 24] for more details.)

$T_0$  and  $T_C$  can be expressed by means of  $\gamma$  and  $\alpha$ , respectively, since

$$\begin{aligned} g(T_0(x, y)z, w) &= g(y, z)g(x, w) - g(x, z)g(y, w) \\ &= (g_{jk}g_{il} - g_{ik}g_{jl})x^i y^j z^k w^l \\ &= 3\gamma(g)_{ijkl}x^i y^j z^k w^l \end{aligned}$$

and

$$\begin{aligned} g(T_C(x, y)z, w) &= g(y, C \cdot z)g(C \cdot x, w) - g(x, C \cdot z)g(C \cdot y, w) - \\ &\quad 2g(x, C \cdot y)g(C \cdot z, w) \\ &= g_{js}y^j C^s_k z^k g_{il}C^t_i x^i w^l - g_{is}x^i C^s_k z^k g_{tl}C^t_j y^j w^l - \\ &\quad 2g_{is}x^i C^s_j y^j g_{tl}C^t_k z^k w^l \\ &= \{C_{jk}C_{li} - C_{ik}C_{lj} - 2C_{ij}C_{lk}\}x^i y^j z^k w^l \\ &= \{-C_{jk}C_{il} + C_{ik}C_{jl} + 2C_{ij}C_{kl}\}x^i y^j z^k w^l \\ &= 3\alpha(C)_{ijkl}x^i y^j z^k w^l. \end{aligned}$$

Thus we obtain

$$(1.12) \quad T = 3\lambda_0\gamma(g) + 3\sum_{i=1}^r \lambda_i \alpha(C_i)$$

from (1.11). It was conjectured that any algebraic Osserman curvature tensor has a structure given in (1.11). This is true for all known examples of Osserman tensors in the case of a positive definite metric  $g$  (see [24]). But for indefinite metrics there exist counter examples (see Remark 6.7).

In the present paper we show that every algebraic curvature tensor  $T$  has a representation

$$(1.13) \quad T = \sum_{S \in \mathcal{S}} \epsilon_S \gamma(S) + \sum_{A \in \mathcal{A}} \epsilon_A \alpha(A) \quad , \quad \epsilon_S, \epsilon_A \in \{1, -1\},$$

where  $\mathcal{S}$  and  $\mathcal{A}$  are finite sets of symmetric or skew-symmetric tensors of order 2, respectively. (See Theorem 3.1.) To prove (1.13) we use the connection between tensors and the representation theory of symmetric groups. We discussed the formula (1.13) with P. B. Gilkey. In the course of this discussion P. B. Gilkey found another proof of (1.13) which is based on tensor algebra (see [26, pp. 41–44]). More precisely, P. B. Gilkey could show the stronger result that every algebraic curvature tensor has a representation

$$(1.14) \quad T = \sum_{S \in \mathcal{S}} \epsilon_S \gamma(S) \quad , \quad \epsilon_S \in \{1, -1\},$$

as well as a representation

$$(1.15) \quad T = \sum_{A \in \mathcal{A}} \epsilon_A \alpha(A) \quad , \quad \epsilon_A \in \{1, -1\}.$$



Obviously, a Young frame of  $r$  is characterized by a partition  $\lambda = (\lambda_1, \dots, \lambda_l) \vdash r$  of  $r$ . A *Young tableau*  $t$  of a partition  $\lambda \vdash r$  is the Young frame corresponding to  $\lambda$  which was fulfilled by the numbers  $1, 2, \dots, r$  in any order.

$$t = \begin{array}{|c|c|c|c|c|} \hline 11 & 2 & 5 & 4 & 12 \\ \hline 9 & 6 & 16 & 15 & \\ \hline 8 & 14 & 1 & 7 & \\ \hline 13 & 3 & & & \\ \hline 10 & & & & \\ \hline \end{array}$$

If a Young tableau  $t$  of a partition  $\lambda \vdash r$  is given, then the *Young symmetrizer*  $y_t$  of  $t$  is defined by<sup>1</sup>

$$(2.6) \quad y_t := \sum_{p \in \mathcal{H}_t} \sum_{q \in \mathcal{V}_t} \text{sign}(q) p \circ q$$

where  $\mathcal{H}_t, \mathcal{V}_t$  are the groups of the *horizontal* or *vertical permutations* of  $t$  which only permute numbers within rows or columns of  $t$ , respectively. The Young symmetrizers of  $\mathbb{K}[\mathcal{S}_r]$  are essentially idempotent and define decompositions

$$(2.7) \quad \mathbb{K}[\mathcal{S}_r] = \bigoplus_{\lambda \vdash r} \bigoplus_{t \in \mathcal{ST}_\lambda} \mathbb{K}[\mathcal{S}_r] \cdot y_t \quad , \quad \mathbb{K}[\mathcal{S}_r] = \bigoplus_{\lambda \vdash r} \bigoplus_{t \in \mathcal{ST}_\lambda} y_t \cdot \mathbb{K}[\mathcal{S}_r]$$

of  $\mathbb{K}[\mathcal{S}_r]$  into minimal left or right ideals. In (2.7), the symbol  $\mathcal{ST}_\lambda$  denotes the set of all standard tableaux of the partition  $\lambda$ . Standard tableaux are Young tableaux in which the entries of every row and every column form an increasing number sequence.<sup>2</sup>

S.A. Fulling, R.C. King, B.G.Wybourne and C.J. Cummins showed in [17] that the symmetry classes of the Riemannian curvature tensor  $R$  and its covariant derivatives  $\nabla^{(u)}R$  are generated by special Young symmetrizers. To this end they assumed that the covariant derivatives  $\nabla_{s_1} \nabla_{s_2} \dots \nabla_{s_u} R_{ijkl}$  are symmetric in  $s_1, \dots, s_u$  for  $u \geq 2$ . This is possible if one uses Ricci identities and neglect derivatives of orders smaller than  $u$ . Under this assumption one has

**Theorem 2.3.** *Consider the Levi-Civita connection  $\nabla$  of a pseudo-Riemannian metric  $g$ . For  $u \geq 0$  the Riemann tensor and its covariant derivatives  $\nabla^{(u)}R$  fulfil*

$$(2.8) \quad e_t^* \nabla^{(u)}R = \nabla^{(u)}R$$

where  $e_t := y_t(u+1)/(2 \cdot (u+3)!)$  is an idempotent which is formed from the Young symmetrizer  $y_t$  of the standard tableau

$$(2.9) \quad t = \begin{array}{|c|c|c|c|c|} \hline 1 & 3 & 5 & \dots & \dots & (u+4) \\ \hline 2 & 4 & & & & \\ \hline \end{array}.$$

<sup>1</sup>We use the convention  $(p \circ q)(i) := p(q(i))$  for the product of two permutations  $p, q$ .

<sup>2</sup>About Young symmetrizers and Young tableaux see for instance [4, 5, 17, 19, 28, 29, 32, 33, 34, 35, 40, 41]. In particular, properties of Young symmetrizers in the case  $\mathbb{K} \neq \mathbb{C}$  are described in [34].

The star " $*$ " denotes the mapping

$$(2.10) \quad * : a = \sum_{p \in \mathcal{S}_r} a(p) p \mapsto a^* := \sum_{p \in \mathcal{S}_r} a(p) p^{-1}.$$

A proof of this result of [17] can be found in [12], too. The proof needs only the symmetry properties (1.3), the identities Bianchi I and Bianchi II and Ricci identities. Thus Theorem 2.3 is a statement about the algebraic curvature tensors and "algebraic" tensors that possess the symmetry properties of the covariant derivatives of the Riemann tensor. In particular, a tensor  $T \in \mathcal{T}_4V$  is an algebraic curvature tensor iff  $T$  satisfies

$$(2.11) \quad y_t^* T = 12 T$$

where  $y_t$  is the Young symmetrizer of the standard tableau

$$(2.12) \quad t = \begin{array}{|c|c|} \hline 1 & 3 \\ \hline 2 & 4 \\ \hline \end{array}.$$

### 3. A STRUCTURE THEOREM FOR ALGEBRAIC CURVATURE TENSORS

Now we will show that every algebraic curvature tensor can be built from certain finite sets of symmetric and skew-symmetric tensors from  $\mathcal{T}_2V$ .

**Theorem 3.1.** *For every algebraic curvature tensor  $T \in \mathcal{T}_4V$  there exist finite sets  $\mathcal{S}$  and  $\mathcal{A}$  of symmetric and alternating tensors  $S, A \in \mathcal{T}_2V$ , respectively, such that*

$$(3.1) \quad T = \sum_{S \in \mathcal{S}} \epsilon_S \gamma(S) + \sum_{A \in \mathcal{A}} \epsilon_A \alpha(A) \quad , \quad \epsilon_S, \epsilon_A \in \{1, -1\}.$$

*Proof.* Let  $T$  be an algebraic curvature tensor. Because of the symmetry  $T_{klij} = T_{ijkl}$ , every algebraic curvature tensor  $T$  fulfils  $fT = T$ , where  $f$  is the idempotent  $f := \frac{1}{2}(\text{id} + (13)(24)) \in \mathbb{K}[\mathcal{S}_4]$ . Thus every algebraic curvature tensor belongs to the symmetry class  $\mathcal{T}_{\mathfrak{r}}$  of the right ideal  $\mathfrak{r} := f \cdot \mathbb{K}[\mathcal{S}_4]$  and has a representation  $T = fT'$  with  $T' \in \mathcal{T}_4V$ .

Since an arbitrary tensor  $T' \in \mathcal{T}_4V$  can be written as a finite sum of decomposable tensors  $a_1 \otimes a_2 \otimes a_3 \otimes a_4 \in \mathcal{T}_4V$ , we can use a representation

$$(3.2) \quad T' = \sum_{(M,N) \in \mathcal{P}} M \otimes N$$

for a  $T' \in \mathcal{T}_4V$ , where  $\mathcal{P}$  is a finite set of pairs  $(M, N)$  of tensors  $M, N \in \mathcal{T}_2V$ . From (3.2) and  $T = fT'$  we obtain

$$\begin{aligned} T = fT' &= \frac{1}{2} \sum_{(M,N) \in \mathcal{P}} (M \otimes N + N \otimes M) \\ &= \frac{1}{2} \sum_{(M,N) \in \mathcal{P}} ((M + N) \otimes (M + N) - M \otimes M - N \otimes N) . \end{aligned}$$

Thus every tensor  $T \in \mathcal{T}_\tau$  has a representation

$$(3.3) \quad T = \frac{1}{2} \sum_{M' \in \mathcal{M}'} z_{M'} M' \otimes M' \quad , \quad z_{M'} \in \mathbb{Z}$$

with a certain finite set  $\mathcal{M}'$  of tensors from  $\mathcal{T}_2V$ . If we carry out a transformation  $M' \mapsto M := \sqrt{|z_{M'}|/2} M'$  in every summand of (3.3), we obtain

$$(3.4) \quad T = \sum_{M \in \mathcal{M}} \epsilon_M M \otimes M \quad , \quad \epsilon_M \in \{1, -1\}$$

from (3.3).

Since every matrix  $M$  has a unique decomposition into a symmetric matrix  $S$  and a skew-symmetric matrix  $A$ , i.e.  $M = S + A$ , (3.4) leads to

$$(3.5) \quad T = \sum_{S+A \in \mathcal{M}} \epsilon_{S+A} (S \otimes S + S \otimes A + A \otimes S + A \otimes A) .$$

Now, let  $y_t$  be the Young symmetrizer of the standard tableau (2.12). If we apply (2.11) to the representation (3.5) of an algebraic curvature tensor  $T$ , we obtain

$$(3.6) \quad T = \frac{1}{12} y_t^* T = \frac{1}{12} \sum_{S+A \in \mathcal{M}} \epsilon_{S+A} y_t^* (S \otimes S + S \otimes A + A \otimes S + A \otimes A) .$$

Then a calculation<sup>3</sup> yields

$$\begin{aligned} \frac{1}{12} y_t^*(S \otimes S) &= \gamma(S) & , & & \frac{1}{12} y_t^*(A \otimes S) &= 0 , \\ \frac{1}{12} y_t^*(A \otimes A) &= \alpha(A) & , & & \frac{1}{12} y_t^*(S \otimes A) &= 0 , \end{aligned}$$

and if we set  $\epsilon_S := \epsilon_{S+A}$ ,  $\epsilon_A := \epsilon_{S+A}$ , we obtain (3.1).  $\square$

**Corollary 3.2.** *If  $\mathcal{S} \subseteq \mathcal{T}_2V$  and  $\mathcal{A} \subseteq \mathcal{T}_2V$  are finite sets of symmetric or alternating tensors, respectively, then*

$$(3.7) \quad T := \sum_{S \in \mathcal{S}} \epsilon_S \gamma(S) + \sum_{A \in \mathcal{A}} \epsilon_A \alpha(A) \quad , \quad \epsilon_S, \epsilon_A \in \{1, -1\}$$

*is an algebraic curvature tensor.*

*Proof.* Every summand of (3.7) is an algebraic curvature tensor since  $\gamma(S) = \frac{1}{12} y_t^*(S \otimes S)$  and  $\alpha(A) = \frac{1}{12} y_t^*(A \otimes A)$ .  $\square$

<sup>3</sup>This calculation can be done by hand or by a cooperation of the Mathematica package PERMS [14] with one of the tensor packages Ricci [31] or MathTensor [8]. See [9].

## 4. VERIFICATION BY THE LITTLEWOOD–RICHARDSON RULE AND PLETHYSMS

We can verify the results about  $y_t^*(S \otimes S)$ ,  $y_t^*(S \otimes A)$ ,  $y_t^*(A \otimes S)$  and  $y_t^*(A \otimes A)$  by means of the *Littlewood–Richardson rule*<sup>4</sup> and *plethysms*<sup>5</sup>. To this end we form group ring elements from tensors (see B. Fiedler [13, Sec.III.1] and [10, 11, 15, 16]).

**Definition 4.1.** Any tensor  $T \in \mathcal{T}_r V$  and any  $r$ -tuple  $b := (v_1, \dots, v_r) \in V^r$  of vectors from  $V$  induce a function  $T_b : \mathcal{S}_r \rightarrow \mathbb{K}$  according to the rule

$$(4.1) \quad T_b(p) := T(v_{p(1)}, \dots, v_{p(r)}) \quad , \quad p \in \mathcal{S}_r .$$

We identify this function with the group ring element  $\sum_{p \in \mathcal{S}_r} T_b(p) p \in \mathbb{K}[\mathcal{S}_r]$ , which we denote by  $T_b$ , too.

These  $T_b$  fulfil<sup>6</sup>

**Corollary 4.2.** *If  $T \in \mathcal{T}_r V$ ,  $b \in V^r$  and  $a \in \mathbb{K}[\mathcal{S}_r]$ , then*

$$(4.2) \quad (aT)_b = T_b \cdot a^* .$$

If a tensor  $T$  belongs to a certain symmetry class, then its  $T_b$  lie in a certain left ideal<sup>6</sup>.

**Proposition 4.3.** *Let  $e \in \mathbb{K}[\mathcal{S}_r]$  be an idempotent. Then a  $T \in \mathcal{T}_r V$  fulfils the condition  $eT = T$  iff  $T_b \in \mathfrak{l} := \mathbb{K}[\mathcal{S}_r] \cdot e^*$  for all  $b \in V^r$ , i.e. all  $T_b$  of  $T$  lie in the left ideal  $\mathfrak{l}$  generated by  $e^*$ .*

Obviously, the  $S_b$  and  $A_b$  of symmetric/skew-symmetric tensors of order 2 lie in the left ideals

$$(4.3) \quad \forall b \in V^2 : S_b \in \mathfrak{l}_1 := \mathbb{K}[\mathcal{S}_2] \cdot e_1 \quad , \quad e_1 := \frac{1}{2} (\text{id} + (12))$$

$$(4.4) \quad \forall b \in V^2 : A_b \in \mathfrak{l}_2 := \mathbb{K}[\mathcal{S}_2] \cdot e_2 \quad , \quad e_2 := \frac{1}{2} (\text{id} - (12))$$

$e_1$  and  $e_2$  are the (normalized) Young symmetrizers of the standard tableaux

$$t_1 := \boxed{1} \boxed{2} \quad , \quad t_2 := \begin{array}{|c|} \hline 1 \\ \hline 2 \\ \hline \end{array}$$

and the left ideals  $\mathfrak{l}_1$  and  $\mathfrak{l}_2$  are 1-dimensional, minimal left ideals, which belong to the equivalence classes of minimal left ideals characterized by the partitions  $\lambda_1 = (2)$  and  $\lambda_2 = (1^2)$ , respectively. Then the group ring elements  $(A \otimes S)_b$ ,  $(S \otimes A)_b$ ,  $(S \otimes S)_b$  and  $(A \otimes A)_b$  belong to left ideals which are characterized by Littlewood–Richardson products  $[2][1^2]$  or plethysms  $[2] \odot [2]$ ,  $[1^2] \odot [2]$  (see

<sup>4</sup>See D. E. Littlewood [32, pp. 94–96], A. Kerber [29, Vol.240/p. 84], G. D. James and A. Kerber [28, p. 93], A. Kerber [30, Sec.5.5], I. G. Macdonald [33, Chap. I, Sec. 9], W. Fulton and J. Harris [18, pp. 455–456], S. A. Fulling, R. C. King, B. G. Wybourne and C. J. Cummins [17]. See also B. Fiedler [13, Sec.II.5].

<sup>5</sup>See A. Kerber [29], G. D. James and A. Kerber [28], A. Kerber [30], F. Sanger [39], D. E. Littlewood [32], I. G. Macdonald [33], S. A. Fulling, R. C. King, B. G. Wybourne and C. J. Cummins [17]. See also B. Fiedler [13, Sec.II.6].

<sup>6</sup>See B. Fiedler [13, Sec.III.3.1] and [10, 11, 15, 16].



B. Fiedler [13, Sec.III.3.2] or [15, Sec.4.2], [16]). Especially, we obtain the following structures for these ideals:

$(S \otimes S)_b$	$[2] \odot [2] \sim [4] + [2^2]$	$\mathfrak{l}^{(1)} = \mathfrak{l}_{[4]}^{(1)} \oplus \mathfrak{l}_{[2^2]}^{(1)}$
$(S \otimes A)_b$	$[2][1^2] \sim [3\ 1] + [2\ 1^2]$	$\mathfrak{l}^{(2)} = \mathfrak{l}_{[3\ 1]}^{(2)} \oplus \mathfrak{l}_{[2\ 1^2]}^{(2)}$
$(A \otimes S)_b$	$[2][1^2] \sim [3\ 1] + [2\ 1^2]$	$\mathfrak{l}^{(3)} = \mathfrak{l}_{[3\ 1]}^{(3)} \oplus \mathfrak{l}_{[2\ 1^2]}^{(3)}$
$(A \otimes A)_b$	$[1^2] \odot [2] \sim [2^2] + [1^4]$	$\mathfrak{l}^{(4)} = \mathfrak{l}_{[2^2]}^{(4)} \oplus \mathfrak{l}_{[1^4]}^{(4)}$

The relation  $[2][1^2] \sim [3\ 1] + [2\ 1^2]$  can be determined by the Littlewood–Richardson rule. The relation  $[2] \odot [2] \sim [4] + [2^2]$  follows from the formula

$$(4.5) \quad [2] \odot [n] \sim \sum_{\lambda \vdash n} [2\ \lambda]$$

(see G.D. James and A. Kerber [28, p. 224]) and  $[1^2] \odot [2] \sim [2^2] + [1^4]$  is a consequence of (4.5) and

$$(4.6) \quad [\mu] \odot [\nu] \sim \sum_{\lambda \vdash m \cdot n} m_\lambda [\lambda'] \Leftrightarrow [\mu'] \odot [\nu] \sim \sum_{\lambda \vdash m \cdot n} m_\lambda [\lambda]$$

$$\mu \vdash m, \nu \vdash n, m, n, m_\lambda \in \mathbb{N}, m \text{ even}.$$

(See F. Sanger [39, pp. 10–11]. See also B. Fiedler [13, Corr.II.6.8].) In (4.6)  $\lambda'$  denotes the associated or transposed partition of a partition  $\lambda$ . All formulas of the above table can be determined by the **Mathematica** package **Perms** [14], too.

Every of the left ideals  $\mathfrak{l}^{(k)}$ ,  $k = 1, 2, 3, 4$ , decomposes into two minimal left ideals  $\mathfrak{l}^{(k)} = \mathfrak{l}_{[\mu_k]}^{(k)} \oplus \mathfrak{l}_{[\nu_k]}^{(k)}$ , where the partitions  $\mu_k$  and  $\nu_k$  characterize the equivalence class of minimal left ideals to which such an ideal  $\mathfrak{l}_{[\mu_k]}^{(k)}$ ,  $\mathfrak{l}_{[\nu_k]}^{(k)}$  belongs.

Now the formula (4.2) yields that the group ring elements  $(y_t^*(S \otimes S))_b$ ,  $(y_t^*(A \otimes S))_b$ ,  $(y_t^*(S \otimes A))_b$  and  $(y_t^*(A \otimes A))_b$  are contained in one of the left ideals  $\mathfrak{l}^{(k)} \cdot y_t$ . Since the Young symmetrizer  $y_t$  of (2.12) generates a minimal left ideal of the equivalence class of  $\lambda = (2^2) \vdash 4$ , we obtain  $\mathfrak{l}_{[\mu]}^{(k)} \cdot y_t = \{0\}$  if  $\mu \neq (2^2)$ . This leads to  $\mathfrak{l}^{(2)} \cdot y_t = \{0\}$ ,  $\mathfrak{l}^{(3)} \cdot y_t = \{0\}$  and  $y_t^*(A \otimes S) = 0$ ,  $y_t^*(S \otimes A)$ . Furthermore, we obtain  $\mathfrak{l}_{[4]}^{(1)} \cdot y_t = \{0\}$ ,  $\mathfrak{l}_{[1^4]}^{(4)} \cdot y_t = \{0\}$ . At most the products  $\mathfrak{l}_{[2^2]}^{(1)} \cdot y_t$ ,  $\mathfrak{l}_{[2^2]}^{(4)} \cdot y_t$  are candidates for non-vanishing results. We have already seen in the proof of Theorem 3.1 that these cases really lead to non-vanishing expressions (1.6) and (1.7).

## 5. A NEW PROOF OF THE STRONGER RESULT OF P. B. GILKEY

Now we give a new proof of the following result of P. B. Gilkey (see [26, pp. 41–44]).

**Theorem 5.1.** *For every algebraic curvature tensor  $T \in \mathcal{T}_4V$  there exist finite sets  $\mathcal{S}$  and  $\mathcal{A}$  of symmetric and alternating tensors  $S, A \in \mathcal{T}_2V$ , respectively, such that*

$T$  has a representation

$$(5.1) \quad T = \sum_{S \in \mathcal{S}} \epsilon_S \gamma(S) \quad , \quad \epsilon_S \in \{1, -1\},$$

as well as a representation

$$(5.2) \quad T = \sum_{A \in \mathcal{A}} \epsilon_A \alpha(A) \quad , \quad \epsilon_A \in \{1, -1\}.$$

*Proof.* We consider the following group ring elements

$$(5.3) \quad \sigma_\epsilon := y_t^* \cdot f \cdot \xi_\epsilon \quad , \quad \epsilon \in \{1, -1\},$$

where  $y_t$  is the Young symmetrizer of the Young tableau (2.12) and

$$(5.4) \quad f := \text{id} + (13)(24)$$

$$(5.5) \quad \xi_\epsilon := (\text{id} + \epsilon(12)) \cdot (\text{id} + \epsilon(34)) \quad , \quad \epsilon \in \{1, -1\}.$$

A calculation (by means of PERMS [14]) shows that  $\sigma_1 \neq 0$  and  $\sigma_{-1} \neq 0$  (see [9]).

The right ideals  $\sigma_\epsilon \cdot \mathbb{K}[\mathcal{S}_4]$  are non-vanishing subideals of the right ideal  $\mathfrak{r} = y_t^* \cdot \mathbb{K}[\mathcal{S}_4]$  which defines the symmetry class  $\mathcal{T}_{\mathfrak{r}}$  of algebraic curvature tensors. Since  $\mathfrak{r}$  is a minimal right ideal, we obtain  $\mathfrak{r} = \sigma_1 \cdot \mathbb{K}[\mathcal{S}_4] = \sigma_{-1} \cdot \mathbb{K}[\mathcal{S}_4]$ , i.e.  $\sigma_1$  and  $\sigma_{-1}$  are generating elements of  $\mathfrak{r}$ , too.

A tensor  $T \in \mathcal{T}_4V$  is an algebraic curvature tensor iff there exist  $a \in \mathfrak{r}$  and  $T' \in \mathcal{T}_4V$  such that  $T = aT'$ . Since further every  $a \in \mathfrak{r}$  has a representation  $a = \sigma_\epsilon \cdot x_\epsilon$ ,  $\epsilon = 1, -1$ , a tensor  $T \in \mathcal{T}_4V$  is an algebraic curvature tensor iff there is a tensor  $T' \in \mathcal{T}_4V$  such that  $T = \sigma_1 T'$  or  $T = \sigma_{-1} T'$ .

Let us consider the case  $\epsilon = 1$ . We obtain all algebraic curvature tensors if we form  $T = \sigma_1 T'$ ,  $T' \in \mathcal{T}_4V$ . As in the proof of Theorem 3.1 we can use a representation

$$(5.6) \quad T' = \sum_{(M,N) \in \mathcal{P}} M \otimes N$$

for a  $T' \in \mathcal{T}_4V$ , where  $\mathcal{P}$  is a finite set of pairs  $(M, N)$  of tensors  $M, N \in \mathcal{T}_2V$ . From (5.6) we obtain

$$(5.7) \quad \xi_1 T' = \sum_{(S', S'') \in \tilde{\mathcal{P}}} S' \otimes S''$$

where  $\tilde{\mathcal{P}}$  is the finite set of pairs  $(S', S'')$  of the symmetrized tensors  $S' = M + M^t$ ,  $S'' = N + N^t$ ,  $(M, N) \in \mathcal{P}$ . Further the application of the symmetry operator  $f$

to  $\xi_1 T'$  yields

$$\begin{aligned}
f(\xi_1 T') &= \sum_{(S', S'') \in \hat{\mathcal{P}}} (S' \otimes S'' + S'' \otimes S') \\
&= \sum_{(S', S'') \in \hat{\mathcal{P}}} ((S' + S'') \otimes (S' + S'') - S' \otimes S' - S'' \otimes S'') \\
&= \sum_{S''' \in \mathcal{S}'''} z_{S'''} S''' \otimes S''' \quad , \quad z_{S'''} \in \mathbb{Z} \\
(5.8) \quad &= \sum_{S \in \mathcal{S}} \epsilon_S S \otimes S \quad , \quad \epsilon_S \in \{1, -1\}.
\end{aligned}$$

The last step includes a transformation  $S''' \mapsto S := \sqrt{|z_{S'''}|} S'''$ . The application of  $y_t^*$  finishes the proof of (5.1):

$$T = \sigma_1 T' = (y_t^* \cdot f \cdot \xi_1) T' = \sum_{S \in \mathcal{S}} \epsilon_S y_t^*(S \otimes S) = 12 \sum_{S \in \mathcal{S}} \epsilon_S \gamma(S).$$

The proof of (5.2) can be carried out in exactly the same way. We have only to replace (5.7) by

$$(5.9) \quad \xi_{-1} T' = \sum_{(A', A'') \in \hat{\mathcal{P}}} A' \otimes A''$$

where  $\hat{\mathcal{P}}$  is the finite set of pairs  $(A', A'')$  of the skew-symmetrized tensors  $A' = M - M^t$ ,  $A'' = N - N^t$ ,  $(M, N) \in \mathcal{P}$ .  $\square$

**Remark 5.2.** The new generating elements  $\sigma_1, \sigma_{-1}$  of the right ideal  $\mathfrak{r}$  do not automatically lead to new generating idempotents of  $\mathfrak{r}$ . A calculation [9] by means of PERMS [14] yields

$$\begin{aligned}
y_t^* \cdot y_t^* &= 12 y_t^* \\
y_t^* \cdot \sigma_1 &= 12 \sigma_1 \\
\sigma_1 \cdot y_t^* &= 0 \\
y_t^* \cdot \sigma_{-1} &= 12 \sigma_{-1} \\
\sigma_{-1} \cdot y_t^* &= 96 y_t^* \\
\sigma_1 \cdot \sigma_1 &= 0 \\
\sigma_{-1} \cdot \sigma_{-1} &= 96 \sigma_{-1} \\
\sigma_{-1} \cdot \sigma_1 &= 96 \sigma_1 \\
\sigma_1 \cdot \sigma_{-1} &= 0.
\end{aligned}$$

Thus  $\sigma_{-1}$  is an essentially idempotent element from which a generating idempotent of  $\mathfrak{r}$  can be determined. However,  $\sigma_1$  is a nilpotent generating element of  $\mathfrak{r}$ .

6. SOME APPLICATIONS OF  $\alpha$  AND  $\gamma$ 

We finish our paper with some remarks which arise from suggestions and hints of P. B. Gilkey.

In [26, pp. 191–193] P. B. Gilkey presents a family of algebraic curvature tensors which is a generalization of the family (1.11) in the case of a metric  $g$  with an arbitrary signature  $(p, q)$ . This new family can be used to show that there are *Jordan Osserman algebraic curvature tensors*  $T$  for which  $J_T$  can have an arbitrary complicated Jordan normal form. We give a short view of the construction of Gilkey.

We consider a metric  $g \in \mathcal{T}_2V$  of signature  $(p, q)$  on  $V$ .

**Definition 6.1.** An algebraic curvature tensor  $T \in \mathcal{T}_4V$  is called *spacelike Jordan Osserman* (respectively *timelike Jordan Osserman*) if the Jordan normal form of the Jacobi operator  $J_T(x)$  is constant on  $S^+(V) := \{x \in V \mid g(x, x) = +1\}$  (respectively  $S^-(V) := \{x \in V \mid g(x, x) = -1\}$ ).  $T$  is called *Jordan Osserman* if  $T$  is both spacelike Jordan Osserman and timelike Jordan Osserman.

Now we generalize the Clifford commutation relation (1.8)

**Definition 6.2.** Let  $V$  and  $W$  be vector spaces which possess metrics with signature  $(p, q)$  and  $(r, s)$ , respectively, where  $r \leq p$  and  $s \leq q$ . We say that  $V$  admits a *Clifford module structure*  $\text{Cliff}(W)$  if there exist skew-symmetric linear transformations  $C_i : V \rightarrow V$  for  $1 \leq i \leq r + s$  so that

- (1)  $C_i C_j + C_j C_i = 0$  for  $i \neq j$ .
- (2)  $C_i^2 = \text{Id}$  for  $r$  values of  $i$ .
- (3)  $C_i^2 = -\text{Id}$  for  $s$  values of  $i$ .

Note that a linear map  $C : V \rightarrow V$  is called *skew-symmetric* if

$$\forall x, y \in V : g(Cx, y) = -g(x, Cy).$$

Now the following Lemma holds

**Lemma 6.3.** *Let  $V$  and  $W$  possess metrics of signature  $(p, q)$  and  $(r, s)$ , respectively ( $r \leq p, s \leq q$ ). Assume that  $V$  admits a  $\text{Cliff}(W)$  module structure. Let  $J : W \rightarrow W$  be a self-adjoint linear map of  $W$ . Then there exists a Jordan Osserman algebraic curvature tensor  $T$  on  $V$  so that the Jacobi operator  $J_T(x)$  is conjugate to  $g(x, x)J \oplus 0$  for every  $x \in S^\pm(V)$ , where  $g$  is the metric of  $V$ .*

In Gilkey's proof of Lemma 6.3 the above Jordan Osserman algebraic curvature tensor  $T$  is given by the formula

$$(6.1) \quad T = \sum_i c_i \alpha(A_i) + \frac{1}{2} \sum_{i \neq j} c_{ij} \alpha(A_i + A_j)$$

where  $c_i$  and  $c_{ij} = c_{ji}$  are constants. The  $A_i$  are tensors of order 2 on  $V$  defined by  $A_i(x, y) := g(C_i x, y)$ . Obviously the  $A_i$  are skew-symmetric.

The expression  $J \oplus 0$  is defined as follows: Under the assumptions of Lemma 6.3 we can find an isometry between  $V$  and  $W \oplus \mathbb{R}^{(p-r, q-s)}$ . Then  $J \oplus 0$  is defined by

setting

$$(J \oplus 0)(w \oplus z) := Jw \oplus 0.$$

The sum (6.1) is a representation (5.2) of an algebraic curvature tensor which is formed by means of  $\alpha$  from skew-symmetric tensors  $A_i \in \mathcal{T}_2V$ .

Now the following Theorem by Gilkey tells us that we can construct Jordan Osserman algebraic curvature tensors with arbitrarily complicated Jordan normal form if we greatly increase the dimension by suitable powers of 2.

**Theorem 6.4.** *Let  $J : W \rightarrow W$  be an arbitrary linear map of a vector space  $W$  of dimension  $m$ . Then there exist  $l = l(m)$  and a Jordan Osserman algebraic curvature tensor  $T$  on  $V = \mathbb{R}^{(2^l, 2^l)}$  so that  $J_T(x)$  is conjugate to  $\pm J \oplus 0$  if  $x \in S^\pm(V)$ .*

If we choose a suitable  $J$  we can generate a Jacobi operator  $J_T(x)$  with an arbitrarily complicated Jordan normal form. In particular we can construct non-diagonalizable Jacobi operators. Thus the family (6.1) is different from the family (1.11) since the Jacobi operators of the tensors (1.11) are defined on a vector space  $V$  with a positive definite metric and can be diagonalized as self-adjoint operators on  $V$ .

In [26, p. 191] P. B. Gilkey presents also a second generalization of (1.11), (1.12) in the case of certain indefinite metrics. The algebraic curvature tensors of this generalization have a structure

$$(6.2) \quad T = \lambda_0 \gamma(g) + \sum_i \lambda_i \alpha(A_i) \quad , \quad \lambda_0, \lambda_i = \text{const.}$$

which is very similar to (1.12). The  $A_i \in \mathcal{T}_2V$  are alternating tensors whose corresponding skew-symmetric maps  $C_i : V \rightarrow V$  satisfy the Clifford commutation relations (1.8). The Jacobi operators of these tensors (6.2) are diagonalizable.

In a further hint P. B. Gilkey pointed out that there is a simple possibility to generate examples of Osserman algebraic curvature tensors of the form  $\gamma(S)$  or  $\alpha(A)$  ( $S, A \in \mathcal{T}_2V$  symmetric or alternating) if we have a pseudo-Riemannian metric  $g$  of indefinite signature  $(p, q)$ . In particular, we can obtain in this way Osserman algebraic curvature tensors not of the form given in (1.11) or (6.2) (see Remark 6.7).

**Proposition 6.5.** *Assume that  $V$  has a metric  $g$  with signature  $(p, q)$ .*

(1) *If  $p, q \geq 1$  and  $C : V \rightarrow V$  is a symmetric map of  $V$  with  $C^2 = 0$ , then*

$$(6.3) \quad T := \gamma(S) \quad , \quad S(x, y) := g(Cx, y) \quad \forall x, y \in V$$

*is an Osserman algebraic curvature tensor.*

(2) *If  $p, q \geq 2$  and  $C : V \rightarrow V$  is a skew-symmetric map of  $V$  with  $C^2 = 0$ , then*

$$(6.4) \quad T := \alpha(A) \quad , \quad A(x, y) := g(Cx, y) \quad \forall x, y \in V$$

*is an Osserman algebraic curvature tensor.*

- (3) *The Jacobi operators  $J_T(x)$  of (6.3) and (6.4) are nilpotent of order 2, i.e.  $J_T(x)^2 = 0$  for all  $x \in V$ .*

Note that the tensors  $S, A \in \mathcal{T}_2V$  defined by (6.3), (6.4), respectively, are symmetric or skew-symmetric, respectively.

To prove Proposition 6.5 we show first the existence of the above nilpotent symmetric or skew-symmetric maps  $C$ . Furthermore we show that such maps  $C$  do not exist in the positive definite case.

**Lemma 6.6.** (1) *If  $M$  is a symmetric or skew-symmetric  $m \times m$ -matrix with  $M^2 = 0$ , then  $M = 0$ .*

- (2) *Let  $p, q \geq 2$ ,  $m = p + q$ , and  $F$  be the  $m \times m$ -diagonal matrix*

$$(6.5) \quad F := \text{diag}(\underbrace{1, \dots, 1}_p, \underbrace{-1, \dots, -1}_q)$$

*Then we can find non-vanishing symmetric or skew-symmetric  $m \times m$ -matrices  $S, A$ , respectively, such that*

$$(6.6) \quad (S \cdot F)^2 = 0 \quad , \quad (A \cdot F)^2 = 0.$$

*The assertion about a symmetric matrix  $S$  holds also for  $p, q \geq 1$ .*

*Proof. Ad (1):* If a matrix  $M = (m_{ij})$  is symmetric or skew-symmetric, we have  $M^T = \epsilon M$ ,  $\epsilon \in \{1, -1\}$ . Then  $M^2 = 0$  leads to  $\epsilon M \cdot M^T = 0$  and

$$(6.7) \quad 0 = \text{trace } M \cdot M^T = \sum_{i,j} m_{ij}^2.$$

From (6.7) we obtain  $m_{ij} = 0$  and  $M = 0$ .

**Ad (2):** The existence of skew-symmetric matrices  $A$  with  $(A \cdot F)^2 = 0$  is guaranteed by an example of P. B. Gilkey [26, p. 186] which was constructed for  $p, q \geq 2$ .

Let us now consider symmetric matrices under  $p, q \geq 1$ . If we write  $F$  as block matrix

$$F = \begin{pmatrix} \ddots & & & & & \\ & 1 & 0 & & & \\ & 0 & -1 & & & \\ & & & \ddots & & \\ & & & & \ddots & \\ & & & & & \ddots \end{pmatrix}$$

and choose a corresponding symmetric block matrix

$$S = \begin{pmatrix} 0 & & & & & \\ & 1 & 1 & & & \\ & 1 & 1 & & & \\ & & & & & \\ & & & & & \\ & & & & & 0 \end{pmatrix}$$

then we obtain

$$S \cdot F = \begin{pmatrix} 0 & & & \\ & 1 & -1 & \\ & 1 & -1 & \\ & & & 0 \end{pmatrix}.$$

Obviously, the matrix  $S \cdot F$  satisfies  $(S \cdot F)^2 = 0$ .  $\square$

A relation  $B(x, y) = g(Cx, y)$  between a linear map  $C : V \rightarrow V$  and a tensor  $B \in \mathcal{T}_2V$  leads to relations

$$(6.8) \quad B_{ij} = C_i^k g_{kj} \quad \text{or} \quad C_i^j = B_{ik} g^{kj}$$

for the coordinates of  $B$ ,  $C$ , and  $g$ . If we use an orthonormal basis of  $V$  then we have  $(g^{ij}) = \text{diag}(1, \dots, 1, -1, \dots, -1)$  and the matrix  $(B_{ik} g^{kj})$  is a matrix product of the type  $B \cdot F$  considered in Lemma 6.6. Thus statement (2) of Lemma 6.6 guarantees the existence of symmetric or skew-symmetric maps  $C$  with  $C^2 = 0$  in the above pseudo-Riemannian settings. Furthermore we see from statement (1) of Lemma 6.6 that such maps do not exist if  $g$  is positive definite.

Now we can prove Proposition 6.5.

*Proof.* For a symmetric map  $C : V \rightarrow V$  the definitions (1.5), (1.6) and  $S(x, w) = g(Cx, w)$  lead to

$$g(J_{\gamma(S)}(x)y, w) = \frac{1}{3}(S(y, w)S(x, x) - S(x, y)S(x, w))$$

and

$$(6.9) \quad J_{\gamma(S)}(x)y = \frac{1}{3}(g(Cx, x)Cy - g(Cy, x)Cx).$$

If now  $C^2 = 0$ , then we obtain  $CJ_{\gamma(S)}(x)y = \frac{1}{3}(g(Cx, x)C^2y - g(Cy, x)C^2x) = 0$  and  $J_{\gamma(S)}(x)^2y = \frac{1}{3}(g(Cx, x)CJ_{\gamma(S)}(x)y - g(CJ_{\gamma(S)}(x)y, x)Cx) = 0$ , i.e.  $J_{\gamma(S)}(x)$  is nilpotent of order 2 for all  $x \in V$ . But then all eigenvalues of  $J_{\gamma(S)}(x)$  are equal to zero, i.e.  $\gamma(S)$  is Osserman.

For a skew-symmetric map  $C : V \rightarrow V$  we obtain from (1.5), (1.7),  $A(x, w) = g(Cx, w)$  and  $A(x, x) = 0$

$$\begin{aligned} g(J_{\alpha(A)}(x)y, w) &= \frac{1}{3}(2A(y, x)A(x, w) + A(y, x)A(x, w) - A(y, w)A(x, x)) \\ &= A(y, x)A(x, w) \end{aligned}$$

and

$$J_{\alpha(A)}(x)y = g(Cy, x)Cx.$$

Again, the condition  $C^2 = 0$  leads to  $CJ_{\alpha(A)}(x)y = 0$  and  $J_{\alpha(A)}(x)^2y = 0$  for all  $x \in V$ . Consequently all eigenvalues of  $J_{\alpha(A)}(x)$  vanish for all  $x \in V$  and  $\alpha(A)$  is Osserman.  $\square$

**Remark 6.7.** If we use a representation of (1.11) or (6.2) in which we arrange the skew-symmetric maps  $C_i$  as in (1.9) then we will see that the Jacobi operator of (1.11) or (6.2) has the eigenvalues

$$0 \quad , \quad \lambda_0 \quad , \quad \lambda_0 - 3\lambda_i .$$

on the (pseudo)-unit sphere (see [24] and [26, p. 191]). The vanishing of all these eigenvalues leads to  $T = 0$  for (1.11) and (6.2). Proposition 6.5, however, yields examples of non-vanishing Osserman algebraic curvature tensors  $\gamma(S)$ ,  $\alpha(A)$  whose Jacobi operators have only the eigenvalues zero. Consequently the examples of Proposition 6.5 can not be transformed into a representation (1.11) or (6.2).

Moreover, the Jacobi operators of the algebraic curvature tensors (1.11) and (6.2) are diagonalizable (see [24] and [26, p. 191]). Since the Jacobi operators  $J_{\gamma(S)}$ ,  $J_{\alpha(A)}$  from Proposition 6.5 have only the eigenvalue zero, a Jacobi operator  $J_{\gamma(S)}$ ,  $J_{\alpha(A)}$  according to Proposition 6.5 is equal to zero if it is diagonalizable. Thus every algebraic curvature tensor  $\gamma(S)$ ,  $\alpha(A)$  from Proposition 6.5, which has a non-vanishing Jacobi operator, is different from every algebraic curvature tensor with a diagonalizable Jacobi operator.

**Remark 6.8.** Osserman tensors (6.3) for metrics  $g$  with Lorentzian signature  $(1, q)$  are a special case. It is known that an Osserman algebraic curvature tensor has constant sectional curvature if the metric  $g$  has Lorentzian signature  $(1, q)$ ,  $q \geq 1$  (see [1, 20]). The Jacobi operator of an algebraic curvature tensor with constant sectional curvature is diagonalizable. Thus we obtain from Remark 6.7 that every algebraic curvature tensor  $\gamma(S)$  from Proposition 6.5 has a Jacobi operator  $J_{\gamma(S)} = 0$  if  $p = 1$  or  $q = 1$ . We verify this fact.

**Lemma 6.9.** *Let  $F$  be the  $m \times m$ -diagonal matrix*

$$F := \text{diag}(1, -1, \dots, -1).$$

- (1) *If  $S \neq 0$  is a symmetric  $m \times m$ -matrix with  $(S \cdot F)^2 = 0$  then there exists an invertible matrix  $D$  and a  $\lambda \neq 0$  such that*

$$(6.10) \quad D \cdot S \cdot D^{-1} = \begin{pmatrix} \lambda & \pm\lambda & & \\ \pm\lambda & \lambda & & \\ & & & \\ & & & 0 \end{pmatrix} \quad \text{and} \quad D \cdot F \cdot D^{-1} = F.$$

- (2) *A skew-symmetric  $m \times m$ -matrix  $A$  with  $(A \cdot F)^2 = 0$  vanishes.*

*Proof. Ad (1):* We write a symmetric  $m \times m$ -matrix  $S$  as block matrix

$$(6.11) \quad S = \begin{pmatrix} a & b \\ b^T & \tilde{S} \end{pmatrix},$$

where  $\tilde{S}$  is a symmetric  $(m-1) \times (m-1)$ -matrix. Then we can transform  $\tilde{S}$  into a diagonal matrix by a conjugation of  $S$  by a matrix

$$D = \begin{pmatrix} 1 & 0 \\ 0^T & \tilde{D} \end{pmatrix}$$



where  $\tilde{D}$  is a suitable orthogonal  $(m-1) \times (m-1)$ -matrix. Obviously it holds  $D \cdot F \cdot D^{-1} = F$  for such a  $D$ .

From (6.11) we obtain

$$S \cdot F = \begin{pmatrix} a & -b \\ b^T & -\tilde{S} \end{pmatrix}, \quad (S \cdot F)^2 = \begin{pmatrix} a^2 - b \cdot b^T & -ab + b \cdot \tilde{S} \\ ab^T - \tilde{S} \cdot b^T & -b^T \cdot b + \tilde{S}^2 \end{pmatrix}.$$

Assume that  $\tilde{S} = 0$ . Then the condition  $(S \cdot F)^2 = 0$  leads to the relations  $a^2 - b \cdot b^T = 0$  and  $ab = 0$  from which  $S = 0$  follows. However this is a contradiction to  $S \neq 0$ .

In case of  $\tilde{S} \neq 0$  the matrix  $\tilde{S}$  has rank 1 because of  $b^T \cdot b - \tilde{S}^2 = 0$ . Then  $\tilde{S}$  has one and only one non-vanishing eigenvalue  $\lambda$ . Furthermore we obtain  $b \neq 0$  from  $b^T \cdot b - \tilde{S}^2 = 0$ .

Now the condition  $ab^T - \tilde{S} \cdot b^T = 0$  means that  $b^T$  is an eigenvector of  $\tilde{S}$  which belongs to the eigenvalue  $a$ . The value  $a = 0$  is impossible since then  $a^2 - b \cdot b^T = 0$  yields  $b = 0$ . Thus we obtain  $a = \lambda \neq 0$ .

If we place  $\lambda \neq 0$  in the left upper corner of  $\tilde{S}$ , then  $b$  is an  $(m-1)$ -tuple  $b = (\mu, 0, \dots, 0)$  and  $a^2 - b \cdot b^T = 0$  yields  $\mu = \pm\lambda$ .

**Ad (2):** We write a skew-symmetric  $m \times m$ -matrix  $A$  as block matrix

$$(6.12) \quad A = \begin{pmatrix} 0 & b \\ -b^T & \tilde{A} \end{pmatrix},$$

where  $\tilde{A}$  is a skew-symmetric  $(m-1) \times (m-1)$ -matrix. From (6.12) we obtain

$$A \cdot F = \begin{pmatrix} 0 & -b \\ -b^T & -\tilde{A} \end{pmatrix}, \quad (A \cdot F)^2 = \begin{pmatrix} b \cdot b^T & b \cdot \tilde{A} \\ \tilde{A} \cdot b^T & b^T \cdot b + \tilde{A}^2 \end{pmatrix}.$$

Now the condition  $(A \cdot F)^2 = 0$  yields  $b \cdot b^T = 0$ , i.e.  $b = 0$ . But then  $\tilde{A}$  is a skew-symmetric matrix which has to fulfil  $\tilde{A}^2 = 0$ . From Lemma 6.6 we obtain  $\tilde{A} = 0$ .  $\square$

Now let  $g$  be a Lorentzian metric with signature  $(1, q)$  and  $T = \gamma(S)$  be an Osserman algebraic curvature tensor according to Proposition 6.5. Then  $S$  can be transformed into a form (6.10) and the corresponding symmetric map

$$C := S \cdot F = \begin{pmatrix} \lambda & \mp\lambda & \\ \pm\lambda & -\lambda & \\ & & 0 \end{pmatrix}$$

has rank 1. Let us consider the Jacobi operator (6.9). If  $Cx = 0$  then  $J_{\gamma(S)}(x) = 0$ . If  $Cx \neq 0$  then the vectors  $Cx$  and  $Cy$  are proportional since  $C$  has a 1-dimensional range. Again we obtain  $J_{\gamma(S)}(x) = 0$ . Thus  $J_{\gamma(S)}(x) \equiv 0$  for every  $T = \gamma(S)$  according to Proposition 6.5 in the Lorentzian setting. Statement (2) of Lemma 6.9 tells us that a non-vanishing Osserman tensor  $T = \alpha(A)$  of the type (6.4) does not exist in the case of a Lorentzian signature.

**Acknowledgements.** I would like to thank Prof. P. B. Gilkey for important and helpful discussions and for valuable suggestions for future investigations.

## REFERENCES

- [1] N. Blažić, N. Bokan and P. B. Gilkey. A note on Osserman Lorentzian manifolds. *Bull. London Math. Soc.*, 29:227–230, 1997.
- [2] N. Blažić, N. Bokan, P. B. Gilkey and Z. Rakić. Pseudo-Riemannian Osserman manifolds. *Balkan J. Geom. Appl.*, 2(2):1–12, 1997.
- [3] N. Blažić, N. Bokan and Z. Rakić. Nondiagonalizable timelike (spacelike) Osserman (2,2) manifolds. *Saitama Math. J.*, 16:15–22, 1998.
- [4] H. Boerner. *Darstellungen von Gruppen*, volume 74 of *Die Grundlehren der mathematischen Wissenschaften in Einzeldarstellungen*. Springer-Verlag, Berlin, Göttingen, Heidelberg, 1955.
- [5] H. Boerner. *Representations of Groups*. North-Holland Publishing Company, Amsterdam, 2. revised edition, 1970.
- [6] A. Bonome, P. Castro, E. García-Río, L. Hervella and R. Vázquez-Lorenzo. Nonsymmetric Osserman indefinite Kähler manifolds. *Proc. Amer. Math. Soc.*, 126:2763–2769, 1998.
- [7] Q.-S. Chi. A curvature characterization of certain locally rank-one symmetric spaces. *J. Differential Geom.*, 28:187–202, 1988.
- [8] S. M. Christensen and L. Parker. *MathTensor: A System for Doing Tensor Analysis by Computer*. Addison-Wesley, Reading, Mass., Menlo Park, Ca., New York, Don Mills, Ont., Wokingham, UK, Amsterdam, Bonn, Sydney, Singapore, Tokyo, Madrid, San Juan, Milan, Paris, 1994.
- [9] B. Fiedler. Examples of calculations by means of PERMS. Mathematica notebooks. Internet <http://home.t-online.de/home/Bernd.Fiedler.RoschStr.Leipzig/pnbks.htm>.
- [10] B. Fiedler. A use of ideal decomposition in the computer algebra of tensor expressions. *Z. Anal. Anw.*, 16(1):145–164, 1997.
- [11] B. Fiedler. An algorithm for the decomposition of ideals of the group ring of a symmetric group. In Adalbert Kerber, editor, *Actes 39<sup>e</sup> Séminaire Lotharingien de Combinatoire, Thurnau, 1997*, Publ. I.R.M.A. Strasbourg. Institut de Recherche Mathématique Avancée, Université Louis Pasteur et C.N.R.S. (URA 01), 1998. Electronically published: <http://www.mat.univie.ac.at/~slc>. B39e, 26 pp.
- [12] B. Fiedler. A characterization of the dependence of the Riemannian metric on the curvature tensor by Young symmetrizers. *Z. Anal. Anw.*, 17(1):135–157, 1998.
- [13] B. Fiedler. *An Algorithm for the Decomposition of Ideals of Semi-Simple Rings and its Application to Symbolic Tensor Calculations by Computer*. Habilitationsschrift, Universität Leipzig, Leipzig, Germany, November 1999. Fakultät für Mathematik und Informatik.
- [14] B. Fiedler. *PERMS 2.1 (15.1.1999)*. Mathematisches Institut, Universität Leipzig, Leipzig, 1999. Will be sent in to MathSource, Wolfram Research Inc.
- [15] B. Fiedler. Characterization of tensor symmetries by group ring subspaces and computation of normal forms of tensor coordinates. In A. Betten, A. Kohnert, R. Laue, and A. Wassermann, editors, *Algebraic Combinatorics and Applications. Proceedings of the Euroconference, ALCOMA, Gößweinstein, Germany, September 12–19, 1999*, pages 118–133, Berlin, 2001. Springer-Verlag.
- [16] B. Fiedler. Ideal decompositions and computation of tensor normal forms. In *Séminaire Lotharingien de Combinatoire*, 2001. El. published: <http://www.mat.univie.ac.at/~slc>. B45g, 16 pp. ArXiv: <http://arXiv.org/abs/math.CO/0211156>.
- [17] S. A. Fulling, R. C. King, B. G. Wybourne and C. J. Cummins. Normal forms for tensor polynomials: I. The Riemann tensor. *Class. Quantum Grav.*, 9:1151–1197, 1992.
- [18] W. Fulton and J. Harris. *Representation Theory: A First Course*, volume 129 of *Graduate Texts in Mathematics, Readings in Mathematics*. Springer-Verlag, New York, Berlin, Heidelberg, London, Paris, Tokyo, Hong Kong, Barcelona, Budapest, 1991.
- [19] W. Fulton. *Young Tableaux*. Number 35 in London Mathematical Society Student Texts. Cambridge University Press, Cambridge, New York, Melbourne, 1997.

- [20] E. García-Riό, D. N. Kupeli and M. E. Vázquez-Abal. On a problem of Osserman in Lorentzian geometry. *Differential Geom. Appl.*, 7:85–100, 1997.
- [21] E. García-Riό, D. N. Kupeli, M. E. Vázquez-Abal and R. Vázquez-Lorenzo. Affine Osserman connections and their Riemann extensions. *Differential Geom. Appl.*, 11:145–153, 1999.
- [22] E. García-Riό, M. E. Vázquez-Abal and R. Vázquez-Lorenzo. Nonsymmetric Osserman pseudo-Riemannian manifolds. *Proc. Amer. Math. Soc.*, 126:2771–2778, 1998.
- [23] E. García-Riό, D. N. Kupeli and Ramon Vázquez-Lorenzo. *Osserman Manifolds in Semi-Riemannian Geometry*, volume 1777 of *Lecture notes in Mathematics*. Springer-Verlag, Berlin, 2002.
- [24] P. B. Gilkey. Geometric properties of the curvature operator. In W. H. Chen, A.-M. Li, U. Simon, L. Verstraelen, C. P. Wang and M. Wiehe, editors, *Proceedings of Beijing Conference "Geometry and Topology of Submanifolds"*, volume 10, Singapore. World Scientific. March 2000. To appear.
- [25] P. B. Gilkey. Relating algebraic properties of the curvature tensor to geometry. *Novi Sad J. Math.*, 29:109–119, 1999.
- [26] P. B. Gilkey. *Geometric Properties of Natural Operators Defined by the Riemann Curvature Tensor*. World Scientific Publishing Co., Singapore, New Jersey, London, Hong Kong, 2001.
- [27] P. B. Gilkey, J. V. Leahy and H. Sadofsky. Riemannian manifolds whose skew-symmetric curvature tensor has constant eigenvalues. *Indiana Univ. Math. J.*, 48(2):615–634, 1999. Preprint: Research paper Nr. 125 on <http://darkwing.uoregon.edu/~gilkey/>.
- [28] G. D. James and A. Kerber. *The Representation Theory of the Symmetric Group*, volume 16 of *Encyclopedia of Mathematics and its Applications*. Addison-Wesley Publishing Company, Reading, Mass., London, Amsterdam, Don Mills, Ont., Sidney, Tokyo, 1981.
- [29] A. Kerber. *Representations of Permutation Groups*, volume 240, 495 of *Lecture Notes in Mathematics*. Springer-Verlag, Berlin, Heidelberg, New York, 1971, 1975.
- [30] A. Kerber. *Algebraic combinatorics via finite group actions*. BI-Wiss.-Verl., Mannheim, Wien, Zürich, 1991.
- [31] J. M. Lee, D. Lear, J. Roth, J. Coskey and L. Nave. *Ricci. A Mathematica package for doing tensor calculations in differential geometry. User's Manual. Version 1.32*. Department of Mathematics, Box 354350, University of Washington, Seattle, WA 98195-4350, 1992 – 1998. Ricci's home page: <http://www.math.washington.edu/~lee/Ricci/>.
- [32] D. E. Littlewood. *The Theory of Group Characters and Matrix Representations of Groups*. Clarendon Press, Oxford, 2. edition, 1950.
- [33] I. G. Macdonald. *Symmetric Functions and Hall Polynomials*. Clarendon Press, Oxford, 1979.
- [34] W. Müller. *Darstellungstheorie von endlichen Gruppen*. Teubner Studienbücher Mathematik. B. G. Teubner, Stuttgart, 1980.
- [35] M. A. Naimark and A. I. Štern. *Theory of Group Representations*, volume 246 of *Grundlehren der Mathematischen Wissenschaften*. Springer-Verlag, Berlin, Heidelberg, New York, 1982.
- [36] Y. Nikolayevsky. Osserman conjecture in dimension  $n \neq 8, 16$ . Preprint, 2002. ArXiv: <http://arXiv.org/abs/math.DG/0204258>.
- [37] R. Osserman. Curvature in the eighties. *Amer. Math. Monthly*, 97:731–756, 1990.
- [38] Z. Rakić. An example of rank two symmetric Osserman space. *Bull. Austral. Math. Soc.*, 56:517–521, 1997.
- [39] F. Sānger. Plethysmen von irreduziblen Darstellungen symmetrischer Gruppen. Dissertation, Rheinisch-Westfälische Technische Hochschule, Mathematisch-Naturwissenschaftliche Fakultät, Aachen, 1980.
- [40] B. L. van der Waerden. *Algebra*, volume I, II. Springer-Verlag, Berlin, Heidelberg, New York, London, Paris, Tokyo, Hong Kong, Barcelona, Budapest, 9., 6. edition, 1993.
- [41] H. Weyl. *The Classical Groups, their Invariants and Representations*. Princeton University Press, Princeton, New Jersey, 1939.

BERND FIEDLER, MATHEMATISCHES INSTITUT, UNIVERSITÄT LEIPZIG, AUGUSTUSPLATZ  
10/11, D-04109 LEIPZIG, GERMANY

*URL:* <http://home.t-online.de/home/Bernd.Fiedler.RoschStr.Leipzig/>

*E-mail address:* Bernd.Fiedler.RoschStr.Leipzig@t-online.de