

Determination of $|V_{ub}|$ from Measurements of the Electron and Neutrino Momenta in Inclusive Semileptonic B Decays

B. Aubert,¹ R. Barate,¹ D. Boutigny,¹ F. Couderc,¹ Y. Karyotakis,¹ J. P. Lees,¹ V. Poireau,¹ V. Tisserand,¹ A. Zghiche,¹ E. Grauges,² A. Palano,³ M. Pappagallo,³ A. Pompili,³ J. C. Chen,⁴ N. D. Qi,⁴ G. Rong,⁴ P. Wang,⁴ Y. S. Zhu,⁴ G. Eigen,⁵ I. Ofte,⁵ B. Stugu,⁵ G. S. Abrams,⁶ M. Battaglia,⁶ A. B. Breon,⁶ D. N. Brown,⁶ J. Button-Shafer,⁶ R. N. Cahn,⁶ E. Charles,⁶ C. T. Day,⁶ M. S. Gill,⁶ A. V. Gritsan,⁶ Y. Groysman,⁶ R. G. Jacobsen,⁶ R. W. Kadel,⁶ J. Kadyk,⁶ L. T. Kerth,⁶ Yu. G. Kolomensky,⁶ G. Kukartsev,⁶ G. Lynch,⁶ L. M. Mir,⁶ P. J. Oddone,⁶ T. J. Orimoto,⁶ M. Pripstein,⁶ N. A. Roe,⁶ M. T. Ronan,⁶ W. A. Wenzel,⁶ M. Barrett,⁷ K. E. Ford,⁷ T. J. Harrison,⁷ A. J. Hart,⁷ C. M. Hawkes,⁷ S. E. Morgan,⁷ A. T. Watson,⁷ M. Fritsch,⁸ K. Goetzen,⁸ T. Held,⁸ H. Koch,⁸ B. Lewandowski,⁸ M. Pelizaeus,⁸ K. Peters,⁸ T. Schroeder,⁸ M. Steinke,⁸ J. T. Boyd,⁹ J. P. Burke,⁹ N. Chevalier,⁹ W. N. Cottingham,⁹ M. P. Kelly,⁹ T. Cuhadar-Donszelmann,¹⁰ B. G. Fulsom,¹⁰ C. Hearty,¹⁰ N. S. Knecht,¹⁰ T. S. Mattison,¹⁰ J. A. McKenna,¹⁰ A. Khan,¹¹ P. Kyberd,¹¹ M. Saleem,¹¹ L. Teodorescu,¹¹ A. E. Blinov,¹² V. E. Blinov,¹² A. D. Bukin,¹² V. P. Druzhinin,¹² V. B. Golubev,¹² E. A. Kravchenko,¹² A. P. Onuchin,¹² S. I. Serednyakov,¹² Yu. I. Skovpen,¹² E. P. Solodov,¹² A. N. Yushkov,¹² D. Best,¹³ M. Bondioli,¹³ M. Bruinsma,¹³ M. Chao,¹³ I. Eschrich,¹³ D. Kirkby,¹³ A. J. Lankford,¹³ M. Mandelkern,¹³ R. K. Mommsen,¹³ W. Roethel,¹³ D. P. Stoker,¹³ C. Buchanan,¹⁴ B. L. Hartfiel,¹⁴ A. J. R. Weinstein,¹⁴ S. D. Foulkes,¹⁵ J. W. Gary,¹⁵ O. Long,¹⁵ B. C. Shen,¹⁵ K. Wang,¹⁵ L. Zhang,¹⁵ D. del Re,¹⁶ H. K. Hadavand,¹⁶ E. J. Hill,¹⁶ D. B. MacFarlane,¹⁶ H. P. Paar,¹⁶ S. Rahatlou,¹⁶ V. Sharma,¹⁶ J. W. Berryhill,¹⁷ C. Campagnari,¹⁷ A. Cunha,¹⁷ B. Dahmes,¹⁷ T. M. Hong,¹⁷ M. A. Mazur,¹⁷ J. D. Richman,¹⁷ W. Verkerke,¹⁷ T. W. Beck,¹⁸ A. M. Eisner,¹⁸ C. J. Flacco,¹⁸ C. A. Heusch,¹⁸ J. Kroseberg,¹⁸ W. S. Lockman,¹⁸ G. Nesom,¹⁸ T. Schalk,¹⁸ B. A. Schumm,¹⁸ A. Seiden,¹⁸ P. Spradlin,¹⁸ D. C. Williams,¹⁸ M. G. Wilson,¹⁸ J. Albert,¹⁹ E. Chen,¹⁹ G. P. Dubois-Felsmann,¹⁹ A. Dvoretzki,¹⁹ D. G. Hitlin,¹⁹ I. Narsky,¹⁹ T. Piatenko,¹⁹ F. C. Porter,¹⁹ A. Ryd,¹⁹ A. Samuel,¹⁹ R. Andreassen,²⁰ S. Jayatilleke,²⁰ G. Mancinelli,²⁰ B. T. Meadows,²⁰ M. D. Sokoloff,²⁰ F. Blanc,²¹ P. Bloom,²¹ S. Chen,²¹ W. T. Ford,²¹ U. Nauenberg,²¹ A. Olivas,²¹ P. Rankin,²¹ W. O. Ruddick,²¹ J. G. Smith,²¹ K. A. Ulmer,²¹ S. R. Wagner,²¹ J. Zhang,²¹ A. Chen,²² E. A. Eckhart,²² A. Soffer,²² W. H. Toki,²² R. J. Wilson,²² Q. Zeng,²² D. Altenburg,²³ E. Feltresi,²³ A. Hauke,²³ B. Spaan,²³ T. Brandt,²⁴ J. Brose,²⁴ M. Dickopp,²⁴ V. Klose,²⁴ H. M. Lacker,²⁴ R. Nogowski,²⁴ S. Otto,²⁴ A. Petzold,²⁴ G. Schott,²⁴ J. Schubert,²⁴ K. R. Schubert,²⁴ R. Schwierz,²⁴ J. E. Sundermann,²⁴ D. Bernard,²⁵ G. R. Bonneaud,²⁵ P. Grenier,²⁵ S. Schrenk,²⁵ Ch. Thiebaux,²⁵ G. Vasileiadis,²⁵ M. Verderi,²⁵ D. J. Bard,²⁶ P. J. Clark,²⁶ W. Gradl,²⁶ F. Muheim,²⁶ S. Playfer,²⁶ Y. Xie,²⁶ M. Andreotti,²⁷ V. Azzolini,²⁷ D. Bettoni,²⁷ C. Bozzi,²⁷ R. Calabrese,²⁷ G. Cibinetto,²⁷ E. Luppi,²⁷ M. Negrini,²⁷ L. Piemontese,²⁷ F. Anulli,²⁸ R. Baldini-Ferrolli,²⁸ A. Calcaterra,²⁸ R. de Sangro,²⁸ G. Finocchiaro,²⁸ P. Patteri,²⁸ I. M. Peruzzi,²⁸ * M. Piccolo,²⁸ A. Zallo,²⁸ A. Buzzo,²⁹ R. Capra,²⁹ R. Contri,²⁹ M. Lo Vetere,²⁹ M. Macri,²⁹ M. R. Monge,²⁹ S. Passaggio,²⁹ C. Patrignani,²⁹ E. Robutti,²⁹ A. Santroni,²⁹ S. Tosi,²⁹ S. Bailey,³⁰ G. Brandenburg,³⁰ K. S. Chaisanguanthum,³⁰ M. Morii,³⁰ E. Won,³⁰ J. Wu,³⁰ R. S. Dubitzky,³¹ U. Langenegger,³¹ J. Marks,³¹ S. Schenk,³¹ U. Uwer,³¹ W. Bhimji,³² D. A. Bowerman,³² P. D. Dauncey,³² U. Egede,³² R. L. Flack,³² J. R. Gaillard,³² G. W. Morton,³² J. A. Nash,³² M. B. Nikolich,³² G. P. Taylor,³² W. P. Vazquez,³² M. J. Charles,³³ W. F. Mader,³³ U. Mallik,³³ A. K. Mohapatra,³³ J. Cochran,³⁴ H. B. Crawley,³⁴ V. Eyges,³⁴ W. T. Meyer,³⁴ S. Prell,³⁴ E. I. Rosenberg,³⁴ A. E. Rubin,³⁴ J. Yi,³⁴ N. Arnaud,³⁵ M. Davier,³⁵ X. Giroux,³⁵ G. Grosdidier,³⁵ A. Höcker,³⁵ F. Le Diberder,³⁵ V. Lepeltier,³⁵ A. M. Lutz,³⁵ A. Oyanguren,³⁵ T. C. Petersen,³⁵ M. Pierini,³⁵ S. Plaszczynski,³⁵ S. Rodier,³⁵ P. Roudeau,³⁵ M. H. Schune,³⁵ A. Stocchi,³⁵ G. Wormser,³⁵ C. H. Cheng,³⁶ D. J. Lange,³⁶ M. C. Simani,³⁶ D. M. Wright,³⁶ A. J. Bevan,³⁷ C. A. Chavez,³⁷ J. P. Coleman,³⁷ I. J. Forster,³⁷ J. R. Fry,³⁷ E. Gabathuler,³⁷ R. Gamet,³⁷ K. A. George,³⁷ D. E. Hutchcroft,³⁷ R. J. Parry,³⁷ D. J. Payne,³⁷ K. C. Schofield,³⁷ C. Touramanis,³⁷ C. M. Cormack,³⁸ F. Di Lodovico,³⁸ R. Sacco,³⁸ C. L. Brown,³⁹ G. Cowan,³⁹ H. U. Flaecher,³⁹ M. G. Green,³⁹ D. A. Hopkins,³⁹ P. S. Jackson,³⁹ T. R. McMahon,³⁹ S. Ricciardi,³⁹ F. Salvatore,³⁹ D. Brown,⁴⁰ C. L. Davis,⁴⁰ J. Allison,⁴¹ N. R. Barlow,⁴¹ R. J. Barlow,⁴¹ M. C. Hodgkinson,⁴¹ G. D. Lafferty,⁴¹ M. T. Naisbit,⁴¹ J. C. Williams,⁴¹ C. Chen,⁴² A. Farbin,⁴² W. D. Hulsbergen,⁴² A. Jawahery,⁴² D. Kovalskyi,⁴² C. K. Lae,⁴² V. Lillard,⁴² D. A. Roberts,⁴² G. Simi,⁴² G. Blaylock,⁴³ C. Dallapiccola,⁴³ S. S. Hertzbach,⁴³

R. Kofler,⁴³ V. B. Koptchev,⁴³ X. Li,⁴³ T. B. Moore,⁴³ S. Saremi,⁴³ H. Staengle,⁴³ S. Willocq,⁴³ R. Cowan,⁴⁴ K. Koeneke,⁴⁴ G. Sciolla,⁴⁴ S. J. Sekula,⁴⁴ M. Spitznagel,⁴⁴ F. Taylor,⁴⁴ R. K. Yamamoto,⁴⁴ H. Kim,⁴⁵ P. M. Patel,⁴⁵ S. H. Robertson,⁴⁵ A. Lazzaro,⁴⁶ V. Lombardo,⁴⁶ F. Palombo,⁴⁶ J. M. Bauer,⁴⁷ L. Cremaldi,⁴⁷ V. Eschenburg,⁴⁷ R. Godang,⁴⁷ R. Kroeger,⁴⁷ J. Reidy,⁴⁷ D. A. Sanders,⁴⁷ D. J. Summers,⁴⁷ H. W. Zhao,⁴⁷ S. Brunet,⁴⁸ D. Côté,⁴⁸ P. Taras,⁴⁸ B. Viaud,⁴⁸ H. Nicholson,⁴⁹ N. Cavallo,^{50,†} G. De Nardo,⁵⁰ F. Fabozzi,^{50,†} C. Gatto,⁵⁰ L. Lista,⁵⁰ D. Monorchio,⁵⁰ P. Paolucci,⁵⁰ D. Piccolo,⁵⁰ C. Sciacca,⁵⁰ M. Baak,⁵¹ H. Bulten,⁵¹ G. Raven,⁵¹ H. L. Snoek,⁵¹ L. Wilden,⁵¹ C. P. Jessop,⁵² J. M. LoSecco,⁵² T. Allmendinger,⁵³ G. Benelli,⁵³ K. K. Gan,⁵³ K. Honscheid,⁵³ D. Hufnagel,⁵³ P. D. Jackson,⁵³ H. Kagan,⁵³ R. Kass,⁵³ T. Pulliam,⁵³ A. M. Rahimi,⁵³ R. Ter-Antonyan,⁵³ Q. K. Wong,⁵³ J. Brau,⁵⁴ R. Frey,⁵⁴ O. Igonkina,⁵⁴ M. Lu,⁵⁴ C. T. Potter,⁵⁴ N. B. Sinev,⁵⁴ D. Strom,⁵⁴ J. Strube,⁵⁴ E. Torrence,⁵⁴ A. Dorigo,⁵⁵ F. Galeazzi,⁵⁵ M. Margoni,⁵⁵ M. Morandin,⁵⁵ M. Posocco,⁵⁵ M. Rotondo,⁵⁵ F. Simonetto,⁵⁵ R. Stroili,⁵⁵ C. Voci,⁵⁵ M. Benayoun,⁵⁶ H. Briand,⁵⁶ J. Chauveau,⁵⁶ P. David,⁵⁶ L. Del Buono,⁵⁶ Ch. de la Vaissière,⁵⁶ O. Hamon,⁵⁶ M. J. J. John,⁵⁶ Ph. Leruste,⁵⁶ J. Malclès,⁵⁶ J. Ocariz,⁵⁶ L. Roos,⁵⁶ G. Therin,⁵⁶ P. K. Behera,⁵⁷ L. Gladney,⁵⁷ Q. H. Guo,⁵⁷ J. Panetta,⁵⁷ M. Biasini,⁵⁸ R. Covarelli,⁵⁸ S. Pacetti,⁵⁸ M. Pioppi,⁵⁸ C. Angelini,⁵⁹ G. Batignani,⁵⁹ S. Bettarini,⁵⁹ F. Bucci,⁵⁹ G. Calderini,⁵⁹ M. Carpinelli,⁵⁹ R. Cenci,⁵⁹ F. Forti,⁵⁹ M. A. Giorgi,⁵⁹ A. Lusiani,⁵⁹ G. Marchiori,⁵⁹ M. Morganti,⁵⁹ N. Neri,⁵⁹ E. Paoloni,⁵⁹ M. Rama,⁵⁹ G. Rizzo,⁵⁹ J. Walsh,⁵⁹ M. Haire,⁶⁰ D. Judd,⁶⁰ D. E. Wagoner,⁶⁰ J. Biesiada,⁶¹ N. Danielson,⁶¹ P. Elmer,⁶¹ Y. P. Lau,⁶¹ C. Lu,⁶¹ J. Olsen,⁶¹ A. J. S. Smith,⁶¹ A. V. Telnov,⁶¹ F. Bellini,⁶² G. Cavoto,⁶² A. D'Orazio,⁶² E. Di Marco,⁶² R. Faccini,⁶² F. Ferrarotto,⁶² F. Ferroni,⁶² M. Gaspero,⁶² L. Li Gioi,⁶² M. A. Mazzoni,⁶² S. Morganti,⁶² G. Piredda,⁶² F. Polci,⁶² F. Safai Tehrani,⁶² C. Voena,⁶² H. Schröder,⁶³ G. Wagner,⁶³ R. Waldi,⁶³ T. Adye,⁶⁴ N. De Groot,⁶⁴ B. Franek,⁶⁴ G. P. Gopal,⁶⁴ E. O. Olaiya,⁶⁴ F. F. Wilson,⁶⁴ R. Aleksan,⁶⁵ S. Emery,⁶⁵ A. Gaidot,⁶⁵ S. F. Ganzhur,⁶⁵ P.-F. Giraud,⁶⁵ G. Graziani,⁶⁵ G. Hamel de Monchenault,⁶⁵ W. Kozanecki,⁶⁵ M. Legendre,⁶⁵ G. W. London,⁶⁵ B. Mayer,⁶⁵ G. Vasseur,⁶⁵ Ch. Yèche,⁶⁵ M. Zito,⁶⁵ M. V. Purohit,⁶⁶ A. W. Weidemann,⁶⁶ J. R. Wilson,⁶⁶ F. X. Yumiceva,⁶⁶ T. Abe,⁶⁷ M. T. Allen,⁶⁷ D. Aston,⁶⁷ R. Bartoldus,⁶⁷ N. Berger,⁶⁷ A. M. Boyarski,⁶⁷ O. L. Buchmueller,⁶⁷ R. Claus,⁶⁷ M. R. Convery,⁶⁷ M. Cristinziani,⁶⁷ J. C. Dingfelder,⁶⁷ D. Dong,⁶⁷ J. Dorfan,⁶⁷ D. Dujmic,⁶⁷ W. Dunwoodie,⁶⁷ S. Fan,⁶⁷ R. C. Field,⁶⁷ T. Glanzman,⁶⁷ S. J. Gowdy,⁶⁷ T. Hadig,⁶⁷ V. Halyo,⁶⁷ C. Hast,⁶⁷ T. Hryn'ova,⁶⁷ W. R. Innes,⁶⁷ M. H. Kelsey,⁶⁷ P. Kim,⁶⁷ M. L. Kocian,⁶⁷ D. W. G. S. Leith,⁶⁷ J. Libby,⁶⁷ S. Luitz,⁶⁷ V. Luth,⁶⁷ H. L. Lynch,⁶⁷ H. Marsiske,⁶⁷ S. Menke,⁶⁷ R. Messner,⁶⁷ D. R. Muller,⁶⁷ C. P. O'Grady,⁶⁷ V. E. Ozcan,⁶⁷ A. Perazzo,⁶⁷ M. Perl,⁶⁷ B. N. Ratcliff,⁶⁷ A. Roodman,⁶⁷ A. A. Salnikov,⁶⁷ R. H. Schindler,⁶⁷ J. Schwiening,⁶⁷ A. Snyder,⁶⁷ J. Stelzer,⁶⁷ D. Su,⁶⁷ M. K. Sullivan,⁶⁷ K. Suzuki,⁶⁷ S. Swain,⁶⁷ J. M. Thompson,⁶⁷ J. Va'vra,⁶⁷ M. Weaver,⁶⁷ W. J. Wisniewski,⁶⁷ M. Wittgen,⁶⁷ D. H. Wright,⁶⁷ A. K. Yarritu,⁶⁷ K. Yi,⁶⁷ C. C. Young,⁶⁷ P. R. Burchat,⁶⁸ A. J. Edwards,⁶⁸ S. A. Majewski,⁶⁸ B. A. Petersen,⁶⁸ C. Roat,⁶⁸ M. Ahmed,⁶⁹ S. Ahmed,⁶⁹ M. S. Alam,⁶⁹ J. A. Ernst,⁶⁹ M. A. Saeed,⁶⁹ F. R. Wappler,⁶⁹ S. B. Zain,⁶⁹ W. Bugg,⁷⁰ M. Krishnamurthy,⁷⁰ S. M. Spanier,⁷⁰ R. Eckmann,⁷¹ J. L. Ritchie,⁷¹ A. Satpathy,⁷¹ R. F. Schwitters,⁷¹ J. M. Izen,⁷² I. Kitayama,⁷² X. C. Lou,⁷² S. Ye,⁷² F. Bianchi,⁷³ M. Bona,⁷³ F. Gallo,⁷³ D. Gamba,⁷³ M. Bomben,⁷⁴ L. Bosisio,⁷⁴ C. Cartaro,⁷⁴ F. Cossutti,⁷⁴ G. Della Ricca,⁷⁴ S. Dittongo,⁷⁴ S. Grancagnolo,⁷⁴ L. Lanceri,⁷⁴ L. Vitale,⁷⁴ F. Martinez-Vidal,⁷⁵ R. S. Panvini,^{76,‡} Sw. Banerjee,⁷⁷ B. Bhuyan,⁷⁷ C. M. Brown,⁷⁷ D. Fortin,⁷⁷ K. Hamano,⁷⁷ R. Kowalewski,⁷⁷ J. M. Roney,⁷⁷ R. J. Sobie,⁷⁷ J. J. Back,⁷⁸ P. F. Harrison,⁷⁸ T. E. Latham,⁷⁸ G. B. Mohanty,⁷⁸ H. R. Band,⁷⁹ X. Chen,⁷⁹ B. Cheng,⁷⁹ S. Dasu,⁷⁹ M. Datta,⁷⁹ A. M. Eichenbaum,⁷⁹ K. T. Flood,⁷⁹ M. Graham,⁷⁹ J. J. Hollar,⁷⁹ J. R. Johnson,⁷⁹ P. E. Kutter,⁷⁹ H. Li,⁷⁹ R. Liu,⁷⁹ B. Mellado,⁷⁹ A. Mihalyi,⁷⁹ Y. Pan,⁷⁹ R. Prepost,⁷⁹ P. Tan,⁷⁹ J. H. von Wimmersperg-Toeller,⁷⁹ S. L. Wu,⁷⁹ Z. Yu,⁷⁹ and H. Neal⁸⁰

(The BABAR Collaboration)

¹Laboratoire de Physique des Particules, F-74941 Annecy-le-Vieux, France

²IFAE, Universitat Autònoma de Barcelona, E-08193 Bellaterra, Barcelona, Spain

³Università di Bari, Dipartimento di Fisica and INFN, I-70126 Bari, Italy

⁴Institute of High Energy Physics, Beijing 100039, China

⁵University of Bergen, Inst. of Physics, N-5007 Bergen, Norway

⁶Lawrence Berkeley National Laboratory and University of California, Berkeley, California 94720, USA

⁷University of Birmingham, Birmingham, B15 2TT, United Kingdom

⁸Ruhr Universität Bochum, Institut für Experimentalphysik 1, D-44780 Bochum, Germany

⁹University of Bristol, Bristol BS8 1TL, United Kingdom

¹⁰University of British Columbia, Vancouver, British Columbia, Canada V6T 1Z1

¹¹Brunel University, Uxbridge, Middlesex UB8 3PH, United Kingdom

¹²Budker Institute of Nuclear Physics, Novosibirsk 630090, Russia

¹³University of California at Irvine, Irvine, California 92697, USA

- ¹⁴University of California at Los Angeles, Los Angeles, California 90024, USA
¹⁵University of California at Riverside, Riverside, California 92521, USA
¹⁶University of California at San Diego, La Jolla, California 92093, USA
¹⁷University of California at Santa Barbara, Santa Barbara, California 93106, USA
¹⁸University of California at Santa Cruz, Institute for Particle Physics, Santa Cruz, California 95064, USA
¹⁹California Institute of Technology, Pasadena, California 91125, USA
²⁰University of Cincinnati, Cincinnati, Ohio 45221, USA
²¹University of Colorado, Boulder, Colorado 80309, USA
²²Colorado State University, Fort Collins, Colorado 80523, USA
²³Universität Dortmund, Institut für Physik, D-44221 Dortmund, Germany
²⁴Technische Universität Dresden, Institut für Kern- und Teilchenphysik, D-01062 Dresden, Germany
²⁵Ecole Polytechnique, LLR, F-91128 Palaiseau, France
²⁶University of Edinburgh, Edinburgh EH9 3JZ, United Kingdom
²⁷Università di Ferrara, Dipartimento di Fisica and INFN, I-44100 Ferrara, Italy
²⁸Laboratori Nazionali di Frascati dell'INFN, I-00044 Frascati, Italy
²⁹Università di Genova, Dipartimento di Fisica and INFN, I-16146 Genova, Italy
³⁰Harvard University, Cambridge, Massachusetts 02138, USA
³¹Universität Heidelberg, Physikalisches Institut, Philosophenweg 12, D-69120 Heidelberg, Germany
³²Imperial College London, London, SW7 2AZ, United Kingdom
³³University of Iowa, Iowa City, Iowa 52242, USA
³⁴Iowa State University, Ames, Iowa 50011-3160, USA
³⁵Laboratoire de l'Accélérateur Linéaire, F-91898 Orsay, France
³⁶Lawrence Livermore National Laboratory, Livermore, California 94550, USA
³⁷University of Liverpool, Liverpool L69 7ZE, United Kingdom
³⁸Queen Mary, University of London, E1 4NS, United Kingdom
³⁹University of London, Royal Holloway and Bedford New College, Egham, Surrey TW20 0EX, United Kingdom
⁴⁰University of Louisville, Louisville, Kentucky 40292, USA
⁴¹University of Manchester, Manchester M13 9PL, United Kingdom
⁴²University of Maryland, College Park, Maryland 20742, USA
⁴³University of Massachusetts, Amherst, Massachusetts 01003, USA
⁴⁴Massachusetts Institute of Technology, Laboratory for Nuclear Science, Cambridge, Massachusetts 02139, USA
⁴⁵McGill University, Montréal, Quebec, Canada H3A 2T8
⁴⁶Università di Milano, Dipartimento di Fisica and INFN, I-20133 Milano, Italy
⁴⁷University of Mississippi, University, Mississippi 38677, USA
⁴⁸Université de Montréal, Laboratoire René J. A. Lévesque, Montréal, Quebec, Canada H3C 3J7
⁴⁹Mount Holyoke College, South Hadley, Massachusetts 01075, USA
⁵⁰Università di Napoli Federico II, Dipartimento di Scienze Fisiche and INFN, I-80126, Napoli, Italy
⁵¹NIKHEF, National Institute for Nuclear Physics and High Energy Physics, NL-1009 DB Amsterdam, The Netherlands
⁵²University of Notre Dame, Notre Dame, Indiana 46556, USA
⁵³Ohio State University, Columbus, Ohio 43210, USA
⁵⁴University of Oregon, Eugene, Oregon 97403, USA
⁵⁵Università di Padova, Dipartimento di Fisica and INFN, I-35131 Padova, Italy
⁵⁶Universités Paris VI et VII, Laboratoire de Physique Nucléaire et de Hautes Energies, F-75252 Paris, France
⁵⁷University of Pennsylvania, Philadelphia, Pennsylvania 19104, USA
⁵⁸Università di Perugia, Dipartimento di Fisica and INFN, I-06100 Perugia, Italy
⁵⁹Università di Pisa, Dipartimento di Fisica, Scuola Normale Superiore and INFN, I-56127 Pisa, Italy
⁶⁰Prairie View A&M University, Prairie View, Texas 77446, USA
⁶¹Princeton University, Princeton, New Jersey 08544, USA
⁶²Università di Roma La Sapienza, Dipartimento di Fisica and INFN, I-00185 Roma, Italy
⁶³Universität Rostock, D-18051 Rostock, Germany
⁶⁴Rutherford Appleton Laboratory, Chilton, Didcot, Oxon, OX11 0QX, United Kingdom
⁶⁵DSM/Dapnia, CEA/Saclay, F-91191 Gif-sur-Yvette, France
⁶⁶University of South Carolina, Columbia, South Carolina 29208, USA
⁶⁷Stanford Linear Accelerator Center, Stanford, California 94309, USA
⁶⁸Stanford University, Stanford, California 94305-4060, USA
⁶⁹State University of New York, Albany, New York 12222, USA
⁷⁰University of Tennessee, Knoxville, Tennessee 37996, USA
⁷¹University of Texas at Austin, Austin, Texas 78712, USA
⁷²University of Texas at Dallas, Richardson, Texas 75083, USA
⁷³Università di Torino, Dipartimento di Fisica Sperimentale and INFN, I-10125 Torino, Italy
⁷⁴Università di Trieste, Dipartimento di Fisica and INFN, I-34127 Trieste, Italy
⁷⁵IFIC, Universitat de Valencia-CSIC, E-46071 Valencia, Spain
⁷⁶Vanderbilt University, Nashville, Tennessee 37235, USA
⁷⁷University of Victoria, Victoria, British Columbia, Canada V8W 3P6

⁷⁸*Department of Physics, University of Warwick, Coventry CV4 7AL, United Kingdom*

⁷⁹*University of Wisconsin, Madison, Wisconsin 53706, USA*

⁸⁰*Yale University, New Haven, Connecticut 06511, USA*

(Dated: June 9, 2006)

We present a determination of the CKM matrix element $|V_{ub}|$ based on the analysis of semileptonic B decays from a sample of 88 million $\Upsilon(4S)$ decays collected with the BABAR detector at the PEP-II e^+e^- storage ring. Charmless semileptonic B decays are selected using measurements of the electron energy and the invariant mass squared of the electron-neutrino pair. We obtain $|V_{ub}| = (4.41 \pm 0.30 \text{ }^{+0.65}_{-0.47} \pm 0.28) \times 10^{-3}$, where the errors represent experimental uncertainties, heavy quark parameter uncertainties, and theoretical uncertainties, respectively.

Note: this document includes corrections to the original publication of this work [1]. These corrections, which affect the calculated efficiencies and quantities derived from them, will appear in an erratum.

PACS numbers: 13.20.He, 12.15.Hh, 14.40.Nd

The study of the weak interactions of quarks has played a crucial role in the development of the Standard Model (SM), which embodies our understanding of the fundamental interactions. The increasingly precise measurements of CP asymmetries in B decays allow stringent experimental tests of the SM mechanism for CP violation via the complex phase in the Cabibbo-Kobayashi-Maskawa (CKM) matrix [2]. Improved determinations of $|V_{ub}|$, the coupling strength of the b quark to the u quark, will improve the sensitivity of these tests.

Two observables have been used to determine $|V_{ub}|$ from inclusive semileptonic B decays: the endpoint of the lepton momentum spectrum [3] and the mass of the accompanying hadronic system [4]. In this paper, semileptonic $\bar{B} \rightarrow X_u e \bar{\nu}$ decays are selected using a novel approach based on simultaneous requirements for the electron energy, E_e , and the invariant mass squared of the $e\bar{\nu}$ pair, q^2 [5]. The neutrino 4-momentum is reconstructed from the visible 4-momentum and knowledge of the e^+e^- initial state. The dominant charm background is suppressed by selecting a region of the q^2 - E_e phase space where correctly reconstructed $\bar{B} \rightarrow X_c e \bar{\nu}$ events are kinematically excluded. Background contamination in the signal region is due to resolution effects and is evaluated in Monte Carlo (MC) simulations. Theoretical calculations are applied to the measured $\bar{B} \rightarrow X_u e \bar{\nu}$ partial rate to determine $|V_{ub}|$, the precision of which is limited mostly by our current knowledge of the b -quark mass, m_b .

The data used in this analysis were collected with the BABAR detector [6] at the PEP-II asymmetric-energy e^+e^- storage ring. The data set consists of 88.4 million $B\bar{B}$ pairs collected at the $\Upsilon(4S)$ resonance, corresponding to an integrated luminosity of 81.4 fb^{-1} at $\sqrt{s} = 10.58 \text{ GeV}$. An additional 9.6 fb^{-1} of data were collected at center-of-mass energies 20 MeV below the $B\bar{B}$ threshold. Off-resonance data are used to subtract the non- $B\bar{B}$ contributions from the data collected at the $\Upsilon(4S)$ resonance. To do so, the off-resonance data are scaled according to the integrated luminosity and the energy dependence of the QED cross-section, and

the particles are boosted to the $\Upsilon(4S)$ resonance energy. Throughout this paper, all kinematic variables are given in the $\Upsilon(4S)$ rest frame unless stated otherwise.

The simulation of charmless semileptonic B decays used in optimizing the analysis and determining reconstruction efficiencies is based on the Heavy Quark Expansion (HQE) including $\mathcal{O}(\alpha_S)$ corrections [7]. This calculation produces a continuous spectrum of hadronic masses, m_X . Subsequent hadronization is simulated using JETSET down to $2m_\pi$ [8]. Decays to low-mass hadrons (π , η , ρ , ω , η') are simulated separately using the ISGW2 model [9], and mixed with the non-resonant states so that the m_X , q^2 and E_e spectral distributions correspond as closely as possible to the HQE calculation.

Hadronic events containing an identified electron with energy $2.1 \text{ GeV} < E_e < 2.8 \text{ GeV}$ are selected. Radiative Bhabha events rejected using the criteria given in Ref. [10] and electrons from $J/\psi \rightarrow e^+e^-$ decays are vetoed. The total visible 4-momentum, p_{vis} , is determined using charged tracks emanating from the collision point, identified pairs of charged tracks from $K_S^0 \rightarrow \pi^+\pi^-$, $\Lambda \rightarrow p\pi^-$ and $\gamma \rightarrow e^+e^-$, and energy deposits in the electromagnetic calorimeter. Each charged particle is assigned a mass hypothesis based on particle identification information. Calorimeter clusters unassociated with a charged track and with a lateral energy spread consistent with electromagnetic showers are treated as photons.

Additional requirements are made to improve the quality of the neutrino reconstruction and suppress contributions from $e^+e^- \rightarrow q\bar{q}$ continuum events. We form the missing 4-momentum, $p_{\text{miss}} = p_{e^+e^-} - p_{\text{vis}}$, where $p_{e^+e^-}$ is the 4-momentum of the initial state. For each event we require (1) no additional identified e or μ ; (2) $-0.95 < \cos \theta_{\text{miss}} < 0.8$, where θ_{miss} is the polar angle of the missing 3-momentum; (3) $0.0 \text{ GeV} < E_{\text{miss}} - |\mathbf{p}_{\text{miss}}| < 0.8 \text{ GeV}$, where E_{miss} is the missing energy in the event; (4) $|\mathbf{p}_{\text{miss}}| < 2.5 \text{ GeV}$ and (5) $|\cos \theta_T| < 0.75$, where θ_T is the angle between the electron momentum and the thrust vector of the remaining particles in the event.

The measured $|\mathbf{p}_{\text{miss}}|$ differs from the true neutrino momentum due to additional particles that escape de-

tection. Therefore, a bias correction, $\mathbf{p}_\nu = \mathbf{p}_{\text{miss}}(0.804 - 0.078/|\mathbf{p}_{\text{miss}}|)$, is derived from the simulation. Since the resolution on $|\mathbf{p}_{\text{miss}}|$ is superior to that of E_{miss} , we set $p_\nu = (|\mathbf{p}_\nu|, |\mathbf{p}_\nu|)$ and $q^2 = (p_e + p_\nu)^2$. Defining $\eta_\pm = \sqrt{(1 \pm \beta)/(1 \mp \beta)}$, where $\beta \simeq 0.06$ is the velocity of the B meson in the $\Upsilon(4S)$ frame, the maximum kinematically allowed hadronic mass squared for a given E_e and q^2 is $s_h^{\text{max}} = m_B^2 + q^2 - 2m_B(E_e\eta_- + q^2\eta_+/4E_e)$ for $\pm 2E_e > \pm\sqrt{q^2}\eta_\pm$, and $s_h^{\text{max}} = m_B^2 + q^2 - 2m_B\sqrt{q^2}$ otherwise. We require $s_h^{\text{max}} < 3.5 \text{ GeV}^2 \approx m_{D^0}^2$; no $\bar{B} \rightarrow X_c e \bar{\nu}$ decays can have values of s_h^{max} below this limit before accounting for resolution. The requirements on E_e and s_h^{max} and criteria (1)–(5) were chosen to minimize the total (experimental and theoretical) expected uncertainty $\sigma(|V_{ub}|)/|V_{ub}|$.

The quality of the neutrino reconstruction is evaluated using a control sample ($De\bar{\nu}$) consisting of the decays $\bar{B} \rightarrow D^0 e \bar{\nu}(X)$, where kinematic criteria result in the X system typically being no more than a π or γ from a $D^* \rightarrow D^0 X$ transition. The D^0 is reconstructed in the $K^-\pi^+$ decay mode and we require $|\mathbf{p}_{D^0}| > 0.5 \text{ GeV}$ and $E_e > 1.4 \text{ GeV}$. The $D^0 e$ combination must satisfy $-2.5 < \cos\theta_{B,De} < 1.1$, where $\cos\theta_{B,De} = (2E_B E_{De} - m_B^2 - m_{De}^2)/(2|\mathbf{p}_B||\mathbf{p}_{De}|)$ is the cosine of the angle between the vector momenta of the \bar{B} and the $D^0 e$ system assuming the only missing particle in the \bar{B} decay was a single neutrino. After the combinatorial background is subtracted using D^0 mass sidebands, the selected sample consists primarily ($\simeq 95\%$) of $\bar{B} \rightarrow D^0 e \bar{\nu}$ and $\bar{B} \rightarrow D^* e \bar{\nu}$ decays. The control sample selection makes no requirements on the other B in the event, and can therefore be used to study the impact of the modeling of the other B on the neutrino reconstruction. Since the unreconstructed X system in the $\bar{B} \rightarrow D^0 e \bar{\nu}(X)$ decays carries away little energy, a good estimate (r.m.s. $\sim 0.2 \text{ GeV}$) of the neutrino energy can be obtained from the known \bar{B} energy and the measured D^0 and e energies, E_ν^{De} . A second estimate of the neutrino energy is constructed from the visible momentum as described previously. Subtracting the first estimate from the second gives the distribution shown in Fig. 1, where the criteria (1)–(5) described above have been imposed. We find good agreement between data and MC; the average (r.m.s.) is 0.066 GeV (0.366 GeV) for data and 0.072 GeV (0.365 GeV) for simulated events.

The $De\bar{\nu}$ control sample is also used to improve the modeling of the $\bar{B} \rightarrow X_c e \bar{\nu}$ decays. After relaxing the $\cos\theta_{B,De}$ requirements and subtracting continuum and combinatorial backgrounds, we perform a binned χ^2 fit to the $De\bar{\nu}$ sample in the variables $|\mathbf{p}_D|$, E_e and $\cos\theta_{B,De}$. The fit determines scale factors for the MC components $\bar{B} \rightarrow De\bar{\nu}$, $\bar{B} \rightarrow D^* e \bar{\nu}$ and other contributions (85% of which are decays to D^{**} states), while keeping the total $\bar{B} \rightarrow X_c e \bar{\nu}$ branching fraction fixed to the measured value [11]. The fit increases the $\bar{B} \rightarrow De\bar{\nu}$ and $\bar{B} \rightarrow D^* e \bar{\nu}$ branching fractions to 2.29% and 6.02% (2.48%




FIG. 1: The difference between the two neutrino energy estimates described in the text for continuum-subtracted data and simulated $B\bar{B}$ events for the $De\bar{\nu}$ control sample.

and 6.52%) for neutral (charged) B mesons, respectively, while decreasing the remaining contributions. By design, these revised branching fractions respect isospin symmetry and are used in the determination of the background.

Two control samples are used to reduce the sensitivity of the efficiency and background estimates to details of the simulation: the $De\bar{\nu}$ control sample described above, but with $E_e > 2.0 \text{ GeV}$; and events satisfying the normal selection criteria but having $s_h^{\text{max}} > 4.25 \text{ GeV}^2$, a sample with $< 5\%$ signal decays. Efficiencies $\epsilon_{De\bar{\nu}}^{\text{data}}$ and $\epsilon_{De\bar{\nu}}^{\text{MC}}$ are calculated separately in data and MC as the ratio of $De\bar{\nu}$ candidates satisfying criteria (1)–(5) to the total $De\bar{\nu}$ sample. The $\bar{B} \rightarrow X_u e \bar{\nu}$ signal efficiency is multiplied by the ratio of these efficiencies to reduce sensitivity to details of the simulation. The $s_h^{\text{max}} > 4.25 \text{ GeV}^2$ sideband region is used to normalize the simulated s_h^{max} distribution to the data, reducing sensitivity to background normalization uncertainties.

We determine a partial branching fraction $\Delta\mathcal{B}(\tilde{E}, \tilde{s}_h^{\text{max}}) = \mathcal{B}(\bar{B} \rightarrow X_u e \bar{\nu})f_u$, unfolded for detector effects. The acceptance, f_u , is the fraction of $\bar{B} \rightarrow X_u e \bar{\nu}$ decays in the region of interest, $\tilde{E}_e > 2.0 \text{ GeV}$ and $\tilde{s}_h^{\text{max}} < 3.5 \text{ GeV}^2$, where \tilde{E}_e and \tilde{s}_h^{max} are the true (generated) values in the B meson rest frame. Slightly lower values are accepted for \tilde{E}_e than for E_e to account for the boost of the B meson and to increase f_u . The efficiency times acceptance for $\bar{B} \rightarrow X_u e \bar{\nu}$ decays can be written as $\epsilon_u = \epsilon_{\text{sig}} f_u + \epsilon_{\overline{\text{sig}}}(1 - f_u)$, where ϵ_{sig} ($\epsilon_{\overline{\text{sig}}}$) is the efficiency for an event inside (outside) the region of interest to be reconstructed and pass our selection criteria. We calculate the partial branching fraction as follows:

$$\Delta\mathcal{B} = \frac{N_{\text{cand}} - M_{\text{bkg}} \frac{N_{\text{side}}}{M_{\text{side}}}}{2 N_{B\bar{B}} \frac{\epsilon_{De\bar{\nu}}^{\text{data}}}{\epsilon_{De\bar{\nu}}^{\text{MC}}} \epsilon_{\text{sig}}} \left[1 + \frac{1 - f_u \epsilon_{\overline{\text{sig}}}}{f_u \epsilon_{\text{sig}}} \right]^{-1}, \quad (1)$$

where N_{cand} and N_{side} refer to the number of candidates in the signal and s_h^{max} sideband regions of the data, M_{bkg} and M_{side} refer to background in the signal re-

gion and the yield in the sideband region in simulated events and $2N_{B\bar{B}}$ is the number of B mesons produced from $\Upsilon(4S) \rightarrow B\bar{B}$ decays. Since the resulting ratio of $\epsilon_{\text{sig}}/\epsilon_{\text{sig}}$ is small, $\Delta\mathcal{B}$ depends only weakly on the model used to determine f_u .

Fig. 2 shows the electron energy and s_h^{max} distributions after cuts have been applied to all variables except the one being displayed. The discrepancy observed between data and MC for $E_e < 1.95$ GeV is covered by the systematic error on the $\bar{B} \rightarrow X_c e \bar{\nu}$ modeling. The yields and efficiencies are given in Table I. We find

$$\Delta\mathcal{B}(2.0, 3.5) = (4.41 \pm 0.42 \pm 0.42) \times 10^{-4} \quad (2)$$

where the uncertainties are statistical and systematic, respectively. Alternative values of $\Delta\mathcal{B}$ are obtained using different electron energy requirements: $\Delta\mathcal{B}(1.9, 3.5) = (5.29 \pm 0.44 \pm 0.72) \times 10^{-4}$ and $\Delta\mathcal{B}(2.1, 3.5) = (3.68 \pm 0.43 \pm 0.36) \times 10^{-4}$.




FIG. 2: The electron energy, E_e , and s_h^{max} spectra in the $\Upsilon(4S)$ frame for continuum-subtracted data and simulated $B\bar{B}$ events satisfying all selection requirements except for the variable shown. The arrows denote the signal (and sideband) region in E_e and s_h^{max} . Note that $\tilde{E}_e \neq E_e$ (see text).

Systematic uncertainties are assigned for the modeling of the signal $\bar{B} \rightarrow X_u e \bar{\nu}$ decays, background and detector response. The leading sources of uncertainty are listed in Table II. Uncertainties from the simulation of charged particle tracking, neutral reconstruction, charged particle identification, and the energy deposition by K_L^0 were evaluated from studies comparing data and simulation. Radiation in the decay process was simulated using

TABLE I: Yields and efficiencies from data and simulation. All uncertainties are statistical except for $N_{B\bar{B}}$ where systematics are included. Efficiencies are quoted in units of 10^{-4} .

| | | | | | |
|-------------------|-------------------|--|-------------------------|------------------------------|---------|
| N_{cand} | N_{side} | $\epsilon_{D e \bar{\nu}}^{\text{data}}$ | $N_{B\bar{B}} (10^6)$ | | |
| 5130 ± 150 | 6152 ± 130 | 902 ± 39 | 88.35 ± 0.97 | | |
| M_{bkg} | M_{side} | $\epsilon_{D e \bar{\nu}}^{\text{MC}}$ | ϵ_{sig} | $\epsilon_{\text{sig}}^{-1}$ | f_u |
| 3176 ± 35 | 6423 ± 49 | 906 ± 19 | 256 ± 3 | 2.7 ± 0.2 | 0.174 |

PHOTOS [12]; comparisons with the analytical result of Ref. [13] were used to assess the systematic uncertainty. The uncertainty due to bremsstrahlung in the detector was evaluated using the method of Ref. [11]. The uncertainty in modeling the background was first evaluated by varying the total $\bar{B} \rightarrow X_c e \bar{\nu}$, $\bar{B} \rightarrow D e \bar{\nu}$ and $\bar{B} \rightarrow D^* e \bar{\nu}$ rates within their measured range. Furthermore, the form factors for $\bar{B} \rightarrow D e \bar{\nu}$ [14] and $\bar{B} \rightarrow D^* e \bar{\nu}$ [15] were varied within their uncertainties, and the composition of the D^{**} states was modified to include only narrow resonances, broad resonances, or Goity-Roberts decays [16]; the effect of these variations is reduced by the fit to the $D e \bar{\nu}$ control sample. The modeling of D decays was varied based on the measurements reported in Ref. [17]; the variation in the $D \rightarrow K_L^0 X$ branching fractions dominates the uncertainty. The modeling of $\bar{B} \rightarrow X_u e \bar{\nu}$ decays is sensitive to the resonance structure at low mass. The branching fractions of $\bar{B} \rightarrow (\pi, \rho, \omega, \eta, \eta') e \bar{\nu}$ were varied as follows: π : $\pm 30\%$; ρ : $\pm 30\%$; ω : $\pm 40\%$; simultaneously η and η' : $\pm 100\%$.

TABLE II: Uncertainties on $|V_{ub}|$ and $\Delta\mathcal{B}$.

| Source | $\sigma(V_{ub})/ V_{ub} $ (%) | $\sigma(\Delta\mathcal{B})/\Delta\mathcal{B}$ (%) |
|--|---------------------------------|---|
| Tracking | ± 0.8 | ± 1.5 |
| Neutrals | ± 1.7 | ± 3.4 |
| Electron ID | ± 0.5 | ± 1.0 |
| Hadron ID | ± 1.0 | ± 2.0 |
| Bremsstrahlung | ± 1.0 | ± 2.0 |
| K_L^0 | ± 1.3 | ± 2.6 |
| $N_{B\bar{B}}$ | ± 0.6 | ± 1.1 |
| Radiation | ± 1.9 | ± 3.8 |
| $\bar{B} \rightarrow X_c e \bar{\nu}$ modeling | ± 2.5 | ± 5.0 |
| $\bar{B} \rightarrow X_u e \bar{\nu}$ resonances | ± 2.2 | ± 4.4 |
| Statistical | ± 4.7 | ± 9.3 |
| Total experimental | ± 6.7 | ± 13.3 |
| Heavy quark parameters | $^{+14.6}_{-10.6}$ | ± 1.5 |
| Theoretical | ± 6.3 | |

We extract $|V_{ub}| = [\Delta\mathcal{B}/(\Delta\zeta \tau_B)]^{1/2}$ using $\tau_B = 1.604 \pm 0.023$ ps [17]. The normalized partial rate, $\Delta\zeta$, computed in units of $\Delta\Gamma/|V_{ub}|^2$, is taken from Ref. [18], in which the leading terms in the HQE of the $\bar{B} \rightarrow X_u e \bar{\nu}$ spectra are computed at next-to-leading order, and power corrections are included at $\mathcal{O}(\alpha_S)$ for the lead-

ing shape function (SF) and at tree level for subleading SFs. The values used for the heavy quark parameters, $m_b = 4.61 \pm 0.08 \text{ GeV}$ and $\mu_\pi^2 = 0.15 \pm 0.07 \text{ GeV}^2$, with a correlation coefficient of -0.4 , are based on fits to $\bar{B} \rightarrow X_c \ell \bar{\nu}_\ell$ moments [19], translated to the shape-function scheme of Ref. [20].

We find $|V_{ub}| = (4.41 \pm 0.30_{-0.47}^{+0.65} \pm 0.28) \times 10^{-3}$ for $\tilde{E}_e > 2.0 \text{ GeV}$, where the errors represent experimental, heavy quark parameters, and theoretical uncertainties, respectively. The latter include estimates of the effects of subleading SFs [21], variations in the matching scales used in the calculation, and weak annihilation [22]. No uncertainty is assigned for possible quark-hadron duality violation. The determination of $|V_{ub}|$ is limited primarily by our knowledge of m_b . An approximate dependence is $|V_{ub}(m_b)| = |V_{ub}(m_0)|(1 + 7(m_b - m_0)/m_0)$, where $m_0 = 4.61 \text{ GeV}$. The sensitivity to higher moments of the SF is weak; the change in $|V_{ub}|$ when varying μ_π^2 from 0.03 to 0.35 GeV^2 with m_b fixed is 2% , and the impact of using alternative SF parameterizations [23] is $< 2\%$. The overall precision on the above result surpasses that of Refs. [3] and [4], but is comparable to determinations of $|V_{ub}|$ which have become available while this paper was nearing completion [24].

ACKNOWLEDGMENTS

We are grateful for the excellent luminosity and machine conditions provided by our PEP-II colleagues, and for the substantial dedicated effort from the computing organizations that support *BABAR*. The collaborating institutions wish to thank SLAC for its support and kind hospitality. This work is supported by DOE and NSF (USA), NSERC (Canada), IHEP (China), CEA and CNRS-IN2P3 (France), BMBF and DFG (Germany), INFN (Italy), FOM (The Netherlands), NFR (Norway), MIST (Russia), and PPARC (United Kingdom). Individuals have received support from CONACyT (Mexico), Marie Curie EIF (European Union), the A. P. Sloan Foundation, the Research Corporation, and the Alexander von Humboldt Foundation. Finally, we would like to thank the many theorists with whom we have had valuable discussions, and further thank M. Neubert, B. Lange and G. Paz for making available for our use a computer code implementing their calculations.

* Also with Università di Perugia, Dipartimento di Fisica, Perugia, Italy

- † Also with Università della Basilicata, Potenza, Italy
‡ Deceased
- [1] B. Aubert *et al.* (*BABAR* Collab.), Phys. Rev. Lett. **95**, 111801 (2005).
[2] M. Kobayashi and T. Maskawa, Prog. Theor. Phys. **49**, 652 (1973).
[3] R. Fulton *et al.* (CLEO Collab.), Phys. Rev. Lett. **64**, 16 (1990); J. Bartelt *et al.*, Phys. Rev. Lett. **71**, 4111 (1993); A. Bornheim *et al.*, Phys. Rev. Lett. **88**, 231803 (2002); H. Albrecht *et al.* (ARGUS Collab.), Phys. Lett. B **234**, 409 (1990); and Phys. Lett. B **255**, 297 (1991).
[4] B. Aubert *et al.* (*BABAR* Collab.), Phys. Rev. Lett. **92**, 071802 (2004); H. Kakuno *et al.* (BELLE Collab.), Phys. Rev. Lett. **92**, 101801 (2004).
[5] R. Kowalewski and S. Menke, Phys. Lett. B **541**, 29 (2002). We have modified the definition of s_h^{max} from that given in this reference.
[6] B. Aubert *et al.* (*BABAR* Collab.), Nucl. Instr. Methods Phys. Res., Sect. A **479**, 1 (2002).
[7] F. De Fazio and M. Neubert, JHEP **06**, 017 (1999).
[8] T. Sjöstrand, Comput. Phys. Commun. **82**, 74 (1994).
[9] N. Isgur, D. Scora, B. Grinstein and M. Wise, Phys. Rev. D **39**, 799 (1989).
[10] B. Aubert *et al.* (*BABAR* Collab.), Phys. Rev. D **67**, 031101 (2003).
[11] B. Aubert *et al.* (*BABAR* Collab.), Phys. Rev. D **69**, 111104 (2004).
[12] E. Richter-Was, Phys. Lett. B **303**, 163 (1993).
[13] E.S. Ginsberg, Phys. Rev. **142**, 1035 (1966).
[14] J. Bartelt *et al.* (CLEO Collab.), Phys. Rev. Lett. **82**, 3746 (1999).
[15] B. Aubert *et al.* (*BABAR* Collab.), “Measurement of the $B \rightarrow D^* \text{ Form Factors in the Semileptonic Decay } \bar{B} \rightarrow D^{*+} e^- \bar{\nu}$ ”, Contribution to the 2004 International Conference on High Energy Physics, hep-ex/409047 (2004).
[16] J. L. Goity and W. Roberts, Phys. Rev. D **51**, 3459 (1995).
[17] S. Eidelman *et al.*, Phys. Lett. B **592**, 1 (2004).
[18] B.O. Lange, M. Neubert and G. Paz, “Theory of Charmless Inclusive B Decays and the Extraction of $|V_{ub}|$ ”, hep-ph/0504071 (2005).
[19] B. Aubert *et al.* (*BABAR* Collab.), Phys. Rev. Lett. **93**, 011803 (2004).
[20] M. Neubert, Phys. Lett. B **612**, 13 (2005).
[21] S.W. Bosch, M. Neubert and G. Paz, JHEP **0411**, 073 (2004); M. Neubert, “Impact of four-quark shape functions on inclusive B decay spectra”, hep-ph/0411027 (2004).
[22] T.O. Meyer, “Limits on weak annihilation in inclusive charmless semileptonic B decays”, Cornell Univ. THESIS 05-1 (2005).
[23] A. Limosani and T. Nozaki, “Extraction of the b -quark shape function parameters using the Belle $B \rightarrow X_s \gamma$ photon energy spectrum”, hep-ex/0407052 (2004).
[24] The Heavy Flavor Averaging Group (EPS 2005 update): <http://www.slac.stanford.edu/xorg/hfag/>.