

Full Length Research Paper

Determining water content and other impurities in *Siparuna guianensis* Aublet essential oil using differential scanning calorimetry

Augustus Caesar Franke Portella^{1*}, Marilda Munaro², Chrystian de Assis Siqueira¹, Kleber Franke Portella², Magnólia de Mendonca Lopes¹, Alex Sander Rodrigues Cangussu¹ and Raimundo Wagner de Souza Aguiar¹

¹Universidade Federal do Tocantins, Departamento de Ciências Exatas, Campus Universitário de Gurupi, Gurupi, Tocantins, Brazil.

²Departamento de Tecnologia de Materiais, Instituto de Tecnologia para o Desenvolvimento, LACTEC, CP 19067, 81531-980 Curitiba – PR, Brazil.

Received 19 March, 2015; Accepted 31 August, 2015

Siparuna guianensis Aublet is a predominant species in the Brazilian Cerrado. Some studies found that its essential oil has properties that could be useful for manufacturing new products. Its quality depends on physicochemical properties, since its degradation, as well as water content and other volatile materials may cause several changes in its features. The differential scanning calorimetry (DSC) was used to determine the water content in the *S. guianensis* essential oil in order to evaluate its thermal parameters. The method is based on the hypothesis that the desolvation enthalpy (ΔH_{desol}) needed for removing n water moles from the essential oil is approximately the enthalpy needed for melting the same n water moles ($n\Delta H_f$). Thus, the current study considers the dissociation enthalpy (ΔH_{diss}) to be negligible. The total number of moles was calculated from the molecular mass value of the main essential oil components mentioned in the literature. The DSC curve in nitrogen atmosphere indicated the melting temperature of 1.6°C ($\Delta H=18.95 \text{ Jg}^{-1}$). Based on the dehydration enthalpies demonstrated in the DSC curve, it was possible to infer that the calculations on the number of water moles and other constant volatile materials found in the *S. guianensis* essential oil were compatible with Karl Fischer Titration (KFT) method. This method is commonly used to determine the water content in substances.

Key words: *Siparuna guianensis*, differential scanning calorimetry, thermogravimetric analyses.

INTRODUCTION

The physicochemical properties of the essential oils are crucial in the development of new products. Featuring the active ingredient may optimize quality parameters, ensure effectiveness and decrease the presence of water, impurities and/or degradation compounds. They may lead to undesired effects on the oil's composition. Water is a polar molecule and it may strongly interact

with ions or polar molecules of ion-dipole or dipole-dipole interactions, respectively. Since the hydrogen bonding energy between water molecules in solid state does not significantly differ from the bonding energy of hydrogen molecules of water in liquid state water molecules grouping may occur on the ice surface even before a complete monolayer is formed (Brown, 2005). Thermal

analysis refers to a group of techniques in which a property of the sample is measured against time or temperature, whereas the temperature of the sample, in a specified atmosphere, is heated or cooled at fixed rate of temperature change or hold at constant temperature (Giron, 2002). The differential scanning calorimetry (DSC) and the thermogravimetric analysis (TG) are the most used analytical techniques in the development of different studies on a wide variety of pharmaceutical materials (Souza et al., 2012). Thermal analysis methods, especially DSC and TG, have been used to feature the study, development and quality control of medications in general (Shamsipur, 2013).


This work applied these techniques to feature moisture contents in the study on the essential oil from *Siparuna guianensis*. It is a plant widely used in the Northern and Northeastern Brazilian regions due to its ethnobotanical properties such as antipyretic and decongestant and to its importance to healthcare. DSC and TG techniques are mainly applied to feature the molecules used in the development of bioproducts (Shen et al., 2010). Therefore, determining moisture content and testing the stability and decomposition kinetics of some valuable parameters regarding the thermal behavior of drugs and medication compounds, such as decomposition rates, possible mechanisms, and values of some thermokinetic and thermodynamic parameters, became necessary (Santos, 2004; Dantas, 2006).

Using DSC to determine essential oil moisture has some advantages in comparison to other methods, such as the relatively short analysis time and the use of small amounts of sample (Sandor, 2005). Thus, the DSC technique allows detecting a fraction in the total amount of a given substance, such as water molecules. It also allows quantifying the components from other volatile materials according to the dehydration and vaporization enthalpies (Khankari et al., 1992). Therefore, the current study aimed to determine the thermal featuring parameters and the number of water moles per mole in the *S. guianensis* essential oil.



Calculation methods

The hydrodistillation method used to extract essential oil is able to modify the water content in the samples. Terpenic and sesquiterpenic compounds easily oxidize under the influence of air, light and humidity due to their unsaturated character (Ganlim et al., 2002). Thus, it is necessary to accurately measure the water content in the tested samples before analyzing the pure essential oils.

It is done in order to obtain conclusive results. The current study suggests using an oil dehydration process similar to the hydrate desolvation performed by Khankari et al. (1992). In that case, the endothermic desolvation included steps that begin with the breaking up of solvate bonds up to solvent vaporization. The crystal structure reorganizes itself by breaking up and forming intermolecular forces after the desolvation. The structural rearrangement is part of the process (Chadha et al., 2012). According to the current study, the dehydration process is given by:



ΔH_{desol} corresponds to the enthalpy of desolvation (J / g hydrate) obtained from the area in DSC. The whole process can be divided into the following steps:



With

$$\Delta H_{desol} = \Delta H_{diss} + n\Delta H_f \quad 4$$

ΔH_{diss} is the water/oil dissociation heat. ΔH_f is the water fusion heat after its dissociation, and is similar to the vaporization hypothesis by Khankari et al. (1992), the current study considers ΔH_{diss} negligible.

$$\Delta H_{diss} \approx 0 \quad 5$$

which results in

$$\Delta H_{desolv} \approx n\Delta H_f \quad 6$$

The equation does not apply to the desolvation process in which the ΔH_{diss} dissociation heat is not negligible. The calculation performed to determine the number of water moles (n) per mole of anhydride is based on the premise that if one does not know the hydrate molecular weight value, it is necessary to determine the enthalpy change per mass unit (Δh_f) instead of per

*Corresponding author. E-mail: portella@uft.edu.br.

Abbreviations: DSC, Differential scanning calorimetry; KFT, Karl Fischer titration; TG, thermogravimetric analysis.

Author(s) agree that this article remains permanently open access under the terms of the [Creative Commons Attribution License 4.0 International License](https://creativecommons.org/licenses/by/4.0/)

mole (ΔH_f)

$$\Delta H_{desolv} = \Delta h_{desolv} \cdot (M_s + nM_w) \quad 7$$

$$\Delta H_f = \Delta h_f \cdot M_w \quad 8$$

Δh_f is the melting enthalpy of water (333 J/g, Stark and Wallace, 1976); M_s is the average molecular weight of the major components of *S. guianensis* essential oil identified by the CG-MS ($120.77 \pm 12 \text{ g mol}^{-1}$); and M_w is the molecular weight of water ($18.016 \text{ g mol}^{-1}$). Equation 9 was formed by grouping Equations 5, 6 and 7:

$$n = \frac{\Delta h_d M_s}{(\Delta h_l - \Delta h_d) M_w} \quad 9$$

The stoichiometric number, n , represents the number of water moles per oil mole.

MATERIALS AND METHODS

Oil extraction

The harvesting of *S. guianensis* Aublet leaves was performed in a Cerrado fragment in Formoso do Araguaia, in Northern Brazil, Latitude: $11^\circ 47' 48''$ S Longitude: $49^\circ 31' 44''$ W. The botanical material exsiccates were deposited at Universidade Federal do Tocantins - Campus Porto Nacional (register 10.298). The samples were ground, and their mass was analyzed ($25 \pm 5 \text{ g}$) and stored at 4°C for later use. Subsequently, they were placed in a volumetric flask containing 500 ml distilled water in each extraction. The adjusted methodology defined the ideal time to extract the essential oil in 80 min distillation (after the first sign of boiling). The essential oil was extracted according to the steam distillation method - using Clevenger type apparatus - and it was expressed based on its fresh mass (Sandor, 2005). All experiments were performed in triplicate and standard deviations were calculated.

Differential scanning calorimetry (DSC)

A 25.78 mg essential oil sample extracted from *S. guianensis* leaves was subjected to DSC analysis using Netzsch apparatus, model STA 449 F3 Jupiter, in N_2 atmosphere at the flow of 50 ml / min under temperature ranging from -30 to 40°C , with heating rate $\beta = 10^\circ\text{C}/\text{min}$. The sample mass (%) and temperature ($^\circ\text{C}$) were measured at 1.2 s intervals for 36 min.

Simultaneous thermal analysis (DSC-TG)

Essential oil samples of approximately 32.3 mg extracted from *S. guianensis* leaves were subjected to thermal analysis and the thermogravimetric curves (TG) were obtained using Netzsch apparatus, model STA 449 F3 Jupiter, in N_2 atmosphere, at the flow of 50 ml / min under temperature ranging from 39.15 to 400°C , with heating rate $\beta = 2, 3$ and $5^\circ\text{C}/\text{min}$. The sample mass (%) and temperature ($^\circ\text{C}$) were measured at 1.2 s intervals for 36 min.

Analysis in Karl Fischer titrator

Karl Fischer method

The water content of the essential oil was determined according to Karl Fischer method in order to compare the results; Equation (10):

$$\% \text{H}_2\text{O} = (100 \text{ E.V}) / m \quad 10$$

Where E = Water equivalent to the Karl Fischer reagent (g/mL); V = volume of Karl Fischer's solution spent in the oil sample titration (mL); and m = Sample weight (g).

The measurements were performed in a Mettler Toledo Karl Fischer titrator, DL31 model. The principle of this method is the titration of a conventional Karl Fischer reagent in an anhydrous methanol solvent in the presence of *S. guianensis* essential oil. The same procedure was carried out in triplicate. Regarding this titration test, the oil samples were measured on an analytical scale, using specific glazing. The values were expressed as % of humidity and converted into moles.

RESULTS AND DISCUSSION

Thermoanalytical featuring of the *S. guianensis* essential oil

Figure 1 shows the DSC curve. As it can be seen, the mass loss, due to water sublimation in *S. guianensis* essential oil, happened at the temperature interval from -3.1 to 5.9°C , with maximum peak at 1.9°C , thus totaling the area of 18.95 jg^{-1} . Many studies use these methodologies as alternatives in the quality control of pharmaceutical materials (Atkins, 2002; ASTM, 2003; Sandor, 2005).

Using DSC to determine the water content and other impurities in the *S. guianensis* essential oil

The non-linearity theory explains the determination of water content and other impurities by means of DSC. In other words, the area where the event normally starts is derived from processes such as the vaporization, oxidation or the decomposition of the oil itself as well as the impurities themselves. The endothermic transition of approximately 1.6°C may be evidence that the *S. guianensis* essential oil presents water of hydration in its composition. The stoichiometry of water molecules may be calculated at each stage by using the vaporization hypothesis. The results are listed in Table 1. As for the analysis in the Karl Fischer titrator, the essential oil was titrated in triplicate and presented mean moisture of 0.352 ± 0.005 water moles per anhydrous mole. The presence of moisture in oils may negatively influence the transesterification process (for the production of bioproducts) by disabling the basic catalysts and releasing water molecules, thus reducing yield (Simões et al., 2004). Table 2 shows the values obtained during the analyses of samples with mass almost identical to that of




Figure 1. DSC curves of *S. guianensis* essential oil obtained in N_2 atmosphere (50 mL min^{-1}) at $10 \text{ }^\circ\text{C min}^{-1}$.

Table 1. Water distribution between both endothermic dissolutions of *S. guianensis* essential oil.

Peak temperature ($^\circ\text{C}$)	ΔH of the desolvation endotherm (J/g)	Water in the energy state determined by using DSC in the vaporization hypothesis, Eq. 9 (moles/mole of anhydrous \pm EP)	Water moles from Karl Fischer titration (KFT \pm EP)
1.6	18.95	0.404 ± 0.04	
Total water moles		0.404 ± 0.04	0.352 ± 0.005

Table 2. Measurements and parameters used from DTA curves.

β ($^\circ\text{C min}^{-1}$)	T_M (K)	$1/T$ (K^{-1})	$\ln(\beta/T_M^2)$
2	367.64	0.002720	-11.1211
3	375.15	0.002666	-10.7560
5	390.47	0.002561	-10.3253

the *S. guianensis* essential oil, at three different heating rates (β): 2, 3 and 5°C min^{-1} (Figure 2). The values mentioned in the T_M column are, in this case, those read in the respective maximum peaks of DTA curves, as they were determined by the method developed by Kissinger (Kissinger, 1957). Figure 3 shows the values presented in graphic form, with the linear regression, the straight-line equation and the correlation coefficient R^2 . The straight-line equation (Equation 11) is

$$Y = -4885.5x + 2.207 \quad 11$$

The Activation Energy value E_a , may be calculated using parameters such as angular coefficient of 4885.5 and gas constant of $8.314 \text{ KJ mol}^{-1}$,

$$E_a = -(-4885.5 \times 8.314) / 1000 = 40.61 \text{ kJ mol}^{-1} \quad 12$$

The E_a calculated for the *S. guianensis* essential oil was $40.61 \text{ kJ mol}^{-1}$. From the linear coefficient (Equations 13-16):

$$2.207 = \ln(A/R/E_a) \quad 13$$

$$e^{2.207} = A \cdot 8.314 / 40.61 \times 10^3 \quad 14$$

$$A = e^{2.207} \times 40.61 \times 10^3 / 8.314 \quad 15$$

$$A = 4.44 \times 10^4 \text{ min}^{-1} \quad 16$$

Therefore, based on the data analysis by DTG, it is




Figure 2. DTG curves obtained from the analyses of 25.78 mg essential oil, at different heating rates.




Figure 3. Determination of kinetic parameters according to Kissinger's method, using DTA data.

possible to highlight that the Arrhenius Equation (Equation 17) for the overall conversion of the *S. guianensis* essential oil can be described by:

$$K(T) = A e^{\frac{-E_a}{RT}} \tag{17}$$

$$K = 4.44 \times 10^4 e^{\frac{-40.61KJ}{RT}} \tag{18}$$

Equation 17 adjusts itself to the experimental data at

temperature ranging from 35 to 400°C. It highlights the hypothesis that melting is the prevalent phenomenon, although there are parallel ones, since the activation energy was relatively low (Martins, 2008).

Kinetic study of *S. guianensis* decomposition

The stability (in days) of the *S. guianensis* essential oil was calculated based on Equation (18), using 35°C, (lower limit of the analyzed range). The estimated thermal

stability time of approximately 33 days or 1 month was obtained starting from the analysis date. Guimaraes et al. (2008) used other analysis techniques and found 50% myrcene concentration loss in 120 days. Myrcene is the major component found in *S. guianensis* essential oil.

Conclusion

The thermoanalytical techniques used in the current study provided the parameters of the *S. guianensis* essential oil. They are important to determine water content and other impurities during the quality control of this oil. The DSC found water content similar to that found by Karl Fischer technique. DSC was considered to be a faster and more reliable technique because the oil melting is not followed by material decomposition.

Conflict of interests

The author(s) did not declare any conflict of interest.

REFERENCES

- ASTM E 928-03, (2003). Standard Test Method for Purity by Differential Scanning Calorimetry", West Conshohocken, USA.
- Atkins PW (2002). Kurzlehrbuch Physikalische Chemie. 3rd ed. Weinheim, Germany: Wiley-VCH.
- Brown TL (2005). Química a Ciência Central. 9 ed. São Paulo: Pearson-Prentice Hall.
- Chadha R, Kuhad A, Arora P, Kishor S (2012). Characterisation and evaluation of pharmaceutical solvates of Atorvastatin calcium by thermoanalytical and spectroscopic studies. Chem. Cent. J. 6:114.
- Dantas MB (2006). Obtenção, Caracterização e Estudo Termoanalítico de Biodiesel de Milho (*Zea mays* L.). João Pessoa, UFPB, Dissertação de Mestrado. 114.
- Ganlim CD, Dutta NK, Roy CN, Kehoe D, Matisonss J (2002). Evaluation of Kinetic Parameters of Thermal Oxidative Decomposition of Base Oils by Conventional, Isothermal and Modulated TGA, and Pressure DSC. Thermochim. Acta 392: 357-369.
- Giron DJ (2002). Applications of thermal analysis and coupled techniques in pharmaceutical industry. J. Therm. Anal. Calorim. 68:335-57.
- Guimarães LG, Cardoso MDG, Zacaroni LM, de Lima RK, Pimentel FA, de Moraes AR (2008). Influência da luz e da temperatura sobre a oxidação do óleo essencial de capim-limão (*Cymbopogon citratus* (D.C.) Stapf). Quím. Nova [online]. 31(6):1476-1480.
- Khankari RK, Law D, Grant DJW (1992). Determination of water content in pharmaceutical hydrates by differential scanning calorimetry. Int. J. Pharm. 82:117-127.
- Kissinger HE (1957). Reaction Kinetics in Differential Thermal Analysis. Anal. Chem. 29(11):1702-1706.
- Martins JBL (2008). Equilíbrio e cinética química. Físico-Química Teórica. Publicação interna do Laboratório de Química Computacional. 18 f. Instituto de Química, Universidade de Brasília. Brasília, DF.
- Sandor G (2005). The sacred cow: the questionable role of assay methods in characterizing the quality of bulk pharmaceuticals. J. Pharm. Biomed. Anal. 5(36):931-937.
- Santos JCO (2004). Estudo Termoanalítico e Cinético da Degradação Térmica de Oleos Lubrificantes Automotivos. João Pessoa, Programa de Pós-graduação em Química, UFPB, Tese de Doutorado.
- Shen L, Worrel E, Patel M (2010). Biofuels. Bioproducts Biorefining 4:25-40.
- Simões CMO, Schenkel EP, Gosmann G, Mello JCP, Mentz LA, Petrovicck PR (2004). Farmacognosia: da planta ao medicamento, 5^a ed., Editora da UFSC: Porto Alegre.
- Souza SPMC, Moraes FE, Santos EV, Silva ML, Martinez-Huitle CA, Fernandes NS (2012). Determinação do teor de cálcio em comprimido à base de lactato de cálcio utilizado no tratamento da osteoporose. Quím. Nova 7(35):1355-1359.
- Stark JG, Wallace HG (1976). Chemistry Data Book, S1 Edn. John Murray, London. P 122.