



Development of a coupling-phase SCAR marker linked to the powdery mildew resistance gene 'er1' in pea (*Pisum sativum* L.)

Rakesh K. Srivastava, Shashi K. Mishra, Ashok K. Singh and Trilochan Mohapatra

DOI: <http://dx.doi.org/10.1007/s10681-012-0650-z>

This is author version post print archived in the official Institutional Repository of ICRISAT www.icrisat.org

Development of a coupling-phase SCAR marker linked to the powdery mildew resistance gene 'er1' in pea (*Pisum sativum* L.)

Rakesh K Srivastava*^{1,2}. Shashi K Mishra^{2,3} . Ashok K Singh² . Trilochan Mohapatra⁴

(1) **Rakesh K. Srivastava**

*Corresponding author

Present address:

Building 305, Global Theme- Biotechnology,

International Crops Research Institute for the Semi-Arid Tropics (ICRISAT)

Patancheru, Hyderabad- 502324, Andhra Pradesh, India

E-mail: r.k.srivastava@cgiar.org

Phone: +91 40 30713433

Telefax: +91 40 30713074

(2) Division of Genetics,

Indian Agricultural Research Institute (IARI),

Pusa Campus, New Delhi- 110012, India.

(3) Last address (late):

Division of Germplasm Evaluation,

National Bureau of Plant Genetic Resources (NBPGR)

Pusa Campus, New Delhi- 110012, India.

- (4) National Research Centre on Plant Biotechnology (NRCPB),
Indian Agricultural Research Institute (IARI),
Pusa Campus, New Delhi- 110012, India.
E-mail: tm@nrcpb.org

Abstract

Pea powdery mildew is one of the major constraints in pea production worldwide, causing severe seed yield and quality loss. The resistance is governed by a single recessive gene *er1* in majority of resistant cultivars, but *er2* and *Er3* have also been reported. The objective of the study was to find out tightly linked SCAR markers to *er1* gene using NILs. A total of 620 RAPD markers were screened for length polymorphism between seven sets of NILs. The 880 bp polymorphic band of the tightly linked RAPD marker OPX 04₈₈₀ was cloned, sequenced and a SCAR marker ScOPX 04₈₈₀ was developed. In a population of completely classified 208 F₂ plants (supported by phenotypic data from 208 F_{2:3} and 4,390 F_{3:4} families) ScOPX 04₈₈₀ was linked at 0.6 cM in coupling phase with *er1* gene in the order ScOPX 04₈₈₀-*er1*-ScOPD 10₆₅₀. ScOPX 04₈₈₀ will correctly differentiate homozygous resistant plants from the susceptible accessions with more than 99% accuracy. In combination with repulsion phase marker ScOPD 10₆₅₀, ScOPX 04₈₈₀ can help in an error free marker-assisted selection (MAS).

Introduction

The pea powdery mildew is caused by an obligate ecto-parasitic biotrophic fungus *Erysiphe pisi* var. *pisi* D.C. (Braun 1980; Cook and Fox 1992), and has been known as one of the major constraints in pea production globally. Dixon (1978) identified powdery mildew as the greatest threat to the crop of dry pea. This disease occurs worldwide (Kraft and Pflieger 2001; Dixon 1978), and in India it is the most serious disease of pea. Powdery mildew alone can reduce 25-50% of seed yield (Kumar and Singh 1981; Reiling 1984) and also the seed quality of pea (Rathi and Tripathi 1994). Dry weather conditions favour the disease development, but do not assure an epidemic. In India, due to severe economic consequence of powdery mildew attack, All India Coordinated Pulse Improvement Project (AICPIP) made a decision in 1982 that no powdery mildew susceptible line would be tested in the coordinated trials (Sharma 1997).

Hammarlund (1925) reported four recessive genes for powdery mildew of field pea, but it was not confirmed by further studies. Later, Harland (1948) isolated some resistant plants from local Peruvian material. His F₂ and F₃ data perfectly fitted to 3:1 segregation ratio, susceptible being dominant over resistance. He designated this gene as *er*. The resistance was controlled by a single gene in homozygous recessive condition. This was later supported by Pierce (1948), Yarnell (1962), Cousin (1965), Marx (1971), Saxena et al. (1975), Singh et al. (1983), Mishra and Shukla (1984), Gupta (1987), Sarala (1993), Timmerman et al. (1994), Rakshit (1997), Vaid and Tyagi (1997), Janila and Sharma (2004), and Srivastava and Mishra (2004). However, the inheritance of powdery mildew has not been totally unambiguous. Several workers have reported digenic segregation (Heringa et al. 1969; Sokhi et al. 1979;

Kumar and Singh 1981; and Tiwari et al. 1997). In these studies genes governing resistance to *E. pisi* were reported to be controlled by recessive genes *er1* and *er2*. However, *er2* has not been confirmed in the present sets of NILs during our earlier studies. There has been limited mention of this gene in linkage studies. Katoch et al (2010) recently reported that they have mapped *er2* to LGIII. Overwhelming evidence favors one-gene (*er1*) control of powdery mildew resistance in pea (Sharma 2003; Sharma and Yadav 2003, Srivastava and Mishra 2004). This may be partly due to the nature of resistance governed by *er2*. While *er1* offers complete resistance against the powdery mildew pathogen by preventing pathogen penetration, *er2* mediated resistance is primarily based on reduced penetration rate complimented by post-penetration hypersensitive response leading to cell death (Fondevilla et al. 2006; Katoch et al. 2010).

Development of molecular markers for *er1* gene was thought to be essential for pyramiding it with *er2* and *Er3* in single cultivar, or in a backcrossing program to convert a susceptible variety into resistant one. This is a durable gene and since the last five decades there has been no report of the resistance break-down conferred by *er1*. Scores of attempts were made in the past to find molecular markers linked to the *er1* gene. Many workers tried to develop random amplified polymorphic DNA (RAPD) and amplified fragment length polymorphism (AFLP) derived sequence characterized amplified regions (SCAR) markers. Timmerman et al. (1994) developed a RAPD-derived SCAR marker ScOPD 10₆₅₀ in Canadian germplasm. This marker was reported to be linked at 2.1 cM in repulsion phase. Subsequent validation studies in two different mapping populations placed ScOPD 10₆₅₀ at 3.6 cM (Rakshit, 1997) and 3.4 cM (Janila and Sharma 2004) from *er1*. Both reported distances are large for efficient marker-assisted selection (MAS). The other two closely linked SCAR markers, ScOPO 18₁₂₀₀ and ScOPE 16₁₆₀₀ reported by Tiwari et al. (1998) were not found to be polymorphic by Janila and Sharma (2004), and did not differentiate resistant and susceptible lines of Indian origin. Several new marker systems have been reported to be linked to various powdery mildew resistance genes in pea as *er1* (Ek et al. 2005; Pereira et al. 2010), *er2* (Katoch et al. 2010) and *Er3* (Fondevilla et al. 2008).

Herein we describe our experiment that led to development of a tightly linked SCAR marker to the powdery mildew resistance gene (*er1*) in pea, using advanced generation (BC₉) near-isogenic lines (NILs).

Material and Methods

Plant materials

NILs, developed at the Division of Genetics, Indian Agricultural Research Institute (IARI), New Delhi, India, through nine cycles of backcrossing with their respective recurrent parents, were used in the present study (Table 1). The parents for generation of mapping population were selected based on their pedigree and sources of resistance (PMR)/susceptibility (PMS) to powdery mildew.

Crosses and Phenotypic Scoring:

A set of seven crosses were attempted during winter of 2000, at I.A.R.I, New Delhi involving NILs viz., HFP 4 (PMS)^{Bonneville#} × HFP 4 (PMR); HFP 4 (PMS)^{PG 3} × HFP 4 (PMR); PG 3 (PMR)^{HFP 4} × PG 3 (PMS); DMR 11 (PMS)^{KPSD 1} × DMR 11 (PMR); T 163 (PMR)^{DMR 11} × T 163 (PMS); L 116 (PMR)^{DMR 11} × L 116 (PMS) and Pusa 10 (PMR)^{DMR 11} × Pusa 10 (PMS). About 50-60 buds per cross were emasculated to get sufficient number of F₁ seeds. It was assumed that there are no reciprocal differences for powdery mildew resistance (Singh 1984). The PMR (*er1er1*) line was chosen as female parent in all the crosses. The crossed as well as selfed seeds from both parents were harvested. Simultaneously, young unopened leaves and growing stipules were collected from each parent of all the seven crosses for parental polymorphism study.

Disease phenotyping and zygosity testing for *er1* locus in the mapping population

Plots of five meter row length with 65 cm row spacing and 30 cm spacing between plans were planted in the field. To get error free disease reaction for the powdery mildew, all the seven F₂ populations were

#Superscript indicates the donor parent for powdery mildew susceptible (PMS) or resistant (PMR) trait.

sown in late December, 2001. This facilitated natural epidemic of the disease at IARI, New Delhi. A single row or five row plot of susceptible variety L116 was planted every fourth row and on the boarder (five rows) of each population. Dusting of the infected plant material from L116 was also done on alternate days from 30 days after sowing till maturity for uniform spread of the inoculum. The infected foliage surface was totally covered with a white powdery mass of the fungus and the infection spread to all aerial parts of the plant, including stems and pods. Tissue beneath the infected areas in the susceptible plants turned brown, followed by the production of black fruiting bodies (cleistothecia). The resistant plants were free from infection or infection was localized in very small patches only on the foliage (stipules and leaves), but it never spread to the stem, peduncle and pods. The plants with stem, peduncle and pods free from the infection symptoms were categorized as resistant, while plants with infection on

leaves, stem, stipules and pods were categorized as susceptible. Disease phenotyping was conducted according to Janila and Sharma (2004), and Srivastava and Mishra (2004). There were clear cut differences between resistant and susceptible plants with no ambiguity, based on the scoring criteria mentioned above.

The entire F_2 population of the cross PG 3 (PMR)^{HFP 4} × PG 3 (PMS) was advanced to F_4 generation through F_3 by harvesting single plants separately. Unequivocal phenotyping was obtained by screening of the F_3 and F_4 families. This testing dissected the heterozygosity/homogygosity of the susceptible plants at *er1* locus and also ruled out disease escape of any susceptible plants. The F_3 families were screened at the hot-spot location for powdery mildew during 2003 at off-season nursery Dalang Maidan, Lahaul Spithi, Himachal Pradesh State, India. All the 208 F_2 plants were represented as 208 F_3 families. The population was grown in two-row plots of five m length comprising at least 20 plants per family along with resistant (HFP 4) and susceptible (PG 3) parents at every fifth row. A total of 4,390 plants in the F_3 population were screened. Seeds were harvested from each of the F_3 families to raise F_4 families that were screened again at I.A.R.I New Delhi during winter 2003/04. Presence of significant number of resistant plants in a particular F_3/F_4 family was taken to infer heterozygosity at *er1* locus for the corresponding F_2 plant.

DNA isolation and quantification

The DNA was extracted from individual 208 F_2 plants, from unopened leaves and young stipules using the cetyl methylammonium bromide (CTAB) protocol (Doyle and Doyle 1990) with few modifications as use of 1% PVP and 4% mercaptoethanol in the extraction process. DNA was also isolated from the PMR (15 plants) and PMS (15 plants) pairs of NILs, and was bulked together as resistant and susceptible bulks, respectively. Quantification was performed by running DNA samples on 0.8% agarose gels along with known quantity of uncut λ DNA. Quantification was also made based on UV absorbance at 260 nm using a spectrophotometer. The ratio of absorbance at 260 to 280 nm gave the purity of extracted DNA. The samples were considered pure, if the ratio was between 1.8 and 2.0.

PCR analysis

We used RAPD marker system for SCAR marker development. A total of 620 random, 10-mer RAPD primers procured from Operon Technologies, Inc. (Alameda, CA, USA) were used for establishing

polymorphism among seven pairs of NILs. These primers included OPA 01-20, OPB 01-20 to OPZ 01-20, OPAA 01-20 to OPAC 01-20, OPAE 01-20 and OPAK 1-20. RAPD analysis was done as conducted in Williams et al. (1990). The 25 µl reaction mixture contained 200 µM of dNTP mix (dATP, dTTP, dGTP, dCTP, all in precisely equal molar concentration), 0.2 µM of a given 10-mer primer, 0.5 units of Taq polymerase assay buffer (Genetix) and 25 ng of genomic DNA in 1X Taq polymerase assay buffer (Genetix). The reaction was carried out in Perkin Elmer 9600 thermal cycler (Barnstead Thermolyne Corp., Iowa, USA) programmed for one cycle at 92 °C for 4 minutes for initial template denaturation, followed by 44 cycles at 92 °C (denaturation) for 30 seconds, 37 °C (annealing) for 30 seconds and 72 °C (template extension) for 1 minute, and one cycle at 72 °C for 7 minutes for final template extension. The amplified fragments of DNA were resolved by electrophoresis on 1.4% agarose gel stained with ethidium bromide, and photographed under transmitted UV light (302 nm) using gel documentation system. The primers exhibiting polymorphism between near-isogenic parents were repeated four times to select only reproducible primers. Such reproducible primers were used to screen the F₂ population and analyzed for their segregation patterns.

In addition to RAPDs, previously reported SCAR markers were also used to screen parents. Three SCAR primers viz., ScOPD 10₆₅₀, ScOPO 18₁₂₀₀ and ScOPE 16₁₆₀₀ were synthesized. The primer ScOPD 10₆₅₀ was reported to amplify a 650-bp band in the resistant cultivars (Timmerman et al. 1994). The primers ScOPO 18₁₂₀₀ and ScOPE 16₁₆₀₀ were reported to amplify 1200 and 1600 bp bands in susceptible and resistant parents, respectively (Tiwari et al. 1998). All three SCAR markers were studied for polymorphism. PCR reactions were performed as discussed above for RAPD analysis with suitable modifications in the annealing temperatures.

Linkage map Construction

Polymorphic markers identified by NIL marker analysis were used to screen the entire 208 F₂ mapping population of the PG 3 (PMR)^{HFP 4} × PG 3 (PMS) cross. A linkage map using the candidate markers and the already reported polymorphic marker was constructed using computer programme MAPMAKER/EXP Ver. 3.0 (Lander et al. 1987), using Kosambi's mapping function (Kosambi 1944).

SCAR Marker Development

The parent PG 3 was used to amplify the most closely linked RAPD band of 880 bp using the primer OPX 04₈₈₀. The band of interest was precisely demarcated by giving a brief exposure to long wave UV. The band was then sliced out using a sterile scalpel. The eluted band was purified using Millipore filter

kit obtained from M/S Millipore Inc., USA, following the manufacturer's instructions. The purified band was quantified on 0.8% agarose gel. It was then used as a template for its re-amplification using the same primer. TA cloning strategy was employed using pGEM[®]-T Easy Vector system. The standard reaction consisted of 2X rapid ligation buffer- 5µl, pGEM[®]-T Easy Vector -1µl (50ng); PCR product:1-3 µl; T₄ DNA ligase- 5 µl; deionized water to a final volume of 10 µl. The insert to vector molar ratio was kept between 3:1 and 8.2:1. The reaction mixture was mixed by pipetting, and incubated at 4°C overnight. Transformation was carried out using high efficiency competent cells (DH-5α strain of *Escherichia coli*) following the protocol described by Sambrook and Russell (2001). The recombinant plasmid DNA from *E. coli* was isolated using Qiagen's QIAprep[®] Miniprep kit following the manufacturer's directions. About 500 ng (1 µl) of plasmid was treated with 6 units of *Eco*RI enzyme and digested for 1 hour at 37°C. The insert size was then checked on 1.4% agarose gel, by comparing with DNA molecular weight marker. Sequencing of the recombinant plasmids was done by taking one representative sample from each of the four classes of clones. Sequencing was performed on megaBACE-1000 sequencing machine, using M13 universal primers, at the Rice Genome Sequencing Project, I.A.R.I, New Delhi.

Designing of the SCAR primers

The sequence data generated from all four classes of clones were subjected to BLAST search (Basic Local Alignment Search Tool; <http://www.ncbi.nih.gov/BLAST/>). This programme was used for alignment of unique sequence and for retrieving vector and the 10-mer RAPD primer sequences. The primers were designed using the software "Primer 3" available at the URL http://frodo.wi.mit.edu/cgi-bin/primer3/primer3_www.cgi and "Web Primer" from the URL <http://genome-www2.stanford.edu/cgi-bin/SGD/web-primer> for construction of suitable sets of primers. The primers were synthesized by M/S Qiagen Operon, Cologne, Germany. Each primer was supplied as a lyophilized salt with quantity ranging between 450-570 µg (13-19 OD). The primers were re-suspended in 1 ml of sterile T₁₀E₁ buffer at 4°C for 48 hours aliquoted in different tubes and stored at -20°C.

Standardization of SCAR reaction

Based on the average T_m of the forward and reverse primers of each set, an annealing temperature gradient was set up. The annealing temperature gradient ranged between -5 and +6 of the respective average T_m for different sets of primers. The PCR reaction conditions in 25 µl of reaction mixture were maintained as follows: 200 µmoles of dNTPmix (dATP, dTTP, dGTP and dCTP in exactly equal concentration) (Promega); 25 µg of each primer; 1 unit Taq polymerase (Promega) and 25ng of genomic

DNA in 1X Taq polymerase assay buffer (Promega). Amplification was carried out in 35 cycles of 3 minutes of initial denaturation at 92°C, 1 minute for primer annealing at 58-67°C for different sets of SCAR primers, 2 minutes of template extension at 72°C, followed by 34 cycles of 1 minute of denaturation at 92°C; 1 minute of annealing at 58-67°C depending upon SCAR primer, 2 minutes at 72°C. One additional profile of 72°C for 7 minutes was given for final extension. The reactions were carried out in Biometra and Perkin Elmer 9600 thermal-cyclers. The amplified products were run on 1.2% agarose gel and photographed in Gel-documentation system.

Results

Screening for polymorphism among parents

All seven sets of NILs were screened pair-wise to detect polymorphism for RAPD markers. Out of 620 random 10-mer RAPD primers used (OPA 01-20, OPB 01-20 to OPZ 01-20, OPAA 01-20 to OPAC 01-20, OPAE 01-20 and OPAK 1-20), nine polymorphic primers were identified between resistant and susceptible NILs. However, based on stringent reproducibility conditions (four times), three primers retained polymorphism, out of which only OPX 04 produced a bright polymorphic band (880bp) and retained polymorphism in the NIL set PG 3 (PMR)^{HFP4} × PG 3 (PMS). The RAPD primer OPX 04 (5'-CCGCTACCGA-3') amplified a fragment of about 880 bp in the susceptible parent PG 3 (Figure 1). The other two polymorphic primers did not produce bright and thick polymorphic band as OPX 04.

Earlier reported *erI*-linked molecular markers *viz.*, ScOPE 16₁₆₀₀, OPL 06₁₉₀₀ and ScOPO 18₁₂₀₀ by Tiwari et al. (1998); and OPU 17₁₀₀₀ and OPU 02₁₁₀₀ by Janila and Sharma (2004) did not reveal polymorphism between the parents since the plant materials used in these studies were different. Out of seven sets of NILs used to study polymorphism, only one set namely PG 3 (PMR)^{HFP4} × PG 3 (PMS) was simultaneously differentiated by all the three polymorphic primers. The other NIL sets were either not differentiated by all the three polymorphic primers, or differentiated only by OPX 04.

SCAR marker development

Initial analysis revealed that the RAPD marker OPX 04₈₈₀ and the SCAR marker ScOPD 10₆₅₀ belonged to same linkage group in relation to *erI* gene. Hence OPX 04₈₈₀ was converted to a SCAR marker. The polymorphic linked band of 880 bp, generated by the RAPD primer OPX 04 was amplified from susceptible parent PG 3, gel extracted and cloned. A total of 29 positive clones were picked up for insert check and subjected to *EcoRI* digestion. Eight clones showed insert of the right size *i.e.*, 880 bp. Within such clones, four different types of banding pattern were observed (Figure 2). One representative sample

from each of the four types was sequenced for primer designing and sequence homology search. Each of the four classes of clones gave four different types of sequence information. BLAST search did not yield any homology with the deposited sequences in the gene bank. Also none of the sequences shared any distinct types of repeat sequences.

Designing, synthesis and amplification check of the primers

Four Primer sets were designed from the sequence information of all the four classes of clones, using “Primer 3” and “Web Primer” software. The sense and antisense sequence details of all the four primer sets are given in Table 2. Out of the four primer sets, two (Class I & II) sets failed to distinguish between the two parents and amplified a band of 880 bp in both the parents, while Class IV primer set did not yield any amplification product. One primer set designated as ScOPX 04₈₈₀ obtained from the Class III clone, correctly distinguished between resistant and susceptible parent (annealing temperature of 65.5°C). This 21 mer sense (5'-CCGCTACCGATGTTATGTTTG-3') and 20 mer antisense (5'-CCGCTACCGAACTGGTTGGA-3') SCAR primer amplified a band of 880 bp (similar to its polymorphic RAPD counterpart OPX 04₈₈₀) in the susceptible parent PG 3, while no amplification was observed in the resistant isogenic parent PG 3 (PMR)^{HFP 4}. This marker, thus, behaved as a dominant coupling phase SCAR marker.

Determination of the linkage relationship of the new SCAR marker

The SCAR marker ScOPX 04₈₈₀ segregated in precisely the same fashion as the 880 bp band of the RAPD primer OPX 04. It exhibited only one recombination (plant no. 9) with *erl* in the population of 208 F₂ plants (Figure 3). MAPMAKER analysis placed ScOPX 04₈₈₀ linked to *erl* gene at 0.6 cM, in coupling phase, while ScOPD 10₆₅₀ was placed at 2.2 cM from *erl* and 2.8 cM from ScOPX 04₈₈₀ flanking the *erl* gene in repulsion phase (Figure 4). The segregation pattern for these three molecular markers, and also the *erl* gene was in agreement with 3:1 ratio with non-significant χ^2 value with P ranging between 0.88 and 1.00 (Table 3). The presence of the amplified DNA band was taken as dominant over its absence as all these markers amplified their respective bands in the F₁ hybrids. As mentioned earlier, powdery mildew resistance was controlled by monogenic recessive gene in several studies. In the present investigation also, the F₂ mapping population always segregated in a good fit ratio of 3 PMS : 1 PMR in all the crosses involving NILs.

Discussion

RAPD markers have been successfully used in the recent past to develop more reliable sequence tagged SCAR markers in crops such as oilseed rape (*Brassica napus* L.) (Mikolajczyk et al. 2008; Saal and Struss 2005), grapes (*Vitis vinifera* L.) (Kim et al. 2008; Akkurt et al. 2007), *Amaranthus* (Ray and Chandra 2009), safflower (*Carthamus tinctorius* L.) (Hamdan et al. 2008), papaya (*Carica papaya* L.) (Giovanni and Víctor 2007), *Pandanus fascicularis* (Vinod et al. 2007), faba bean (*Vicia faba* L.) (Gutierrez et al. 2007), common bean (*Phaseolus vulgaris* L.) (Park et al. 2008), sorghum [*Sorghum bicolor* (L.) Moench] (Singh et al. 2006) and wheat (*Triticum aestivum* L.) (Gupta et al. 2006). Recently Jones et al. (2009) have re-emphasized role of SCAR markers in tracking economically important genes.

This study elucidates successful use of advance generation NILs for fine gene tagging in plants. This has also been emphasized by several other workers (Martin et al. 1991; Young et al. 1988), as a more efficient tool in tagging genes with scorable phenotype than bulked segregant analysis (BSA) proposed by Michelmore et al. (1991). Bi-parental mapping populations generated by crossing NILs are more precise in mapping the target gene because it allows recombination to take place in the regions close to the gene of interest, since the rest of the genome would be homogenous. Although polymorphism between pairs of NILs may be low, but a marker, if found will have high probability to be closely linked to the gene of interest.

In the present study, seven pairs of NILs were used to screen for polymorphism. On the basis of highest number of polymorphic markers, one NIL pair was selected for development and use of mapping population for identification of molecular markers linked to the *er1* gene. This NIL pair was developed by crossing HFP 4, carrying *er1* allele in homozygous form, to the susceptible recurrent parent PG 3. In the ninth generation of back-crossing it was expected that *Er1/Er1* locus was replaced with *er/er*, while the rest of the genome was the same (>99.9%) as that of recurrent parent PG 3. The converted NIL was resistant to the powdery mildew (as HFP 4) and was designated as PG 3 (PMR)^{HFP 4}.

SCAR marker development

Sequencing of multiple clones of the 880bp informative amplicon obtained with the random primer OPX 04 revealed presence of four different types of clones. No sequence homology between Class I and IV clones (Figure 2) might be due to amplification of non-target regions of the genome by the RAPD primer or co-migration. Such random 10-mer sequences have a probability to find unlinked regions of the genome once every 1.048 Mb. It was also unusual not to find any repetitive elements in these sequences.

This may be due to small sequence length (sampling error) and/or due to sequences from the functional region of the genome, known to carry very little repetitive elements.

Lack of polymorphism for Class I, II and IV SCAR primers was on expected lines. Although RAPD primer sequence was present in Class I and II primers, there was no sequence homology to the *Er1* locus, and may represent sequence complementarity with regions unlinked to the *Er1* locus. Class IV SCAR primer set was a clear case of co-migration due to non-complimentary amplification as no RAPD primer sequence could be located. This is a common phenomenon observed with RAPD primers due to their low annealing temperature (37° C in the presents study) and short primer length.

On the other hand, polymorphism and co-segregation with the powdery mildew disease for Class III SCAR primer sets derived from Class III clones may be due to presence of sequence homology to the *Er1* locus.

Mapping of the *er1* gene with SCAR markers

Comparable segregation pattern and band sizes of SCAR and RAPD markers may be due to inclusion of RAPD primer sequence at the 5' ends of the sense and antisense primers. It is worth mentioning that ScOPX 04₈₈₀ appeared at a distance of zero cM from *er1* gene in multipoint analysis, whereas pair wise analysis (two point cross) placed it at 0.6 cM. In multipoint analysis, the lone recombinant of ScOPX 04₈₈₀ was taken as an error (with “error detection on” command in MAPMAKER) and, therefore, was shown at zero cM distance (data not presented). Hence, the map distance of 0.6 cM obtained from pair-wise analysis appears to be more realistic (Table 4). The composite linkage map is presented in Figure 4. In this map ScOPD10₆₅₀ maps at 2.2 cM, very close to the 2.1 cM map distance reported by Timmerman et al. (1994), but is different from Rakshit (1997) and Janila and Sharma (2004) who reported ScOPD10₆₅₀ at distance of 3.7 and 3.4 cM, respectively, from *er1*. Such minor discrepancies in map distance may arise due to differences in genetic architecture. In fact markers mapped in a particular cross rarely give the same map distance in other crosses. In present investigation SCAR markers ScOPE 16₁₆₀₀ and ScOPO 18₁₂₀₀ reported by Tiwari et al. (1998), and RAPD markers OPU 17₁₀₀₀ and OPU 02₁₁₀₀ reported by Janila and Sharma (2004) did not reveal polymorphism. This may be attributed to differences in the primer binding sites due to evolution of the lines at different geographical locations under varying kinds of selection pressure. The implications of such results may warrant development of molecular markers which are anchored to the polymorphic regions of the genome which remain conserved by

pedigree and origin. The SCAR marker developed in the present study need to be verified in the other materials with different genetic background for its suitability in gene pyramiding and MAS.

ScOPX 04₈₈₀ as a dominant coupling phase marker can precisely identify *erlerl* plants with more than 99% precision. On the other hand, ScOPD 10₆₅₀ would amplify in susceptible carriers as well, thereby limiting its use in germplasm screening. Therefore, the major limitation of a less tightly linked repulsion phase SCAR marker ScOPD 10₆₅₀ can easily be overcome by the use of new tightly linked SCAR marker ScOPX 04₈₈₀ identified in the present investigation. Extremely reliable marker-assisted selection (MAS) can be carried out by combining the two flanking SCAR markers ScOPX 04₈₈₀ and ScOPD 10₆₅₀ for BC_nF₁ and germplasm screening with almost 100% accuracy. Presence of two bands (of 880 bp and 650 bp size) will indicate heterozygosity at *ErI* locus, while only one band of 880 bp will indicate homozygous resistant plants.

RAPD-derived SCARs can circumvent the problems associated with the SCARs generated from AFLP bands that typically range between 50-400 bp. Such short sequences in many cases do not reveal polymorphism upon conversion to SCAR, and may need inverse PCR (I-PCR) or PCR walking strategy to generate polymorphism (Negi et al. 2000). It is easier to convert RAPD markers to SCAR markers as the former generate fragments in the size range of 500-1,500 bp (Barret et al. 1998), obviating the need for I-PCR or PCR walking. The same has been demonstrated successfully in the present study. The SCAR marker linked to the *erl* gene developed in the present study, and the other reported marker systems could be used for pyramiding genes for a more broad based durable resistance against powdery mildew pathogen.

We need to better understand *erl* gene, which is thought to evolve from loss-of-function mutation of a plant-specific Mildew Resistance Locus O1 (*PsMLO1*) (Humphry et al. 2011) under natural selection pressure. We also need to understand how a recessive gene coding for a defective/malfunctioning/nonfunctional gene product, initiates a cascade of metabolic pathways that result in a near-perfect protection of the pea from *E. pisi* globally.

Acknowledgements

The authors thank Dr. N.K. Singh, National Research Centre on Plant Biotechnology (NRCPB), New Delhi, India for his help and valuable suggestions in the course of the study. Rakesh K. Srivastava

gratefully acknowledges the SRF grant from the Council of Scientific and Industrial Research (CSIR), India; and I.A.R.I fellowship from the Indian Agricultural Research Institute, India.

References

- Akkurt M, Welter L, Maul E, Töpfer R, Zyprian E (2007) Development of SCAR markers linked to powdery mildew (*Uncinula necator*) resistance in grapevine (*Vitis vinifera* L. and *Vitis* sp.) Mol Breed 19: 103–111
- Barret P, Delourme R, Foisset N, Renard M (1998) Development of a SCAR (sequence characterized amplified region) marker for molecular tagging of the dwarf BREIZH (*Bzh*) gene in *Brassica napus* L. Theor Appl Genet 97: 828–833
- Braun V (1980) A monograph of *Erysiphales* (powdery mildews). Nova Hedwigia (Suppl.) no. 89. Stuttgart, Germany
- Cook RTA, Fox, RTU (1992) *Erysiphe pisi* var. *pisi* of faba bean and other legumes in Britain. Plant Pathol 41: 506–512
- Cousin R (1965) Resistance to powdery mildew in pea. Ann. Amélior. Plantes 15: 93–97
- Dixon GR (1978) Powdery mildew of vegetable and allied crops. In: DM Spencer (ed) The Powdery Mildews. Academic Press, London, pp 475–524
- Doyle JJ, Doyle JL (1990) Isolation of plant DNA from fresh tissue. Focus 12: 13–15
- Ek M, Eklund M, von Post R, Dayteg C, Henriksson T, Weibull P, Ceplitis A, Isaac P, Tuveesson S (2005) Microsatellite markers for powdery mildew resistance in pea (*Pisum sativum* L.). Hereditas 142: 86–89. doi:10.1111/j.1601-5223.2005.01906.x
- Fondevilla S, Moreno MT, Carver TLW, Rubiales D (2006) Macroscopic and histological characterisation of genes *er1* and *er2* for powdery mildew resistance in pea. Eur J Plant Pathol 115: 309–321. doi:10.1007/s10658-006-9015-6
- Fondevilla S, Rubiales D, Moreno MT, Torres AM (2008) Identification and validation of RAPD and SCAR markers linked to the gene *Er3* conferring resistance to *Erysiphe pisi* DC in pea. Mol Breed 22: 93–200. doi:10.1007/s11032-008-9166-6
- Giovanni Chaves-Bedoya, Víctor N (2007) SCAR marker for the sex types determination in Colombian genotypes of *Carica papaya*. Euphytica 153: 215–220
- Gupta MD (1987) Inheritance of powdery mildew resistance in pea (*Pisum sativum* L.). Ph.D. Thesis. Indian Agricultural Research Institute, New Delhi, India


- Gupta S, Charpe A, Koul S Haque QMR, Prabhu KV (2006) Development and Validation of SCAR Markers Co-Segregating with an *Agropyron Elongatum* Derived Leaf Rust Resistance Gene *Lr24* in Wheat. *Euphytica* 150: 233–240
- Gutierrez N, Avila CM, Rodriguez-Suarez C, Moreno1 MT, Torres1 AM (2007) Development of SCAR markers linked to a gene controlling absence of tannins in faba bean. *Mol Breed* 19: 305–314
- Hamdan YAS, Velasco L, Pérez-Vich B (2008) Development of SCAR markers linked to male sterility and very high linoleic acid content in safflower. *Mol Breed* 22: 385–393
- Hammarlund C (1925) Zur Genetik, Biologic und Physiologic einiger *Erysiphaeecen*. *Hereditas* 6: 1–126
- Harland SC (1948) Inheritance of immunity to powdery mildew in Peruvian forms of *Pisum sativum*. *Heredity* 2:263–269
- Heringa RJ, Norel A van, Tazelaar MF (1969) Resistance to powdery mildew (*E. Polygoni* D.C.) in peas (*Pisum sativum* L.). *Euphytica* 18: 163–169
- Humphry M, Reinstädler A, Ivanov S, Bisseling T, Panstruga R (2011) Durable broad-spectrum powdery mildew resistance in pea *er1* plants is conferred by natural loss-of-function mutations in *PsMLO1*. *Molecular Plant Pathology*. doi: 10.1111/j.1364-3703.2011.00718.x
- Janila P, Sharma B (2004) RAPD and SCAR markers for powdery mildew resistance gene *er* in pea. *Plant Breed* 123: 271–274 doi:10.1111/j.1439-0523.2004.00963.x
- Jones N, Ougham H, Thomas H, Pašakinskienė I (2009) Markers and mapping revisited: finding your gene. *New Phytologist*. DOI 10.1111/j.1469-8137.2009.02933.x
- Katoch V, Sharma S, Pathania S, Banayal DK, Sharma SK, Rathour R (2010) Molecular Mapping of pea powdery mildew resistance gene *er2* to pea linkage group III. *Mol Breed* 25:229–237
- Kim GH, Yun HK, Choi C, Park JH, Jung YJ, Park KS, Dane F, Kang KK (2008) Identification of AFLP and RAPD markers linked to anthracnose resistance in grapes and their conversion to SCAR markers. *Plant Breed* 127: 418–423
- Kosambi DD (1944) The estimation of map units from recombination values. *Ann Eugen Lond* 12: 172–175
- Kraft JM, Pflieger FL (2001) *Compendium of pea diseases and pests*. 2nd ed. APS Press, Saint Paul, MN
- Kumar H, Singh RB (1981) Genetic analysis of adult plant resistance to powdery mildew in pea (*Pisum sativum* L.). *Euphytica* 30: 147–151. doi:10.1007/BF00033671
- Lander E, Green P, Abrahamson J, Barlow A, Daly MJ, Lincoln SE, Newberg LA (1987) MAPMAKER: An interactive computer package for constructing primary genetic linkage maps of experimental and natural populations. *Genomics* 1: 174–181

- Martin GB, Williams JGK, Tanksley SD (1991) Rapid identification of markers linked to *Pseudomonas* resistance gene in tomato by using random primers and near-isogenic lines. *Proc Natl Acad Sci USA* 88: 2336–2340
- Marx GA (1971) New linkage relations for chromosome III of *Pisum sativum*. *Pisum Newsletter* 3: 18–19
- Michelmore RW, Paran I, Kesseli RV (1991) Identification of markers linked to disease-resistance genes by bulked segregant analysis: A rapid method to detect markers in specific genomic regions by using segregating populations. *Proc. Natl. Acad. Sci. USA* 88:9828–9832
- Mikolajczyk K, Dabert M, Nowakowska J, Podkowinski J, Poplawska W, Bartkowiak-Broda I (2008) Conversion of the RAPD OPC02₁₁₅₀ marker of the *Rfo* restorer gene into a SCAR marker for rapid selection of oilseed rape. *Plant Breed* 127: 647–649
- Mishra SP, Shukla P (1984) Inheritance of powdery mildew resistance in pea. *Z. Pflanzenzuchtg*, 93: 251–254
- Negi MS, Devic M, Delseny M, Lakshmikumaran M (2000) Identification of AFLP fragments linked to seed coat colour in *Brassica juncea* and conversion to SCAR marker for rapid selection. *Theor Appl Genet* 101: 146–152
- Park SO, Steadman JR, Coyne DP, Crosby KM (2008) Development of a Coupling-Phase SCAR marker linked to the *Ur-7* Rust Resistance Gene and Its Occurrence in Diverse Common Bean Lines. *Crop Sci* 48: 357–363
- Pereira G, Marques C, Ribeiro R, Formiga S, Dâmaso M, Sousa TM, Farinhó M, Leitão JM (2010) Identification of DNA markers linked to an induced mutated gene conferring resistance to powdery mildew in pea (*Pisum sativum* L.). *Euphytica* 171: 327–335
- Pierce WH (1948) Resistance to powdery mildew in peas. *Phytopathology* 38: 21
- Rakshit S (1997) Biochemical and molecular analysis of powdery mildew resistance in pea (*Pisum sativum* L.). Ph.D. thesis, I.A.R.I., New Delhi
- Rathi AS, Tripathi NN (1994) Assessment of growth reduction and yield losses in peas (*Pisum sativum* L.) due to powdery mildew disease caused by *Erysiphe polygoni* D.C. *Crop Reis* 8(2): 372–376
- Ray T, Chandra S (2009) Genetic Diversity of *Amaranthus* Species from the Indo-Gangetic Plains Revealed by RAPD Analysis Leading to the Development of Ecotype-Specific SCAR Marker. *Journal of Heredity* 100: 338–347
- Reiling TP (1984) Powdery mildews. In: Hagedorn DJ (ed) *Compendium of pea diseases*. American Phytopathological Society, St. Paul, MN, pp 21–22

- Saal B, Struss D (2005) RGA- and RAPD-derived SCAR markers for a *Brassica* B-genome introgression conferring resistance to blackleg in oilseed rape. *Theor Appl Genet* 111: 281–290
- Sambrook J, Russell DW (2001) *Molecular Cloning: A Laboratory Manual*. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, New York
- Sarala K (1993) Linkage studies in pea (*Pisum sativum* L.) with reference to *er* gene for powdery mildew resistance and other genes. Ph.D. Thesis, Indian Agricultural Research Institute, New Delhi, India
- Saxena JK, Tripathi RM, Srivastava RL (1975) Powdery mildew resistance in pea (*Pisum sativum* L.). *Curr Sci* 44: 746
- Sharma B (1997) Other pulses. In: Paroda RS, Chadha KL (eds) *50 Years of Crop Science Research in India*. Indian Council of Agricultural Research, pp 279–317
- Sharma B (2003) The *Pisum* genus has only one recessive gene for powdery mildew resistance. *Pisum Genet* 35: 22–27
- Sharma B, Yadav Y (2003) *Pisum fulvum* carries a recessive gene for powdery mildew resistance. *Pisum Genet* 35: 30
- Singh M, Chaudhary K, Boora K (2006) RAPD-based SCAR marker SCA 12 linked to recessive gene conferring resistance to anthracnose in sorghum [*Sorghum bicolor* (L.) Moench]. *Theor Appl Genet* 114: 187–192
- Singh MP (1984) Genetics of resistance to powdery mildew (*E. polygoni* DC), combining ability and heterosis in peas (*Pisum sativum* L.). Ph.D. Thesis. GB Pant Univ of Agric and Tech Pantnagar (Nainital), India, pp 169
- Singh RB, Singh MN, Singh UP Singh RM (1983) Inheritance of resistance to powdery mildew in pea and its use in breeding. *Indian J Agric Sci* 53: 855–856
- Sokhi SS, Jhooty JS, Bains SS (1979) Resistance in pea against powdery mildew. *Indian Phytopathol* 32: 571–574
- Srivastava RK, Mishra SK (2004) Inheritance of powdery mildew resistance using near-isogenic lines (NILs) in pea (*Pisum sativum* L.). *Indian J Genet* 64(4): 303–305
- Timmerman GM, Frew TJ, Weeden NF (1994) Linkage analysis of *er 1* a recessive *Pisum sativum* gene for resistance to powdery mildew fungus (*Erysiphe pisi* D.C.). *Theor Appl Genet* 88: 1050–1055
- Tiwari KR, Penner GA, Warkentin TD (1997) Inheritance of powdery mildew resistance in pea. *Can J Plant Sci* 77: 307–310
- Tiwari KR, Penner GA, Warkentin TD (1998) Identification of coupling and repulsion phase RAPD markers for powdery mildew resistance gene *er-1* in pea. *Genome* 41: 440–444
- Vaid A, Tyagi PD (1997) Genetics of powdery mildew resistance in pea. *Euphytica* 96: 203–206


- Vinod MS, Raghavan, Prashanth S, George S, Parida A (2007) Identification of a sex-specific SCAR marker in dioecious *Pandanus fascicularis* L. (Pandanaceae). *Genome* 50: 834–839
- Williams JGK, Kubelik AR, Livak KJ, Rafalski JA, Tingey SV (1990) DNA polymorphisms amplified by arbitrary primers are useful as genetic markers. *Nucl Acids Res* 18: 6531–6535
- Yarnell SH (1962) Cytogenetics of vegetable crops. III. Legumes. *Bot Rev* 28: 467–537
- Young D, Zamin M, Ganae W, Tanksley SD (1988) Use of isogenic lines and simultaneously probing to identify DNA markers tightly linked to the *Tm-2a* gene in tomato. *Genetics* 120: 579–585

Figure 1. RAPD marker OPX 04 showing 880 bp polymorphic band among parents of the mapping population.



S= Susceptible parent PG 3
 R= Resistant parent PG 3 (PMR)^{HFP 4}
 L= 100 bp plus DNA molecular weight marker

Figure 2. Insert size check (880bp) for the pGEM[®]-T Easy vector with *Eco*RI digestion



The clones 8, 9, 13, 16, 18, 21, 26 and 27 are showing insert of the right size i.e. 880 bp. Within these clone types, there are four distinct classes of clones. Clones 8 and 9 form class I; 13 and 18 from class II; 16 and 21 from class III; and 26 and 27 from class IV.

Figure 3. Comparative segregation pattern of the RAPD (OPX₈₈₀) derived SCAR marker ScOPX₈₈₀ in F₂ mapping population of the cross PG 3 (PMR)^{HFP 4} × PG 3 (PMS)




Figure 4. Linkage map of *er1* gene with SCAR and RAPD markers

ScOPX 04₈₈₀ (coupling phase)

0.6cM

ScOPD 10₆₅₀ (repulsion phase)

Loglikelihood = -158.10






Table 1. Near-isogenic lines, their donor and recipient parents used in the present investigation

S.No.	Near-isogenic line	PMS/PMR donor parent	Recipient parent
1.	HFP 4 (PMS) ^{Bonneville}	Bonneville (PMS)	HFP 4 (PMR)
2.	HFP 4 (PMS) ^{PG 3}	PG 3 (PMS)	HFP 4 (PMR)
3.	PG 3 (PMR) ^{HFP 4}	HFP 4 (PMR)	PG 3 (PMS)
4.	DMR 11 (PMS) ^{KPSD 1}	KPSD 1 (PMS)	DMR 11 (PMR)
5.	T 163 (PMR) ^{DMR 11}	DMR 11 (PMR)	T 163 (PMS)
6.	L 116 (PMR) ^{DMR 11}	DMR 11 (PMR)	L 116 (PMS)
7.	Pusa 10 (PMR) ^{DMR 11}	DMR 11 (PMR)	Pusa 10 (PMS)

PMR= Powdery Mildew Resistant, PMS= Powdery Mildew Susceptible.

Near-isogenic lines are written with the PMS/PMR donor source as indicated in superscripts.

Table 2. DNA Sequence information of different SCAR primers tested and the amplification pattern on parents of the mapping population

Class	Primer Sequence (5' → 3')	Length	Annealing Temperature		Amplification		Band Size
					S	R	
Ia.	<u>CCGCTACCGA</u> AGTATTGCAAG (Sense)	21	50-68 ⁰ C		+	+	880 bp
Ib.	<u>CCGCTACCGA</u> TTGATAGATAT (Antisense)	21	50-68 ⁰ C		+	+	880 bp
IIa.	<u>CCGCTACCGA</u> TAAATAGATTGCC (Sense)	22	55-69 ⁰ C		+	+	880 bp
IIb.	<u>CCGCTACCGA</u> GTTTTATCATTAA (Antisense)	22	55-69 ⁰ C		+	+	880 bp
IIIa.	<u>CCGCTACCGA</u> TGTTATGTTTG (Sense)	21	65.5 ⁰ C		+	--	880 bp
IIIb.	<u>CCGCTACCGA</u> ACTGGTTGGA (Antisense)	20	65.5 ⁰ C		+	--	880 bp
IVa.	TAGGGTGACACCGCAGTGAC (Sense)	20	53-69 ⁰ C		--	--	No Band
IVb.	TCATGCACGAGCCATGTATTC (Antisense)	21	53-69 ⁰ C		--	--	No Band

S= Susceptible parent PG 3 (PMS); R= Resistant parent PG 3 (PMR)^{HFP4}; Class III SCAR primer, designated as ScOPX 04₈₈₀ was synthesized from sequence information of class III clone (Figure 3)

Table 3. F₂ segregation of molecular markers in the cross PG 3 (PMR)^{HFP 4} × PG 3 (PMS)

Marker	Cross			No. of F ₂ plants			χ^2 (3:1)	P
	[PG 3 (PMR) ^{HFP 4}] × [PG 3(PMS)]			with the marker				
				Present (S)	Absent (R)	Total		
<i>er1</i>	<i>er1er1</i>	<i>Er1er1</i>	<i>Er1Er1</i>	155	53	208	0.026	0.88
ScOPX 04 ₈₈₀	-	+	+	156	52	208	0.000	1.00
OPX 04 ₈₈₀	-	+	+	156	52	208	0.000	1.00
ScOPD 10 ₆₅₀	+	+	-	155	53	208	0.026	0.88

Table 4. Pair-wise joint segregation analysis of molecular markers and *er1* gene

Marker pair	Pair-wise distance (cM)	Pair-wise LOD value
<i>er1</i> - Sc OPX 04 ₈₈₀	0.6 cM	48.34
<i>er1</i> - OPX 04 ₈₈₀	0.6 cM	48.34
<i>er1</i> - ScOPD 10 ₆₅₀	2.2 cM	37.12
ScOPX 04 ₈₈₀ –ScOPD 10 ₆₅₀	2.8 cM	3.79

*****END*****