



## Original Article

## Development of a rapid and simple HPLC-UV method for determination of gallic acid in *Schinopsis brasiliensis*



Felipe H.A. Fernandes<sup>a</sup>, Rayanne S. de A. Batista<sup>b</sup>, Francinalva D. de Medeiros<sup>a</sup>, Fábio S. Santos<sup>b</sup>, Ana C.D. Medeiros<sup>a,\*</sup>

<sup>a</sup> Laboratório de Desenvolvimento e Ensaio de Medicamentos, Centro de Ciências Biológicas e da Saúde, Universidade Estadual da Paraíba, Campina Grande, PB, Brazil

<sup>b</sup> Laboratórios Unificados de Desenvolvimento e Ensaio de Medicamentos, Departamento de Ciências Farmacêuticas, Centro de Ciências da Saúde, Universidade Federal da Paraíba, João Pessoa, PB, Brazil

## ARTICLE INFO

## Article history:

Received 17 December 2014

Accepted 19 May 2015

Available online 11 June 2015

## Keywords:

Gallic acid

Chemical marker

Validation

*Schinopsis brasiliensis*

## ABSTRACT

The aim of this work was to develop and validate an analytical method for the identification of the chemical marker of *Schinopsis brasiliensis* Engl., Anacardiaceae. It would determine the total polyphenols and flavonoid content by spectrophotometric methodology in the dried extract of plant. The chromatographic profiles of *S. brasiliensis* were determined using HPLC-UV. The liquid chromatography method was conducted on a Phenomenex Gemini NX C18 column (250 × 4.6 mm, 5 μm). The mobile phase consisted of 0.05% orthophosphoric acid: methanol. The flow rate was 1 ml/min and effluents were monitored at 271 nm. The retention time for gallic acid was 8.5 min. The described method has the advantage of being both rapid and easy. Hence it can be applied for routine quality control analysis of herbal preparation containing *S. brasiliensis*.

© 2015 Published by Elsevier Editora Ltda. on behalf of Sociedade Brasileira de Farmacognosia.

## Introduction

Polyphenols are a group of secondary metabolites present in different plants. In general polyphenols are classified into four classes: phenolic acids, flavonoids, lignans and stilbenes. These are very important to the pharmaceutical, cosmetic and food industries, mainly for their antioxidant action and properties (Gharras, 2009).

*Schinopsis brasiliensis* Engl., Anacardiaceae, is a plant which belongs to the Anacardiaceae family, known as “baraúna” or “bráuna,” and can be found in the Brazilian semi-arid regions. Their shells have ethnopharmacological uses for the treatment of diarrhea and coughs, and can also be used as an antiseptic and analgesic (Chaves et al., 2013). Several studies have demonstrated the antimicrobial activity of *S. brasiliensis* against *Pseudomonas aeruginosa*, *Staphylococcus aureus*, *Escherichia coli*, *Klebsiella pneumoniae* and *Candida albicans* (Chaves et al., 2011; Saraiva et al., 2011; Silva et al., 2012); *Salmonella typhimurium* (Saraiva et al., 2011); *Streptococcus oralis*, *Streptococcus mutans*, *Streptococcus parasanguinis* and *Enterococcus faecalis* (Silva et al., 2012). Other studies included multi-drug

resistant strains for tetracycline, gentamicin and oxacillin (Saraiva et al., 2011, 2013) and erythromycin (Silva et al., 2015).

The current list of compounds found in *S. brasiliensis* leaves includes the presence of tannins and polyphenols, such as methyl gallate, gallic acid, ellagic acid and 5,6,7,8,3',4'-hexahydroxy-flavonol (Saraiva et al., 2011). New alkylphenol methyl 6-eicosanoyl-2-hydroxy-4-methoxybenzoate and steroid 5α,8α-epidioxy ergosta-6,22-dien-3β-ol are both compounds isolated from the bark of *S. brasiliensis* (Cardoso et al., 2005). The essential oil extracted from the leaves of *S. brasiliensis* has a good amount of myrcene and low amounts of other compounds such as β-caryophyllene, eucalyptol and guaiol (Donati et al., 2014).

One important aspect for the quality control of herbal medicine products is the determination of the chemical marker, which consists of a constituent (or groups of constituents) that is chemically defined and present in the plant product. These constituents may or may not be related to their pharmacological activity (Marques et al., 2013). While several different analytical techniques can be used to identify and quantify these substances, we used a high performance liquid chromatography (HPLC), which is the method most commonly used by the various official standards (Zollner and Schwarz, 2013).

The aim of this work was to develop and validate an analytical method using HPLC for the identification and quantification of the chemical markers of *S. brasiliensis*.

\* Corresponding author.

E-mail: [anaclaudia@uepb.edu.br](mailto:anaclaudia@uepb.edu.br) (A.C.D. Medeiros).

## Materials and methods

### Raw material

The stem bark of *Schinopsis brasiliensis* Engl., Anacardiaceae, was collected in July 2013 from the region of Campina Grande, state of Paraíba (7°13'50" S, 35°52'52" W). The voucher specimen was prepared and identified by Professor Jayme Coelho de Moraes Herbarium, Universidade Federal da Paraíba, under the EAN-14049.

The herbal drugs were dried at 40 °C using an air oven. After drying, the material was reduced in a knife mill (10 mesh outlet sieve) to a fine powder. The hydroalcoholic extract was obtained by maceration of the powder in a water:ethanol (30:70, v/v) solvent system for 72 h and then dried using Spray Dryer (LabPlant®), with onset temperature of 140 °C and under 3 ml/min flow rate. Colloidal silicon dioxide (Aerosil®200) was used as a drying agent at 20% (w/w). The drying extract was weighted, protected from light and stored in closed flasks at room temperature.

### Determination of total polyphenols

The Folin–Ciocalteu method was used to determine the total polyphenols, using gallic acid as a standard (Chaves et al., 2013). A sample was dissolved in water with 0.5 ml of Folin–Ciocalteu solution (Sigma–Aldrich) and left at rest for 2 min. Then it was added to 1 ml of sodium carbonate solution 20% (w/w) and was left to stand for 10 min. The calibration curve was prepared with a gallic acid standard in different concentrations. The samples were measured using a UV–Vis Spectrophotometer Shimadzu UV-Mini 1240, with a wavelength of 757 nm. The results were expressed as micrograms of gallic acid/mg of extract (µg GAE/mg extract).

### Determination of total flavonoids

The aluminum chloride method was used to determine the flavonoid content using a quercetin standard (Chaves et al., 2013). Initially an aliquot of the extract was dissolved in methanol. Then the extract was mixed with an aluminum chloride solution (2% in methanol) in a ratio of 1:1 and left to stand for 10 min. The calibration curve was prepared with a quercetin standard in different concentrations. We used a UV–Vis Spectrophotometer Shimadzu UV-Mini 1240, with a wavelength of 415 nm to analyze our sample. The results were expressed as micrograms of quercetin/mg of the extract (µg QE/mg extract).

### Chemical marker

Chromatography analysis for the quantification of the chemical marker gallic acid in *S. brasiliensis* was conducted on a Shimadzu® HPLC model LC-10 controlled by Class VP Software and with quaternary pumps (LC-10AD), an on-line degasser (DGL-14A), and an injection valve (Rheodyne® LLC) with a 20 µl loop. The equipment included a UV–Vis detector (SPD-10A). The column used was a Phenomenex RP-Gemini NX C18 (250 mm × 4.6 mm, 5 µm) and it was protected by a pre-column Gemini NX C18 (10 mm × 4 mm).

The chromatographic separation was carried out using a mobile phase with phosphoric acid: water 0.05% as solvent A and methanol as solvent B at a flow rate of 1 ml/min. The gradient program was as follows: 90–10% B (10 min), 70–30% B (3 min), 40–60% B (5 min), 60–40% B (3 min), 80–20% B (3 min) and 90–10% B (6 min). The peaks were detected at 271 nm and were identified by comparing the retention time with standard gallic acid.

The validation of the analytical method has been made while taking into account the figures of merit such as precision, accuracy, linearity, limit of detection and limit of quantification (Marques et al., 2013; Ribani et al., 2004; Pedrosa and Salgado, 2014).

**Table 1**

Polyphenol and flavonoid contents of *Schinopsis brasiliensis*.

	Total polyphenols (µg/mg extract GAE)	Total flavonoids (µg/mg extract QE)
Concentration	24.52	1.43
%	15.08	1.44

The linearity was evaluated from the calibration curve, found = with a standard solution of gallic acid with a concentration between 2.5 and 15 µg/ml, and obtained the linear regression equation  $y = ax + b$ . According to Ribani et al. (2004), verification of the data fit can be observed by the correlation coefficient and determination. The Official Compendium recommends a value greater than 0.99.

## Results and discussion

The total polyphenol content is quite high when compared to the total flavonoids (Table 1). Both these classes of compounds are present in various medicinal plants and are related to antioxidants, antimicrobial, and anti-inflammatory activities (Corradini et al., 2011). Although we used a quantitative method to determine these compounds, this technique cannot predict the composition of each polyphenol, since it may possibly also measure non-phenolic compounds (Roby et al., 2013).

The polyphenols were present in greater quantity in the plant when we analyzed the hydroalcoholic extract of *S. brasiliensis* using gallic acid as standard. Gallic acid (3,4,5-trihydroxybenzoic acid) is an important polyphenol present in plants and is a compound with important applications in the synthesis of the drug Trimethoprim, as a food preservative, and an antioxidant in oils and products rich in lipids (Battestin et al., 2004).

In the preliminary analysis, we detected the presence of these compound classes in large quantities. Based on these results, gallic acid was chosen as a marker and we proceeded to quantify and validate the methods by means of the standard addition. For this test, we used high performance liquid chromatography with a UV–Vis detector, as it is the method most suitable for this type of compounds (Arapitsas, 2012).

Fig. 1 was obtained from the chromatogram of the extract of *S. brasiliensis*, which shows the peak of gallic acid and its standard. The retention time was 8.5 min, in addition to peaks present in other compounds of the phytocomplex. Good linearity ( $R^2 = 0.9984$ ) was achieved within the investigated ranges in the extract and was 99.5% of the explainable variance (Fig. 2).

Different studies are described in the literature (Arapitsas, 2012) analyzing gallic acid as the major compound; however, the difference between them is the type of mobile phase used. In a study of gallic acid by *Dendrophthoe falcata* L. f., Deshmukh and Prabhu (2011) found a retention time of approximately 12 min, using a mobile phase of water acidified with phosphoric acid (0.1%) and acetonitrile in the ratio of 4:6 and a flow rate 1 ml/min. In another work with *Schinus terebinthifolius* Raddi, gallic acid was quantified using two phases: one with water:methanol and the other with water:acetonitrile. In both cases, the water was acidified with formic acid to pH 2.7. The retention times under both conditions were 5 min, although the phase with acetonitrile was slightly shorter (Carvalho et al., 2009).

The choice of mobile phase (0.05% phosphoric acid and methanol) was determined using experiments described in the literature, which report that the acidified water prevents the ionization of gallic acid, causing it to have affinity for the stationary phase. Generally, the most widely used method is a reverse-phase column C-18. Although gallic acid absorbs at various wavelengths between 201 and 280 nm, we chose 271 nm due to its higher absorption and




Fig. 1. Chromatograms (HPLC/UV) of extract of *Schinopsis brasiliensis* (A) and gallic acid (B) in 271 nm.




Fig. 2. Calibration plot of gallic acid standard in 271 nm.

better resolution of the peaks in these conditions (Arapitsas, 2012; De Souza et al., 2002; Lopes et al., 2009).

Through the parameters of the calibration curve it was possible to determine with 95% confidence the limit of detection (LOD=1.22 µg/ml) and the limit of quantification (LOQ=1.82 µg/ml). The accuracy of the method was evaluated by repeatability and intermediate precision (Table 2).

Table 2  
Results of precision for the two analytes.

Analytes	Area peak	RSD (%)
Gallic acid (10 µg/ml)	623,253 ± 14,409.03	2.31
Extract (day 1)	616,653.8 ± 17,097.11	2.77
Extract (day 2)	627,397 ± 24,066	3.83

Table 3

Accuracy of three recoveries with different concentration to extract of *Schinopsis brasiliensis*.

	75%	100%	125%
Recovery (%)	92.83	96.87	111.20
RSD (%)	1.65	0.72	0.65

The accuracy of the method was determined by analysis of three fortifications by adding standard gallic acid (Table 3), and it was determined that the method performs within acceptable limits between 80% and 120% (Lima et al., 2010). According Betz et al., the accuracy test through recovery of the matrix is crucial to the development of methodologies for herbal products. The recovery test consists of adding a standard of known concentration after the extraction process in the matrix (Betz et al., 2011).

## Conclusions

In this study, a simple, accurate and reliable method was developed to determine the chemical marker of *S. brasiliensis* by using HPLC-UV. The results demonstrated that the method showed good linearity, recovery and reproducibility and low limits of detection and quantification. Thus, it is suitable to be used in the quantification of gallic acid in herbal medicine produced with dried extract of *S. brasiliensis*.

## Conflicts of interest

The authors declare no conflicts of interest.

## Acknowledgements

This study was supported by Fapesq, Propesq/UEPB, Capes and CNPq (Process No.: 562957/2010-3), Brazil for scholarships and

research fellowships. The authors especially wish to give thanks to Professor Ivan Coelho Dantas (*in memoriam*) for indicating the variety of plant to be studied in this work.

## References

- Arapitsas, P., 2012. Hydrolyzable tannin analysis in food. *Food Chem.* 135, 1708–1717.
- Battestin, V., Matsuda, L.K., Macedo, G.A., 2004. Fonte e aplicações de taninos e tanases em alimentos. *Aliment. Nutr.* 15, 63–72.
- Betz, J.M., Brown, P.N., Roman, M.C., 2011. Accuracy, precision, and reliability of chemical measurements in natural products research. *Fitoterapia* 82, 44–52.
- Cardoso, M.P., David, J.M., David, J.P., 2005. A new alkylphenol from *Schinopsis brasiliensis* Engl. *Nat. Prod. Res.* 19, 431–433.
- Carvalho, M.G., Freire, F.D., Raffin, F.N., Aragão, C.F.S., Moura, T.F.A., 2009. LC determination of gallic acid in preparations derived from *Schinus terebinthifolius* Raddi. *Chromatographia* 29, S249–S253.
- Chaves, T.P., Santana, C.P., Vêras, G., Brandão, D.O., Felismino, D.C., Medeiros, A.C.D., Trovão, D.M.B.M., 2013. Seasonal variation in the production of secondary metabolites and antimicrobial activity of two plant species used in Brazilian traditional medicine. *Afr. J. Biotechnol.* 12, 847–853.
- Chaves, T.P., Dantas, I.C., Felismino, D.C., Vieira, K.V.M., Clementino, E.L.C., Costa, L.S., 2011. Atividade antimicrobiana das folhas de *Schinopsis brasiliensis* Engler. *Biofar* 5, 11–17.
- Corradini, E., Foglia, P., Giansanti, P., Gubbiotti, R., Samperi, R., Laganá, A., 2011. Flavonoids: chemical properties and analytical methodologies of identification and quantitation in foods and plants. *Nat. Prod. Res.* 25, 469–495.
- De Souza, T.P., Holzschuh, M.H., Lionço, M.I., Gonzáles-Órtega, G., Petrovick, P.R., 2002. Validation of a LC method for the analysis of phenolic compounds from aqueous extract of *Phyllanthus niruri* aerial parts. *J. Pharm. Biomed.* 30, 351–356.
- Deshmukh, H., Prabhu, P.J., 2011. Development of RP-HPLC method for qualitative analysis of active ingredient (Gallic acid) from stem bark of *Dendrophthoe falcata* Linn. *Int. J. Pharm. Sci. Drug Res.* 3, 146–149.
- Donati, M., Mondin, A., Chen, Z., Miranda, F.M., Nascimento Júnior, B.B., Schirato, G., Pastori, P., Frolidi, G., 2014. Radical scavenging and antimicrobial activities of *Croton zehntneri*, *Pterodon emarginatus* and *Schinopsis brasiliensis* essential oils and their major constituents: estragole, trans-anethole,  $\beta$ -caryophyllene and myrcene. *Nat. Prod. Res.* 29, 939–946.
- Gharras, H.E., 2009. Polyphenols: food sources, properties and applications – a review. *Int. J. Food Sci. Technol.* 44, 2512–2518.
- Lima, L.L.A., Schuler, A., Guerra, N., Pereira, G.E., Lima, T.C.A., Rocha, H., 2010. Otimização e validação de método para determinação de ácidos orgânicos em vinhos por cromatografia líquida de alta eficiência. *Quim. Nova* 33, 1186–1189.
- Lopes, G.C., Sanches, A.C.C., Toledo, C.E.M., Isler, A.C., Mello, J.C.P., 2009. Determinação quantitativa de taninos em três espécies de *Stryphnodendron* por cromatografia líquida de alta eficiência. *Braz. J. Pharm. Sci.* 45, 135–143.
- Marques, G.S., Leão, W.F., Lyra, M.A.M., Peixoto, M.S., Monteiro, R.P.M., Rolim, L.A., Xavier, H.S., Rolim Neto, P.J., Soares, L.A.L., 2013. Comparative evaluation of UV/VIS and HPLC analytical methodologies applied for quantification of flavonoids from leaves of *Bauhinia forficata*. *Rev. Bras. Farmacogn.* 23, 51–57.
- Pedroso, T.M., Salgado, H.R.N., 2014. Validation of analytical methodology for quantification of cefazolin sodium pharmaceutical dosage form by high performance liquid chromatography to be applied for quality control in pharmaceutical industry. *Braz. J. Pharm. Sci.* 50, 213–223.
- Ribani, M., Botolli, C.B.G., Collins, C.H., Jardim, I.C.S.F., Melo, L.F.C., 2004. Validação de métodos cromatográficos e eletroforéticos. *Quim. Nova* 27, 771–780.
- Roby, M.H.H., Sarhan, M.A., Selim, K.A., Khalel, K.I., 2013. Evaluation of antioxidant activity, total phenols and phenolic compounds in thyme (*Thymus vulgaris* L.), sage (*Salvia officinalis* L.), and marjoram (*Origanum majorana* L.) extracts. *Ind. Crop. Prod.* 43, 827–831.
- Saraiva, A.M., Castro, R.H.A., Cordeiro, R.P., Peixoto Sobrinho, T.J.S., Castro, V.T.N.A., Amorim, E.L.C., Xavier, H.S., Pisciotano, M.N.C., 2011. In vitro evaluation of antioxidant, antimicrobial and toxicity properties of extracts of *Schinopsis brasiliensis* Engler (Anacardiaceae). *Afr. J. Pharm. Pharmacol.* 5, 1724–1731.
- Saraiva, A.M., Saraiva, C.L., Cordeiro, R.P., Soares, R.R., Xavier, H.R., Caetano, N., 2013. Atividade antimicrobiana e sinérgica das frações das folhas de *Schinopsis brasiliensis* Engler frente a clones multirresistentes de *Staphylococcus aureus*. *Rev. Bras. Plant. Med.* 15, 199–207.
- Silva, M.S.P., Brandão, D.O., Chaves, T.P., Formiga Filho, A.L.N., Costa, E.M.M.B., Santos, V.L., Medeiros, A.C.D., 2012. Study bioprospecting of medicinal plant extracts of the semiarid northeast: contribution to the control of oral microorganisms. *Evid.-Based Complement. Altern. Med.*, <http://dx.doi.org/10.1155/2012/681207>.
- Silva, K.M.A., Chaves, T.P., Santos, R.L., Brandão, D.O., Fernandes, F.H.A., Ramos Júnior, F.J.L., Santos, V.L., Felismino, D.C., Medeiros, A.C.D., 2015. Modulation of the erythromycin resistance in *Staphylococcus aureus* by ethanolic extracts of *Ximelia americana* L. and *Schinopsis brasiliensis* Engler. *Bol. Latinoam. Caribe.* 14, 92–98.
- Zollner, T., Schwarz, M., 2013. Herbal reference standards: applications, definitions and regulatory requirements. *Rev. Bras. Farmacogn.* 23, 1–21.