

# Development of a Specific Polymerase Chain Reaction-Based Assay for the Identification of *Fusarium oxysporum* f. sp. *ciceris* and Its Pathogenic Races 0, 1A, 5, and 6

María del Mar Jiménez-Gasco and Rafael M. Jiménez-Díaz

First author: Departamento de Protección de Cultivos, Instituto de Agricultura Sostenible (IAS), Consejo Superior de Investigaciones Científicas (CSIC), Apartado 4084, 14080 Córdoba, Spain; and second author: Departamento de Protección de Cultivos, IAS-CSIC, and Escuela Técnica Superior de Ingenieros Agrónomos y de Montes, Universidad de Córdoba, Apartado 3048, 14080 Córdoba, Spain.  
Accepted for publication 13 September 2002.

## ABSTRACT

Jiménez-Gasco, M. M., and Jiménez-Díaz, R. M. 2003. Development of a specific polymerase chain reaction-based assay for the identification of *Fusarium oxysporum* f. sp. *ciceris* and its pathogenic races 0, 1A, 5, and 6. *Phytopathology* 93:200-209.

Specific primers and polymerase chain reaction (PCR) assays that identify *Fusarium oxysporum* f. sp. *ciceris* and each of the *F. oxysporum* f. sp. *ciceris* pathogenic races 0, 1A, 5, and 6 were developed. *F. oxysporum* f. sp. *ciceris*- and race-specific random amplified polymorphic DNA (RAPD) markers identified in a previous study were cloned and sequenced, and sequence characterized amplified region (SCAR) primers for specific PCR were developed. Each cloned RAPD marker was characterized by Southern hybridization analysis of *Eco*RI-digested genomic DNA of a subset of *F. oxysporum* f. sp. *ciceris* and nonpathogenic *F. oxysporum* isolates. All except two cloned RAPD markers consisted of DNA sequences that were found highly repetitive in the genome of all *F. oxysporum* f. sp. *ciceris* races. *F. oxysporum* f. sp. *ciceris* isolates representing eight reported races from a wide geographic range, nonpathogenic *F. oxysporum* isolates, isolates of *F. oxysporum* f.

spp. *lycopersici*, *melonis*, *niveum*, *phaseoli*, and *pisi*, and isolates of 47 different *Fusarium* spp. were tested using the SCAR markers developed. The specific primer pairs amplified a single 1,503-bp product from all *F. oxysporum* f. sp. *ciceris* isolates; and single 900- and 1,000-bp products were selectively amplified from race 0 and race 6 isolates, respectively. The specificity of these amplifications was confirmed by hybridization analysis of the PCR products. A race 5-specific identification assay was developed using a touchdown-PCR procedure. A joint use of race 0- and race 6-specific SCAR primers in a single-PCR reaction together with a PCR assay using the race 6-specific primer pair correctly identified race 1A isolates for which no RAPD marker had been found previously. All the PCR assays described herein detected up to 0.1 ng of fungal genomic DNA. The specific SCAR primers and PCR assays developed in this study clearly identify and differentiate isolates of *F. oxysporum* f. sp. *ciceris* and of each of its pathogenic races 0, 1A, 5, and 6.

*Additional keywords:* *Cicer arietinum*, diagnostics, DNA hybridizations, DNA markers, Fusarium wilt of chickpea, repetitive DNA.

Accurate and rapid identification of pathogens is necessary for appropriate management of plant diseases. In particular, phenotypic and genetic characterization of the pathogenic variants of the plant pathogens prevalent in an area is required for efficient disease management. *Fusarium oxysporum* Schlechtend.:Fr. is an anamorphic soilborne fungus that includes both nonpathogenic and plant-pathogenic isolates. Pathogenic forms of this fungus cause vascular and cortical rot diseases in many agricultural crops, and have been classified into formae speciales based on their host specificity (26). Also, variation in virulence among host cultivars can be found within isolates of a forma specialis, leading to the designation of pathotypes and pathogenic races (26).

*F. oxysporum* f. sp. *ciceris* (Padwick) Matuo & K. Sato, the agent of Fusarium wilt of chickpea (*Cicer arietinum* L.), exhibits great diversity. Two pathotypes, yellowing and wilting, have been differentiated by the disease syndrome induced in the plant in pathogenicity tests (34). The yellowing pathotype induces progressive foliar yellowing with vascular discoloration, followed by plant death within 40 days of inoculation. The wilting pathotype induces severe chlorosis and flaccidity, vascular discoloration, and plant death within 20 days after inoculation. In addition to variation in symptom type, there are eight races of *F. oxysporum* f. sp.

*ciceris* (races 0, 1A, 1B/C, 2, 3, 4, 5, and 6), which are identified by reactions on a set of differential chickpea cultivars (12,15). Races 0 and 1B/C induce the yellowing syndrome (yellowing pathotype), whereas races 1A, 2, 3, 4, 5, and 6 induce the wilting syndrome (wilting pathotype) (15). The eight races have distinct geographic distributions. Races 2, 3, and 4 have been reported only in India (12), whereas races 0, 1B/C, 5, and 6 are found mainly in the Mediterranean region and in the United States (California) (11,15,17). Race 1A has been reported in India (12), California, and the Mediterranean region (15,17).

Fusarium wilt of chickpea is managed primarily by the use of resistant cultivars (14). This makes the identification of pathogenic races of *F. oxysporum* f. sp. *ciceris* in a given area important for disease resistance breeding and for the efficient use of resistant cultivars. In the absence of alternative methods, *F. oxysporum* isolates from chickpea are characterized into *F. oxysporum* f. sp. *ciceris* and further into pathogenic races. Determination of races in this pathogen is conceptually simple but costly in time (50 to 60 days), facilities, and resources (15,34). Furthermore, nonpathogenic *F. oxysporum* isolates infecting the chickpea root cortex may be identified as *F. oxysporum* f. sp. *ciceris* due to the variability inherent to biological pathotyping (17,19). Therefore, new methods for the rapid, consistent, reliable, and reproducible identification of the pathogen and its pathogenic races are needed.

Polymerase chain reaction (PCR) has been widely and successfully employed for the identification of important plant-pathogenic

Corresponding author: R. M. Jiménez-Díaz; E-mail address: agljdir@uco.es

Publication no. P-2002-1213-01R  
© 2003 The American Phytopathological Society

fungi (13,23). In many cases, primers for these uses were based on DNA sequence polymorphisms existing within highly conserved regions of the nuclear ribosomal DNA, such as the internal transcribed spacer or the intergenic spacer region (35). Although this strategy proved successful for species identification (3), the above regions do not normally reveal sufficient polymorphism for distinguishing subspecific groups such as formae speciales or pathogenic races. An alternative strategy in the development of markers for infraspecific taxa is based on the isolation and sequencing of distinct fragments of random amplified polymorphic DNA (RAPD), and the use of these sequences to design PCR primers that specifically amplify selected markers. This approach of using sequence characterized amplified regions (SCARs) was first applied by Paran and Michelmore (29), and since then has been very effective for the intraspecific identification of a diversity of plant pathogens (20,24).

In a previous study, we analyzed a worldwide collection of *F. oxysporum* f. sp. *ciceris* isolates representative of all pathogenic races, as well as nonpathogenic *F. oxysporum* isolates (17). Pathogenic isolates were unambiguously typed as *F. oxysporum* f. sp. *ciceris* and pathogenic races 0, 1B/C, 5, and 6 by distinct RAPD markers. The applicability of those markers for diagnostic purposes was confirmed by biological and molecular "blind trials" and found reproducible in repeated amplification experiments (17). However, the reliability of RAPD markers might be influenced by the source and procedure used for DNA isolation, the occurrence of contaminants, the amplification of different DNA sequences of the same size, the thermocycler, reaction conditions, and other factors. This limits the use of the RAPD markers for general diagnostics among many laboratories. To overcome these difficulties, there is a need to develop improved molecular assays and markers for the identification of the pathogen and its pathogenic races.

The aim of the present study was to develop a PCR assay that could selectively discriminate isolates of *F. oxysporum* f. sp. *ciceris* from nonpathogenic *F. oxysporum* from chickpea, other *F. oxysporum* formae speciales and *Fusarium* spp., and identify the races of the pathogen that are prevalent in the Mediterranean region. Our approach was based in that *F. oxysporum* f. sp. *ciceris* is a strictly asexually reproducing, highly clonal fungus (16,30) in which genetic recombination does not occur and for which genetic markers can be used to identify races. For our purpose, we used SCARs to generate specific PCR primers based on informative RAPD markers and optimized conditions for assays (17). This process required the cloning, sequencing, and characterization of the RAPD markers of the races.

## MATERIALS AND METHODS

**Fungal isolates.** A large, systematic collection of fungal isolates was used in this study, consisting of 76 *F. oxysporum* f. sp. *ciceris* isolates representative of the eight described races and from a wide geographic distribution of the pathogen, 20 chickpea isolates of *F. oxysporum* from roots of healthy plants and nonpathogenic to chickpea, 15 isolates from five different *F. oxysporum* formae speciales, and 58 isolates of 47 different *Fusarium* spp. (Table 1). The isolates of *F. oxysporum* f. sp. *ciceris* races were characterized in previous studies (15, 17,19).

All 96 *F. oxysporum* isolates from chickpea (including pathogenic and nonpathogenic ones) and five isolates of other *F. oxysporum* formae speciales were from the culture collection of Departamento de Protección de Cultivos, Instituto de Agricultura Sostenible, CSIC, Córdoba, Spain. An isolate of *F. oxysporum* f. sp. *lycopersici* was a gift from D. Fravel, U.S. Department of Agriculture-Agricultural Research Service (USDA-ARS), Beltsville, MD, and eight *F. oxysporum* f. sp. *pisi* isolates were provided by W. Boge, USDA-ARS, Prosser, WA. Isolates of other

*Fusarium* spp. were obtained from J. F. Leslie, Department of Plant Pathology, Kansas University, Manhattan. Monoconidial cultures of isolates were stored in sterile soil in test tubes at 4°C (26). Active cultures of the isolates were subsequently obtained by placing small aliquots of the soil culture onto a plate of fresh potato dextrose agar (PDA) (250 g of unpeeled potatoes, 20 g of agar, and 20 g of glucose per liter of distilled water) and incubating for 4 days at 25°C and a 12-h photoperiod of fluorescent and near-UV light at 36  $\mu\text{E m}^{-2} \text{s}^{-1}$ . Mycelia for DNA extraction were produced by the following methods. (i) For large-scale production of mycelium, 1 ml of conidial suspension ( $5 \times 10^6$  conidia per ml) was transferred to 250-ml flasks containing 100 ml of potato dextrose broth (250 g of unpeeled potatoes and 20 g of glucose per liter of distilled water). Cultures were incubated on an orbital shaker at 125 rpm, 25°C, and 12 h of light for 4 days. Mycelia were harvested by filtration through sterile cloth, washed with sterile water, lyophilized, and stored at -20°C. (ii) For small-scale production of mycelium, a small piece from an active fungal culture on PDA was placed onto a sterile cellophane film layered over a PDA plate and incubated at 25°C and 12 h of light for 4 to 6 days. Mycelia were harvested from the cellophane surface, lyophilized, and stored at -20°C.

**DNA extraction.** Genomic DNA of the fungal isolates was purified from 50 mg of lyophilized ground mycelia as described by Raeder and Broda (31), with slight modifications. Ground mycelium was suspended in an extraction buffer (200 mM Tris-HCl [pH 8.5], 25 mM NaCl, 25 mM EDTA, and 0.5% sodium dodecyl sulfate). The mixture was homogenized with equilibrated phenol/chloroform/isoamyl alcohol (25:24:1), and centrifuged for 1 h at 12,000  $\times g$  (4°C). The supernatant was treated with RNAase and extracted with chloroform/isoamyl alcohol (24:1), followed by centrifugation (15 min at 12,000  $\times g$  and 4°C). DNA was precipitated by adding cold isopropanol, resuspended in sterile water, and stored at -20°C. Aliquots of the DNA samples were analyzed on 0.7% agarose gels in 1 $\times$  Tris-acetate-EDTA (TAE) buffer to estimate the concentration and quality of the extracted DNA. For PCR reactions, samples were diluted to a final concentration of 25 to 50 ng/ $\mu\text{l}$  in sterile water.

**Cloning of RAPD fragments.** The following RAPD markers of *F. oxysporum* f. sp. *ciceris* and of its pathogenic races 0, 1B/C, 5, and 6 (17) were cloned: *F. oxysporum* f. sp. *ciceris*, 1.5 kb (OPF-12, 5'-ACGGTACCAG-3'); race 0, 0.39 kb (OPF-12) and 0.9 kb (OPI-09, 5'-TGGAGAGCAG-3'); race 1B/C, 0.53 kb (OPI-09) and 1.1 kb (OPF-10, 5'-GGAAGCTTGG-3'); race 5, 0.9 kb (OPF-10); and race 6, 1.0 and 1.4 kb (OPI-09) (Operon Technology, Alameda, CA). RAPD reactions were carried out as described previously (17). RAPD products were separated by electrophoresis in 1.5% agarose gels in 1 $\times$  TAE buffer, stained with ethidium bromide, and visualized under UV light. The 0.1-kb DNA ladder XIV size marker was used for electrophoresis (Roche Diagnostics, Mannheim, Germany). The RAPD markers of *F. oxysporum* f. sp. *ciceris* or its pathogenic races were excised from the agarose gel, and DNA was purified using the Qiaex gel extraction kit (Qiagen, Hilden, Germany). The purified RAPD products were cloned into the vectors pGem-T (Promega, Madison, WI) or pCR2.1 TOPO (TOPO TA, Invitrogen, Groningen, Netherlands) according to the manufacturer's instructions and used to transform competent *Escherichia coli* cells. For each cloned RAPD marker, plasmid DNA was released from 30 to 50 transformed *E. coli* colonies by the alkaline lysis method (33). A RAPD marker may comprise DNA fragments of the same size but differing in sequence; therefore, we aimed to ensure that DNA inserts carried by selected bacterial clones corresponded to the putative correct RAPD marker. Therefore, insert DNAs were digested with *Apa*I, *Bam*HI, *Eco*RI, *Pst*I, *Sac*I, and *Xba*I endonucleases (Pharmacia LKB, Upsala, Sweden) and the restriction analyses identified a number of different DNA sequences. Thereafter, we hybridized blotted RAPD patterns associated to *F.*

*oxysporum* f. sp. *ciceris* and its pathogenic races 0, 1B/C, 5, and 6 using each DNA insert as a probe to verify that the insert hybridized only to the race from which the SCAR was developed.

**Hybridization assays.** Three types of hybridization assays were carried out in this study using each of the cloned RAPD markers as a probe: (i) hybridization of blotted RAPD patterns, (ii) hybridization of fungal genomic DNA digested with the restriction enzyme *EcoRI* (Southern blot), and (iii) hybridization of

blotted specific PCR products. For hybridization analyses of either RAPD patterns or specific PCR products, amplification products (25 µl) were separated in 1.5% agarose gels and transferred onto nylon Zeta-Probe blotting membranes (Bio-Rad Laboratory, S.A., Madrid, Spain) by capillarity in alkaline conditions (33). Cloned RAPD markers used as probes were labeled using digoxigenin-3-O-methylcarbonyl-amino-caproyl-5-(3-aminoallyl)-uridine-5'-triphosphate (DIG-11-dUTP) by random-

TABLE 1. Isolates of *Fusarium* spp. evaluated by polymerase chain reaction (PCR) with sequence characterized amplified region (SCAR) primers, with reference number, geographic origin, and race classification

Isolate reference <sup>a</sup>	Geographic origin <sup>b</sup>	Isolate reference <sup>a</sup>
<i>Fusarium oxysporum</i> f. sp. <i>ciceris</i>		
Race 0		
Foc-7802, -7952, -8207, -82108, -82113, -8503, -8733, -8912, -9018 PV1, -9018 JG62, -9032, -90111, -91100, -91015, -91108, -91114	Spain	<i>F. acuminatum</i> 11442 <i>F. acutatum</i> 10769 <i>F. armeniacum</i> 11623 <i>F. andiyazi</i> 4647
Foc-9601, -9603, -9604, -9605, -9606, -T3	Tunisia	<i>F. anthophilum</i> 11560
Foc-cc21C, -cc62R, -cc63K	Israel	<i>F. avenaceum</i> 11440
Foc-L96-5, -L96-6, -L96-7, -L96-9, -L96-10, -L96-11	Lebanon	<i>F. babinda</i> 11478
Foc-Sy 96-14-1, -Sy 96-18-3, -Sy 96-19-1	Syria	<i>F. begoniae</i> 10767
Foc-T 96-1-3	Turkey	<i>F. brevicatenulatum</i> 10756 <i>F. bulbicola</i> 10759
Race 1B/C		
Foc-USA 3-1JG62, -1987-W17	United States	<i>F. chlamydosporum</i> 11397
Foc-9602	Tunisia	<i>F. circinatum</i> 10766, H-10847
Foc-Sy 96-10-1	Syria	<i>F. circinatum</i> H-10850
Foc-T 96-1-1, -T 96-1-2, T -96-2-1, -T 96-2-2	Turkey	<i>F. compactum</i> 11624
Foc-cc22D	Israel	<i>F. concentricum</i> 10765 <i>F. crookwellense</i> 11451
Race 1A		
Foc-7989	India	<i>F. culmorum</i> 11427
Foc-9168	Morocco	<i>F. decemcellulare</i> 11411
Foc-8272	Spain	<i>F. denticulatum</i> 10763 <i>F. dimerum</i> 11425
Race 2		
Foc-8605, -1992 R2N	India	<i>F. equiseti</i> 11439
Race 3		
Foc-8606, -1992 R3N	India	<i>F. globosum</i> 11554 <i>F. guttiforme</i> 10764
Race 4		
Foc-8607, -1992 R4N	India	<i>F. lactis</i> 10757 <i>F. lateritium</i> 11403
Race 5		
Foc-8012, -8257, -8408, -8508, -9035, -9094 PV1, -9094 JG62	Spain	<i>F. longipes</i> 11428 <i>F. nelsonii</i> 11564 <i>F. nisikadoi</i> 10758
Foc-USA1-1JG62, -USA W6-1, -USA14201	United States	<i>F. nygamai</i> 5111 <i>F. phyllophilum</i> 10768
Race 6		
Foc-8905, -8924, -9023, -9027 PV1, -9093 PV1	Spain	<i>F. poae</i> 11470
Foc-9164, -9166, -9170	Morocco	<i>F. polyphialidicum</i> 11414
Foc-Tonini	United States	<i>F. proliferatum</i> 11558, D-4853, D-4854
Foc-9620, -9622, -9628, -9632	Israel	<i>F. pseudoanthophilum</i> 10755 <i>F. pseudocircinatum</i> 10761
<i>F. oxysporum</i> f. sp. <i>lycopersici</i> Fol-PV, -325-3	Unknown	<i>F. pseudograminearum</i> 11435
<i>F. oxysporum</i> f. sp. <i>melonis</i> Fom-8701, -9016	Unknown	<i>F. pseudonygamai</i> 10762
<i>F. oxysporum</i> f. sp. <i>niveum</i> Fon-8805, -8822	Unknown	<i>F. ramigenum</i> 10670
<i>F. oxysporum</i> f. sp. <i>phaseoli</i> Fop-DR85	Unknown	<i>F. sacchari</i> 278, B-3852, B-3853 <i>F. scirpi</i> 11409
<i>F. oxysporum</i> f. sp. <i>pisi</i>		
Race 1		
F-51, -22	Unknown	<i>F. semisectum</i> 11432
Race 2		
F-35, -42, -81	Unknown	<i>F. solani</i> 11420 <i>F. sporotrichioides</i> 11552
Race 6		
F-4A, -207, -241	Unknown	<i>F. subglutinans</i> 11544, E-990, E-2192 <i>F. thapsinum</i> F-4093, F-4094 <i>F. torulosum</i> 11419 <i>F. tricinctum</i> 11566
<i>F. oxysporum</i>		
Nonpathogenic		
Fo-8250, -9009, -9081, -90101, -90105, -91117	Spain	<i>F. verticillioides</i> 11556, A-149, A-999
Fo-9169	Morocco	
Fo-Fsp4, -Fsp7, -Fsp8, -Fsp9	Algeria	
Fo-420, -422, -425, -442, -448, -457, -506, -526, -C4P	Italy	

<sup>a</sup> Race of *Fusarium oxysporum* f. sp. *ciceris* was determined by pathogenicity tests on differential chickpea lines before PCR assays (15,17,20). Isolates of *F. oxysporum* f. sp. *lycopersici* Fol-PV, *melonis*, *niveum*, and *phaseoli* were from the fungal culture collection of Departamento de Protección de Cultivos, Instituto de Agricultura Sostenible (IAS-CSIC), Córdoba, Spain. Isolate of *F. oxysporum* f. sp. *lycopersici* Fol-325-3 was provided by D. Fravel, United States Department of Agriculture-Agricultural Research Service (USDA-ARS) Vegetable Development Laboratory, Beltsville, MD. Isolates of *F. oxysporum* f. sp. *pisi* were provided by W. Boge, USDA-ARS Pacific West Area, Prosser, WA. Isolates of other *Fusarium* spp. were obtained from J. F. Leslie, Department of Plant Pathology, Kansas State University, Manhattan. Geographic origin of these latter isolates was not provided. *F. oxysporum* isolates listed as nonpathogenic are not pathogenic to chickpea.

<sup>b</sup> Isolates from Algeria, Italy, Morocco, Spain, and the United States (California) were obtained from the fungal culture collection of Departamento de Protección de Cultivos, IAS-CSIC, Córdoba, Spain. *F. oxysporum* f. sp. *ciceris* isolates from India were provided by M. P. Haware, ICRISAT, Hyderabad, India; isolates from Tunisia were provided by M. H. Halila, Institute Nationale de la Recherche Agronomique, Ariana, Tunisia; isolates from Israel were provided by J. Katan, Department of Plant Pathology and Microbiology, The Hebrew University of Jerusalem, Israel; isolates from Lebanon, Syria, and Turkey were provided by C. Akem, ICARDA, Aleppo, Syria; unknown = geographic origin not known.

primed labeling (Roche Diagnostics) according to the manufacturer's instructions.

For Southern blots, 3 to 5 µg of genomic DNA was digested with *EcoRI*. The restriction fragments were separated by electrophoresis in 0.7% agarose gels and transferred to nylon membranes using a vacuum transfer system (VacuGene XL, VacuPump, Pharmacia LKB). Hybridization conditions were as described by Sambrook et al. (33) using high-astringency temperature (68°C). Homologous sequences in the fungal genome were detected by the DIG Luminescent Detection kit (Roche Diagnostics) according to the manufacturer's instructions.

**Sequencing of cloned RAPD markers and design of specific SCAR primers.** Cloned RAPD inserts were sequenced at the DNA Sequencing Facility at Cornell University (Ithaca, NY), or at Newbiotechnic, S.A. (Sevilla, Spain), using universal M13 and internal sequencing primers. A search for sequence similarities was performed with BLASTN and BLASTX programs of NCBI network service (1). We designed a PCR primer pair so as to correspond to sequences between 18 to 30 nucleotides that were identified at both ends of the insert. These primers included part of or the full nucleotide sequence of the Operon primer that amplified the RAPD marker originally (29). Primers were designed using the computer program PrimerSelect 3.11 (DNASar, Lasergen, Madison, WI). Sequences for the specific primers designed in this study are shown in Table 2 and were synthesized by GENSET (Paris).

**Specific PCR reactions.** The PCR reaction mixture (25 µl) consisted of 2.5 µl of 10× reaction buffer (166 mM (NH<sub>4</sub>)<sub>2</sub>SO<sub>4</sub>, 670 mM Tris-HCl [pH 8.0, 25°C], Tween 20), 0.2 µM of each primer, 200 µM of each dNTP, 0.64 units of *EcoTaq* DNA

Polymerase (EcoGen, Madrid, Spain), 1.0 mM MgCl<sub>2</sub>, and 10 to 25 ng of fungal DNA. Amplifications were performed in Perkin-Elmer 2400 and 9600 thermocyclers (Perkin-Elmer, Norwalk, CT). All reactions were repeated at least twice and always included a positive control of a known template DNA and a negative control with no DNA. Cycling profiles consisted of an initial step of 2 min at 94°C, 25 or 28 cycles of 30 s at 94°C, 1 min of annealing temperature, and 30 s at 72°C, followed by a final step of 4 min at 72°C. Annealing temperatures were 58°C for the *F. oxysporum* f. sp. *ciceris*-specific primer pair (Foc0-12f/Foc0-12r), and 61°C for specific primer pairs of races 0, 1A/6, and 6 of the pathogen (Table 2). For the race 5-specific primer pair (FocR5-L10f/FocR5-L10r), we used a touchdown-PCR procedure (6) to ensure specificity of the amplification product. For this procedure, the annealing temperature was 71°C for the first PCR cycle and decreased by 1°C per cycle for the next 10 cycles until an optimal annealing temperature of 61°C was reached. Then, there were 15 PCR cycles at the annealing temperature of 61°C, as described previously. PCR amplification products were separated and visualized as described above for the RAPD reactions.

## RESULTS

**Cloning and hybridization of RAPD markers.** Jiménez-Gasco et al. (17) identified the following specific RAPD markers for *F. oxysporum* f. sp. *ciceris* and each of the pathogenic races: *F. oxysporum* f. sp. *ciceris*, 1.5 kb (OPF-12); race 0, 0.39 kb (OPF-12) and 0.9 kb (OPI-09); race 1B/C, 0.53 kb (OPI-09) and 1.1 kb (OPF-10); race 5, 0.9 kb (OPF-10); and race 6, 1.0 and 1.4 kb

TABLE 2. GenBank accession numbers of sequence characterized amplified regions (SCARs), characteristics of recombinant DNA bacterial plasmids, SCAR primers, and amplification conditions for specific polymerase chain reaction assays developed for the identification of *Fusarium oxysporum* f. sp. *ciceris* and races 0, 1A, 1B/C, 5, and 6

Bacterial clone or primer	Isolate	Insert <sup>a</sup>	Accession number	Sequence (5'-3') <sup>b</sup>	Product, race <sup>c</sup>	Conditions <sup>d</sup>
<b>Clone<sup>e</sup></b>						
FocC19	Foc-7802	0.39 kb-R0-OPF-12	AF491867	...	...	...
FocD33	Foc-USA 3-1 JG62	1.1 kb-R1B/C-OPF-10	AF491868	...	...	...
FocL10	Foc-9035	0.9 kb-R5-OPF-10	AF491869	...	...	...
FocM15	Foc-9018 JG62	0.9 kb-R0-OPI-09	AF491870, AF492450	...	...	...
FocN5	Foc-USA 3-1 JG62	0.53 kb-R1B/C-OPI-09	AF492012	...	...	...
FocO2	Foc-9093 PV1	1.0 kb-R6-OPI-09	AF492013	...	...	...
FocP18	Foc-9093 PV1	1.4 kb-R6-OPI-09	AF492014, AF492450	...	...	...
Foc0-12	Foc-9018 JG62	1.5 kb-Foc-OPF-12	AF492451	...	...	...
<b>Primer<sup>f</sup></b>						
Foc0-12f	...	...	...	GGCGTTTCGCAGCCTTACAATGAAG	1,503-bp,	58°C,
Foc0-12r	...	...	...	<u>G</u> ACTCCTTTTCCCGAGGTAGGTCAGAT	f. sp. <i>ciceris</i>	28 cycles
FocR0-M15f	...	...	...	<u>GGAGAGCAGGACAGCAAAGACTA</u>	≈900-bp, race 0	61°C,
FocR0-M15r	...	...	...	<u>GGAGAGCAGCTACCTTAGATACACC</u>		28 cycles
FocR1B/C-N5f	...	...	...	<u>GAGAGCAGGGTCAGCGTAGATAG</u>	553-bp,	61°C,
FocR1B/C-N5r	...	...	...	<u>GCAGCAGAAGAGGAAGAAAATGTA</u>	race 1B/C	28 cycles
FocR5-L10f	...	...	...	<u>GGAAGCTTGGCATGACATAC</u>	938-bp, race 5	71-61°C,
FocR5-L10r	...	...	...	<u>AAGCTTGGGCACCCTCTT</u>		touchdown
FocR6-O2f	...	...	...	<u>GAGCAGTCAATGGCAATGG</u>	1,000-bp, race 6	61°C,
FocR6-O2r	...	...	...	<u>AGAGCAGGGTCAGCGTAGATA</u>		28 cycles
FocR6-P18f	...	...	...	<u>GGAGAGCAGTAGAGTTACAGCAGTATT</u>	≈1,400-bp,	61°C,
FocR0-M15r	...	...	...	<u>GGAGAGCAGCTACCTTAGATACACC</u>	races 1A, 6	25 cycles

<sup>a</sup> Approximate molecular weight of the cloned random amplified polymorphic DNA (RAPD) markers of races (R0 = race 0, R1B/C = race 1B/C, R5 = race 5, R6 = race 6) and *F. oxysporum* f. sp. *ciceris*, and arbitrary OPERON primer that originated it, respectively.

<sup>b</sup> Underlined nucleotides indicate sequences of the corresponding arbitrary primers.

<sup>c</sup> Polymerase chain reaction (PCR) product and race from which amplified.

<sup>d</sup> Amplification conditions: annealing temperature and number of PCR cycles. Cycles were as follows: 2 min at 94°C; 25 or 28 cycles of 30 s at 94°C, 1 min of annealing temperature, and 30 s at 72°C; and a final step of 4 min at 72°C. Touchdown PCR (6) consisted of 2 min at 94°C; 10 cycles at decreasing annealing temperature (71 to 61°C); 15 cycles of 30 s at 94°C, 1 min at 61°C, and 30 s at 72°C; followed by 4 min at 72°C.

<sup>e</sup> RAPD markers of *F. oxysporum* f. sp. *ciceris* and pathogenic races 0, 1B/C, 5, and 6 developed in a previous study (17) were cloned using commercial bacterial vectors pGem-T (Promega), and pCR2.1 (TOPO TA Cloning Kit, Invitrogen).

<sup>f</sup> Primers nomenclature: race of *F. oxysporum* f. sp. *ciceris* identified and bacterial clone from which the sequences were designed.

(OPI-09). Cloning of these RAPD marker bands into a bacterial vector yielded three to six types of different DNA inserts from a cloned RAPD band. These DNA inserts were distinguished because of slight differences in molecular size, and the distinct restriction patterns of inserts digested with *Apa*I, *Bam*HI, *Eco*RI, *Pst*I, *Sac*I, and *Xba*I endonucleases. Hybridization of the RAPD patterns that contained the marker DNA band for *F. oxysporum* f. sp. *ciceris* and for its races 0, 1B/C, 5, and 6, using as probes each of DNA inserts derived from cloning of the corresponding RAPD marker, allowed to identify the relevant marker sequences. In all cases, only one of the inserts hybridized to the corresponding RAPD marker. Other insert DNAs hybridized to RAPD products of similar molecular weight that were amplified from all the races tested but were not distinguishable in the agarose gels following electrophoretic separation. The above results allowed selection of bacterial clones carrying inserts corresponding to each of *F. oxysporum* f. sp. *ciceris* and races 0, 1B/C, 5, and 6 (Table 2).


Inserts from clones FocD33, FocL10, FocN5, FocO2, and Foc0-12 hybridized strongly with the single RAPD marker band amplified from isolates of the appropriate race, and this signal corresponded in size to that of the progenitor RAPD fragment (Fig. 1). Therefore, these inserts were appropriate for the identification of the corresponding *F. oxysporum* f. sp. *ciceris* race. Conversely, clone FocC19 derived from the race 0 RAPD marker

amplified by OPF-12 hybridized to this fragment from all race 0 isolates as expected, but it also hybridized to DNA bands amplified by this primer from other *F. oxysporum* f. sp. *ciceris* races as well as from *F. oxysporum* isolates nonpathogenic to chickpea. However, none of the FocC19-hybridized amplicons of above were distinguishable after separation of the RAPD products in agarose gels. Therefore, the insert carried by clone FocC19 was not suitable for identification of race 0 and was disregarded for further investigations. Finally, inserts from clones derived from race 0 (FocM15) and race 6 (FocP18) marker bands amplified by primer OPI-09 (Table 2) produced a faint cross-hybridization signal when probed against the RAPD patterns generated by OPI-09 from isolates of these races. However, these clones did not hybridize with any other amplicon. Subsequent nucleotide sequence analysis showed that these two insert DNAs share a short identical sequence.

Southern hybridizations of *Eco*RI-digested genomic DNA from isolates of *F. oxysporum* f. sp. *ciceris* races using the inserts as probes allowed characterization of the following insert DNAs as moderate to highly repetitive (20 to 30 bands, 1.5 to 12.0 kb in size) DNAs (8): FocD33, FocM15, FocN5, FocO2, and FocP18. The genome of each of the races tested shared homologies to all insert DNAs of above, but the hybridization patterns showed race-specific polymorphisms (i.e., for every insert DNA of above the hybridization patterns showed higher similarity among isolates of a race compared with that with isolates of different races). Similarly, hybridization of *Eco*RI-digested genomic DNA from nonpathogenic *F. oxysporum* isolates Fo-420, -8250, -9009, -9081, -90105, and -9169 using the inserts from clones FocD33, FocM15, FocN5, FocO2, and FocP18 (Table 2) as probes indicated that the genomes of these isolates, except for isolate Fo-9009, also comprised homologous sequences to the insert DNAs listed above. Isolate Fo-9009 genomic DNA showed no homology to any of the insert DNAs (Table 2). Homologies in nonpathogenic *F. oxysporum* occurred in a smaller number of copies and displayed completely different hybridization patterns compared with those in *F. oxysporum* f. sp. *ciceris* (Fig. 2).

Clones FocL10 and Foc0-12 did not exhibit repetitive copies. They hybridized only to one to four bands in the *Eco*RI-digested DNA of *F. oxysporum* f. sp. *ciceris* isolates, depending upon the race, and to one band in that of nonpathogenic *F. oxysporum* isolates (only clone FocL-10) (Fig. 3). Contrary to the rest of the DNA sequences, the insert from Foc0-12 did not hybridize with any fragments from the *Eco*RI-digested genomic DNA of nonpathogenic *F. oxysporum* isolates. This insert was selected for the identification of *F. oxysporum* f. sp. *ciceris*.


**Sequencing of cloned RAPD markers and design of SCAR primers specific for *F. oxysporum* f. sp. *ciceris* and its pathogenic races 0, 1B/C, 5, and 6.** The complete (clones FocD33, FocL10, FocN5, FocO2, and Foc0-12) or partial (5' and 3' ends; clones FocM15 and FocP18) sequences of insert DNAs were obtained and compared with the GenBank database for the identification of sequence similarity to published gene sequences (1). The insert from clone FocO2 showed 45% nucleotide similarity with a transposase-like protein identified in *F. oxysporum* (28). The inserts from clones FocM15 and FocP18 shared a fragment of identical nucleotide sequence, which was identified as a portion of the *impala* transposon reported in *F. oxysporum* (21) in a BLAST search. Also, a BLAST search showed similarity of the Foc10L sequence to several transport proteins. The complete and partial sequences were deposited in the GenBank nucleotide sequence database (accession numbers, Table 2). From the complete and partial sequences of the clones we designed seven specific primer pairs for the identification of *F. oxysporum* f. sp. *ciceris* (primer pair Foc0-12f/Foc0-12r) and of the pathogenic races 0 (primer pair FocR0-M15f/FocR0-M15r), 1B/C (primer pairs FocR1B/C-N5f/FocR1B/C-N5r and FocR1B/C-D33f/FocR1B/C-D33r), 5 (primer pair FocR5-L10f/FocR5-L10r), and 6 (primer pairs



**Fig. 1.** A, Random amplified polymorphic DNAs (RAPDs) generated by primer OPF-10 using genomic DNA of isolates of *Fusarium oxysporum* f. sp. *ciceris* and of nonpathogenic *F. oxysporum*; and B, hybridization and chemiluminescence detection results using the same gel as in A and the DNA insert FocD33 derived from the cloned race 1B/C-RAPD marker of 1.1 kb as a probe. Numbers on the left are the molecular weights (kb) of the 0.1-kb ladder XIV (Roche Diagnostics) (lane M). Lanes correspond to the following isolates: lanes 1 to 3, race 0 Foc-8207, -9018 JG62, and -7952; lanes 4 to 5, race 1B/C Foc-USA 3-1 JG62, and -1987 W17; lane 6, race 1A Foc-7989; lane 7, race 6 Foc-9166; lanes 8 to 9, race 2 Foc-8605 and -1992R2N; lanes 10 to 11, race 3 Foc-8606 and -1992R3N; lanes 12 to 13, race 4 Foc-8607 and -1992R4N; lanes 14 to 16, race 5 Foc-8012, -9035, and -USA 1-1 JG62; lanes 17 to 18, race 6 Foc-9023 and -Tonini; and lanes 19 to 20, nonpathogenic Fo-9009 and -90105.

FocR6-O2f/FocR6-O2r and FocR6-P18f/FocR6-P18r) by specific PCR assays. Primer designation and sequences, target races, and PCR products under optimized amplification conditions are presented in Table 2.


**Identification of *F. oxysporum* f. sp. *ciceris* and its pathogenic races 0, 1A, 5, and 6 by SCAR primers.** Primers derived from sequences of inserts in clones FocM15, FocN5, FocL10, and FocO2 amplified only a single PCR product from genomic DNA of isolates of each of races 0, 1B/C, 5, and 6, respectively (Fig. 4). None of these products were amplified when DNA of isolates of *F. oxysporum* f. sp. *ciceris* races other than the target races, and of nonpathogenic isolates of *F. oxysporum*, were used as template in specific PCR assays (Fig. 4). Similarly, the primer pair Foc0-12f/Foc0-12r selectively amplified only a single 1,503-bp product from the genomic DNA of a large number of *F. oxysporum* f. sp. *ciceris* isolates representative of all races of the pathogen and of a wide geographic range. This primer pair was designed for the specific identification of the forma specialis *ciceris*. Conversely, SCAR primer pairs based on sequences from the inserts in clones



**Fig. 2.** Southern blot hybridization and chemiluminescence detection results of *EcoRI*-digested genomic DNA of representative isolates of *Fusarium oxysporum* f. sp. *ciceris* and nonpathogenic *F. oxysporum*. Hybridization probe was the insert DNA FocO2 derived from the cloned race 0 random amplified polymorphic DNA marker of 1.0 kb. Numbers on the left are approximate molecular weights (kb). Lanes correspond to the following isolates: lanes 1 to 2, race 0 Foc-9018 PV1 and -T3; lane 3, race 3 Foc-8606; lane 4, race 4 Foc-8607; lanes 5 to 6, race 5 Foc-USA 1-1 JG62 and -USA W6-1; lanes 7 to 9, race 6 Foc-9164, -9023, and -9093 PV1; lanes 10 to 13, nonpathogenic Fo-8250, -9081, -9169, and -420. Lane C is a positive hybridization control (DNA of the corresponding hybridization probe).

FocD33 and FocP18 amplified a single PCR product from all races of *F. oxysporum* f. sp. *ciceris*, although these primers were originally designed for the specific identification of races 1B/C and 6, respectively. This result was unexpected because hybridization assays of RAPD patterns from the eight pathogen races using the above insert DNAs as probes showed hybridization signals only for races 1B/C and 6, and negative results otherwise.

To verify that heterologous amplifications do not occur in the specific PCR assays, specific PCR products from isolates representative of the eight pathogenic races of *F. oxysporum* f. sp. *ciceris*, *F. oxysporum* nonpathogenic to chickpea, other *F. oxysporum* formae speciales, and other *Fusarium* spp. (Table 1) were hybridized using the corresponding insert DNAs as probes. Specific PCR assays using single DNA from isolates of the above and primer pairs FocR0-M15f/FocR0-M15r, FocR1B/C-N5f/FocR1B/C-N5r, and FocR6-O2f/FocR6-O2r yielded the predicted single products only from DNA of races 0, 1B/C, and 6, respectively. Furthermore, only these amplification products yielded a signal in the hybridizations using the corresponding insert DNA as probes. There were no hybridization signals detectable when these later probes were used for hybridization of the putative PCR products from isolates of other *F. oxysporum* f. sp. *ciceris* races, nonpathogenic *F. oxysporum*, other *F. oxysporum* formae speciales, and other *Fusarium* spp. Results were similar with primer pair Foc0-12f/Foc0-12r (Fig. 5). Hybridization of the insert from clone FocL10 to the specific-PCR products yielded by primer pair FocR5-L10f/FocR5-L10r from the same fungal DNAs mentioned above indicated that amplification occurred for isolates of the target race 5 but also for isolates of other races. However, the hybridization signals produced by the nonspecific amplification products were of much less intensity compared with that produced by race 5 isolates. This difficulty was overcome by touchdown PCR (6), which increased the specificity of the amplification to the point that only the race 5-diagnostic amplicon hybridized to the insert from clone FocL10 (Fig. 4B). These results indicated that the SCAR primer pairs designed are effective for the specific identification of *F. oxysporum* f. sp. *ciceris* and of its pathogenic races 0, 1B/C, 5, and 6 by specific- and touchdown-PCR assays. In addition, PCR assays using primers FocR6-P18f



**Fig. 3.** Southern blot hybridization and chemiluminescence detection results of *EcoRI*-digested genomic DNA of representative isolates of *Fusarium oxysporum* f. sp. *ciceris* and nonpathogenic *F. oxysporum*. Hybridization probe was insert DNA FocL10 derived from the cloned race 5 random amplified polymorphic DNA marker of 0.9 kb. Numbers on the left are approximate molecular weights (kb). Lanes correspond to the following isolates: lanes 1 to 2, race 0 Foc-8207 and -82018; lane 3, race 1B/C Foc-1987 W-17; lane 4, race 2 Foc-1992R2N; lane 5, race 4 Foc-1992R4N; lanes 6 to 8, race 1A Foc-7989, -9166, and -9027 PV1; lanes 9 to 13, race 5 Foc-8012, -USA 1-1 JG62, -USA W6-1, -9035, and -9094 JG62; lanes 14 and 16, race 6 Foc-Tonini and -9093 PV1; lanes 15, 17, and 18, nonpathogenic Fo-8250, -9009, and -90105. Lane C is a positive hybridization control (DNA of the corresponding hybridization probe).


and FocR0-M15r yielded a single product of approximately 1,400-bp only from genomic DNA of isolates of *F. oxysporum* f. sp. *ciceris* races R1A and 6 (Fig. 4D). Therefore, a positive result from this assay strategy, together with a negative result from PCR assay using the race 6-specific primers pair FocR6-O2f/FocR6-O2r, will allow the identification of race 1A isolates, albeit in a two-step process.

The specificity and reliability of the SCAR primers developed in this study for the identification of *F. oxysporum* f. sp. *ciceris* and its races 0, 1B/C, 1A, 5, and 6 was further challenged by additional PCR assays using a large collection of *Fusarium* isolates (Table 1). No cross-reactions or amplification of additional fragments were observed for isolates of other *F. oxysporum* formae speciales, other *Fusarium* spp., and nonpathogenic *F. oxysporum*. Furthermore, the specific PCR assays correctly identified isolates of *F. oxysporum* f. sp. *ciceris* and assigned them to the correct pathogenic race. Nevertheless, these PCR assays proved problematic for some of the tested isolates. Primer pair FocR1B/C-N5f/FocR1B/C-N5r yielded a single PCR product of 553-bp from race 1B/C isolates originating from California that was not amplified from isolates of the same race of other geographic origin (i.e., Israel, Syria, Tunisia, or Turkey). Races 2, 3, and 4 were beyond the scope of this work. These three races have been reported only in India (12) and only two isolates of each of them were made available to us. Results from two isolates were not considered appropriate for the aims of our research. Finally, we tested the PCR assays developed in this study using a range of DNA from 200 to 10<sup>-5</sup> ng to determine the minimum amount of

fungal genomic DNA that could be detected. Every primer pair designed in this study detected a minimum amount of 0.1 ng of fungal DNA (Fig. 6).

## DISCUSSION

Phenotypic and genetic characterization of *F. oxysporum* f. sp. *ciceris* races is important for the efficient management of Fusarium wilt through use of resistant cultivars in chickpea-growing areas. This study demonstrated that SCAR primers developed from RAPD markers (17) can be used to unambiguously identify *F. oxysporum* f. sp. *ciceris* races 0, 1A, 5, and 6, which are prevalent in the Mediterranean Basin (17). Also, these SCAR primers proved useful in discriminating *F. oxysporum* f. sp. *ciceris* from other diverse formae speciales of this species, other *Fusarium* spp., and nonpathogenic *F. oxysporum*. The SCAR primer pairs amplified a single diagnostic PCR product from all isolates of the target forma specialis and races irrespective of the geographic origin. This is the first report of PCR-based identification of *F. oxysporum* f. sp. *ciceris*. Although races 0, 5, and 6 can be characterized by the RAPD markers previously reported (17), the SCAR primers and PCR assays are free of problems associated with RAPD assays that reduce its applicability for diagnosis of those races and enhance the possibility of fast, extensive, and reliable discrimination of the pathogen and races. An additional benefit from the present study concerns the molecular identification of race 1A of the pathogen, for which no RAPD marker was found previously (17). This can be achieved through two independent



**Fig. 4.** Agarose gels showing amplification products from polymerase chain reaction (PCR) using genomic DNA from isolates of *Fusarium oxysporum* f. sp. *ciceris* and of nonpathogenic *F. oxysporum* and sequence-characterized amplified region (SCAR) primers designed for races **A**, **0**, **B**, **5**, and **C**, **6**. **D**, PCR using primer pair FocR6-P18f/FocR0-M15r, which amplifies a 1.4-kb product from genomic DNA of race 1A and race 6 isolates. Numbers on the left are the molecular weights (kb) of the 0.1-kb ladder XIV (Roche Diagnostics) (lane M). Lanes correspond to the following isolates: **A** and **C**, lanes 1 to 3, race 0 Foc-7802, -82113, and -91108; lanes 4 to 5, race 1B/C Foc-USA 3-1 JG62 and -1987 W17; lane 6, race 2 Foc-8605; lane 7, race 3 Foc-8606; lane 8, race 4 Foc-8607; lane 9, race 1A Foc-7989; lanes 10 to 14, race 6 Foc-9166, -9164, -Tonini, -9093 JG62, and -9023; lanes 15 to 16, race 5 Foc-8012 and -USA14201; lanes 17 to 19, nonpathogenic Fo-90101, -90105, and -9169; **B**, lanes 1 to 3, race 0 Foc-7802, -82113, and -91108; lanes 4 to 5, race 1B/C Foc-USA 3-1 JG62, and -1987 W17; lane 6, race 2 Foc-8605; lane 7, race 3 Foc-8606; lane 8, race 4 Foc-8607; lane 9, race 1A Foc-7989; lanes 10 to 12, race 6 Foc-9166, -Tonini, and -9023; lanes 13 to 17, race 5 Foc-8012, -9035, -USA W6-1, USA 1-1 JG62, and -USA14201; lanes 18 to 19, nonpathogenic Fo-90101 and -9081; **D**, lanes 1 to 2, race 1A isolates Foc-7989 and -9168; lanes 3 to 12, race 6 Foc-9164, -9166, -9027 PV1, -Tonini, -9093 JG62, -9023, -9170, -8272, -8905, and -8924; lane 13, race 0 Foc-7802; lane 14, race 1B/C Foc-USA 3-1 JG62; lane 15, race 2 Foc-8605; lane 16, race 3 Foc-8606; lane 17, race 4 Foc-8607; lane 18, race 5 Foc-8012; and lane 19, nonpathogenic Fo-90105.



PCR assays, one using the SCAR primers which amplified a single PCR product from races 1A and 6, and another assay using the race 6-specific primers. Races 1A and 6 showed close virulence patterns on chickpea differentials (11,15) and high genetic similarities by RAPD analyses (17). This could explain the occurrence of a common specific PCR marker.

An unexpected, though interesting result in our study concerns the SCAR primer pair FocN5-R1B/Cf/FocN5-R1B/Cr. These primers were developed for the identification of race 1B/C and amplified a single PCR product from race 1B/C isolates from California but not from those from other geographic origins (i.e., Israel, Syria, Tunisia, and Turkey). Races 0 and 1B/C induce the yellowing syndrome in chickpea and share high genetic similarity, as indicated by cluster analysis of RAPD amplifications (17). The limitations of pathogenicity assays may have led to misclassification of race 0 and race 1B/C isolates (19). Therefore, additional work, including new pathogenicity tests, must be done to clarify the race identity of the above isolates as well as to ascertain the possibility that significant genetic diversity may exist among isolates of race 1B/C from different geographic origin.


Our study illustrates the need for extreme caution in the process of selecting the appropriate cloned DNA fragments, a crucial step in the design of race-specific SCAR primers from RAPD markers. The methodology used had a critical impact on the possibility of achieving our research goals; therefore, it deserves special men-

tion. Restriction analyses of a large number of DNA inserts derived from each cloning experiment showed nonhomologous DNA sequences within each RAPD marker. Thereafter, hybridization of RAPD profiles using each resulting insert as a probe showed that many sequences also were amplified in pathogen races other than the target race, which allowed us to identify clones containing the race-specific DNA inserts and discard the nonspecific ones. Had these precautions not been taken, inadequate sequences for primer design might have been chosen, which would have resulted in the development of nonspecific primers and PCR assays.


It often is assumed that DNA fragments of the same molecular size amplified in RAPD assays represent homologous sequences. This assumption is not necessarily correct and homology of RAPD bands needs to be demonstrated using Southern analyses (32). Using this procedure, we herein verified that the RAPD bands amplified from isolates of the target form *specialis* or belonging to the same race were homologous. This strengthens the suitability of our approach for developing race-specific SCAR primers and PCR assays from informative RAPD fragments.

Repetitive DNA sequences have been used before to successfully identify subpopulations within fungal species, including *F. oxysporum* (4,7,25). The efficiency of such repetitive sequences for generating SCAR markers has been demonstrated previously for the detection of *Peronospora tabacina* by specific PCR (36). Although sequences derived from the RAPD marker of a particular *F. oxysporum* f. sp. *ciceris* race were present in a high-copy number in the genome of all the races of the pathogen, we were able to design SCAR primers that yielded race-specific PCR products. Thus, the single 900- and 1,000-bp products of primer pairs FocR0-M15f/FocR0-M15r and FocR6-O2f/FocR6-O2r that identify races 0 and 6, respectively, were proven specific by hybridization experiments of the blotted PCR products. Primer pair FocR5-L10f/FocR5-L10r was based on a single- or low-copy DNA fragment. These primers yielded a single 900-bp product that was diagnostic for race 5. Although this product also was amplified from other *F. oxysporum* f. sp. *ciceris* races, that was to a much lesser extent because it could be detectable only by hybridization of the PCR products. A touchdown-PCR procedure significantly improved the intensity of the amplification signal and the specificity of this assay, making it suitable for the selective identification of race 5 (6).

That DNA sequences from all RAPD markers studied herein were present in the genome of the eight pathogen races, irrespective of geographic origin, was not surprising. Previous studies showed that races of *F. oxysporum* f. sp. *ciceris* share the same mitochondrial DNA restriction fragment length polymorphism



**Fig. 5. A**, Agarose gel showing the 1.5-kb product diagnostic for *Fusarium oxysporum* f. sp. *ciceris* using genomic DNA from *F. oxysporum* f. sp. *ciceris* isolates and isolates of nonpathogenic *F. oxysporum*, and sequence-characterized amplified region (SCAR) primer pair Foc0-12f/Foc0-12r; and **B**, hybridization and chemiluminescence detection results using the same gel as in **A** and the insert DNA Foc0-12 derived from the cloned random amplified polymorphic DNA marker of 1.5 kb. Numbers on the left are the molecular weights (kb) of the 0.1-kb ladder XIV (Roche Diagnostics) (lane M). Lanes 1 to 8 correspond to *F. oxysporum* f. sp. *ciceris* isolates Foc-7802 (race 0), -USA 3-1 JG62 (race 1B/C), -7989 (race 1A), -8605 (race 2), -8606 (race 3), -8607 (race 4), -8012 (race 5), and -9023 (race 6). Lanes 9 to 12 correspond to nonpathogenic *F. oxysporum* isolates Fo-8250, -9081, -90105, and -506. Lanes 13 and 14 correspond to isolates *F. oxysporum* f. sp. *melonis* 9016 and *F. oxysporum* f. sp. *niveum* 8805. Lanes 15 to 19 correspond to isolates *F. acuminatum* 11442, *F. avenaceum* 11440, *F. lactis* 10757, *F. proliferatum* 11558, and *F. solani* 11420.



**Fig. 6.** Agarose gel showing the sensitivity of polymerase chain reaction (PCR) using genomic DNA of *Fusarium oxysporum* f. sp. *ciceris* and sequence-characterized amplified region (SCAR) primer pair Foc0-12f/Foc0-12r: Amplification of decreasing amount of race 0 isolate Foc-7802 DNA ranging from 200 to  $10^{-5}$  ng. Numbers on the left correspond to the molecular weights (kb) of the 0.1-kb ladder XIV (Roche Diagnostics) (lane M). Lanes CR0 and CNP, amplification controls for race 0 isolate Foc-7802 and nonpathogenic *F. oxysporum* isolate Fo-90101 DNAs, respectively. Lane C, control reaction with no template DNA.



pattern (30) and belong to a single vegetative compatibility group (27). Additionally, sequence analyses of five nuclear genes and further gene genealogy analysis of the translation elongation factor 1 $\alpha$  indicated a monophyletic origin of *F. oxysporum* f. sp. *ciceris* (16). Therefore, isolates of this strictly asexually reproducing fungus probably derived from a small founder population or single individual that acquired pathogenicity to *Cicer* spp., and populations that further developed likely comprise clonal individuals that share a common genetic background (9).

Numerous studies have demonstrated the utility of RAPD methodology for correlating molecular markers to pathogenic races in *F. oxysporum* (2,10,22). However, the number of studies that subsequently have converted race-associated RAPD markers into SCARs and have developed specific primers for race identification is more limited (5). Our results proved that this strategy is highly efficient, for it allowed the unambiguous differentiation of races within a monophyletic forma specialis of *F. oxysporum*, as well as the discrimination of the forma specialis itself from other *F. oxysporum* and *Fusarium* spp. Also, our results have important practical applications for diagnosis, epidemiology, and management of Fusarium wilt of chickpea. Thus, the SCAR primers specific for *F. oxysporum* f. sp. *ciceris* provide a useful tool for easily discriminating nonpathogenic *F. oxysporum* isolates that can be recovered from virus-infected chickpea (18,34), as well as from Fusarium wilt-affected chickpea (34). Similarly, the race-specific SCAR primers provide an attractive alternative for the analysis of race structure in *F. oxysporum* f. sp. *ciceris* populations compared with the use of pathogenicity tests. The race-specific PCR products of these SCAR primers should facilitate the early detection of introduced races, as well as detect changes in the relative race frequencies that may occur as a response to the use of resistant cultivars. Finally, all SCAR primer pairs developed were able to detect 0.1 ng of fungal DNA using the amplification conditions described herein. Such amounts of fungal DNA can be obtained easily from several natural substrates that harbor the pathogen. Relatively straightforward adaptations of this technique could be envisioned for the detection of races of *F. oxysporum* f. sp. *ciceris* in infested soil or in infected plants (20), which will find additional practical applications in the epidemiology and management of Fusarium wilt of chickpea.

## ACKNOWLEDGMENTS

This research was supported by grants AGF97-1479 and AGF98-0878 from Comisión Interministerial de Ciencia y Tecnología (CICYT) of Spain. We thank G. Contreras for technical support; C. Akem, W. Boge, D. Fravel, M. H. Halila, J. Katan, and J. F. Leslie for providing fungal isolates; and E. Manias for critical reading of the manuscript.

## LITERATURE CITED

- Altschul, S. F., Madden, T. L., Schäffer, A. A., Zhang, J., Zhang, Z., Miller, W., and Lipman, D. J. 1997. Gapped BLAST and PSI-BLAST: A new generation of protein database search programs. *Nucleic Acids Res.* 25:3389-3402.
- Assigbetse, K. B., Fernandez, D., Dubois, M. P., and Geiger, J. P. 1994. Differentiation of *Fusarium oxysporum* f. sp. *vasinfectum* races on cotton by random amplified polymorphic DNA (RAPD) analysis. *Phytopathology* 84:622-626.
- Bunting, T. E., Plumley, K. A., Clarke, B. B., and Hillman, B. I. 1996. Identification of *Magnaporthe poae* by PCR and examination of its relationship to other fungi by analysis of their nuclear rDNA ITS-1 regions. *Phytopathology* 86:398-404.
- Chiocchetti, A., Bernardo, I., Daboussi, M. J., Garibaldi, A., Gullino, M. L., Langin, T., and Migheli, Q. 1999. Detection of *Fusarium oxysporum* f. sp. *dianthi* in carnation tissue by PCR amplification of transposon insertions. *Phytopathology* 89:1169-1175.
- de Haan, L. A. M., Numansen, A., Roebroek, E. J. A., and van Doorn, J. 2000. PCR detection of *Fusarium oxysporum* f. sp. *gladioli* race 1, causal agent of *Gladiolus* yellows disease, from infected corms. *Plant Pathol.* 49:89-100.
- Don, R. H., Cox, P. T., Wainwright, B. J., Baker, K., and Mattick, J. S. 1991. 'Touch-down' PCR to circumvent spurious priming during gene amplification. *Nucleic Acids Res.* 19:4008.
- Edel, V., Steinberg, C., Avelange, I., Laguerre, G., and Alabouvette, C. 1995. Comparison of three methods for the characterization of *Fusarium oxysporum* strains. *Phytopathology* 85:579-585.
- Goodwin, S. B., Drench, A., and Fry, W. E. 1992. Cloning and genetic analysis of two highly polymorphic, moderately repetitive nuclear DNAs from *Phytophthora infestans*. *Curr. Genet.* 22:107-115.
- Gordon, T. R., and Martyn, R. D. 1997. The evolutionary biology of *Fusarium oxysporum*. *Annu. Rev. Phytopathol.* 35:111-128.
- Grajal-Martín, M. J., Simon, C. J., and Muehlbauer, F. J. 1993. Use of random amplified polymorphic DNA (RAPD) to characterize race 2 of *Fusarium oxysporum* f. sp. *pisi*. *Phytopathology* 83:612-614.
- Halila, H. M., and Strange, R. N. 1996. Identification of the causal agent of wilt of chickpea in Tunisia as *Fusarium oxysporum* f. sp. *ciceris* race 0. *Phytopathol. Mediterr.* 35:67-74.
- Haware, M. P., and Nene, Y. L. 1982. Races of *Fusarium oxysporum* f. sp. *ciceris*. *Plant Dis.* 66:809-810.
- Henson, J. M., and French, R. 1993. The polymerase chain reaction and plant disease diagnosis. *Annu. Rev. Phytopathol.* 31:81-109.
- Jalali, B. L., and Chand, H. 1992. Chickpea wilt. Pages 420-444 in: *Plant Diseases of International Importance. Vol. 1. Diseases of Cereals and Pulses.* U. S. Singh, A. N. Mukhopadhyay, J. Kumar, and H. S. Chaube, eds. Prentice Hall, Englewood Cliffs, NJ.
- Jiménez-Díaz, R. M., Alcalá-Jiménez, A. R., Hervás, A., and Traperocas, J. L. 1993. Pathogenic variability and hosts resistance in the *Fusarium oxysporum* f. sp. *ciceris/Cicer arietinum* pathosystem. Pages 87-94 in: *Proc. 3rd Eur. Semin. Fusarium Mycotoxins, Taxonomy, Pathogenicity and Host Resistance.* Hodowska Róslin Aklimatyazacja i Nasiennictwo. Plant Breeding and Acclimatization Institute, Radzików, Poland.
- Jiménez-Gasco, M. M., Milgroom, M. G., and Jiménez-Díaz, R. M. 2002. Gene genealogies support *Fusarium oxysporum* f. sp. *ciceris* as a monophyletic group. *Plant Pathol.* 51:72-77.
- Jiménez-Gasco, M. M., Pérez-Artés, E., and Jiménez-Díaz, R. M. 2001. Identification of pathogenic races 0, 1B/C, 5, and 6 of *Fusarium oxysporum* f. sp. *ciceris* with random amplified polymorphic DNA (RAPD). *Eur. J. Plant Pathol.* 107:237-248.
- Kaiser, W. J., Klein, R. E., Larsen, R. C., and Wyatt, S. D. 1993. Chickpea wilt incited by pea streak carlavirus. *Plant Dis.* 77:922-926.
- Kelly, A. G., Alcalá-Jiménez, A. R., Bainbridge, B. W., Heale, J. B., Pérez-Artés, E., and Jiménez-Díaz, R. M. 1994. Use of genetic fingerprinting and random amplified polymorphic DNA to characterize pathotypes of *Fusarium oxysporum* f. sp. *ciceris* infecting chickpea. *Phytopathology* 84:1293-1298.
- Kelly, A. G., Bainbridge, B. W., Heale, J. B., Pérez-Artés, E., and Jiménez-Díaz, R. M. 1998. In planta-polymerase-chain-reaction detection of the wilt-inducing pathotype of *Fusarium oxysporum* f. sp. *ciceris* in chickpea (*Cicer arietinum* L.). *Physiol. Mol. Plant Pathol.* 52:397-409.
- Langin, T., Capy, P., and Daboussi, M. J. 1995. The transposable element *impala*, a fungal member of the *Tc1-mariner* superfamily. *Mol. Gen. Genet.* 246:19-28.
- Manulis, S., Kogan, N., Reuven, M., and Ben Yehet, Y. 1994. Use of the RAPD technique for identification of *Fusarium oxysporum* f. sp. *dianthi* from carnation. *Phytopathology* 84:98-101.
- Martin, R. R., James, D., and Levensque, C. A. 2000. Impacts of molecular diagnostic technologies on plant disease management. *Annu. Rev. Phytopathol.* 38:207-239.
- McDermott, J. M., Brändle, U., Dutly, F., Haemmerli, U. A., Keller, S., Müller, K. E., and Wolfe, M. S. 1994. Genetic variation in powdery mildew of barley: Development of RAPD, SCAR and VNTR markers. *Phytopathology* 84:1316-1321.
- Namiki, F., Shiomi, T., Kayamura, T., and Tsuge, T. 1994. Characterization of the formae speciales of *Fusarium oxysporum* causing wilts of cucurbits by DNA fingerprinting with nuclear repetitive DNA sequences. *Appl. Environ. Microbiol.* 60:2684-2691.
- Nelson, P. E., Tousson, T. A., and Cook, R. J. 1981. *Fusarium: Diseases, Biology and Taxonomy.* The Pennsylvania University Press, University Park.
- Nogales-Moncada, A. M. 1997. Compatibilidad Vegetativa en *Fusarium oxysporum* f. sp. *ciceris* y *Fusarium oxysporum* f. sp. *melonis*, Agentes, Respectivamente de las Fusariosis Vasculares del Garbanzo y del Melón. Ph. D. Thesis. University of Córdoba, Córdoba, Spain.
- Okuda, M., Ikeda, K., Namiki, F., Nishi, K., and Tsuge, T. 1998. *Tfo1*: An Ac-like transposon from the plant pathogenic fungus *Fusarium oxysporum*. *Mol. Gen. Genet.* 258:599-607.
- Paran, I., and Michelmore, R. W. 1993. Development of reliable PCR-

- based markers linked to downy mildew resistance genes in lettuce. *Theor. Appl. Genet.* 85:985-993.
30. Pérez-Artés, E., Roncero, M. I. G., and Jiménez-Díaz, R. M. 1995. Restriction fragment length polymorphism analysis of the mitochondrial DNA of *Fusarium oxysporum* f. sp. *ciceris*. *J. Phytopathol.* 143:105-109.
  31. Raeder, U., and Broda, P. 1985. Rapid DNA preparation from filamentous fungi. *Lett. Appl. Microbiol.* 1:17-20.
  32. Rieseberg, L. H. 1996. Homology among RAPD fragments in interspecific comparisons. *Mol. Ecol.* 5:99-105.
  33. Sambrook, J., Fritsch, E. F., and Maniatis, T. 1989. *Molecular Cloning: A Laboratory Manual*, 2nd ed. Cold Spring Harbor Laboratory, Cold Spring Harbor, NY.
  34. Trapero-Casas, A., and Jiménez-Díaz, R. M. 1985. Fungal wilt and root rot diseases of chickpea in southern Spain. *Phytopathology* 75:1146-1151.
  35. Ward, E. 1994. Use of polymerase chain reaction for identifying plant pathogens. Page 143-160 in: *Ecology of Plant Pathogens*. J. P. Blakeman and B. Williamson, eds. CAB International, Oxon, UK.
  36. Wigglesworth, M. D., Nesmith, W. C., Schardl, C. L., Li, D., and Siegel, M. R. 1994. Use of specific repetitive sequences in *Peronospora tabacina* for early detection of the tobacco blue mold pathogen. *Phytopathology* 84:425-430.