



Tehran University of Medical
Sciences Publication
<http://tums.ac.ir>

Iran J Parasitol

Open access Journal at
<http://ijpa.tums.ac.ir>



Iranian Society of Parasitology
<http://isp.tums.ac.ir>

Original Article

Development of an Indirect ELISA Using Different Fragments of Recombinant Ncgra7 for Detection of *Neospora caninum* Infection in Cattle and Water Buffalo

*Hossein HAMIDINEJAT¹, Massoud Reza SEIFI ABAD SHAPOURI¹, Mohammad Mehdi NAMAVARI², Parviz SHAYAN³, Marzieh KEFAYAT⁴

1. Dept. of Pathobiology, Veterinary Faculty, Shahid Chamran University of Ahvaz, Ahvaz, Iran
2. Razi Vaccine and Serum Research Institute, Shiraz Branch, Shiraz, Iran
3. Dept. of Parasitology, Veterinary Faculty, University of Tehran, Tehran, Iran
4. Dept. of Parasitology, Shahid Chamran University of Ahvaz, Ahvaz, Iran

Received 09 Jul 2014
Accepted 21 Nov 2014

Keywords:

Neospora caninum,
ELISA,
NcGRA7,
Recombinant antigen,
Cattle,
Water buffalo,
Iran

***Correspondence**

Email:

hamidinejat@yahoo.com

Abstract

Background: Dense granules are immunodominant proteins for the standardization of immunodiagnostic procedures to detect neosporosis. In the presented study different fragment of a dense-granule protein was evaluated for serodiagnosis of *Neospora caninum* in cattle and water buffalo.

Methods: NcGRA7, from *N. caninum* tachyzoites was amplified. PCR product and pMAL-c2X plasmid were digested with *EcoRI* restriction enzyme and expressed in *Escherichia coli* to evaluate its competence for detection of anti- *N. caninum* antibodies with ELISA in comparison with commercial IDEXX ELISA. Furthermore, 230 sera of presumably healthy cattle and water buffaloes (108 cattle and 122 water buffaloes) were analyzed by both tests to determine the agreement of these two procedures.

Results: Sensitivities and specificities of NcGRA7-based ELISA were 94.64% and 90.38% respectively using sera of cattle, but were 98.57% and 86.54% in the case of buffaloes respectively. A good correlation between the results of IDEXX ELISA and ELISA based on recombinant NcGRA7 for detecting *N. caninum* antibodies was appeared. Analyzing by Mc Nemar's showed that NcGRA7-based ELISA has acceptable capability to differentiate the positive results in comparison with IDEXX ELISA.

Conclusion: NcGRA7-based ELISA considering utilized new fragment of genomic DNA is a good tool for serodiagnosis of anti- *N. caninum* antibodies for screening and epidemiological purposes on cattle herd and water buffaloes as well.

Introduction

Cyst-forming coccidian *Neospora caninum* (Apicomplexa: Sarcocystidae) is an important intracellular protozoan parasite and causes neosporosis that related to congenital birth defects in cattle and water buffalo (*Bubalus bubalis*) worldwide (1, 2). The parasite was first recognized in 1984 from Norwegian dogs when Bjerkas et al. reported unidentified protozoan parasites belong to sporozoa (3).

Confirmed definitive hosts of *N. caninum* are dogs and coyotes, but other wild canids may play a role as definitive hosts. Multitude of mammalian species including cattle, water buffalo, sheep, goat, deer, horse and canids are defined as intermediate hosts of the *N. caninum*, however researches revealed that the range of these hosts is actually numerous (4-7). This infection can cause serious problems especially when occur simultaneously with other infectious diseases. For instance, 9.4% of dogs in Meshkin-Shahr, which infected with *N. caninum* had afflicted with *Leishmania infantum* infection (8). Transmission of *N. caninum* to intermediate hosts occurs either by ingestion of sporulated oocysts via contaminated food and water or vertically from mother to fetus through the placenta (5-9). Abortion and other reproductive failures are the main features of the infection in adult cows and water buffaloes. Abortions originated from *N. caninum* can occur at any time of gestation but the majority of abortions are at 5-6 months of gestation and this situation always leads to serious economic losses (2, 4, 9, 10).

Serologic procedures are routinely and widely used among the available diagnostic tests for detection of *N. caninum* infection. Currently, serologic assays have been developed to determine the anti-*N. caninum* antibodies in cattle including immunoblotting, the indirect fluorescent antibody test (IFAT), direct modified agglutination test (MAT) and ELISA (4,11-14).

For these purposes, using the recombinant immunodominant proteins, for instance surface NcSAG1, NcSRS2 and NcSAG4 bradyzoite stage-specific proteins and the dense-granule protein NcGRA7 of *N. caninum* have been reported as helpful immunogenic targets in the literature (15-20).

Most serodiagnostic tools such as *N. caninum* commercial kits utilize antigens from whole and or sonicated tachyzoites, which always burden production costs. On the other hand, cross-reactions with prevalent and related parasite, *Toxoplasma gondii*, and even existence of host materials affect the specificity of the tests (13, 14, 21). Recombinant antigen based procedures could overcome these drawbacks and it seems that dense-granule antigen, GRA7, which present by both tachyzoite and bradyzoite stages of *N. caninum* has a good competence to detect antibodies in early phase of infection (18, 22, 23).

Water buffaloes (*Bubalus bubalis*) as well as cattle are economically important animals in tropical areas and little is known about the applied epidemiological tools to investigate or to screen of neosporosis in these animals. Hence, we aimed to establish an ELISA based upon different fragments of recombinant NcGRA7 protein in an efficient prokaryotic expression system for diagnosis of *N. caninum* infection in cattle and water buffaloes as well.

Materials and Methods

Parasite

N. caninum tachyzoites, Nc-1 strain, were grown in monolayers of Vero cells cultured in RPMI 1640 medium (Sigma Co., USA) fortified by 2% of fetal calf serum, penicillin (10,000 U/ml), streptomycin (100 mg/ml) and amphotericin B (25 mg/ml) at 37 °C with 5% CO₂ in tissue culture flasks. Tachyzoites were harvested by scraping the monolayer when 80% of the cells were infected and then purified by successive passage of the cell mono-

layer through a 27-gauge needle and centrifuged to remove host cell debris. The supernatant was collected, centrifuged and washed twice in phosphate-buffered saline afterward (PBS, pH 7.2) (15).

DNA extraction and PCR

Tachyzoites were prepared at a concentration of 1×10^8 / ml and extraction of genomic DNA was carried out by commercial kit (Qiagen) according to the kit protocol. The DNA sequences were obtained from the GenBank database and the accession number—U82229 used to design the required primers. Designed and applied forward and reverse primers were 5'- GCGAGAAATTCGCTGGAGACTTGG-3' and 5'-GCGAGAAATTCCTATTCGGTGT-3' respectively. PCR was carried out in a total volume of 50 μ l, containing 5 μ l of DNA template, 1 μ l of each primer at 20 pmol, 1 μ l of 10 mM dNTPs mix, 5 μ l of 109 PFU PCR buffer, 1.5 μ l of 50 mM MgCl₂ and 5u PFU DNA polymerase (BioNeer, Korea). PCR condition was as follow: pre denaturation at 94°C for 5 min, denaturation at 94 °C for 1 min, annealing at 51°C for 1 min, extension at 72 °C for 90s (35 repeats) and final extension at 72°C for 7 min in a thermal cycler. The amplified DNA was visualized on 1.0% agarose gel stained with cyber safe then sequence analysis of PCR product was followed up.

Cloning and expression of NcGRA7

PCR product was purified from gel with the gel extraction kit (Vivantis, Malaysia). The purified PCR product and pMAL-c2X plasmid (New England Biolabs, USA) were digested with *EcoR1* restriction enzymes (Fermentase, Latvian) at 37 °C for 2.5 h and were extracted from agarose gel using the gel extraction kit. Ligation process was applied on digested DNA fragment and plasmid vector at 22 °C for 2 h by T4 DNA ligase (Fermentas, Latvian), followed by overnight incubation at 4 °C. *Escherichia coli* host strain TG1 was transformed by the ligation product using Chung transforming method (24). Transformed colo-

nies appeared on the selective Luria–Bertani (LB) agar, were grown in LB broth containing ampicillin. Extraction of plasmid was performed by Vivantis plasmid extraction kit and presence of GRA7 inserts were confirmed by restriction digestion of recombinant plasmids with *EcoR1*.

Western blotting

After SDS-PAGE, the protein bands were transferred onto nitrocellulose membrane. The membrane was blocked with 0.2% Tween 20 in PBS (137 mM NaCl, 2.7 mM KCl, 4.3 mM Na₂PO₄, 1.4 mM KH₂PO₄ plus 0.05% Tween) at ambient temperature for 2 h. The blots were washed three times with PBST (0.05% Tween 20 in PBS) and incubated with 1/100 dilution of positive bovine serum for 1 h. The membranes washed three times with PBST and incubated with 1/1000 dilution of peroxidase conjugated with rabbit anti-bovine IgG (Sigma, USA) for 1 h. The membranes were washed with PBST then by PBS and soaked in 4-chloro-1-naphthol and hydrogen peroxide afterward (H₂O₂).

Serum samples

Negative and positive sera of cattle and water buffaloes confirmed by two commercial ELISA kits (IDEXX and ID VET, France) and *N. caninum* agglutination test (NAT) as well. Blood samples were also collected utilizing venoject tubes from jugular vein of 230 apparently healthy cattle and water buffaloes from Ahvaz district (108 cattle and 122 water buffaloes). Collected blood samples were centrifuged at 1000×g and the supernatants were frozen at -20°C until the examinations were performed.

ELISA

Checkerboard procedure was conducted to set up the NcGRA7-based ELISA. The cut-off point calculated as average of optical density of negatives plus two standard deviations. Finally, polystyrene microtitre plates (kariz Mehr, Iran) were coated with 0.1 μ g/ μ l of re-

combinant NcGRA7 diluted in 100 µl of 0.1 M carbonate buffer (pH=9.6) as coating buffer and incubated overnight at 4°C. After three times washing with PBS-T (PBS containing 0.05% Tween 20), wells were blocked for 2.5 hours at 37°C using PBS containing 0.2% Tween 20 and washed as mentioned. The sera, negative and positive controls were diluted 1:400 in blocking buffer, added to each well at a volume of 100 µl and incubated at ambient temperature for 30 minutes and washed subsequently. Thereafter, 100 µl/well of horseradish anti-bovine peroxidase conjugate (Sigma, USA) diluted 1:10000 in blocking buffer was added at room temperature for 1 hour. Plates were washed and enzyme activities were revealed by 10 minutes incubation of 100 µl/well with tetramethylbenzidine (TMB) (Sigma, USA) and 3% H₂O₂ as substrate and chromogen. Reactions were stopped with 0.1 M of HCL and at last the plates were read with microtiter plate reader (Tadjhiz Afzar Teb, Iran) at wavelength of 450 nm. For each test a well was coated with purified MBP to cover false reactions of positive sera with this protein.

Due to appearing the existence of cross-reaction with closely related parasite, *T. gondii*, all sera were tested for the presence of *T. gondii* antibodies utilizing the agglutination tests based on the direct agglutination of fixed parasites with sera pre-treated with 2-mercaptoethanol to prevent non-specific IgM agglutination, as described by Dubey and Desmonts (21).

Statistical analyzes

The Kappa statistic test (κ) was used to test the level of agreement between the IDEXX ELISA and ELISA based on recombinant NcGRA7 for detection of *N. caninum* infection in cattle and water buffaloes. Kappa and its 95% Confidence Interval (CI), was used further to measure the degree of agreement between the procedures after taking into account the probability of agreement by chance alone. Strength of agreement based on κ was judged

according to the following guidelines: <0.2=slight; 0.2–0.4=fair; 0.4– 0.6=moderate; 0.6–0.8=good; >0.8=very good. Mc nemar's test was also utilized for comparing the percentage of positive reactions (25).

Results

Cloning, expression and purification of the recombinant NcGRA7

Utilizing mentioned primers, expected PCR product of 600 bp was amplified (Fig.1) and its integrity was assured by sequencing.

Sixty clones appeared on LB agar after expression and subsequent transformation. Screening was done with digesting plasmids by *EcoR1* enzyme. Finally, three clones carried the recombinant plasmid and expressed the MBPNcGRA7 protein. Expression of a protein of about 66 kDa, corresponding to predicted molecular weight of MBPNcGRA7 (GRA7:24 kDa and MBP: 42 kDa) was recognized by SDS-PAGE (Fig.2).

Cloning and expression of NcGRA7 in *E. coli* confirmed with reaction of expressed protein in Western blotting (Fig.3) with a *N. caninum* positive serum.

Serology

The mean value of ELISA OD of the negative sera plus 2 standard deviations gave a cut-off point of 0.355 and 0.384 for determination of existing *N. caninum* specific antibodies in cattle and water buffaloes respectively. Compared with IDEXX ELISA, relative sensitivities and specificities of NcGRA7-based ELISA were 94.64% and 90.38% respectively using sera of cattle, but were 98.57% and 86.54 in the case of buffaloes respectively (Table 1). There was no cross-reaction with related apicomplexan parasite, *T. gondii*.

Kappa statistical test showed a good correlation between the results of IDEXX ELISA and ELISA based on recombinant NcGRA7 for detecting *N. caninum* antibodies in cattle ($\kappa= 0.85$), and also water buffaloes ($\kappa= 0.83$).




Fig.1: The PCR products revealing 600 bp bands




Fig.2: SDS-PAGE electrophoresis on 10% polyacrylamide gel. 1 *E. coli* strain TG1 containing pMAL-c2X as control after induction. 2 *E. coli* extracts expression of pMAL-c2X-GRA7




Fig.3: Western blotting analysis. Lane 1 reveals marker protein and lane 2 reveals the reactivity positive serum with bacteria expressing MBP and MBP-GRA7

Furthermore, Mc Nemar's analyzing revealed that NcGRA7-based ELISA has acceptable competence to discriminate the posi-

tive results in comparison with IDEXX ELISA ($P > 0.05$).

Table1: The sensitivity and specificity of recombinant NcGRA7 ELISA in comparison with IDEXX ELISA

IDEXX ELISA	ELISA(Buffalo)		ELISA(Cattle)		
	Positive	Negative	Positive	Negative	
Positive	69 ^(a)	7 ^(b)	Positive	53 ^(a)	5 ^(b)
Negative	1 ^(c)	45 ^(d)	Negative	3 ^(c)	47 ^(d)
Total	70 ^(a+c)	52 ^(b+d)	Total	56 ^(a+c)	52 ^(b+d)
Sensitivity%	98.57		94.64		
Specificity%	86.54		90.38		

Sensitivity = $a / (a + c)$; Specificity = $d / (b + d)$

Discussion

Neosporosis is one of the serious economic problems in dairy and beef industry worldwide. To date, many researchers have studied on different aspects of *N. caninum* and many studies were reported the prevalence of bovine infection with *N. caninum*. The precise and fast diagnosis of *N. caninum* infections has always a key role on interpreting the prevalence and distribution patterns and subsequently prediction from transmission especially in its vertical form. Since, there is no practical effective vaccine or treatment for neosporosis (26), this ideal goal will lead us to preclude the abortions because of this disease.

On the other point of view, it is also necessary to finding definitive markers detecting specific antibodies of *N. caninum* and decrease cross-reaction with closely related parasite, *Toxoplasma gondii* (4, 27).

Succeeding the evaluation of the first serological test, the IFAT, for detection of antibodies to *N. caninum* (28), numerous ELISA procedures have been defined by many authors applying whole or recombinant antigens (29, 30). In the present study, potential of one recombinantly produced antigen was evaluated for the first time in water buffalo.

Recombinant antigens of *N. caninum* revealed appropriate for improving diagnostic purposes that can be used instead of native antigens, which often have inconstant and unreliable quality (31). Secretary proteins like micronemes, rhoptries, and dense granules released in early stages of infection with *N. caninum*,

just as adhesion onto the host cell surface directly depends on the secretion of these substances from the sporozoites of the parasite. Moreover, dense granules also discharge their content into the parasitophorous vacuole throughout the intracellular stage (32). We predict that these purified and high quality proteins can alternatively detect the early infections when they utilize as antigen in ELISA.

NcGRA7 is an immunodominant antigen in both of tachyzoite and bradyzoite stages (33). Therefore, it sounds applying the recombinant NcGRA7 is a suited and helpful marker for both active and chronic diagnosis of *N. caninum* infection and can serve as a potential vaccine candidate (34). This issue is more considerable when the competence of NcGRA7 for demonstration of infection with *N. caninum* in early stages of parasite life cycle has been reported in bovines (34, 35).

In the present study, we intended to clone and express the NcGRA7 using the suitable and novel plasmid and vector for this purpose. At our knowledge, cloning and expression of NcGRA7 has been performed in different expression systems such as pCMVi-luc1 eukaryotic expression vector or bacterial expression vector, pGEX-4T-3 (31, 36). Here, we successfully cloned and expressed NcGRA7 for the first time with new and different primers which have sites for *EcoRI* restriction enzyme. The pMALc2x used as a vector for gene expression and protein production. This vector has a strong promoter and its protein expression will be high compared to the other vectors. The rate of gene cloning into this vector

and subsequent transformation is high and noticeable, which enables work with this vector to be done more easily than with the other vectors. The presence of *lac* promoter in this vector increases the protein expression and under IPTG induction, this will be enhanced considerably. In addition, the presence of maltose in recombinant expressed protein will facilitate its purification. The NcGRA7 (24 KDa) protein showed a very similar molecular mass for protein expressed by *N. caninum*.

Western blotting revealed that performed cloning and expression with different primers can produce protein that able to bind successfully with *N. caninum* antibodies.

Serological results of current study were compared with widely used commercially available ELISA by IDEXX. Shown by kappa statistical and Mc Nemar's analysis, there was good agreement between NcGRA7-based ELISA and frequently used commercial ELISA to diagnose anti- *N. caninum* antibodies either for cattle or water buffaloes samples. On the other hand, sensitivity and specificity of presented ELISA were nearly comparable with previous reports (34, 37). *N. caninum* has huge antigenic analogy to those of *T. gondii*. The sensitivity of our developed ELISA is more acceptable since there was no cross reactivity with *T. gondii*, by inference sera that did not react with commercial ELISA but identified positive by NcGRA7-based ELISA was not due to cross reactivity with this jointly related and very common protozoa.

Conclusion

We believe that NcGRA7-based ELISA under considered circumstances and high sensitivity and specificity is an appropriate achievement to discriminate between seropositive and seronegative bovines and water buffaloes and is cost-effective simple procedure, do not need skilled technicians for screening and epidemiological purposes.

Acknowledgments

This study was supported by Shahid Chamran University of Ahvaz, Iran, and the authors wish to thank the Vice- Chancellor for Research of Shahid Chamran University for the research opportunity. The authors also declare that there is no conflict of interests.

References

1. Dubey JP, Schares G. Diagnosis of bovine neosporosis. *Vet Parasitol.* 2006; 140: 1–34.
2. Anderson M, Barr B, Rowe J, Conrad P. Neosporosis in dairy cattle. *Jpn J Vet Res.* 2012; 60: 51-4.
3. Bjerkas I, Mohn SF, Presthus J. Unidentified cyst-forming sporozoon causing encephalomyelitis and myositis in dogs. *Z Parasitenkunde* 1984; 70: 271–274.
4. Dubey JP, Lindsay DS. A review of *Neospora caninum* and neosporosis. *Vet Parasitol.* 1996; 67: 1–59.
5. McAllister MM, Dubey JP, Lindsay DS et al. Dogs are definitive hosts of *Neospora caninum*. *Int J Parasitol.* 1998; 28: 1473–1478.
6. Gondim LFP, McAllister MM, Pitt WC, Zemlicka DE. Coyotes (*Canis latrans*) are definitive hosts of *Neospora caninum*. *Int J Parasitol.* 2004; 34: 159–161.
7. Klevar S. Tissue cyst forming coccidian; *Toxoplasma gondii* and *Neospora caninum* as a cause of disease in farm animals. *Acta Vet Scand.* 2007. 49, (Suppl1): S1.
8. Sharifdini M, Mohebbi M, Keshavarz H, Hosseinijad M, Hajjaran H, Akhondi B, Rahimi Foroushani A, Zarei Z, Charehdar S. *Neospora caninum* and *Leishmania infantum* co-infection in domestic dogs (*Canis familiaris*) in Meshkin-Shahr district, Northwestern Iran. *Iran J Arthropod-Borne Dis.* 2011; 1(3): 60–68.
9. Dubey JP. Recent advances in *Neospora* and neosporosis. *Vet Parasitol.* 1999; 84: 349-367.
10. Hall CA, Reichel MP, Ellis JT. *Neospora* abortions in dairy cattle: diagnosis, mode of transmission and control. *Vet Parasitol.* 2005; 128: 231–241.
11. Osawa T, Wastling J, Maley S et al. A multiple antigen ELISA to detect *Neospora*-specific antibodies in bovine sera, bovine foetal fluids,

- ovine and caprine sera. *Vet Parasitol.* 1998; 79: 19–34.
12. Packham AE, Sverlow KW, Conrad PA et al. A modified agglutination test for *Neospora caninum*. Development, optimization, and comparison to the indirect fluorescent-antibody test, and comparison to the indirect fluorescent-antibody test and enzyme-linked immunosorbent assay. *Clin Diagn Lab Immunol.* 1998; 5: 467–473.
 13. Atkinson R, Harper PAW, Reichel MP, Ellis JT. Progress in the serodiagnosis of *Neospora caninum* infections of cattle. *Parasitol Today.* 2000; 3: 110–114.
 14. Silva DAO, Lobato J, Mineo TWP, Mineo JR. Evaluation of serological tests for the diagnosis of *Neospora caninum* infection in dogs: optimization of cut-off titers and inhibition studies of cross reactivity with *Toxoplasma gondii*. *Vet Parasitol.* 2007; 143:234–244.
 15. Howe DK, Crawford AC, Lindsay D, Sibley LD. The p29 and p35 immunodominant antigens of *Neospora caninum* tachyzoites are homologous to the family of surface antigens of *Toxoplasma gondii*. *Infect Immun.* 1998; 66:5322–5328.
 16. Aguado-Martínez A, Ivarez-García A, Fernández-García G et al. Usefulness of rNcGRA7- and rNcSAG4-based ELISA tests for distinguishing primo-infection, recrudescence, and chronic bovine neosporosis. *Vet Parasitol.* 2008; 157: 182–195.
 17. Romano MI. An applied printing immunoassay with recombinant Nc-SAG1 for detection of antibodies to *Neospora caninum* in cattle. *J Vet Diagn Inves.* 2011; 23: 971–976.
 18. Hiasa J, Nishimura M, Itamoto K et al. Enzyme-linked immunosorbent assays based on *Neospora caninum* dense granule protein 7 and profilin for estimating the stage of neosporosis. *Clin Vaccine Immunol.* 2012; 19:411-417.
 19. Jiménez-Ruiz E, Bech-Sabat G, Alvarez-García G et al. Specific antibody responses against *Neospora caninum* recombinant rNcGRA7, rNcSAG4, rNcBSR4 and rNcSRS9 proteins are correlated with virulence in mice. *Parasitology.* 2013; 140:569-79.
 20. Terkawi MA, Kameyama K, Rasul NH et al. Development of an immunochromatographic assay based on dense granule protein 7 for serological detection of *Toxoplasma gondii* infection. *Clin Vaccine Immunol.* 2013; 20:596-601.
 21. Dubey JP, Desmonts G. Serological responses of equids fed *Toxoplasma gondii* oocysts. *Equine Vet J.* 1987; 19: 337-9.
 22. Nigro M, Gutierrez A, Hoffer AM et al. Evaluation of *Toxoplasma gondii* recombinant proteins for the diagnosis of recently acquired toxoplasmosis by an immunoglobulin G analysis. *Diagn Micro Infect Dis.* 2003; 47: 609–613.
 23. Pietkiewicz H, Hiszczynska-Sawicka E, Kur J et al. Usefulness of *Toxoplasma gondii*-specific recombinant antigens in serodiagnosis of human toxoplasmosis. *J Clin Microbiol.* 2004; 42: 1779–1781.
 24. Chung CT, Niemela SL, Miller RH. One-step preparation of competent *Escherichia coli*: transformation and storage of bacterial cells in the same solution. *Biochemistry.* 1989; 86: 2172–2175.
 25. Dohoo IR, Martin SW, Stryhn H. In: Dohoo IR. (Ed.). *Vet Epidemiol Res.* AVC Inc., Charlottetown, 2003; p: 706.
 26. Dubey JP, Schares G, Ortega-Mora LM. Epidemiology and control of neosporosis and *Neospora caninum*. *Clin Microbiol Rev.* 2007; 20: 323–367.
 27. Chahan B, Gaturaga I, Huang X et al. Serodiagnosis of *Neospora caninum* infection in cattle by enzyme-linked immunosorbent assay with recombinant truncated NcSAG1. *Vet Parasitol.* 2003; 118: 177–185.
 28. Dubey JP, Hattel AL, Lindsay DS, Topper MJ. Neonatal *Neospora caninum* infection in dogs: isolation of the causative agent and experimental transmission. *J Am Vet Med Assoc.* 1988; 193: 1259–1263.
 29. Paré J, Hietala SK, Thurmond MC. An enzyme-linked immunosorbent assay (ELISA) for serological diagnosis of *Neospora* sp. infection in cattle. *J Vet Diagn Invest.* 1995; 7:352-599.
 30. Bjorkman C, Alvarez-Garcia G, Conraths FJ et al. *Neospora caninum* IgG avidity tests: an interlaboratory comparison. *Vet Parasitol.* 2006; 140: 273–280.
 31. Hara AO, Liao M, Baticados W, Bannai H et al. Expression of recombinant dense granule protein 7 of *Neospora caninum* and evaluation of its diagnostic potential for canine neosporosis. *J Protozool Res.* 2006; 16: 34-41.

32. Sonda S, Fuchs N, Gottstein B, Hemphill A. Molecular characterization of a novel microneme antigen in *Neospora caninum*. *Mol Biochem Parasitol*. 2000; 108:39–51.
33. Alvarez-Garcia G, Pitarch A, Zaballos A et al. The NcGRA7 gene encodes the immunodominant 17 kDa antigen of *Neospora caninum*. *Parasitology*. 2007; 134: 41–50.
34. Huang P, Liao M, Zhang H, Lee G et al. Dense-granule protein NcGRA7, a new marker for the serodiagnosis of *Neospora caninum* infection in aborting cows DI196. *Clin Vaccine Immunol*. 2007; 14: 1640–1643.
35. Alvarez-Garcia G, Lopez-Perez I, Innes E Colantes-Fernandez et al. Use of an immunodominant P17 antigenic fraction of *Neospora caninum* in detection of antibody response in cattle. *Mem Inst Oswaldo Cruz*. 2006; 101: 529–534.
36. Liddell S, Parker C, Vinyard B et al. Immunization of mice with plasmid DNA coding for NcGRA7 or NcsHSP33 confers partial protection against vertical transmission of *Neospora caninum*. *J Parasitol*. 2003; 89: 496–500.
37. Wilkowsky SE, Bareiro GG, Mon ML et al. An applied printing immunoassay with recombinant Nc-SAG1 for detection of antibodies to *Neospora caninum* in cattle. *J Vet Diagn Invest*. 2011; 23:971-6.