



Title	Development of vaccine strains of H5 and H7 influenza viruses
Author(s)	Soda, Kosuke; Sakoda, Yoshihiro; Isoda, Norikazu; Kajihara, Masahiro; Haraguchi, Yoshinari; Shibuya, Hitomi; Yoshida, Hiromi; Sasaki, Takashi; Sakamoto, Ryuichi; Saijo, Kazue; Hagiwara, Junko; Kida, Hiroshi
Citation	Japanese Journal of Veterinary Research, 55(2-3), 93-98
Issue Date	2008-02-09
DOI	10.14943/jjvr.55.2-3.93
Doc URL	http://hdl.handle.net/2115/32369
Type	bulletin (article)
File Information	55(2-3)_93-98.pdf



[Instructions for use](#)

Development of vaccine strains of H5 and H7 influenza viruses

Kosuke Soda¹⁾, Yoshihiro Sakoda¹⁾, Norikazu Isoda¹⁾, Masahiro Kajihara¹⁾, Yoshinari Haraguchi¹⁾, Hitomi Shibuya¹⁾, Hiromi Yoshida¹⁾, Takashi Sasaki²⁾, Ryuichi Sakamoto³⁾, Kazue Saijo⁴⁾, Junko Hagiwara⁵⁾ and Hiroshi Kida^{1),6)*}

¹⁾Laboratory of Microbiology, Department of Disease Control, Graduate School of Veterinary Medicine, Hokkaido University, Sapporo, 060-0818, Japan

²⁾Avian Biologics Department, Kyoto Biken Laboratories, Inc., Uji, Kyoto, 611-0041, Japan

³⁾Division 2, Second Research Department, The Chemo-Sero-Therapeutic Research Institute, Kikuchi, Kumamoto, 869-1298, Japan

⁴⁾Research Center for Biologicals, The Kitasato Institute, Kitamoto, Saitama, 364-0026, Japan

⁵⁾Research Department, Nippon Institute for Biological Science, Ome, Tokyo, 198-0024, Japan

⁶⁾Research Center for Zoonosis Control, Hokkaido University, Sapporo, 001-0020, Japan

Received for publication, December 27, 2007; accepted, January 30, 2008

Abstract

To establish vaccine strains of H5 and H7 influenza viruses, A/duck/Hokkaido/Vac-1/04 (H5N1) [Vac-1/04 (H5N1)], A/duck/Hokkaido/Vac-3/07 (H5N1) [Vac-3/07 (H5N1)], and A/duck/Hokkaido/Vac-2/04 (H7N7) [Vac-2/04 (H7N7)] were generated from non-pathogenic avian influenza viruses isolated from migratory ducks. Vac-1/04 (H5N1) and Vac-3/07 (H5N1) were generated by genetic reassortment between H5N2 or H5N3 virus as an HA gene provider and H7N1 or H6N1 viruses as an NA gene provider. Vac-2/04 (H7N7) was a genetic reassortant obtained using H7N7 and H9N2 viruses to give high growth character of the H9N2 virus in chicken embryonated eggs. The results of sequence analyses and experimental infections revealed that these H5N1 and H7N7 reassortant viruses were non-pathogenic in chickens and embryos, and had good growth potential in embryonated eggs. These viruses should be useful to develop vaccines against H5 and H7 highly pathogenic avian influenza viruses.

Key Words: genetic reassortant, H5, H7, influenza, vaccine

Introduction

Highly pathogenic avian influenza (HPAI) caused by H5 and H7 influenza viruses have occurred worldwide, leading to serious economic

losses in poultry industries. In Japan, there had been no outbreak since 1925 when HPAI caused by a H7N7 virus occurred in Chiba prefecture¹³⁾. In the beginning of 2004, HPAI outbreaks caused by H5N1 viruses occurred at chicken farms in

*Corresponding author: Hiroshi Kida, Laboratory of Microbiology, Department of Disease Control, Graduate School of Veterinary Medicine, Hokkaido University, Sapporo 060-0818, Japan
Phone: +81-11-706-5207. Fax: +81-11-706-5273. E-mail: kida@vetmed.hokudai.ac.jp

Yamaguchi and Kyoto prefectures^{7,10}. H5N1 infections then reoccurred on poultry farms in Miyazaki and Okayama prefectures in 2007. The depopulation of chickens in the relevant farms and appropriate control measures successfully prevented the spread of HPAI.

Standard countermeasures for the control of HPAI outbreaks are testing and culling of affected chickens. However, when outbreaks spread to a broad area, being uncontrollable, ring vaccination would be an optional measure to reduce the concentration of the virus and, hence, suppress its spread³. In response to the 2004 outbreaks in Japan, commercial vaccines were urgently imported from Mexico as stockpiles. However, these vaccines have been suspected that it might be inadequate to control current outbreaks in Asia because they have been produced from H5N2 viruses of North American lineage, which are antigenically different from those of Eurasian lineage. A domestic vaccine which has a similar antigenicity to viruses circulating in Asia at present has been awaited for stable production and provision in the case of a nation-wide outbreak which cannot be controlled by stamping out strategy.

The aim of the present study is to establish proper seed strains for vaccine manufacturing against the H5 and H7 influenza viruses in Asia. Here, we report the characteristics of non-pathogenic H5N1 and H7N7 viruses generated from parental viruses isolated from ducks, a natural host of influenza A viruses, by genetic reassortment.

Materials and Methods

Viruses: All the parental viruses for generating vaccine candidate strains in this study were isolated in our laboratory from fecal samples of migratory ducks in Mongolia and Japan⁸. The viruses were propagated in ten-day-old embryonated chicken eggs for 48 hours at 35°C and infectious allantoic fluids were used for generating of reassortant viruses.

Generation of genetic reassortant virus: To generate the H5N1 and H7N7 reassortant viruses, parental viruses were mixed and inoculated into the allantoic cavity of embryonated chicken eggs. After incubation at 35°C for 24h or 48h, the allantoic fluids were collected, mixed with chicken antisera (α -A/Duck/Hong Kong/301/78 (H7N2) for A/duck/Hokkaido/Vac-1/04 (H5N1), α -A/turkey/England/73 (H7N3) and α -A/turkey/Massachusetts/3740/65 (H6N2) for A/duck/Hokkaido/Vac-3/07 (H5N1), and α -A/duck/Hokkaido/13/00 (H9N2) for A/duck/Hokkaido/Vac-2/04 (H7N7)) and incubated at room temperature for one hour prior to plaque cloning of the viruses in MDCK cells to neutralize the viruses which have undesired hemagglutinin (HA) and/or neuraminidase (NA) (e.g., H7HA and N2NA in generation of A/duck/Hokkaido/Vac-1/04 (H5N1)). Cloned viruses were subtyped using hemagglutinin-inhibition (HI)¹¹ and neuraminidase-inhibition (NI) tests².

Plaque assay: Purification of viruses by plaque selection method was performed as described previously⁵. Briefly, confluent MDCK cells were incubated one hour at 35°C with ten-fold dilutions of virus. The cells were then washed and overlaid with MEM (MEM, Nissui Pharmaceutical, Tokyo, Japan) containing 0.7% Bacto-agar (Difco, Sparks, MD, U.S.A) and 0.5µg/ml of trypsin and incubated at 35°C. After three days, plaques were picked to be further purified.

Sequencing: Viral RNAs were extracted from infectious allantoic fluids using a commercial kit (TRI LS reagent, Sigma-Aldrich, St. Louis, MO, U.S.A) and reverse-transcribed with the Uni12 primer⁶ and M-MLV Reverse Transcriptase (Invitrogen, Carlsbad, CA, U.S.A). Viral complementary DNA was amplified with the polymerase chain reaction (PCR) with the RNA LA PCR Kit Ver. 1.1 (Takara, Shiga, Japan) and PTC-200 thermal cycler (Biorad, Alfred Novel Drive Hercules, CA, U.S.A). The following primers were used or designed on the basis of published nucleotides sequences of HA genes for RT-PCR in the present study; (BmHA

-1F⁶) 5'-TATTCGTCTCAGGGAGCAAAAGCAGGG G-3', (H5-1695R) 5'-CGATCCATTGGAGCACATCC -3', and (H7-1141R) 5'-CGATCCATTGGAGCACAT CC-3'. For sequencing internal protein genes, gene -specific primer sets were used as described previously⁹. Nucleotide sequences were determined from these RT-PCR products using CEQ2000 automated DNA sequencer (Beckman Coulter, Fullerton, CA, U.S.A) according to Dye Terminator Cycle Sequence Chemistry Protocol (Beckman Coulter). Sequence data were analysed using GENETYX Version 7.0 (Genetyx, Tokyo, Japan).

Pathogenicity studies: Mean death times (MDTs) of chicken embryos were determined as described previously¹⁴. Nine-day-old embryonated chicken eggs were inoculated with ten-fold dilutions of the viruses. Eggs were examined every eight hours by candling; the integrity of the circulatory system and movement of the embryo were the criteria for viability. Any embryos which were dead at 24 hour post-inoculation were discarded as nonspecific deaths and did not figure in the calculation of MDT. The MDT was calculated as the mean time in hours for the minimum dose lethal to kill the embryos. The intracerebral pathogenicity index (ICPI) and the intravenous pathogenicity index (IVPI) tests were performed as described elsewhere¹⁴; each infectious allantoic fluid was inoculated into groups of ten one-day-old chicks and eight six-week-old chickens, via the intracerebral and intravenous route, respectively. In both studies, the birds were observed for clinical symptoms and mortality at 24-hour intervals for a period of eight days for the ICPI test and ten days for the IVPI test. The ICPI and IVPI values were calculated as described¹¹.

Viral Growth Kinetics: Viruses of 10² and 10⁴ 50% egg infectious dose (EID₅₀) (10^{1.5} and 10^{3.5} EID₅₀ for A/duck/Hokkaido/Vac-2/04 (H7N7)) were inoculated into ten-day-old embryonated chicken eggs and incubated at 34°C (35°C for A/duck/Hokkaido/Vac-3/07 (H5N1)). Allantoic fluids were harvested to determine viral titers at different time points (0,

12, 24, 48, and 72 hour). The EID₅₀ and hemagglutination titers of each virus present in the allantoic fluid was determined.

Results

Generation of non-pathogenic H5N1 strains of Eurasian lineage:

Until now, non-pathogenic H5N1 viruses of Eurasian lineage have been rarely isolated from natural hosts¹². To establish H5N1 vaccine candidate strains against H5N1 influenza viruses circulating worldwide, A/duck/Hokkaido/Vac-1/04 (H5N1) [Vac-1/04 (H5N1)] and A/duck/Hokkaido/Vac-3/07 (H5N1) [Vac-3/07 (H5N1)] were generated by the standard genetic reassortant procedure from non-pathogenic viruses of Eurasian lineage derived from waterbirds and origins of the internal proteins of these reassortant viruses were determined (Fig. 1A and 1B). The PB2, PB1, PA, HA, NP, and M gene segments of Vac-1 (H5N1) were derived from A/duck/Mongolia/54/01 (H5N2), and the NA and NS gene segments, from A/duck/Mongolia/47/01 (H7N1) (Fig. 1A). The HA and NS gene segments of Vac-3 (H5N1) were derived from A/duck/Hokkaido/101/04 (H5N3), while the other segments were from A/duck/Hokkaido/262/04 (H6N1) (Fig. 1B). The complete nucleotide sequences of Vac-1 (H5N1) and Vac-3 (H5N1) obtained in the present study have been registered in GenBank (Accession numbers: AB259709-AB259716, and AB355926-AB355933).

Generation of a non-pathogenic H7N7 virus with good growth potential:

Dozens of non-pathogenic H7N7 viruses have been isolated from fecal samples of waterbirds during our surveillance work since 2001 (data not shown). Based on the results of phylogenetic and antigenic analyses (data not shown), A/duck/Mongolia/736/02 (H7N7) [Mon/736 (H7N7)] was selected as a vaccine candidate. However, the growth potential of Mon/736 (H7N7) in chicken embryonated eggs was poor in terms of the HA




Fig. 1. Establishment of candidate strains for vaccines against H5 and H7 highly pathogenic avian influenza viruses by genetic reassortment. Vac-1/04 (H5N1), Vac-3/07 (H5N1), and Vac-2/04 (H7N7) were generated from two non-pathogenic avian influenza viruses isolated from migratory ducks. The origin of viral segments of reassortant viruses is indicated with white and black bands. (A); Vac-1/04 (H5N1) was a reassortant generated with A/duck/Mongolia/54/01 (H5N2) as the HA gene provider and A/duck/Mongolia/47/01 (H7N1) as the NA gene provider. (B); Vac-3/07 (H5N1) was generated with A/duck/Hokkaido/101/04 (H5N3) as the HA gene provider and A/duck/Hokkaido/262/04 (H6N1) as the NA gene provider. (C); Vac-2/04 (H7N7) was generated from Mon/736 (H7N7) as the HA and NA gene provider and Hok/49 (H9N2) to give high growth potential in embryonated eggs.

titer and yield of viral protein. To improve the growth potential of Mon/736 (H7N7), A/duck/Hokkaido/Vac-2/04 (H7N7) [Vac-2/04 (H7N7)] was generated as a genetic reassortant virus between Mon/736 (H7N7) and A/duck/Hokkaido/49/98 (H9N2) [Hok/49 (H9N2)], which possesses good growth potential in embryonated eggs (Fig. 1C). The PB2, PB1, PA, HA, NA, and NS gene segments of Vac-2/04 (H7N7) were derived from Mon/736 (H7N7), and the other segments, from Hok/49 (H9N2) (Fig. 1C).

The complete nucleotide sequences of Vac-2 (H7N7) obtained in the present study have been registered in GenBank (Accession numbers: AB243417-AB243424).

Pathogenicity of vaccine candidate strains:

The pathogenicity of the vaccine candidate strains for chickens, chicks, and chicken embryos was evaluated by measuring the IVPI, ICPI, and MDT. All of the chickens inoculated intravenously with Vac-1 (H5N1), Vac-3 (H5N1), or Vac-2 (H7N7) survived for ten days without showing any clinical signs, and IVPI values were 0.00 (Table 1). The pathogenicity of Vac-1 (H5N1) and Vac-2 (H7N7) for chicks was very low (the ICPI values were 0.10 and 0.20, respectively). Although HAs of all the vaccine candidate strains had typical sequences found in the HA of avirulent type at the cleavage sites as shown in Table 1, half of the chicks inoculated intracerebrally with Vac-3 (H5N1) were dead at two to eight days post-inoculation (ICPI, 0.58), indicating a relatively high pathogenicity for chicks as compared with the other two candidate strains. The MDT of Vac-1 (H5N1), Vac-3 (H5N1), and Vac-2 (H7N7) was 64, 73, and 81 hours, respectively.

Growth character of vaccine candidate strains in chicken embryos:

The growth kinetics of vaccine candidates was determined by using chicken embryos. As shown in Table 2, all the candidates replicated well, and reached plateau by 48 hours. There was no significant difference in peak titers regardless of the in-

Table 1. Pathogenicity of vaccine candidate strains for chickens and chicken embryos

Viruses	Amino acid sequence of HA cleavage site	IVPI ^{a)}	ICPI ^{b)}	MDT ^{c)} (hr)
Vac-1/04 (H5N1)	PQRETR/G	0.00	0.10	64
Vac-3/07 (H5N1)	PQRETR/G	0.00	0.58	73
Vac-2/04 (H7N7)	PEIPKGR/G	0.00	0.20	81

^{a)} Intravenous pathogenicity index

^{b)} Intracerebral pathogenicity index

^{c)} Mean death time

Table 2. Growth kinetics of vaccine candidate strains in chicken embryos^{a)}

Incubation time (hr)	Vac-1/04 (H5N1)				Vac-3/07 (H5N1)				Vac-2/04 (H7N7)			
	10 ^{2.0b)}		10 ^{4.0}		10 ^{2.0}		10 ^{4.0}		10 ^{1.5}		10 ^{3.5}	
	Virus titer	HA titer	Virus titer	HA titer	Virus titer	HA titer	Virus titer	HA titer	Virus titer	HA titer	Virus titer	HA titer
0	≤0.5	<2	≤0.5	<2	≤0.5	<2	≤0.5	<2	≤0.5	<2	≤0.5	<2
12	3.7	<2	4.5	<2	2.3	<2	5.5	<2	3.5	<2	5.1	<2
24	8.7	32	9.5	256	8.3	512	8.4	1024	7.9	4	8.7	64
48	9.3	256	9.5	512	9.3	2048	9.5	2048	9.5	512	9.1	512
72	9.1	512	9.3	256	9.5	2048	9.5	2048	9.1	512	9.3	512

^{a)} The virus titers (logEID₅₀/ml) and HA titers (HAU/ml) were determined from allantoic fluids of chicken embryos inoculated with Vac-1/04 (H5N1), Vac-3/07 (H5N1), and Vac-2/04 (H7N7), respectively.

^{b)} Dose of inoculum given to chicken embryo (EID₅₀/egg)

oculum dose of each vaccine candidate.

Discussion

It is required for influenza vaccine strain to be antigenically similar to endemic virus strains, non-pathogenic for chicken embryos, and high yield in the embryonated eggs. In response to the 2004 outbreaks of HPAI in Japan, the commercial vaccines were emergently imported from Mexico. However, these vaccines were produced using H5N2 viruses of North American lineage, which are antigenically different from current HPAI viruses circulating in countries in Asia, Central East, Europe, and Africa. It is, therefore, required for the stockpile vaccine that is prepared from virus strains of Eurasian lineage, causing the present epidemics. In the present study, we selected the virus strains isolated from feral waterbirds as H5 or H7 HA providers based on the following criteria: (a) antigenically and genetically similar to HPAI virus presently circulating, (b) the latest isolates,

then applied for generation of vaccine candidate strains.

To obtain non-pathogenic H5N1 strains for vaccines, Vac-1 (H5N1) and Vac-3 (H5N1) were generated by genetic reassortment. These two strains showed low pathogenicity in chickens and chicken embryos (Table 1). Additionally, they grew well in embryonated eggs after 48 hours post-inoculation, relatively shorter than their MDT (Table 2), indicating that these virus strains are useful for vaccine production against H5N1 influenza virus infections. Since Vac-3 (H5N1) showed comparatively pathogenic to chicks compared with Vac-1(H5N1) (Table 1), we set Vac-1 (H5N1) as the first candidate strain for H5N1 influenza virus vaccine. The vaccine for chickens by using Vac-1 (H5N1) has been produced in collaboration with vaccine producers and its protective effect is under investigation.

Vac-2 (H7N7) was obtained as a vaccine candidate strain against H7N7 viruses (Fig. 1C). Mon/736 (H7N7), providing the HA and NA gene of Vac-2 (H7N7), showed poor growth potential in chicken

embryos (16HA). To solve this problem, a genetic reassortant was generated using Hok/49 (H9N2), which has high growth potential in eggs. The NP and M gene segments of Vac-2 (H7N7) were derived from Hok/49 (H9N2), indicating that the NP and/or M gene segments were responsible for the growth potential. To determine whether NP and/or M proteins are truly involved in growth potential in chicken embryos or not, further investigation is underway.

References

- 1) Alexander, D. J., Banks, J., Collins, M. S., Manvell, R. J., Frost, K. M., Speidel, E. C. and Aldous, E. W. 1999. Antigenic and genetic characterisation of Newcastle disease viruses isolated from outbreaks in domestic fowl and turkeys in Great Britain during 1997. *Vet. Rec.*, **145**: 417-421.
- 2) Aymard-Henry, M., Coleman, M. T., Dowdle, W. R., Laver, W. G., Schild, G. C. and Webster, R. G. 1973. Influenzavirus neuraminidase and neuraminidase-inhibition test procedures. *Bull. World Health Organ.*, **48**: 199-202.
- 3) Capua, I. and Marangon, S. 2003. The use of vaccination as an option for the control of avian influenza. *World Organization for animal health international committee 71st general session (Paris, 18-23 May 2003)*. http://www.oie.int/eng/avian_influenza/A_71_SG_12_CS3E.pdf
- 4) Capua, I. and Multinelli, F. 2001. A Colour Atlas and Text on Avian Influenza. *Papi Editore, Bologna, Italy*.
- 5) Hayden, F. G., Cote, K. M. and Douglas, R. G., Jr. 1980. Plaque inhibition assay for drug susceptibility testing of influenza viruses. *Antimicrob. Agents Chemother.*, **17**: 865-870.
- 6) Hoffmann, E., Stech, J., Guan, Y., Webster, R. G. and Perez, D. R. 2001. Universal primer set for the full-length amplification of all influenza A viruses. *Arch. Virol.*, **146**: 2275-2289.
- 7) Isoda, N., Sakoda, Y., Kishida, N., Bai, G. R., Matsuda, K., Umemura, T. and Kida, H. 2006. Pathogenicity of a highly pathogenic avian influenza virus, A/chicken/Yamaguchi/7/04 (H5N1) in different species of birds and mammals. *Arch. Virol.*, **151**: 1267-1279.
- 8) Kida, H. and Sakoda, Y. 2006. Library of influenza virus strains for vaccine and diagnostic use against highly pathogenic avian influenza and human pandemics. *Dev. Biol. (Basel)*, **124**: 69-72.
- 9) Liu, J., Okazaki, K., Ozaki, H., Sakoda, Y., Wu, Q., Chen, F. and Kida, H. 2003. H9N2 influenza viruses prevalent in poultry in China are phylogenetically distinct from A/quail/Hong Kong/G1/97 presumed to be the donor of the internal protein genes of the H5N1 Hong Kong/97 virus. *Avian Pathol.*, **32**: 551-560.
- 10) Mase, M., Tsukamoto, K., Imada, T., Imai, K., Tanimura, N., Nakamura, K., Yamamoto, Y., Hitomi, T., Kira, T., Nakai, T., Kiso, M., Horimoto, T., Kawaoka, Y. and Yamaguchi, S. 2005. Characterization of H5N1 influenza A viruses isolated during the 2003-2004 influenza outbreaks in Japan. *Virology*, **332**: 167-176.
- 11) Sever, J. L. 1962. Application of a microtechnique to viral serological investigations. *J. Immunol.*, **88**: 320-329.
- 12) Spackman, E., Swayne, D. E., Suarez, D. L., Senne, D. A., Pedersen, J. C., Killian, M. L., Pasick, J., Handel, K., Pillai, S. P., Lee, C. W., Stallknecht, D., Slemmons, R., Ip, H. S. and Deliberto, T. 2007. Characterization of low-pathogenicity H5N1avian influenza viruses from North America. *J. Virol.*, **81**: 11612-11619.
- 13) Sugimura, T., Ogawa, T., Tanaka, Y. and Kumagai, T. 1981. Antigenic type of fowl plague virus isolated in Japan in 1925. *Natl. Inst. Anim. Health Q. (Tokyo)*, **21**: 104-105.
- 14) Waterson, A. P., Pennington, T. H. and Allan, W. H. 1967. Virulence in Newcastle disease virus. A preliminary study. *Br. Med. Bull.*, **23**: 138-143.