



Developmental Expression of Kv Potassium Channels Segment of Cultured Hippocampal Neurons

Diana Sánchez-Ponce, Javier DeFelipe, Juan José Garrido, Alberto Muñoz

Abstract

Axonal outgrowth and the formation of the axon initial segment (AIS) are early events in the acquisition of neuronal concentration of voltage-dependent sodium and potassium channels. However, the specific ion channel subunits per subdomain vary both during development and among the types of neurons, probably determining their firing characteristics. We characterize the developmental expression of different subfamilies of voltage-gated potassium channels in the AIS, including subunits Kv1.2, Kv2.2 and Kv7.2. In contrast to the early appearance of voltage-gated sodium channels α subunits were tethered at the AIS only after 10 days *in vitro*. Interestingly, we observed different patterns of Kv1.2 to distinct neuronal populations. The accumulation of Kv1.2 and Kv2.2 subunits at the AIS was dependent on ankyrin and the actin cytoskeleton and it was resistant to detergent extraction, as described previously for other AIS proteins. This study further emphasizes the heterogeneity of this structure in different neuronal populations, as proposed previously, and a potential regulation.

Citation: Sánchez-Ponce D, DeFelipe J, Garrido JJ, Muñoz A (2012) Developmental Expression of Kv Potassium Channels in the Axon Initial Segment of Cultured Hippocampal Neurons. PLoS ONE 7(10): e48557. doi:10.1371/journal.pone.0048557

Editor: Stuart E. Dryer, University of Houston, United States of America

Received: July 3, 2012; **Accepted:** September 27, 2012; **Published:** October 31, 2012

Copyright: © 2012 Sánchez-Ponce et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Funding: This work was supported by grants from the Ministerio de Ciencia e Innovación (SAF 2010-18218 to A.M., SAF 2009-09394 to J.D.), the Cajal Blue Brain Project and the Centro de Investigación Biomédica en Red de Enfermedades Neurodegenerativas supported by an FPU fellowship from the Ministerio de Educación (Spain). The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing interests: The authors have declared that no competing interests exist.

Introduction

The non-uniform distribution of specific voltage-gated K^+ (Kv) channels and their restriction to discrete neuronal domains influence neuronal excitability. Indeed, these channels are believed to influence different properties of neurons, including resting potential (AP) firing pattern, transmitter release and synaptic strength. The importance of Kv channels in neuronal excitability is further emphasized by mutations or diseases that disrupt K^+ channel expression, including episodic ataxia and epilepsies [1]. Kv channels are actually complexes made up of four voltage-sensing and pore-forming subunits, each generated from different subfamilies (Kv1–12) [7], [8]. These complexes may assemble with auxiliary β subunits that may influence the expression and function of Kv channels [9], [10], [11]. The axon initial segment (AIS) is a neuronal domain that is densely populated by voltage-gated potassium channels.

critical for input integration and action potential generation [12], [13], [14], [15]. In addition to voltage-gated Na⁺ (V) channels [12], [21], [22], the AIS of neocortical and hippocampal principal cells is characterized by the expression [23], [24], [25], [26]. The distinct subunit composition of these channels confers distinct biophysical properties to them. Together with heterogeneous expression and localization of such channels in the AIS, they are likely to contribute to neuronal populations, and the corresponding differences in AP initiation and/or propagation [17], [18], [19], [20], [2




Figure 1. Developmental expression of potassium channels in cultured hippocampal neurons.

(A) Western blot of Kv1.2, Kv2.2 and Kv7.2 in hippocampal neurons cultured at high density (50,000/cm²) for different developmental stages. (B) Histograms show Kv1.2, Kv2.2 and Kv7.2 expression normalized to actin when quantified densitometrically. Note the delayed onset of Kv1.2 expression as compared with Kv2.2 and Kv7.2. (C) Photomicrograph of hippocampal neurons cultured for 6 DIV and double immunostained for Kv7.2 (green) and MAP2 (red) in a single process emerging from the cell body (arrows). (D) Histogram shows the percentage (mean ± SE) of neurons with Kv7.2 at the AIS at different developmental stages *in vitro*. Scale bar = 16 μm.

doi:10.1371/journal.pone.0048557.g001

Kv1 channels are characterized by low thresholds, as well as rapid activation and slow inactivation kinetics, and they regulate the shape and propagation rate of APs, as well as neurotransmitter release and synaptic efficacy [25], [32], [33], [34]. These channels are also localized at the AISs of different neuronal populations, where they contribute to the inter-spike potential during high frequency firing [44], [45]. Kv7 channels are localized at the AIS [23], [46], [47]. They regulate the M-current. These channels regulate resting potential and AP firing and they are characterized by low-threshold, slow sustained activity and non-inactivation near the AP threshold [47], [52], [53], [54].

Despite the functional importance of the AIS, the timing and the intracellular mechanisms involved in the development of the AIS are poorly understood. A key aim of the present study was to describe the distribution, developmental expression and localization of Kv channels in the AIS of cultured hippocampal neurons, a model commonly used to study the development of neuronal polarity and function. In cultured hippocampal neurons, ankyrin G is one of the earliest markers to be detected at the AIS and it is essential for the tethering of Kv channels to the AIS during development, such as VGSC [56], [57], [58]. Like VGSCs, Kv7 channels (Kv7.2 and Kv7.3 subunits) contain a conserved domain that mediates their targeting to the AIS. Together with the adhesion molecules NrCAM and Neurofascin 186, accumulation of Kv7 channels at the AIS depends on the structural integrity of actin and on the microtubule cytoskeleton in the AIS [64], [65], [66], [67]. Furthermore, the development of hippocampal neurons involves the expression of GABA_A receptor subunits and gephyrin clusters [68], as well as the acquisition of a functional AIS involved in Ca²⁺ regulation and reaches neurochemical maturation during the second week *in vitro* [69], [70]. To date, the developmental expression of Kv2 channels in the AIS, along with the associated trafficking and clustering mechanisms, has not been fully elucidated. We describe the temporal and spatial distribution of Kv1.2 and Kv2.2 subunits during AIS development, and the role of the subunit

results show that Kv1.2 and Kv2.2 expression are mutually exclusive in the AIS of cultured hippocampal neurons, and dependent on ankyrin G, yet independent of actin cytoskeleton integrity.




Figure 2. Kv1.2 is concentrated at the AIS during axonal maturation *in vitro*.


Hippocampal neurons were grown for 1, 3, 6, 8, 10, 13, 15 and 18 days at low density (5,000/cm²), fixed in 4% paraformaldehyde, stained with antibodies against Kv1.2 (green) and 14D4 antibodies (red) to identify the AIS. Note that 14D4 staining is detected at the AIS (A, B) and it is restricted to the AIS as the axon elongates (arrows). Confocal microscopy photomicrographs show neurons cultured for up to 10 DIV (A–E). Staining is light and localized to the soma and neurites. After 10 DIV (F–H), 14D4 staining is observed in the distal AIS. See Figure 3 for quantification. Scale bar = 18 μ m.

doi:10.1371/journal.pone.0048557.g002

Materials and Methods

Neuronal Cultures

Hippocampal neurons were obtained from E17 mouse embryos and prepared as described previously [55]. Mice were housed in a standard animal care facility. Pregnant female mice and embryos were sacrificed by cervical dislocation and decapitation respectively following the European Union Convention ETS123, recently revised as indicated in the Directive 86/609/EEC. In addition all protocols were approved by the local ethics committee (Subcomité de Bioética, CSIC; Institutional review board; IRB 0007851).







Figure 3. Kv2.2 concentration at the AIS increases during axonal maturation *in vitro*.

Confocal microscopy photomicrographs showing representative hippocampal neurons cultured for 1, 3, 6, 8, 10, 14, 16, and 20 (5,000/cm²), fixed in 4% PFA, and stained with antibodies against Kv2.2 (green) and VGSC (red). According to [2008], VGSCs concentrate at the AIS (arrows) after 3 DIV. Note that moderate Kv2.2 immunostaining is localized to the proximal region of the AIS (arrows) at all developmental stages in culture (A–C). After 14 DIV (D–F), Kv2.2 expression is evident in the proximal region of the AIS. Histogram shows the percentage of neurons expressing Kv1.2 (G) and Kv2.2 (H) at *in vitro* (the data represent the mean ± SE from three independent experiments). Scale bar = 18 μm (A–D) and 10 μm (E–F). doi:10.1371/journal.pone.0048557.g003

Briefly, after dissection of the hippocampus, tissue pieces were washed three times in Ca²⁺/Mg²⁺-free HBSS and 0.2% trypsin. The tissue was washed three times in Ca²⁺/Mg²⁺-free HBSS and dissociated with a fire-polished Pasteur pipette. The neurons were resuspended in plating medium (MEM with 10% Horse Serum and 0.6% glucose) and plated on polylysine coated coverslips (low density) for immunostaining, or 50,000 cells per cm² (high density) for Western blots. After 2 hours, the medium was replaced with Neurobasal medium supplemented with B-27 and glutamax-1. To maintain the neurons for 21 days *in vitro*, the coverslips containing astrocyte monolayers that had been cultured in neuronal culture medium for 24 h previously. 1-β-actin was added to the culture after 3 days to prevent astroglial cell growth, and in some cases neurons were treated between 3 and 18 DIV with thapsigargin (Sigma) to impede actin polymerization. For detergent extraction, neurons were maintained in culture for 21 DIV, and then extracted for 15 minutes at 37°C with 1% Triton X-100 in cytoskeletal buffer (2 mM MgCl₂, 10 mM EGTA, 60 mM Pipes [pH 7.0]). In some experiments, the plasmids were introduced into hippocampal neurons by nucleofection prior to plating (Amaxa Biosystems) according to the manufacturer's instructions. Nucleofection was performed using 3 μg of total DNA and the plasmids used for transfection were a GFP plasmid and a shRNA in a pGFP-V-RS plasmid and a shRNA-AnkG (sequence: TCGGATAGGTCCTACACCTTGAACAGAAG) in a pGFP-V-RS plasmid (Amaxa Biosystems, USA). The effects of nucleofection were analyzed at 18 DIV.






Figure 4. Lack of Kv1.2 and Kv2.2 colocalization at the AIS of cultured hippocampal neurons.

A–B and **C–D**: Pairs of representative confocal microscopy photomicrographs of hippocampal neurons cultured (red) and Kv 2.2 (green), and counterstained with DAPI (**A–D**). Note that Kv1.2-expressing AISs (arrows) lack Kv2.2 (**E**). Histogram shows the proportion of neurons expressing Kv1.2, Kv2.2, neither or both at the same AIS at 11–13 independent experiments). Scale bar = 25 μm.

doi:10.1371/journal.pone.0048557.g004

Western Blotting

Protein samples were prepared from hippocampal neurons cultured at high density (50,000/cm²) in control conditions (21 DIV), the plates were washed twice with cold PBS, the neurons were lysed and then homogenized in a buffer containing 100 mM NaF; 1% Triton X-100; 1 mM sodium orthovanadate; 10 mM EDTA; and a complete protease inhibitor cocktail (Roche). The lysates were boiled for 10 minutes, separated by SDS-PAGE on 8% acrylamide gels and transferred to nitrocellulose. The membranes were incubated overnight at 4°C with primary antibodies in blocking solution (PBS, 0.2% Tween-20 and 5% BSA): mouse anti-Kv2.2 (1:500; Alomone, Jerusalem, Israel); mouse anti-Kv7.2 (KCNQ2, 1:500; Neuromab) and mouse anti-ankyrin G (1:500; Neuromab). After washing, the membranes were incubated with the corresponding peroxidase conjugated secondary antibody. The signal was visualized by ECL (Amersham). Densitometry was performed using an imaging densitometer (GS-800, BioRad) and a whole-image background subtraction tool (Quantity One software, BioRad).




Figure 5. AIS resistance to detergent extraction.

Hippocampal neurons cultured for 21 DIV were incubated for 15 min in a buffer containing 0.5% Triton X-100 before being analyzed by confocal microscopy. After detergent extraction, Kv1.2 (**A**) and Kv2.2 (**D**) were still present in the AIS, as indicated by the resistant AIS markers 14D4 (**B**) and ankyrin G (**E**), respectively. Asterisks indicate the location of the neuronal AIS. Scale bar = 25 μm.

doi:10.1371/journal.pone.0048557.g005

Immunocytochemistry

After different times in culture, neurons were fixed in 4% paraformaldehyde for 20 minutes and then washed in PBS with 50 mM NH₄Cl and incubated in blocking buffer for 45 minutes (PBS, 0.22% gelatin and 0.1% Triton X-100). Neurons were incubated for 1 h at room temperature with the primary antibodies to Kv channel subunits diluted in blocking buffer: mouse anti-Kv1.2 (1:250; Neuromab); mouse anti-Kv 7.2 (KCNQ2, 1:200; Neuromab). To identify the AIS we used mouse anti-ankyrin G (1:200; Neuromab) antibodies and rabbit antibodies (14D4) raised against phospho (p-32)-labeled protein. Mouse anti-Tau-1 (1:1,000; Sigma) and mouse anti-phalloidin (1:100; Invitrogen, A-12379) antibodies were used to reveal axonal and neuronal morphology respectively. In some neurons, phalloidin (1:100; Invitrogen, A-12379). The secondary antibodies used were donkey anti-mouse, anti-rabbit or anti-goat (1:1000; Invitrogen, A-12379). After staining coverslips were counterstained with DAPI (1:1000, Calbiochem, San Diego, CA, USA) and mounted in mounting medium (Dako, Ely, UK).

Birmingham, AL, USA). Images were obtained using a DP70 camera attached to an Olympus BX51 fluorescence microscopy (Zeiss 710). Z sections were recorded at 0.2–1- μ m intervals through separate channels and ZEN 2001 composite images from each optical series by combining the images recorded through the different channels. In all cases, the images were processed using Adobe Photoshop CS3 (Adobe Systems Inc., San Jose, CA, USA) and used to generate the figures (Adobe Systems Inc., San Jose, CA, USA). The cell counts in the different experiments were performed using a t-test using Sigma Plot 11.0 software.




Figure 6. AIS Kv channel expression is not dependent on the actin cytoskeleton.

Confocal microscopy photomicrographs show that Kv1.2 (A–F) and Kv2.2 (G–L) accumulation in the AIS is not dependent on the actin cytoskeleton. Cultured hippocampal neurons were exposed to DMSO (control, A–C, G–I) or cytochalasin D (5 μ M; D–F, J–L) from 15 to 17 DIV, double-stained for Kv1.2 (red, A–F) or Kv2.2 (red, G–L), and stained with Alexa 488 phalloidin to reveal F-actin. Note the present and cytochalasin D-treated neurons. Scale bar = 25 μ m (A–F) and 30 μ m (G–L).

doi:10.1371/journal.pone.0048557.g006

Results

Voltage-gated Potassium Channel Expression in the Developing AIS of Cultured Hippocampal Neurons

We first analyzed the expression of voltage-gated potassium channels in hippocampal neurons cultured at high density. The total expression of the different Kv subunits was analyzed in Western blot and the results were normalized to the expression of β -tubulin. Kv1.2 expression was only weakly detected during the first days *in vitro*, yet it increased progressively from about 13 DIV onwards. Kv2.2 expression was clearly evident from the first day in culture and it increased progressively thereafter. The pattern of expression observed for Kv2.2, consistent with previous studies describing the early onset of Kv7.2 expression at the AIS. The expression of ankyrin G [48], which is expressed in the developing AIS from 3 DIV [58].

We next used immunocytochemical analysis to study the localization of these voltage-gated potassium channels in cultured hippocampal neurons. When we studied the localization of Kv7.2 subunit in our cultures (Fig. 1C–D), it was already present in the AIS from 6 DIV onwards, and from 6 DIV onwards, it was detected in ~80% of neurons: 78.64 (\pm 4.38%) at 6 DIV; 80.88 (\pm 1.39%) at 10 DIV; and 84.44 (\pm 2.96%) at 21 DIV. Kv7.2 was distributed homogeneously in the AIS and it was detected along the entire length of the AIS.

development (Fig. 1C). As the expression patterns of Kv7.2 and Kv7.3 at the AIS along with the mechanisms that characterized [48], [50] we subsequently focused on Kv1 and Kv2 channel expression. Accordingly, the results describe Kv1 and Kv2 channels towards the axon lag behind that of sodium channels and Kv7.2 channels.




Figure 7. AIS Kv channel concentration is dependent on ankyrin G.

Confocal microscopy photomicrographs show that interference RNAs against ankyrin G impair the concentrative plating, hippocampal neurons were nucleofected with plasmids expressing scrambled shRNA (A–C, G–I) or ankyrin G shRNA (D–F, J–L) and cultured until 18 DIV. The neurons were then double stained with antibodies against ankyrin G or plkBα (14D4) and Kv1.2 or Kv2.2. Neurons were identified by GFP fluorescence. Ankyrin G, the protein recognized by 14D4 antibodies, Kv1.2 and Kv2.2 were expressed in neurons nucleofected with scrambled shRNA plasmids (A and G) and in non-nucleofected neurons (arrowhead in D–F). However, neurons nucleofected with ankyrin G shRNA (D–E, J–L), no Kv1.2 or Kv2.2 immunostaining was observed at the AIS. Nucleofection did not affect cell somata (K). Scale bar = 25 μm. Arrows indicate AISs and asterisks indicate non-nucleofected neurons.

doi:10.1371/journal.pone.0048557.g007

Kv1.2 Expression in the AIS of Cultured Hippocampal Neurons

To study the expression of low threshold voltage-gated potassium (Kv1) channels during the development of neurons, hippocampal neurons were cultured for different intervals and stained with antibodies raised against the Kv1.2 subunit (Fig. 1). Kv1.2 was expressed in cultured neurons, as identified by 14D4 staining, which co-localizes with the axonal marker SMI-31 [58]. In parallel with axonal growth, 14D4 immunostaining in the AIS, whereas diffuse Kv1.2 staining was observed in neurons, predominantly in the AIS until around 8 DIV (Fig. 2A–E). Consistent with the findings in Western blots, Kv1.2 immunostaining was more prominent in some neurons after 10 DIV (Fig. 2F–H). Indeed, the percentage of neurons exhibiting Kv1.2 staining restricted to the AIS of cultured neurons by 21 DIV (n = 3): 1.46 (±0.25%) at 8 DIV; 5.15 (±0.39%) at 10 DIV; 12.1 (±0.07%) at 13 DIV; and 29.75 (±1.41%) at 21 DIV. Kv1.2 immunostaining was homogeneous, unclustered and mainly concentrated in the region in which 14D4 staining was detected, exhibited Kv1.2 expression at any developmental stage (Fig. 2F–H).

Kv2.2 Expression at the AIS of Cultured Hippocampal Neurons

To study the expression of delayed rectifier voltage-gated potassium channel (Kv2) during the development of the intervals were stained with antibodies directed against the Kv2.2 subunit (Fig. 3), and for sodium channels to identify immunostaining was evident from the initial stages of neuronal differentiation. Indeed, immunocytochemistry reveal processes at all times in culture. In addition, after 14 DIV (Fig. 3D–F) there were intense patches of Kv2.2 immunostaining in contrast to the pattern of Kv1.2 expression observed. The proportion of neurons exhibiting clustered and polarized progressively after 14 DIV, representing over 70% of the cultured neurons at 21 DIV (n = 3): 16.32 (±2.19%) at 11 DIV, 17.32 (±2.19%) at 17 DIV and 74.57 (±7.36%) at 21 DIV.

Mutually Exclusive Kv1.2 and Kv2.2 Expression in the AIS

The differential localization of low-threshold Kv1.2 (distal) and delayed rectifier Kv2.2 potassium channels (proximal) is expressed in a specific region of this structure. Indeed, when we double-stained 21 DIV neurons using antibodies of these subunits at the AIS was mutually exclusive and they were localized in distinct neuronal populations (Fig. 4). 19.47 (±1.85%) of neurons, which in turn exhibited no Kv2.2 immunostaining in the AIS (arrows in Fig. 4). By contrast, the soma and AIS (52.67±1.34%; arrowheads in Fig. 4) no Kv1.2 immunostaining was evident in the AIS. In our experiments, 27.85% ±0.63 of neurons exhibiting Kv1.2/Kv2.2 double immunostaining in the AIS. Moreover, a significant percentage of neurons exhibited Kv1.2/Kv2.2 double immunostaining in the AIS. (27.85% ±0.63; Fig. 4F).

Kv1.2 and Kv2.2 Localization at the AIS is Dependent on Ankyrin G and Independent of the Cytoskeleton

To identify the mechanisms underlying the localization of Kv1.2 and Kv2.2 to the AIS, we first evaluated the resistance property common to other proteins concentrated in the AIS, such as ankyrin G and the protein recognized by 14D4. In 21 DIV hippocampal neurons, Kv1.2 (Fig. 5A) and Kv2.2 (Fig. 5D) expression was still evident in the AIS after extraction with detergent. This suggests that potassium channels in the AIS are associated with scaffolding proteins that are resistant to detergent extraction, such as ankyrin G.

We also assessed whether the localization of Kv1.2 and Kv2.2 in the AIS was dependent on the integrity of the actin cytoskeleton and the structure and function of the AIS [13], [64], and to maintain the structure and neurochemical features of the AIS. We disrupted in neurons by exposing them to cytochalasin D (5 μM) from 15 to 17 DIV, as witnessed by the altered morphology of control neurons (Fig. 6I and L). However, neither the expression nor the distribution of Kv1.2 or Kv2.2 was altered (see arrows in Fig. 6). Hence, the polymerized state of actin microfilaments does not actively influence the distribution of potassium channels at the AIS.

We next investigated the role of ankyrin G in the retention of Kv1.2 and Kv2.2 potassium channels, nucleofecting neurons with ankyrin G shRNA and GFP, and maintaining them in culture until 18 DIV. The absence of ankyrin G expression in neurons transfected with ankyrin G shRNA was verified by ankyrin G immunostaining. No ankyrin G expression was detected in any of the processes of GFP-positive ankyrin G shRNA-nucleofected neurons (Fig. 7J). Moreover, another AIS marker, recognized by 14D4, was absent in ankyrin G shRNA-nucleofected neurons (Fig. 7D), as described previously [72]. By contrast, ankyrin G expression persisted in neurons transfected with scrambled shRNA plasmids (Fig. 7G). In parallel with the loss of ankyrin G and 14D4 staining, no tethering of Kv1.2 and Kv2.2 to the AIS of neurons nucleofected with ankyrin G shRNA, or in any other process emanating from the soma (Fig. 7B, E, I, K), was observed in non-nucleofected neurons (Fig. 2 and 3). Data from three independent experiments showed that the percentage of neurons with clustered and polarized Kv1.2 expression fell significantly (p<0.001), from 16.23 (±2.65%) in scrambled shRNA-nucleofected neurons (total number of nuclei = 106) to 1.89 (±0.97%) in ankyrin G-nucleofected neurons (106 nucleofected neurons). Similarly, the clusters of Kv2.2 immunostaining observed in neurons transfected with scrambled shRNA plasmids (Fig. 7G–I) were absent in ankyrin G-nucleofected neurons (Fig. 7J–L). The mean percentage of neurons exhibiting clustered and polarized Kv2.2 expression fell significantly (p<0.001), from 16.23 (±2.65%) in scrambled shRNA nucleofected neurons (111 nucleofected neurons) to 1.89 (±0.97%) in ankyrin G nucleofected neurons. These results, strongly suggest that Kv1.2 and Kv2.2 tethering and clustering at the AIS is dependent on ankyrin G and independent of the cytoskeleton.

Discussion

The present findings indicate that in contrast to the early expression of voltage-gated sodium channels (VGSC) and Kv2.2 subunits are first tethered at the AIS of cultured hippocampal neurons after 10 days *in vitro* (DIV). Furthermore, Kv1.2 and Kv2.2 are expressed distinctly in the AIS, with each subunit largely restricted to distinct populations of neurons. Our results show that th

AIS is resistant to detergent extraction and like other AIS proteins, it is dependent on the presence of ankyrin G. The subunits in the AIS is not affected by the disruption of the actin cytoskeleton.

AIS Maturation

Cultured hippocampal neurons are widely used as a model to study the development of neuronal polarity [55]. This followed by its subsequent elongation and the development of functionally specialized subdomains, including the AIS and the precise spatial and temporal localization of membrane and cytoskeletal components. Among the first are ankyrin G [56], [57], the protein recognized by 14D4 immunostaining [58], [71] and casein kinase 2 [72], which Ankyrin G is responsible for the accumulation of other structural and functional proteins to the AIS, including VGSC shortly after ankyrin G accumulation in this region [57], [58]. We found that the ankyrin G-dependent targeting and the AIS [48], [49] was a relatively early event in AIS maturation (3–6 DIV), although it occurred after VGSC expression and Kv2 channel expression, which begin to concentrate at the AIS during the second week *in vitro* (10 DIV). The maturation of cultured hippocampal neurons is concomitant with the appearance of gephyrin and GABA_A receptor subunits at the AIS, an organelle, which is involved in Ca²⁺ regulation and reaches neurochemical maturation during the second week *in vitro*. We determined whether these late events in AIS maturation are coordinated with the expression of Kv1 and Kv2 channels in the axonal domain, and to identify the functional consequences of AIS maturation on action potential generation and th

Kv Channel Distribution in the AIS

While the uneven distribution of different Kv channel types is required for proper neuronal function, the specific cell surface channel proteins has not been fully elucidated [73], [74]. We observed a distinct distribution for Kv1, Kv2 and Kv7 channels. No KCNQ channel is expressed in the AIS of different neuronal types, including the rodent adult hippocampus and cerebellum [49], [50], [51]. We found that during the first three weeks of *in vitro* development, the Kv7.2 subunit was homogeneously distributed in the vast majority of hippocampal neurons. This is consistent with the homogeneous distribution of ankyrin G through the AIS. Kv7.2 and Kv7.3 are required for their localization to the AIS [46], [48], [49].

In contrast to Kv7 channels, the Kv1.2 subunit of Kv1 channels was restricted to the distal region of the AIS. AIS compartmentalization and segregation of different VGSCs and the enrichment of the distal AIS with Kv1 channels, and it has been linked with the generation of action potentials in generating and back-propagating APs, respectively [13], [17], [18], [27], [28], [29], [30], [31], [32], [33]. The distal versus the proximal AIS is only observed in certain neuronal types, probably reflecting electrophysiologic differences. These include neocortical pyramidal cells in layer 2/3, interneurons, CA1 pyramidal neurons and retinal ganglion cells, CA1 pyramidal neurons in layer V of the neocortex or in the CA3 region of the hippocampus [17], [18], [28], [35]. In our study, we expressed Kv1.2 at the distal AIS at 21 DIV. This percentage may reflect the proportion of CA1 pyramidal neurons that express Kv1.2 in the AIS of the neuronal population corresponding to the 60% of neurons that do not express Kv1.2.

Kv2 delayed rectifier channels include those comprised of Kv2.1 and Kv2.2 subunits, although they can also form heteromultimers with other subfamilies (Kv5, 6, 8 and 9) [76]. Kv2 channels regulate excitability in hippocampal and cortical neurons rather than the membrane potential repolarization [37], [39], [40], [41], [42], [43], and they are mainly distributed in clusters of around 1.3 μm in diameter dendrites of neocortical and hippocampal neurons [38], [40], [77], [78], [79], [80], [81], [82], [83], [84]. Clusters of Kv2 channels are contact points [85] being also coincident with membrane zones associated with subsurface reticulum cisterns, known as membrane specializations. Kv2.1 clusters overlap with clusters rich in ryanodine receptor Ca²⁺ release channels and the luminal Ca²⁺ binding sites of Kv2 channels in Ca²⁺ regulation [37], [82], [85]. We found that in addition to this somatodendritic domain, Kv2.2 subunits are also expressed in AIS cells, mainly in the proximal AIS. Hence, Kv2.2 subunits may contribute to the maintenance of the AP amplitude in the AIS during high frequency firing, as occurs in neurons of the median nucleus of the trapezoid body [45]. The AIS is composed of stacks of smooth endoplasmic reticulum cisterns. The outermost of these elements is in apposition to IP₃R-expressing microdomains [70]. However, no spatial overlap appears to occur between Kv2.2-expressing AIS microdomains (unpublished observations), suggesting that AIS Kv2 channels are not involved in the IP₃R-mediated

The clustering of Kv2.2 at the AIS described here is consistent with that of the Kv2.1 subunit in hippocampal neurons. Whether Kv2.2 colocalizes with Kv2.1 in the AIS clusters remains unknown. At 21 DIV, Kv2.2 clusters were observed in hippocampal neurons, and a similar proportion of neocortical pyramidal neurons exhibited Kv2.2 somatodendritic immunostaining. The expression of Kv1.2 subunits at the AIS in cultured hippocampal neurons was mutually exclusive. It remains unclear whether the differences in neuronal type within the mature hippocampal formation, or alternatively, a lag in *in vitro* Kv1.2 or Kv2

Mechanisms Mediating AIS Localization of Kv Channels

In recent years, several studies have described mechanisms responsible for the concentration of ion channels at d protein interactions, and have identified amino acid motifs involved in these interactions. However, the functions of and the mechanisms responsible for channel trafficking and clustering at the AIS have yet to be fully characterized

In the AIS, the presence of ankyrin G and its interaction with the actin cytoskeleton through β IV spectrin is critical such as VGSCs, the adhesion molecules neurofascin-186 and NrCAM, and Kv7 potassium channels [16], [46], [48]. Accumulation of the latter occurs through direct binding of Kv7.2 and Kv7.3 subunits to ankyrin G via an ankyrin G [48], [49], [91].

Proteins that form complexes with Kv1 subunits (Kv1.1, Kv1.2 and Kv1.4) include Caspr2, TAG-1 and ADAM22, and localization of Kv1 in the AIS is dependent on the presence of PSD93/Chapsyn-110 and on PDZ domain interaction may also be involved [74], [94]. We found that the accumulation of Kv1.2 subunits in the AIS of cultured hippocampal cytoskeleton and was dependent on the presence of ankyrin G at the AIS, as these subunits were absent in neuro evidence for a direct interaction between ankyrin G and Kv1.2 has been reported [20], our data indicate that the p development, and for the acquisition and/or maintenance of Kv1.2 and Kv2.2 expression at the AIS. This view is in knockdown resulted in the loss of ankyrin G-interacting proteins, such as Na⁺ channels, β IV spectrin and neurofascin casein kinase 2 α , IP₃R1, annexin 6, synaptopodin and α -actinin immunostaining at the AIS [69], [70], [72].

The diverse mechanisms involved in Kv2 channel clustering, including that which occurs at the AIS, remain to be fully dynamic structures, the maintenance and localization of which may depend on the presence of a targeting motif known (PRC) signal [81], or on their interaction with scaffold proteins [95]. Kv2.1 clustering and the voltage-dependence of phosphorylation in response to both neuronal activity-induced Ca²⁺ influx and Ca²⁺ release from internal stores [37] are mobile, although their mean diffusion coefficient is lower than that outside the clusters, suggesting that the clusters cytoskeleton [97], [99]. Kv2.1 clusters are reported to favor cell surface regions not associated with phalloidin-positive depressions in the cortical cytoskeleton corralled by a high density of cortical actin filaments [99], [100]. According hippocampal neurons has been reported to either increase Kv2.1 cluster size [99] or induce complete cluster dissociation more stable than those found in the soma [44]. Moreover, although no measurements of cluster size were performed were not affected by disrupting the actin cytoskeleton with cytochalasin D, which disrupts the diffusion barrier of the asymmetric distribution of other AIS proteins and lipids [64], [65], [101]. Together with the absence of Kv2.2 clusters neurons, these findings suggest that in addition to the actin cytoskeleton, other as yet uncharacterized molecular interactions channel clusters in the AIS.

Author Contributions

Conceived and designed the experiments: AM JGG. Performed the experiments: DS. Analyzed the data: DS AM JGG JD JGG AM. Wrote the paper: DS AM JGG.

References

1. Eunson LH, Rea R, Zuberi SM, Youroukos S, Panayiotopoulos CP, et al. (2000) Clinical, genetic, and experimental channel gene KCNA1 reveal new phenotypic variability. *Ann Neurol* 48: 647–656. doi: 10.1002/1531-8249(200009)48:5<647::AENL210>3.0.CO;2-3
CrossRef • PubMed/NCBI • Google Scholar
2. Manganas LN, Akhtar S, Antonucci DE, Campomanes CR, Dolly JO, et al. (2001) Episodic ataxia type-1 is caused by a distinct folding and intracellular trafficking properties. *J Biol Chem* 276: 49427–49434. doi: 10.1074/jbc.M101010200
CrossRef • PubMed/NCBI • Google Scholar
3. Glasscock E, Yoo JW, Chen TT, Klassen TL, Noebels JL (2010) Kv1.1 potassium channel deficiency reveals a mechanism for sudden unexplained death in epilepsy. *J Neurosci* 30: 5167–5175. doi: 10.1523/JNEUROSCI.4511-10.2010
CrossRef • PubMed/NCBI • Google Scholar
4. Jentsch TJ, Schroeder BC, Kubisch C, Friedrich T, Stein V (2000) Pathophysiology of KCNQ channels: new insights from mouse models. *Epilepsia* 41: 1068–1069. doi: 10.1111/j.1528-1157.2000.tb00302.x
CrossRef • PubMed/NCBI • Google Scholar
5. Misonou H (2010) Homeostatic regulation of neuronal excitability by K(+) channels in normal and diseased neurons. *PLoS One* 5: e11777. doi: 10.1177/1073858409341085

CrossRef • PubMed/NCBI • Google Scholar

6. Nashmi R, Fehlings MG (2001) Mechanisms of axonal dysfunction after spinal cord injury: with an emphasis on potassium channels. *Brain Res Brain Res Rev* 38: 165–191. doi: 10.1016/S0165-0173(01)00134-5.
CrossRef • PubMed/NCBI • Google Scholar
7. Chandy KG (1991) Simplified gene nomenclature. *Nature* 352: 26. doi: 10.1038/352026b0.
CrossRef • PubMed/NCBI • Google Scholar
8. Gutman GA, Chandy KG, Adelman JP, Aiyar J, Bayliss DA, et al. (2003) International Union of Pharmacology classification of potassium channels: potassium channels. *Pharmacol Rev* 55: 583–586. doi: 10.1124/pr.55.4.9.
CrossRef • PubMed/NCBI • Google Scholar
9. Gutman GA, Chandy KG, Grissmer S, Lazdunski M, McKinnon D, et al. (2005) International Union of Pharmacology classification of voltage-gated potassium channels. *Pharmacol Rev* 57: 473–508. doi: 10.1124/pr.57.4.10.
CrossRef • PubMed/NCBI • Google Scholar
10. Yu FH, Yarov-Yarovoy V, Gutman GA, Catterall WA (2005) Overview of molecular relationships in the voltage-gated potassium channel superfamily. *Pharmacol Rev* 57: 387–395. doi: 10.1124/pr.57.4.13.
CrossRef • PubMed/NCBI • Google Scholar
11. Li Y, Um SY, McDonald TV (2006) Voltage-gated potassium channels: regulation by accessory subunits. *Nat Rev Neurosci* 9: 1177–1187. doi: 10.1177/1073858406287717.
CrossRef • PubMed/NCBI • Google Scholar
12. Yu Y, Shu Y, McCormick DA (2008) Cortical action potential backpropagation explains spike threshold variability. *J Neurosci* 28: 7260–7272. doi: 10.1523/JNEUROSCI.1613-08.2008.
CrossRef • PubMed/NCBI • Google Scholar
13. Kole MH, Ilshner SU, Kampa BM, Williams SR, Ruben PC, et al. (2008) Action potential generation requires active propagation into the axon initial segment. *Nat Neurosci* 11: 178–186. doi: 10.1038/nn2040.
CrossRef • PubMed/NCBI • Google Scholar
14. Stuart GJ, Sakmann B (1994) Active propagation of somatic action potentials into neocortical pyramidal cells. *Nat Neurosci* 1: 1038/367069a0. doi: 10.1038/367069a0.
CrossRef • PubMed/NCBI • Google Scholar
15. Stuart G, Schiller J, Sakmann B (1997) Action potential initiation and propagation in rat neocortical pyramidal cells. *J Neurosci* 17: 1469–7793.1997.617ba.x. doi: 10.1111/j.1469-7793.1997.617ba.x.
CrossRef • PubMed/NCBI • Google Scholar
16. Garrido JJ, Giraud P, Carlier E, Fernandes F, Moussif A, et al. (2003) A targeting motif involved in sodium channel localization to the axon initial segment. *Science* 300: 2091–2094. doi: 10.1126/science.1085167.
CrossRef • PubMed/NCBI • Google Scholar
17. Inda MC, DeFelipe J, Munoz A (2006) Voltage-gated ion channels in the axon initial segment of human cortical chandelier cells. *Proc Natl Acad Sci U S A* 103: 2920–2925. doi: 10.1073/pnas.0511197103.
CrossRef • PubMed/NCBI • Google Scholar
18. Lorincz A, Nusser Z (2008) Cell-type-dependent molecular composition of the axon initial segment. *J Neurosci* 28: 4833–4843. doi: 10.1523/JNEUROSCI.4833-08.2008.
CrossRef • PubMed/NCBI • Google Scholar
19. Rasband MN (2010) The axon initial segment and the maintenance of neuronal polarity. *Nat Rev Neurosci* 13: 253–265. doi: 10.1038/nrn2819.
CrossRef • PubMed/NCBI • Google Scholar
20. Rasband MN (2010) Clustered K⁺ channel complexes in axons. *Neurosci Lett* 486: 101–106. doi: 10.1016/j.neulet.2010.07.011.
CrossRef • PubMed/NCBI • Google Scholar
21. Yu Y, Maureira C, Liu X, McCormick D (2010) P/Q and N channels control baseline and spike-triggered calcium transients in axon initial segment. *J Neurosci* 30: 11858–11869. doi: 10.1523/JNEUROSCI.2651-10.2010.
CrossRef • PubMed/NCBI • Google Scholar
22. Bender KJ, Trussell LO (2009) Axon initial segment Ca²⁺ channels influence action potential generation and propagation. *J Neurosci* 29: 10014–10024. doi: 10.1016/j.neuron.2008.12.004.
CrossRef • PubMed/NCBI • Google Scholar
23. Devaux JJ, Kleopa KA, Cooper EC, Scherer SS (2004) KCNQ2 is a nodal K⁺ channel. *J Neurosci* 24: 1231–1240. doi: 10.1523/JNEUROSCI.4833-08.2008.
CrossRef • PubMed/NCBI • Google Scholar

CrossRef • PubMed/NCBI • Google Scholar

24. Ogawa Y, Rasband MN (2008) The functional organization and assembly of the axon initial segment. *Curr Biol* 18: 1016–1026. doi: 10.1016/j.cub.2008.08.008.
CrossRef • PubMed/NCBI • Google Scholar
25. Clark BD, Goldberg EM, Rudy B (2009) Electrogenic tuning of the axon initial segment. *Neuroscientist* 15: 10–18. doi: 10.1177/1043986208325111.
CrossRef • PubMed/NCBI • Google Scholar
26. Duflocq A, Chareyre F, Giovannini M, Couraud F, Davenne M (2011) Characterization of the axon initial segment in a para-AIS and a juxtapara-AIS, organized by protein 4.1B. *BMC Biol* 9: 66. doi: 10.1186/1741-7007-9-66
CrossRef • PubMed/NCBI • Google Scholar
27. Boiko T, Van Wart A, Caldwell JH, Levinson SR, Trimmer JS, et al. (2003) Functional specialization of the axon initial segment: channel targeting. *J Neurosci* 23: 2306–2313. doi: 10.1523/JNEUROSCI.4450-02.2003
CrossRef • PubMed/NCBI • Google Scholar
28. Van Wart A, Matthews G (2006) Expression of sodium channels Nav1.2 and Nav1.6 during postnatal development. *J Neurosci* 26: 1016–1026. doi: 10.1523/JNEUROSCI.2006.05.019.
CrossRef • PubMed/NCBI • Google Scholar
29. Royeck M, Horstmann MT, Remy S, Reitze M, Yaari Y, et al. (2008) Role of axonal Nav1.6 sodium channels in action potential propagation in neurons. *J Neurophysiol* 100: 2361–2380. doi: 10.1152/jn.90332.2008.
CrossRef • PubMed/NCBI • Google Scholar
30. Duflocq A, Le Bras B, Bullier E, Couraud F, Davenne M (2008) Nav1.1 is predominantly expressed in node of Ranvier. *J Neurosci* 28: 180–192. doi: 10.1523/JNEUROSCI.4450-07.2008.
CrossRef • PubMed/NCBI • Google Scholar
31. Hu W, Tian C, Li T, Yang M, Hou H, et al. (2009) Distinct contributions of Nav1.6 and Nav1.2 in action potential propagation in neurons. *J Neurosci* 29: 996–1002. doi: 10.1523/JNEUROSCI.4450-08.2009.
CrossRef • PubMed/NCBI • Google Scholar
32. Kole MH, Letzkus JJ, Stuart GJ (2007) Axon initial segment Kv1 channels control axonal action potential width. *J Neurosci* 27: 633–647. doi: 10.1523/JNEUROSCI.4450-06.2007.
CrossRef • PubMed/NCBI • Google Scholar
33. Shu Y, Duque A, Yu Y, Haider B, McCormick DA (2007) Properties of action-potential initiation in neocortical recordings. *J Neurophysiol* 97: 746–760. doi: 10.1152/jn.00922.2006.
CrossRef • PubMed/NCBI • Google Scholar
34. Bean BP (2007) The action potential in mammalian central neurons. *Nat Rev Neurosci* 8: 451–465. doi: 10.1038/nrn2205.
CrossRef • PubMed/NCBI • Google Scholar
35. Goldberg EM, Clark BD, Zaghera E, Nahmani M, Erisir A, et al. (2008) K⁺ channels at the axon initial segment of neocortical fast-spiking GABAergic interneurons. *Neuron* 58: 387–400. doi: 10.1016/j.neuron.2008.03.003.
CrossRef • PubMed/NCBI • Google Scholar
36. Johnston J, Forsythe ID, Kopp-Scheinflug C (2010) Going native: voltage-gated potassium channels control action potential propagation. *J Physiol* 582: 3187–3200. doi: 10.1113/jphysiol.2010.191973.
CrossRef • PubMed/NCBI • Google Scholar
37. Misonou H, Mohapatra DP, Trimmer JS (2005) Kv2.1: a voltage-gated K⁺ channel critical to dynamic control of action potential propagation. *J Neurosci* 25: 743–752. doi: 10.1523/JNEUROSCI.4450-04.2005.
CrossRef • PubMed/NCBI • Google Scholar
38. Kihira Y, Hermansteyne TO, Misonou H (2010) Formation of heteromeric Kv2 channels in mammalian brain. *J Biol Chem* 285: 10107–10115. doi: 10.1074/jbc.M109.074260.
CrossRef • PubMed/NCBI • Google Scholar
39. Kang J, Huguenard JR, Prince DA (2000) Voltage-gated potassium channels activated during action potential propagation in neocortical pyramidal neurons. *J Neurophysiol* 83: 70–80. doi: 10.1152/jn.2000.83.1.70.
CrossRef • PubMed/NCBI • Google Scholar
40. Bekkers JM (2000) Distribution and activation of voltage-gated potassium channels in cell-attached and outside-out patches of rat pyramidal neurons. *J Physiol* 525 Pt 3: 611–620. doi: 10.1111/j.1469-7793.2000.t01-2-00611.x.
CrossRef • PubMed/NCBI • Google Scholar

41. Colbert CM, Pan E (1999) Arachidonic acid reciprocally alters the availability of transient and sustained de pyramidal neurons. *J Neurosci* 19: 8163–8171.
CrossRef • PubMed/NCBI • Google Scholar
42. Korngreen A, Sakmann B (2000) Voltage-gated K⁺ channels in layer 5 neocortical pyramidal neurones from 525 Pt 3: 621–639. doi: 10.1111/j.1469-7793.2000.00621.x.
CrossRef • PubMed/NCBI • Google Scholar
43. Du J, Haak LL, Phillips-Tansey E, Russell JT, McBain CJ (2000) Frequency-dependent regulation of rat hip channel subunit Kv2.1. *J Physiol* 522 Pt 1: 19–31. doi: 10.1111/j.1469-7793.2000.t01-2-00019.xm.
CrossRef • PubMed/NCBI • Google Scholar
44. Sarmiere PD, Weigle CM, Tamkun MM (2008) The Kv2.1 K⁺ channel targets to the axon initial segment of situ. *BMC Neurosci* 9: 112. doi: 10.1186/1471-2202-9-112.
CrossRef • PubMed/NCBI • Google Scholar
45. Johnston J, Griffin SJ, Baker C, Skrzypiec A, Chernova T, et al. (2008) Initial segment Kv2.2 channels mec frequency action potential firing in medial nucleus of the trapezoid body neurons. *J Physiol* 586: 3493–3505.
CrossRef • PubMed/NCBI • Google Scholar
46. Chung HJ, Jan YN, Jan LY (2006) Polarized axonal surface expression of neuronal KCNQ channels is med KCNQ3 C-terminal domains. *Proc Natl Acad Sci U S A* 103: 8870–8875. doi: 10.1073/pnas.0603376103.
CrossRef • PubMed/NCBI • Google Scholar
47. Schwarz JR, Glassmeier G, Cooper EC, Kao TC, Nodera H, et al. (2006) KCNQ channels mediate IKs, a node of Ranvier. *J Physiol* 573: 17–34. doi: 10.1113/jphysiol.2006.106815.
CrossRef • PubMed/NCBI • Google Scholar
48. Pan Z, Kao T, Horvath Z, Lemos J, Sul JY, et al. (2006) A common ankyrin-G-based mechanism retains K⁺ domains of the axon. *J Neurosci* 26: 2599–2613. doi: 10.1523/JNEUROSCI.4314-05.2006.
CrossRef • PubMed/NCBI • Google Scholar
49. Rasmussen HB, Frokjaer-Jensen C, Jensen CS, Jensen HS, Jorgensen NK, et al. (2007) Requirement of s localization at the axon initial segment. *J Cell Sci* 120: 953–963. doi: 10.1242/jcs.03396.
CrossRef • PubMed/NCBI • Google Scholar
50. Cooper EC (2011) Made for “anchorin”: Kv7.2/7.3 (KCNQ2/KCNQ3) channels and the modulation of neuro Biol 22: 185–192. doi: 10.1016/j.semcd.2010.10.001.
CrossRef • PubMed/NCBI • Google Scholar
51. Klinger F, Gould G, Boehm S, Shapiro MS (2011) Distribution of M-channel subunits KCNQ2 and KCNQ3 in 10.1016/j.neuroimage.2011.07.003.
CrossRef • PubMed/NCBI • Google Scholar
52. Shah MM, Migliore M, Valencia I, Cooper EC, Brown DA (2008) Functional significance of axonal Kv7 char Acad Sci U S A 105: 7869–7874. doi: 10.1073/pnas.0802805105.
CrossRef • PubMed/NCBI • Google Scholar
53. Brown DA, Passmore GM (2009) Neural KCNQ (Kv7) channels. *Br J Pharmacol* 156: 1185–1195.
CrossRef • PubMed/NCBI • Google Scholar
54. Vervaeke K, Gu N, Agdestein C, Hu H, Storm JF (2006) Kv7/KCNQ/M-channels in rat glutamatergic hippoc excitability and transmitter release. *J Physiol* 576: 235–256. doi: 10.1113/jphysiol.2006.111336.
CrossRef • PubMed/NCBI • Google Scholar
55. Banker G, Goslin K (1988) Developments in neuronal cell culture. *Nature* 336: 185–186. doi: 10.1038/336185a0.
CrossRef • PubMed/NCBI • Google Scholar
56. Jenkins SM, Bennett V (2001) Ankyrin-G coordinates assembly of the spectrin-based membrane skeleton, Purkinje neuron initial segments. *J Cell Biol* 155: 739–746. doi: 10.1083/jcb.200109026.
CrossRef • PubMed/NCBI • Google Scholar
57. Boiko T, Vakulenko M, Ewers H, Yap CC, Norden C, et al. (2007) Ankyrin-dependent and -independent me of L1 family members neurofascin and L1/neuron-glia cell adhesion molecule. *J Neurosci* 27: 590–603. doi: 10.1523/JNEUROSCI.4314-05.2006.
CrossRef • PubMed/NCBI • Google Scholar
58. Sanchez-Ponce D, Tapia M, Munoz A, Garrido JJ (2008) New role of IKK alpha/beta phosphorylated I kap

- segment development. *Mol Cell Neurosci* 37: 832–844. doi: 10.1016/j.mcn.2008.01.010.
CrossRef • PubMed/NCBI • Google Scholar
59. Davis JQ, Lambert S, Bennett V (1996) Molecular composition of the node of Ranvier: identification of ank (mucin+/third FNIII domain-) and NrCAM at nodal axon segments. *J Cell Biol* 135: 1355–1367. doi: 10.1083/jcb.135.5.1355.
CrossRef • PubMed/NCBI • Google Scholar
60. Zhou D, Lambert S, Malen PL, Carpenter S, Boland LM, et al. (1998) AnkyrinG is required for clustering of voltage-gated sodium channels at nodes of Ranvier and for normal action potential firing. *J Cell Biol* 143: 1295–1304. doi: 10.1083/jcb.143.5.1295.
CrossRef • PubMed/NCBI • Google Scholar
61. Lemaillet G, Walker B, Lambert S (2003) Identification of a conserved ankyrin-binding motif in the family of ankyrin-like proteins. *J Biol Chem* 278: 27333–27339. doi: 10.1074/jbc.M303327200.
CrossRef • PubMed/NCBI • Google Scholar
62. Hedstrom KL, Xu X, Ogawa Y, Frischknecht R, Seidenbecher CI, et al. (2007) Neurofascin assembles a signaling complex at the node of Ranvier. *J Cell Biol* 178: 875–886. doi: 10.1083/jcb.200705119.
CrossRef • PubMed/NCBI • Google Scholar
63. Hedstrom KL, Ogawa Y, Rasband MN (2008) AnkyrinG is required for maintenance of the axon initial segment. *J Cell Biol* 181: 635–640. doi: 10.1083/jcb.200806112.
CrossRef • PubMed/NCBI • Google Scholar
64. Winckler B, Forscher P, Mellman I (1999) A diffusion barrier maintains distribution of membrane proteins in the axon initial segment. *J Cell Biol* 146: 1038–1046. doi: 10.1083/jcb.146.3.1038.
CrossRef • PubMed/NCBI • Google Scholar
65. Nakada C, Ritchie K, Oba Y, Nakamura M, Hotta Y, et al. (2003) Accumulation of anchored proteins forms a diffusion barrier at the axon initial segment. *Nat Cell Biol* 5: 626–632. doi: 10.1038/ncb1009.
CrossRef • PubMed/NCBI • Google Scholar
66. Nakata T, Hirokawa N (2003) Microtubules provide directional cues for polarized axonal transport through the axon initial segment. *J Cell Biol* 161: 1045–1055. doi: 10.1083/jcb.200302175.
CrossRef • PubMed/NCBI • Google Scholar
67. Tapia M, Wandosell F, Garrido JJ (2010) Impaired function of HDAC6 slows down axonal growth and interferes with the formation of the axon initial segment. *J Neurosci* 30: e12908. doi: 10.1523/JNEUROSCI.0012-10.2010.
CrossRef • PubMed/NCBI • Google Scholar
68. Burkarth N, Kriebel M, Kranz EU, Volkmer H (2007) Neurofascin regulates the formation of gephyrin clusters at the axon hillock of hippocampal neurons. *Mol Cell Neurosci* 36: 59–70. doi: 10.1016/j.mcn.2007.06.001.
CrossRef • PubMed/NCBI • Google Scholar
69. Sanchez-Ponce D, Blazquez-Llorca L, Defelipe J, Garrido JJ, Munoz A (2011c) Colocalization of α -tubulin and AnkyrinG at the Axon Initial Segment. *Cereb Cortex*. doi: 10.1093/cercor/bhr311.
70. Sanchez-Ponce D, DeFelipe J, Garrido JJ, Munoz A (2011b) In vitro maturation of the cisternal organelle in the axon initial segment. *Mol Cell Neurosci* Submitted.
71. Buffington SA, Sobotzik JM, Schultz C, Rasband MN (2012) β -tubulin is not required for axon initial segment formation. *J Cell Biol* 197: 1045–1055. doi: 10.1083/jcb.2012.03.003.
CrossRef • PubMed/NCBI • Google Scholar
72. Sanchez-Ponce D, Munoz A, Garrido JJ (2011) Casein kinase 2 and microtubules control axon initial segment formation. *J Cell Biol* 194: 1045–1055. doi: 10.1083/jcb.2011.09.005.
CrossRef • PubMed/NCBI • Google Scholar
73. Heusser K, Schwappach B (2005) Trafficking of potassium channels. *Curr Opin Neurobiol* 15: 364–369. doi: 10.1016/j.conb.2005.06.001.
CrossRef • PubMed/NCBI • Google Scholar
74. Jensen CS, Rasmussen HB, Misonou H (2011) Neuronal trafficking of voltage-gated potassium channels. *Mol Cell Neurosci* 47: 1045–1055. doi: 10.1016/j.mcn.2011.05.007.
CrossRef • PubMed/NCBI • Google Scholar
75. Palmer LM, Stuart GJ (2006) Site of action potential initiation in layer 5 pyramidal neurons. *J Neurosci* 26: 11523–11533. doi: 10.1523/JNEUROSCI.4812-05.2006.
CrossRef • PubMed/NCBI • Google Scholar

76. Bocksteins E, Raes AL, Van de Vijver G, Bruyns T, Van Bogaert PP, et al. (2009) Kv2.1 and silent Kv subunit cultured small mouse DRG neurons. *Am J Physiol Cell Physiol* 296: C1271–1278. doi: 10.1152/ajpcell.000 CrossRef • PubMed/NCBI • Google Scholar
77. Trimmer JS (1991) Immunological identification and characterization of a delayed rectifier K⁺ channel poly 10764–10768. doi: 10.1073/pnas.88.23.10764. CrossRef • PubMed/NCBI • Google Scholar
78. Hwang PM, Fotuhi M, Bredt DS, Cunningham AM, Snyder SH (1993) Contrasting immunohistochemical localization of the Shab subfamily. *J Neurosci* 13: 1569–1576. CrossRef • PubMed/NCBI • Google Scholar
79. Maletic-Savatic M, Lenn NJ, Trimmer JS (1995) Differential spatiotemporal expression of K⁺ channel poly subunit in situ and in vitro. *J Neurosci* 15: 3840–3851. CrossRef • PubMed/NCBI • Google Scholar
80. Scannevin RH, Murakoshi H, Rhodes KJ, Trimmer JS (1996) Identification of a cytoplasmic domain important for targeting of Kv2.1 K⁺ channel. *J Cell Biol* 135: 1619–1632. doi: 10.1083/jcb.135.6.1619. CrossRef • PubMed/NCBI • Google Scholar
81. Lim ST, Antonucci DE, Scannevin RH, Trimmer JS (2000) A novel targeting signal for proximal clustering of Kv2.1 channels in hippocampal neurons. *J Neurosci* 20: 385–397. doi: 10.1523/JNEUROSCI.0195-04.2004. CrossRef • PubMed/NCBI • Google Scholar
82. Antonucci DE, Lim ST, Vassanelli S, Trimmer JS (2001) Dynamic localization and clustering of dendritic Kv2.1 channels in developing hippocampal neurons. *J Neurosci* 21: 69–81. doi: 10.1523/JNEUROSCI.0195-04.2004. CrossRef • PubMed/NCBI • Google Scholar
83. Guan D, Tkatch T, Surmeier DJ, Armstrong WE, Foehring RC (2007) Kv2 subunits underlie slowly inactivating K⁺ currents in hippocampal neurons. *J Physiol* 581: 941–960. doi: 10.1113/jphysiol.2007.128454. CrossRef • PubMed/NCBI • Google Scholar
84. Murakoshi H, Shi G, Scannevin RH, Trimmer JS (1997) Phosphorylation of the Kv2.1 K⁺ channel alters voltage dependence of activation. *J Neurosci* 17: 821–828. CrossRef • PubMed/NCBI • Google Scholar
85. Du J, Tao-Cheng JH, Zerfas P, McBain CJ (1998) The K⁺ channel, Kv2.1, is apposed to astrocytic processes on the plasma membranes in hippocampal and cortical principal neurons and inhibitory interneurons. *J Neurosci* 18: 37–47. CrossRef • PubMed/NCBI • Google Scholar
86. Berridge MJ (2002) The endoplasmic reticulum: a multifunctional signaling organelle. *Cell Calcium* 32: 235–247. CrossRef • PubMed/NCBI • Google Scholar
87. Paspalas CD, Goldman-Rakic PS (2004) Microdomains for dopamine volume neurotransmission in primate striatum. *J Neurosci* 24: 10152–10161. doi: 10.1523/JNEUROSCI.0195-04.2004. CrossRef • PubMed/NCBI • Google Scholar
88. Kaufmann WA, Ferraguti F, Fukazawa Y, Kasugai Y, Shigemoto R, et al. (2009) Large-conductance calcium-activated potassium channels are clustered at sites of hypolemmal microdomains. *J Comp Neurol* 515: 215–230. doi: 10.1002/ajpa.21000. CrossRef • PubMed/NCBI • Google Scholar
89. Kaufmann WA, Kasugai Y, Ferraguti F, Storm JF (2010) Two distinct pools of large-conductance calcium-activated potassium channels in the plasma membrane of central principal neurons. *J Neurosci* 30: 974–986. doi: 10.1523/JNEUROSCI.0195-04.2004. CrossRef • PubMed/NCBI • Google Scholar
90. Hedstrom KL, Rasband MN (2006) Intrinsic and extrinsic determinants of ion channel localization in neurons. *J Neurosci* 26: 1471–1479. doi: 10.1523/JNEUROSCI.0195-04.2004. CrossRef • PubMed/NCBI • Google Scholar
91. Hill AS, Nishino A, Nakajo K, Zhang G, Fineman JR, et al. (2008) Ion channel clustering at the axon initial segment in early chordates. *PLoS Genet* 4: e1000317. doi: 10.1371/journal.pgen.1000317. CrossRef • PubMed/NCBI • Google Scholar
92. Ogawa Y, Oses-Prieto J, Kim MY, Horresh I, Peles E, et al. (2010) ADAM22, a Kv1 channel-interacting protein, is enriched at the axon initial segment of myelinated axons. *J Neurosci* 30: 1038–1048. doi: 10.1523/JNEUROSCI.0195-04.2004. CrossRef • PubMed/NCBI • Google Scholar

93. Kim E, Niethammer M, Rothschild A, Jan YN, Sheng M (1995) Clustering of Shaker-type K⁺ channels by inorganic guanylate kinases. *Nature* 378: 85–88. doi: 10.1038/378085a0.
CrossRef • PubMed/NCBI • Google Scholar
94. Triller A, Choquet D (2008) New concepts in synaptic biology derived from single-molecule imaging. *Neurosci* 30: 147–157. doi: 10.1016/j.neuron.2008.02.001.
CrossRef • PubMed/NCBI • Google Scholar
95. Surmeier DJ, Foehring R (2004) A mechanism for homeostatic plasticity. *Nat Neurosci* 7: 691–692. doi: 10.1038/nn1260.
CrossRef • PubMed/NCBI • Google Scholar
96. Misonou H, Mohapatra DP, Park EW, Leung V, Zhen D, et al. (2004) Regulation of ion channel localization in dendrites by a cytoskeletal-based perimeter fence. *Neurosci* 7: 711–718. doi: 10.1038/nn1260.
CrossRef • PubMed/NCBI • Google Scholar
97. Tamkun MM, O’Connell K M, Rolig AS (2007) A cytoskeletal-based perimeter fence selectively corrals a subset of Kv2.1 channels. *Sci* 315: 2413–2423. doi: 10.1126/science.1137351.
CrossRef • PubMed/NCBI • Google Scholar
98. Mohapatra DP, Trimmer JS (2006) The Kv2.1 C terminus can autonomously transfer Kv2.1-like phosphorylation, and muscarinic modulation to diverse Kv channels. *J Neurosci* 26: 685–695. doi: 10.1523/JNEUROSCI.1825-06.2006.
CrossRef • PubMed/NCBI • Google Scholar
99. O’Connell KM, Rolig AS, Whitesell JD, Tamkun MM (2006) Kv2.1 potassium channels are retained within dendrites by a perimeter fence. *J Neurosci* 26: 9609–9618. doi: 10.1523/JNEUROSCI.1825-06.2006.
CrossRef • PubMed/NCBI • Google Scholar
100. Weigel AV, Simon B, Tamkun MM, Krapf D (2011) Ergodic and nonergodic processes coexist in the plasma membrane. *Proc Natl Acad Sci U S A* 108: 6438–6443. doi: 10.1073/pnas.1016325108.
CrossRef • PubMed/NCBI • Google Scholar
101. Song AH, Wang D, Chen G, Li Y, Luo J, et al. (2009) A selective filter for cytoplasmic transport at the axon initial segment. *J Cell Biol* 185: 1016–1026. doi: 10.1016/j.jcell.2009.01.016.
CrossRef • PubMed/NCBI • Google Scholar