



# Developmental overnutrition and obesity and type 2 diabetes in offspring

Wei Perng<sup>1,2,3</sup> · Emily Oken<sup>4,5</sup> · Dana Dabelea<sup>1,2,6</sup>

Received: 1 March 2019 / Accepted: 1 May 2019 / Published online: 27 August 2019  
© Springer-Verlag GmbH Germany, part of Springer Nature 2019

## Abstract

Childhood obesity has reached pandemic proportions, and youth-onset type 2 diabetes is following suit. This review summarises the literature on the influence of developmental overnutrition, resulting from maternal diabetes, obesity, maternal dietary intake during pregnancy, excess gestational weight gain, and infant feeding practices, on the aetiology of obesity and type 2 diabetes risk during childhood and adolescence. Key goals of this review are: (1) to summarise evidence to date on consequences of developmental overnutrition; (2) describe shared and distinct biological pathways that may link developmental overnutrition to childhood obesity and youth-onset type 2 diabetes; and (3) to translate current knowledge into clinical and public health strategies that not only target primary prevention in youth, but also encourage primordial prevention during the perinatal period, with the aim of breaking the intergenerational cycle of obesity and diabetes.

**Keywords** Developmental programming · Lifecourse development · Obesity · Overnutrition · Review · Type 1 diabetes · Type 2 diabetes

## Abbreviations

EPOCH Exploring Perinatal Outcomes in Children  
GDM Gestational diabetes mellitus  
GUTS Growing Up Today Study

GWG Gestational weight gain  
HAPO Hyperglycemia and Adverse Pregnancy Outcomes  
HEI Healthy Eating Index

**Electronic supplementary material** The online version of this article (<https://doi.org/10.1007/s00125-019-4914-1>) contains a slide of the figure for download, which is available to authorised users.


✉ Dana Dabelea  
Dana.Dabelea@cuanschutz.edu

- <sup>1</sup> Department of Epidemiology, Colorado School of Public Health, University of Colorado Anschutz Medical Center, Aurora, CO, USA
- <sup>2</sup> Lifecourse Epidemiology of Adiposity and Diabetes (LEAD) Center, Colorado School of Public Health, University of Colorado Anschutz Medical Campus, 13001 East 17th Ave, Box B119, Room W3110, Aurora, CO 80045, USA
- <sup>3</sup> Department of Nutritional Sciences, University of Michigan School of Public Health, Ann Arbor, MI, USA
- <sup>4</sup> Division of Chronic Disease Research Across the Lifecourse (CoRAL), Department of Population Medicine, Harvard Medical School/Harvard Pilgrim Health Care Institute, Boston, MA, USA
- <sup>5</sup> Department of Nutrition, Harvard T.H. Chan School of Public Health, Boston, MA, USA
- <sup>6</sup> Department of Pediatrics, University of Colorado Anschutz Medical Center, Aurora, CO, USA

## Introduction

The obesity pandemic has spared no age group, including young children [1]. Following in its footsteps is youth-onset type 2 diabetes, a novel paediatric condition on the rise in the USA [2] and worldwide [3]. The existence and rise in prevalence of paediatric type 2 diabetes is undoubtedly related to trends in childhood obesity given that excess adiposity is the leading risk factor for type 2 diabetes [4, 5] and emerging evidence suggest that both conditions have origins in utero [6–9]. Little remains known of specific pathways and mechanisms underlying development of youth-onset type 2 diabetes, an important first step to stemming the tide of type 2 diabetes among children and adolescents.

As depicted in Fig. 1, this review expands upon the literature surrounding developmental overnutrition, resulting from maternal diabetes, obesity, diet during pregnancy, and excess gestational weight gain, and infant feeding practices, in relation to childhood obesity and youth-onset type 2 diabetes. We start by summarising and appraising the evidence on



**Fig. 1** Pathways through which exposure to developmental overnutrition during pregnancy (obesity, maternal diabetes, gestational weight gain, diet during pregnancy, and infant feeding) may influence the

development of obesity and type 2 diabetes across the life course. Topics in boxes with a solid border are discussed in depth in this review. This figure is available as a [downloadable slide](#)

consequences of developmental overnutrition and discussing shared and distinct biological pathways that may link developmental overnutrition to obesity and type 2 diabetes in youth. We then translate current knowledge into clinical and public health strategies that not only target primary prevention in youth, but also encourage primordial preventions during the perinatal period, with the aim of breaking the intergenerational cycle of obesity and diabetes [8, 10].

## Developmental overnutrition

### In utero overnutrition

#### Maternal diabetes

Longstanding evidence links maternal diabetes to larger offspring birth size and adiposity across life, and these associations are thought to be driven by maternal fuels: hyperglycaemia and altered lipid and/or amino acid metabolism. While earlier studies evaluated maternal diabetes as a combination of type 1, type 2 and/or gestational diabetes mellitus (GDM), more recent investigations consider diabetes subtypes and degree of hyperglycaemia, which may be more appropriate.

**Maternal diabetes, fetal growth and neonatal adiposity** In the 1950s, Pedersen proposed the fuel-mediated teratogenesis hypothesis, which postulated that intrauterine exposure to hyperglycaemia leads to higher birthweight and future obesity and type 2 diabetes risk [6]. This hypothesis is supported by studies showing that women with pre-existing diabetes and those who develop GDM deliver infants with higher birthweight [11–13] and fat mass [14, 15]. In a study of 195 women with GDM and 220 control individuals [15], mid-

pregnancy fasting glucose was the strongest correlate of newborn fat mass, in comparison with demographic characteristics, family history and maternal anthropometry [15]. Several other studies have since identified associations of maternal hyperglycaemia with offspring adiposity at birth and beyond: (1) maternal glucose levels across all of pregnancy and in the absence of diagnosed diabetes were associated with directly measured neonatal fat mass in the Colorado-based Healthy Start Study ( $n = 804$ ) [16]; (2) mid-pregnancy oral glucose challenge test glucose levels correlated with higher birthweight among 6854 non-diabetic pregnancies in a study conducted in Texas [17]; (3) higher mid-pregnancy oral glucose tolerance test glucose levels were associated with higher birthweight among >25,000 mother–infant pairs in the Hyperglycemia and Adverse Pregnancy Outcomes (HAPO) study [18]; and (4) late pregnancy dysglycaemia among non-GDM pregnancies ( $\text{HbA}_{1c} \geq 39$  mmol/mol at delivery) predicted greater offspring weight gain during early childhood and higher BMI at age 4 years among 898 mother–child pairs in Germany [19]. Together, these findings emphasise the relevance of both degree and timing of maternal hyperglycaemia—even in the absence of frank diabetes—as determinants of offspring adiposity.

Type 1 diabetes, type 2 diabetes and GDM are each associated with altered lipid metabolism [20–22], another fuel-mediated pathway through which maternal diabetes may influence newborn adiposity [23]. In the context of GDM, maternal serum NEFA and/or triacylglycerols are associated with higher birthweight [24–26]. Findings in the general population have been mixed, with some studies suggesting that the relationship between maternal lipids and neonatal outcomes differs by pre-pregnancy weight status [27–30]. In the Healthy Heart Study [16] there was a positive association of NEFA during the second half of pregnancy with birthweight independent of pre-pregnancy BMI, but not with newborn fat

mass, among 804 mother–infant pairs. There was also a positive relationship between late pregnancy cholesterol and newborn fat mass in overweight/obese women, but not among lean women. Such findings emphasise a need to better understand how specific maternal fuels, singly and in combination with other aspects of maternal health and metabolism, may trigger fuel-mediated overnutrition.

Not surprisingly, given the interplay between lipid and amino acid metabolism [31], altered maternal serum amino acid concentrations have been observed in concomitance with maternal hyperglycaemia. In a study of 67 HAPO participants, targeted metabolomics profiling of maternal serum revealed differences in concentrations of metabolites on amino acid and macronutrient degradation pathways between BMI-matched women with high (>90th percentile) vs low fasting glucose (<10th percentile) at 28 weeks' gestation [32]. In light of the interrelationships among biochemical mechanisms of nutrient metabolism, future studies interrogating comprehensive metabolite profiles of maternal blood during pregnancy will shed light on cohesive biological pathways linking maternal hyperglycaemia to offspring health.

**Maternal diabetes and offspring obesity** The relationship between maternal diabetes and offspring adiposity starts at birth and tracks across the life course. In a survey-based analysis of >14,000 US youth in the Growing Up Today Study (GUTS) [33], maternal GDM correlated with 40% higher odds of being overweight during adolescence (OR 1.4, 95% CI 1.2, 1.6). The estimate was attenuated after accounting for maternal pre-pregnancy BMI (OR 1.2, 95% CI 0.8, 1.7). In the Project Viva pre-birth cohort, among 366 boys aged 6–10 years old, the difference in fat mass between those born to mothers with GDM vs those born to normoglycaemic mothers was attenuated from 2.6 kg (95% CI 1.0, 4.2) to 2.0 kg (95% CI 0.4, 3.6) after adjusting for pre-pregnancy BMI [34]. Similarly, in 461 participants of the Exploring Perinatal Outcomes in CHildren (EPOCH) cohort, GDM exposure correlated with higher BMI, waist circumference, visceral and subcutaneous adipose tissue in offspring at age 6–13 years [35]. The investigators noted modest attenuation (i.e. by 14–42%) in estimates of interest after accounting for pre-pregnancy BMI, with some associations retaining statistical significance [35]. A recent investigation of >5000 youth in The Environmental Determinants of Diabetes in the Young (TEDDY) study reported higher odds of being overweight in 5-year-old offspring exposed to maternal GDM (OR 1.48, 95% CI 1.14, 1.92), type 1 diabetes (OR 1.60, 95% CI 1.16, 2.20) and type 2 diabetes (OR 7.39, 95% CI 2.46, 22.23) compared with their unexposed counterparts [36]; adjustment for maternal pre-pregnancy BMI attenuated all three estimates. Similarly, three meta-analyses reported that adjustment for maternal BMI attenuated, but did not completely abolish, associations of GDM with offspring obesity and abnormal

glucose tolerance during childhood [37–39]. Taken together, the evidence suggests an independent effect of exposure to diabetes in utero on future adiposity and type 2 diabetes risk.

We note that although inclusion of maternal pre-pregnancy BMI in regression models exploring associations between GDM and offspring adiposity may partly account for genetic predisposition, such adjustment may also control for shared intrauterine mechanisms that lead to fetal overnutrition, since maternal glucose is elevated among overweight/obese women, even if they do not qualify as having diabetes. Use of appropriate analytical approaches (i.e. inverse probability weighting [40] to balance the distribution of pre-pregnancy BMI among women with and without GDM rather than simple adjustment, which may inadvertently block the effect of shared aetiology) and mechanistic studies, will help to ascertain the extent to which accounting for maternal BMI isolates associations of interest.

**Maternal diabetes and offspring type 2 diabetes** The longitudinal study of Pima Indians in the Gila River Indian Community [41] was one of the first studies to explore associations of in utero exposure to maternal diabetes (combined pre-existing type 2 diabetes and GDM) with subsequent type 2 diabetes in offspring. In this high-risk population, offspring of women with diabetes not only had higher birthweight [41], but also continued on a trajectory of higher weight-for-height through adolescence [42, 43] and had a tenfold greater risk of developing type 2 diabetes in adolescence and young adulthood [10]. Moreover, exposure to maternal diabetes was the single strongest risk factor for youth-onset type 2 diabetes (OR 10.4, 95% CI 4.3, 25.1) and accounted for most of the dramatic increase in youth-onset type 2 diabetes in this population over the last 30 years [44].

Investigations in lower-risk, racially diverse populations support findings from the Pima Indians. In the SEARCH for Diabetes in Youth Case–Control Study, the odds of type 2 diabetes was 7.3 (95% CI 3.2, 16.8) greater in participants whose mothers were diagnosed with diabetes during pregnancy ( $n = 79$ ; with >90% GDM cases) than in their unexposed counterparts ( $n = 190$ ) [45]. In addition, in utero exposure to maternal diabetes in conjunction with obesity contributed to 47% of type 2 diabetes cases in adolescent offspring of various race/ethnicities (non-Hispanic White, African American, Hispanic), suggesting that the transgenerational cycle of diabetes begetting diabetes at increasingly younger ages operates in diverse populations and race/ethnic groups. While such findings also reflect the genetic component of type 2 diabetes, the relationship between in utero exposure to maternal diabetes and future risk of type 2 diabetes is robust to adjustment for paternal diabetes and age at onset of diabetes for either parent [46]. Dabelea et al [47] showed further support for a specific intrauterine effect of maternal diabetes on offspring type 2 diabetes risk above and beyond genetics via a discordant

sibship analysis wherein the sibling born after maternal diagnosis of diabetes had threefold greater odds of type 2 diabetes than those born before diagnosis.

### Pathways linking in utero overnutrition to type 2 diabetes

Although excess adiposity is the leading risk factor for type 2 diabetes in adults [4, 5], whether the relationship between in utero overnutrition and youth-onset type 2 diabetes is a sole consequence of childhood obesity has garnered interest. In the Pima Indian study, acute insulin response to infused glucose was 40% lower in adults whose mothers had diabetes during pregnancy than in those whose mothers developed diabetes after delivery, despite no differences in per cent fat mass [48]. In lower-risk populations, maternal GDM has been associated with precursors of type 2 diabetes in offspring even after accounting for offspring BMI, including higher estimated insulin resistance (HOMA2-IR) among youth in EPOCH [49], and higher fasting glucose among adolescents in the Danish National Birth Cohort [50]. Similarly, in an analysis of 587 mother–offspring dyads in Denmark, Kelstrup et al found that adult offspring exposed to maternal type 1 diabetes or GDM had impaired insulin sensitivity and lower disposition index compared with adult offspring of normoglycaemic women, even after adjustment for BMI [51].

In addition to the effect of overt maternal diabetes on type 2 diabetes in offspring, a recent analysis of 4832 mother–child pairs in the HAPO Follow-Up Study revealed that the entire spectrum of maternal glycaemia (based on maternal fasting glucose levels, as well as plasma glucose levels at 1 h and 2 h post 75 g oral glucose tolerance load) was positively associated with offspring fasting glucose levels and insulin resistance at age 10–14 years, independent of maternal and child BMI and family history of diabetes [9].

Mechanisms underlying the relationship between in utero exposure to maternal diabetes and type 2 diabetes risk in offspring have been gleaned from rodent studies showing a specific detrimental effect of maternal diabetes or hyperglycaemia on offspring pancreatic beta cell development and function [52–54]. In addition, small case–control studies of mother–infant pairs with vs without GDM have noted differential expression of genes encoding the insulin receptor [55] and adiponectin [56] in cord blood, independent of maternal BMI, pointing towards epigenetic modifications of specific genes involved in glycaemic regulation as another mechanistic pathway. Further research to identify the exact mechanisms by which in utero exposure to diabetes influences risk of type 2 diabetes in offspring is needed to develop and implement effective prevention.

### Maternal obesity

Vohr et al reported that maternal pre-pregnancy weight status and gestational weight gain predicted offspring fat mass at birth [57] and age 1 year [58], even among women without

GDM. This positive relationship between maternal weight and offspring adiposity at birth [59] and beyond [49, 60–64] has been confirmed in numerous settings. Moreover, the consequences of pregravid adiposity are detectable across the continuum of maternal BMI and can influence offspring metabolic risk independent of offspring adiposity [64–67].

The concordance between maternal and offspring obesity can stem from genetics [68], shared environment and lifestyle [69, 70], as well as intrauterine mechanisms. The intrauterine effect of maternal obesity on offspring obesity is difficult to isolate in human studies, but animal experiments support biological plausibility. Diet-induced obesity among pregnant rodents altered offspring adipocyte metabolism to favour hypertrophy via epigenetic modifications [71–74]. Other pathways include the effect of maternal obesity on placental function [75], glycaemic regulation [76] and stem cell differentiation [77–81]. We include a brief discussion on mechanistic studies using mesenchymal stem cells in the section ‘Mechanistic studies nested within existing cohorts’; other mechanisms are reviewed elsewhere in this issue.

### Maternal diet during pregnancy

**Macronutrient intake** Studies of maternal macronutrient intake during pregnancy generally indicate that higher energy and carbohydrate intakes and lower protein intakes are associated with higher neonatal adiposity. Specifically, greater carbohydrate intake during late pregnancy was associated with higher neonatal fat mass in a study of 222 Danish mother–child pairs [82], whereas higher protein intake during mid-to-late pregnancy was associated with lower birthweight (Project Viva [83]), neonatal abdominal adiposity (the GUSTO Study [84]) and abdominal fat mass during adolescence (a Danish cohort [85]). In the Healthy Start Study, maternal intake of all energy-providing macronutrients (total fat, saturated fat, unsaturated fat, carbohydrates) except protein was associated with higher neonatal adiposity [86].

**Dietary patterns** The field of nutritional epidemiology recently shifted towards evaluating dietary patterns, rather than individual foods or nutrients, to reflect real-life dietary intake [87]. In the Healthy Start Study, poor diet quality during mid-pregnancy, defined as Healthy Eating Index (HEI) score  $\leq 57$ , corresponded to 0.58% (95% CI: 0.07%, 1.10%) higher fat mass in newborns [88]. A pooled analysis of two cohorts (Project Viva in the USA and the Rhea cohort in Greece) found that adherence to a Mediterranean dietary pattern during mid-pregnancy predicted lower BMI, waist circumference and skinfold thicknesses in offspring across childhood [89]. Yet, a recent study of 721 overweight/obese pregnant women reported no consistent relationship of diet quality indicators, including the HEI, carbohydrate and protein intake and total energy intake, with fetal ultrasound measurements of adiposity at 28–

36 weeks' gestation [90]. While the utility of fetal ultrasounds for assessing neonatal adiposity requires validation, these findings suggest the importance of maternal pre-pregnancy weight status beyond that of diet during pregnancy.

**Gestational weight gain** While gestational weight gain (GWG) is not a dietary factor per se, it is a consequence of dietary intake that contributes to the gestational milieu [91–93], albeit in conjunction with the influence of pre-pregnancy weight status, and the social, genetic and physiological changes that occur during pregnancy, discussed elsewhere [94]. Both higher GWG on a continuous scale, as well as excess GWG according to current Institute of Medicine guidelines [94], have been consistently related to greater offspring adiposity from birth [95, 96] through adulthood [66, 97, 98]. Many of these studies established this association independent of pre-pregnancy BMI and shared environment/lifestyle factors [64, 97, 98]. GWG is also positively correlated with type 2 diabetes-related metabolic biomarkers in offspring, including insulin resistance and adipocytokine profile [99, 100]. In some cases, the metabolic alterations occurred in the absence of offspring obesity [64, 76].

In addition to the impact of total GWG on offspring adiposity and metabolic profile, the timing of weight gain has repercussions. Greater weight gain assessed continuously during early pregnancy has been linked to higher offspring BMI and fat mass during childhood [101, 102], whereas GWG exceeding current guidelines during the second and third trimesters correlated with greater odds of delivering a large-for-gestational age infant [103]. These discrepancies suggest a need to examine associations of GWG timing with growth trajectories rather than weight status at distinct time points.

Of note, while it is tempting to compare the effects of pre-pregnancy BMI with those of GWG on offspring health, as was done in a recent meta-analysis [104], Gillman [105] rightly pointed out that doing so may not be appropriate given that their relationships with offspring health reflect contributions from different factors at different times across a woman's lifespan. GWG occurs during pregnancy, exerting influences on offspring health through the intrauterine environment. On the other hand, pre-pregnancy BMI represents the impact of shared genes and environmental factors between mother and child, as well as a direct influence of the in utero environment and postnatal behaviours.

## Postnatal overnutrition

### Breastfeeding

Exclusive breastfeeding for the first 6 months of life is recommended by the World Health Organization for its protection against infant morbidity [106], benefits to maternal health [106] and potential to reduce childhood obesity [107, 108].

Although some have expressed concern regarding the safety of breastfeeding among diabetic women given the potential for higher levels of insulin and glucose in breast milk [109], studies in diverse populations (e.g. the Pima Indian study [110] and GUTS [111]) reported protective effects of breastfeeding on offspring fat distribution, metabolic traits and type 2 diabetes risk in diabetic and non-diabetic women. Beyond supporting the safety of breastfeeding, Crume et al [112] showed that breastfeeding  $\geq 6$  months mitigated the effects of GDM exposure on adiposity at age 6–13 years, suggesting a specific protective effect of breastfeeding among high-risk offspring.

## Moving from association to causation to inform clinical care and public health practice

### Life course approach

The long latency period between exposure to developmental overnutrition and future obesity/type 2 diabetes necessitates thoughtful study designs and analytical approaches to study disease aetiology, to test mechanisms and to identify opportunities for intervention. Life course epidemiology conceptual models [113] are a valuable tool for these purposes. Conceptual models not only encourage researchers to consider the web of causation among key variables, but also drive the analytical strategy. By accomplishing these tasks, researchers will better understand aetiology and mechanisms, and gain insight into modifiable determinants of type 2 diabetes across the lifespan that both coincide with and occur independently of excess adiposity.

### Clever study designs and analytical strategies

Certain study designs can enhance our ability to draw causal conclusions from observational data. For instance, one could minimise the impact of genetics by comparing effect sizes for maternal vs paternal BMI [114], and/or leverage within- vs between-family comparisons [115] and sibship analyses [47, 116]. In addition, new developments in statistical techniques (i.e. inverse proportional weighting of marginal structural models [117, 118]) can enhance causal inference.

### Mechanistic studies nested within existing cohorts

In the Healthy Start cohort, in vitro studies of mesenchymal stem cells harvested from cord blood in a sample of study participants provided evidence that alterations in  $\beta$ -catenin pathways, expression of genes involved in myocyte growth, amino acid synthesis and oxidative stress link maternal obesity to newborn adiposity and weight gain during infancy [77–81]. Such findings support biological plausibility of

observational findings of the positive correlation between maternal and child obesity in this and other cohorts.

### Randomised clinical trials

Most randomised clinical trials comprising interventions focused on obesogenic conditions during pregnancy have not been effective in mitigating/reducing GDM or excess GWG [119], or preventing macrosomia [119, 120]. The limited success of these trials point towards a need for interventions prior to conception or earlier in pregnancy. Pre-conception interventions may be difficult to implement among first-time mothers given that half of pregnancies are unplanned [121], and early pregnancy interventions may be similarly challenging given that most women seek prenatal care midway through the first trimester. A potential strategy is to focus on women with a history of obesity, GDM and/or excessive GWG for surveillance and interventions prior to the next pregnancy [122]. In light of several recent studies indicating the importance of the first trimester to both maternal [123] and offspring health [101, 102], interventions on maternal behaviours to moderate weight gain during the first trimester seem promising.

There is also room for improvement in the endpoints targeted by pregnancy trials. For example, directly measured neonatal fat mass rather than birthweight is likely to be more relevant to the development of obesity and type 2 diabetes. Furthermore, longer follow-up of health outcomes beyond birth is warranted to better understand the efficacy of pregnancy trials on chronic disease risk prevention.

Finally, following findings that adequate breastfeeding limits the detrimental effects of GDM exposure [112], promoting breastfeeding among diabetic women may help to reduce obesity-related conditions in high-risk offspring, hopefully breaking the intergenerational cycle of disease.

### Summary and conclusions

Youth-onset type 2 diabetes is on the rise, and trends in childhood obesity only partially explain the recent appearance of a condition that was previously confined to adults. Higher maternal BMI entering pregnancy (not simply maternal obesity), maternal hyperglycaemia (even in the absence of overt diabetes), greater GWG (not only excessive GWG according to current guidelines), and greater energy intake during pregnancy are important early-life correlates of excess adiposity and youth-onset type 2 diabetes. Importantly, maternal hyperglycaemia and GDM are associated with precursors of type 2 diabetes (e.g. insulin resistance, reduced disposition index) in offspring, starting as early as late childhood, even after accounting for current body size and/or adiposity, suggesting a specific effect of maternal hyperglycaemia on pancreatic beta cell development and function. Given the

overlapping nature of these aspects of developmental overnutrition, and the shared aetiology of obesity and type 2 diabetes, future studies are warranted to disentangle pathways linking specific aspects of developmental overnutrition to obesity and type 2 diabetes risk. Accomplishment of these tasks will inform timing and targets for early interventions with potential to make measurable impacts on population health.

**Duality of interest** The authors declare that there is no duality of interest associated with this manuscript.

**Contribution statement** WP and DD conceptualised the review. WP wrote the initial draft of the manuscript, incorporated changes suggested by co-authors, and formatted the paper for publication. DD and EO provided critical intellectual feedback. All authors approved the version of the paper to be published.

### References

1. Skinner AC, Ravanbakht SN, Skelton JA, Perrin EM, Armstrong SC (2018) Prevalence of obesity and severe obesity in US children, 1999–2016. *Pediatrics* 141(3):e20173459. <https://doi.org/10.1542/peds.2017-3459>
2. Mayer-Davis EJ, Lawrence JM, Dabelea D et al (2017) Incidence trends of type 1 and type 2 diabetes among youths, 2002–2012. *N Engl J Med* 376(15):1419–1429. <https://doi.org/10.1056/NEJMoal610187>
3. D’Adamo E, Caprio S (2011) Type 2 diabetes in youth: epidemiology and pathophysiology. *Diabetes Care* 34(Suppl 2):S161–S165. <https://doi.org/10.2337/dc11-s212>
4. Chan JM, Rimm EB, Colditz GA, Stampfer MJ, Willett WC (1994) Obesity, fat distribution, and weight gain as risk factors for clinical diabetes in men. *Diabetes Care* 17(9):961–969. <https://doi.org/10.2337/diacare.17.9.961>
5. Carey VJ, Walters EE, Colditz GA et al (1997) Body fat distribution and risk of non-insulin-dependent diabetes mellitus in women. The Nurses’ Health Study. *Am J Epidemiol* 145(7):614–619. <https://doi.org/10.1093/oxfordjournals.aje.a009158>
6. Pedersen J (1971) Diabetes mellitus and pregnancy: present status of the hyperglycaemia–hyperinsulinism theory and the weight of the newborn baby. *Postgrad Med J* 1971(Suppl):66–67
7. Dabelea D, Harrod CS (2013) Role of developmental overnutrition in pediatric obesity and type 2 diabetes. *Nutr Rev* 71(Suppl 1):S62–S67. <https://doi.org/10.1111/nure.12061>
8. Dabelea D, Crume T (2011) Maternal environment and the transgenerational cycle of obesity and diabetes. *Diabetes* 60(7):1849–1855. <https://doi.org/10.2337/db11-0400>
9. Scholtens DM, Kuang A, Lowe LP et al (2019) Hyperglycemia and Adverse Pregnancy Outcome Follow-up Study (HAPO FUS): maternal glycemia and childhood glucose metabolism. *Diabetes Care* 42(3):381–392. <https://doi.org/10.2337/dc18-2021>
10. Dabelea D, Hanson RL, Bennett PH, Roumain J, Knowler WC, Pettitt DJ (1998) Increasing prevalence of Type II diabetes in American Indian children. *Diabetologia* 41(8):904–910. <https://doi.org/10.1007/s001250051006>
11. Berk MA, Mimouni F, Miodovnik M, Hertzberg V, Valuck J (1989) Macrosomia in infants of insulin-dependent diabetic mothers. *Pediatrics* 83(6):1029–1034

12. Kc K, Shakya S, Zhang H (2015) Gestational diabetes mellitus and macrosomia: a literature review. *Ann Nutr Metab* 66(Suppl 2):14–20. <https://doi.org/10.1159/000371628>
13. Ladfors L, Shaat N, Wiberg N, Katarasou A, Berntorp K, Kristensen K (2017) Fetal overgrowth in women with type 1 and type 2 diabetes mellitus. *PLoS One* 12(11):e0187917. <https://doi.org/10.1371/journal.pone.0187917>
14. Lampl M, Jeanty P (2004) Exposure to maternal diabetes is associated with altered fetal growth patterns: a hypothesis regarding metabolic allocation to growth under hyperglycemic-hypoxic conditions. *Am J Hum Biol* 16(3):237–263. <https://doi.org/10.1002/ajhb.20015>
15. Catalano PM, Thomas A, Huston-Presley L, Amini SB (2003) Increased fetal adiposity: a very sensitive marker of abnormal in utero development. *Am J Obstet Gynecol* 189(6):1698–1704. [https://doi.org/10.1016/S0002-9378\(03\)00828-7](https://doi.org/10.1016/S0002-9378(03)00828-7)
16. Crume TL, Shapiro AL, Brinton JT et al (2015) Maternal fuels and metabolic measures during pregnancy and neonatal body composition: The Healthy Start Study. *J Clin Endocrinol Metab* 100(4):1672–1680. <https://doi.org/10.1210/jc.2014-2949>
17. Yogev Y, Langer O, Xenakis EM, Rosenn B (2005) The association between glucose challenge test, obesity and pregnancy outcome in 6390 non-diabetic women. *J Matern Fetal Neonatal Med* 17(1):29–34. <https://doi.org/10.1080/14767050400028766>
18. Metzger BE, Lowe LP, Dyer AR et al (2008) Hyperglycemia and adverse pregnancy outcomes. *N Engl J Med* 358(19):1991–2002. <https://doi.org/10.1056/NEJMoa0707943>
19. Gomes D, von Kries R, Delius M et al (2018) Late-pregnancy dysglycemia in obese pregnancies after negative testing for gestational diabetes and risk of future childhood overweight: an interim analysis from a longitudinal mother–child cohort study. *PLoS Med* 15(10):e1002681. <https://doi.org/10.1371/journal.pmed.1002681>
20. Jones AI, Coleman EL, Husni NR et al (2017) Type 1 diabetes alters lipid handling and metabolism in human fibroblasts and peripheral blood mononuclear cells. *PLoS One* 12(12):e0188474. <https://doi.org/10.1371/journal.pone.0188474>
21. Ji J, Petropavlovskaja M, Khatchadourian A et al (2019) Type 2 diabetes is associated with suppression of autophagy and lipid accumulation in  $\beta$ -cells. *J Cell Mol Med* 23(4):2890–2900. <https://doi.org/10.1111/jcmm.14172>
22. Ryckman K, Spracklen C, Smith C, Robinson J, Saftlas A (2015) Maternal lipid levels during pregnancy and gestational diabetes: a systematic review and meta-analysis. *BJOG* 122(5):643–651. <https://doi.org/10.1111/1471-0528.13261>
23. Tontonoz P, Hu E, Spiegelman BM (1994) Stimulation of adipogenesis in fibroblasts by PPAR $\gamma$ 2, a lipid-activated transcription factor. *Cell* 79(7):1147–1156. [https://doi.org/10.1016/0092-8674\(94\)90006-X](https://doi.org/10.1016/0092-8674(94)90006-X)
24. Schaefer-Graf UM, Meitzner K, Ortega-Senovilla H et al (2011) Differences in the implications of maternal lipids on fetal metabolism and growth between gestational diabetes mellitus and control pregnancies. *Diabet Med* 28(9):1053–1059. <https://doi.org/10.1111/j.1464-5491.2011.03346.x>
25. Schaefer-Graf UM, Graf K, Kulbacka I et al (2008) Maternal lipids as strong determinants of fetal environment and growth in pregnancies with gestational diabetes mellitus. *Diabetes Care* 31(9):1858–1863. <https://doi.org/10.2337/dc08-0039>
26. Simeonova-Krstevska S, Krstevska B, Velkoska-Nakova V et al (2014) Effect of lipid parameters on foetal growth in gestational diabetes mellitus pregnancies. *Pril (Makedon Akad Nauk Umet Odd Med Nauki)* 35(2):131–136
27. Misra VK, Trudeau S, Pemi U (2011) Maternal serum lipids during pregnancy and infant birth weight: the influence of prepregnancy BMI. *Obesity (Silver Spring, Md)* 19(7):1476–1481. <https://doi.org/10.1038/oby.2011.43>
28. Kulkarni SR, Kumaran K, Rao SR et al (2013) Maternal lipids are as important as glucose for fetal growth: findings from the Pune Maternal Nutrition Study. *Diabetes Care* 36(9):2706–2713. <https://doi.org/10.2337/dc12-2445>
29. Harmon KA, Gerard L, Jensen DR et al (2011) Continuous glucose profiles in obese and normal-weight pregnant women on a controlled diet: metabolic determinants of fetal growth. *Diabetes Care* 34(10):2198–2204. <https://doi.org/10.2337/dc11-0723>
30. Clausen T, Burski TK, Oyen N, Godang K, Bollerslev J, Henriksen T (2005) Maternal anthropometric and metabolic factors in the first half of pregnancy and risk of neonatal macrosomia in term pregnancies. A prospective study. *Eur J Endocrinol* 153(6):887–894. <https://doi.org/10.1530/eje.1.02034>
31. Newgard CB (2012) Interplay between lipids and branched-chain amino acids in development of insulin resistance. *Cell Metab* 15(5):606–614. <https://doi.org/10.1016/j.cmet.2012.01.024>
32. Scholtens DM, Muehlbauer MJ, Daya NR et al (2014) Metabolomics reveals broad-scale metabolic perturbations in hyperglycemic mothers during pregnancy. *Diabetes Care* 37(1):158–166. <https://doi.org/10.2337/dc13-0989>
33. Gillman MW, Rifas-Shiman S, Berkey CS, Field AE, Colditz GA (2003) Maternal gestational diabetes, birth weight, and adolescent obesity. *Pediatrics* 111(3):e221–e226. <https://doi.org/10.1542/peds.111.3.e221>
34. Regnault N, Gillman MW, Rifas-Shiman SL, Eggleston E, Oken E (2013) Sex-specific associations of gestational glucose tolerance with childhood body composition. *Diabetes Care* 36(10):3045–3053. <https://doi.org/10.2337/dc13-0333>
35. Crume TL, Ogden L, West NA et al (2011) Association of exposure to diabetes in utero with adiposity and fat distribution in a multiethnic population of youth: the Exploring Perinatal Outcomes among Children (EPOCH) study. *Diabetologia* 54(1):87–92. <https://doi.org/10.1007/s00125-010-1925-3>
36. Pitchika A, Vehik K, Hummel S et al (2018) Associations of maternal diabetes during pregnancy with overweight in offspring: results from the Prospective TEDDY Study. *Obesity (Silver Spring, Md)* 26(9):1457–1466. <https://doi.org/10.1002/oby.22264>
37. Kawasaki M, Arata N, Miyazaki C et al (2018) Obesity and abnormal glucose tolerance in offspring of diabetic mothers: A systematic review and meta-analysis. *PLoS One* 13(1):e0190676. <https://doi.org/10.1371/journal.pone.0190676>
38. Kim SY, England JL, Sharma JA, Njoroge T (2011) Gestational diabetes mellitus and risk of childhood overweight and obesity in offspring: a systematic review. *Exp Diabetes Res* 2011:541308–541309. <https://doi.org/10.1155/2011/541308>
39. Philipps LH, Santhakumaran S, Gale C et al (2011) The diabetic pregnancy and offspring BMI in childhood: a systematic review and meta-analysis. *Diabetologia* 54(8):1957–1966. <https://doi.org/10.1007/s00125-011-2180-y>
40. Hernán MA, Hernández-Díaz S, Robins JM (2004) A structural approach to selection bias. *Epidemiology* 15(5):615–625. <https://doi.org/10.1097/01.ede.0000135174.63482.43>
41. Bennett PH, Rushforth NB, Miller M, LeCompte PM (1976) Epidemiologic studies of diabetes in the Pima Indians. *Recent Prog Horm Res* 32:333–376
42. Pettitt DJ, Baird HR, Aleck KA, Bennett PH, Knowler WC (1983) Excessive obesity in offspring of Pima Indian women with diabetes during pregnancy. *N Engl J Med* 308(5):242–245. <https://doi.org/10.1056/nejm198302033080502>
43. Pettitt DJ, Bennett PH, Knowler WC, Baird HR, Aleck KA (1985) Gestational diabetes mellitus and impaired glucose tolerance during pregnancy. Long-term effects on obesity and glucose tolerance in the offspring. *Diabetes* 34(Suppl 2):119–122. <https://doi.org/10.2337/diab.34.2.S119>
44. Dabelea D, Knowler WC, Pettitt DJ (2000) Effect of diabetes in pregnancy on offspring: follow-up research in the Pima Indians. J

- Matern Fetal Med 9(1):83–88. [https://doi.org/10.1002/\(sici\)1520-6661\(200001/02\)9:1<83::aid-mfm17>3.0.co;2-o](https://doi.org/10.1002/(sici)1520-6661(200001/02)9:1<83::aid-mfm17>3.0.co;2-o)
45. Dabelea D, Mayer-Davis EJ, Lamichhane AP et al (2008) Association of intrauterine exposure to maternal diabetes and obesity with type 2 diabetes in youth: the SEARCH Case-Control Study. *Diabetes Care* 31(7):1422–1426. <https://doi.org/10.2337/dc07-2417>
  46. Pettitt D (1996) Diabetes in subsequent generations. In: Doornhorst A, Hadden DR (eds) *Diabetes and pregnancy: an international approach to diagnosis and management*. John Wiley & Sons, Chichester, pp 367–376
  47. Dabelea D, Hanson RL, Lindsay RS et al (2000) Intrauterine exposure to diabetes conveys risks for type 2 diabetes and obesity: a study of discordant sibships. *Diabetes* 49(12):2208–2211. <https://doi.org/10.2337/diabetes.49.12.2208>
  48. Gautier JF, Wilson C, Weyer C et al (2001) Low acute insulin secretory responses in adult offspring of people with early onset type 2 diabetes. *Diabetes* 50(8):1828–1833. <https://doi.org/10.2337/diabetes.50.8.1828>
  49. Sauder KA, Hockett CW, Ringham BM, Glueck DH, Dabelea D (2017) Fetal overnutrition and offspring insulin resistance and beta-cell function: the Exploring Perinatal Outcomes among Children (EPOCH) study. *Diabet Med* 34(10):1392–1399. <https://doi.org/10.1111/dme.13417>
  50. Grunnet LG, Hansen S, Hjort L et al (2017) Adiposity, dysmetabolic traits, and earlier onset of female puberty in adolescent offspring of women with gestational diabetes mellitus: a clinical study within the Danish National Birth Cohort. *Diabetes Care* 40(12):1746–1755. <https://doi.org/10.2337/dc17-0514>
  51. Kelstrup L, Damm P, Mathiesen ER et al (2013) Insulin resistance and impaired pancreatic beta-cell function in adult offspring of women with diabetes in pregnancy. *J Clin Endocrinol Metab* 98(9):3793–3801. <https://doi.org/10.1210/jc.2013-1536>
  52. Han J, Xu J, Long YS, Epstein PN, Liu YQ (2007) Rat maternal diabetes impairs pancreatic beta-cell function in the offspring. *Am J Physiol Endocrinol Metab* 293(1):E228–E236. <https://doi.org/10.1152/ajpendo.00479.2006>
  53. Aerts L, Van Assche FA (2006) Animal evidence for the transgenerational development of diabetes mellitus. *Int J Biochem Cell Biol* 38(5–6):894–903. <https://doi.org/10.1016/j.biocel.2005.07.006>
  54. Aerts L, Sodoyez-Goffaux F, Sodoyez JC, Malaisse WJ, Van Assche FA (1988) The diabetic intrauterine milieu has a long-lasting effect on insulin secretion by B cells and on insulin uptake by target tissues. *Am J Obstet Gynecol* 159(5):1287–1292. <https://doi.org/10.5555/uri:pii:0002937888904656>
  55. Ott R, Melchior K, Stupin JH et al (2019) Reduced insulin receptor expression and altered DNA methylation in fat tissues and blood of women with GDM and offspring. *J Clin Endocrinol Metab* 104(1):137–149. <https://doi.org/10.1210/jc.2018-01659>
  56. Ott R, Stupin JH, Melchior K et al (2018) Alterations of adiponectin gene expression and DNA methylation in adipose tissues and blood cells are associated with gestational diabetes and neonatal outcome. *Clin Epigenetics* 10(1):131. <https://doi.org/10.1186/s13148-018-0567-z>
  57. Vohr BR, McGarvey ST, Coll CG (1995) Effects of maternal gestational diabetes and adiposity on neonatal adiposity and blood pressure. *Diabetes Care* 18(4):467–475. <https://doi.org/10.2337/diacare.18.4.467>
  58. Vohr BR, McGarvey ST (1997) Growth patterns of large-for-gestational-age and appropriate-for-gestational-age infants of gestational diabetic mothers and control mothers at age 1 year. *Diabetes Care* 20(7):1066–1072. <https://doi.org/10.2337/diacare.20.7.1066>
  59. Starling AP, Brinton JT, Glueck DH et al (2015) Associations of maternal BMI and gestational weight gain with neonatal adiposity in the Healthy Start Study. *Am J Clin Nutr* 101(2):302–309. <https://doi.org/10.3945/ajcn.114.094946>
  60. Sauder KA, Kaar JL, Starling AP, Ringham BM, Glueck DH, Dabelea D (2017) Predictors of infant body composition at 5 months of age: the Healthy Start Study. *J Pediatr* 183:94–99.e91. <https://doi.org/10.1016/j.jpeds.2017.01.014>
  61. Shapiro AL, Schmiede SJ, Brinton JT et al (2015) Testing the fuel-mediated hypothesis: maternal insulin resistance and glucose mediate the association between maternal and neonatal adiposity, the Healthy Start study. *Diabetologia* 58(5):937–941. <https://doi.org/10.1007/s00125-015-3505-z>
  62. Sharp GC, Lawlor DA, Richmond RC et al (2015) Maternal prepregnancy BMI and gestational weight gain, offspring DNA methylation and later offspring adiposity: findings from the Avon Longitudinal Study of Parents and Children. *Int J Epidemiol* 44(4):1288–1304. <https://doi.org/10.1093/ije/dyv042>
  63. Santos S, Monnereau C, Felix JF, Duijts L, Gaillard R, Jaddoe VVW (2018) Maternal body mass index, gestational weight gain, and childhood abdominal, pericardial, and liver fat assessed by magnetic resonance imaging. *Int J Obes* 43(3):581–593. <https://doi.org/10.1038/s41366-018-0186-y>
  64. Peng W, Gillman MW, Mantzoros CS, Oken E (2014) A prospective study of maternal prenatal weight and offspring cardiometabolic health in midchildhood. *Ann Epidemiol* 24(11):793–800.e1. <https://doi.org/10.1016/j.annepidem.2014.08.002>
  65. Lemas DJ, Brinton JT, Shapiro AL, Glueck DH, Friedman JE, Dabelea D (2015) Associations of maternal weight status prior and during pregnancy with neonatal cardiometabolic markers at birth: the Healthy Start study. *Int J Obes* 39(10):1437–1442. <https://doi.org/10.1038/ijo.2015.109>
  66. Hochner H, Friedlander Y, Calderon-Margalit R et al (2012) Associations of maternal prepregnancy body mass index and gestational weight gain with adult offspring cardiometabolic risk factors: the Jerusalem Perinatal Family Follow-up Study. *Circulation* 125(11):1381–1389. <https://doi.org/10.1161/circulationaha.111.070060>
  67. Yu Z, Han S, Zhu J, Sun X, Ji C, Guo X (2013) Pre-pregnancy body mass index in relation to infant birth weight and offspring overweight/obesity: a systematic review and meta-analysis. *PLoS One* 8(4):e61627–e61627. <https://doi.org/10.1371/journal.pone.0061627>
  68. Maes HH, Neale MC, Eaves LJ (1997) Genetic and environmental factors in relative body weight and human adiposity. *Behav Genet* 27(4):325–351. <https://doi.org/10.1023/A:1025635913927>
  69. Scaglioni S, Arrizza C, Vecchi F, Tedeschi S (2011) Determinants of children's eating behavior. *Am J Clin Nutr* 94(6 Suppl):2006s–2011s. <https://doi.org/10.3945/ajcn.110.001685>
  70. Robinson SM (2005) Godfrey KM (2008) feeding practices in pregnancy and infancy: relationship with the development of overweight and obesity in childhood. *Int J Obes* 32(Suppl 6):S4–S10. <https://doi.org/10.1038/ijo.2008.201>
  71. Begum G, Davies A, Stevens A et al (2013) Maternal undernutrition programs tissue-specific epigenetic changes in the glucocorticoid receptor in adult offspring. *Endocrinology*. 154(12):4560–4569. <https://doi.org/10.1210/en.2013-1693>
  72. Samuelsson AM, Matthews PA, Argenton M et al (2008) Diet-induced obesity in female mice leads to offspring hyperphagia, adiposity, hypertension, and insulin resistance: a novel murine model of developmental programming. *Hypertension* 51(2):383–392. <https://doi.org/10.1161/HYPERTENSIONAHA.107.101477>
  73. Bringhenti I, Moraes-Teixeira JA, Cunha MR, Ornellas F, Mandarim-de-Lacerda CA, Aguilu MB (2013) Maternal obesity during the preconception and early life periods alters pancreatic development in early and adult life in male mouse offspring. *PLoS One* 8(1):e55711. <https://doi.org/10.1371/journal.pone.0055711>



74. Howie GJ, Sloboda DM, Kamal T, Vickers MH (2009) Maternal nutritional history predicts obesity in adult offspring independent of postnatal diet. *J Physiol* 587(Pt 4):905–915. <https://doi.org/10.1113/jphysiol.2008.163477>
75. Howell KR, Powell TL (2017) Effects of maternal obesity on placental function and fetal development. *Reproduction* 153(3):R97–R108. <https://doi.org/10.1530/REP-16-0495>
76. Mingrone G, Manco M, Mora MEV et al (2008) Influence of maternal obesity on insulin sensitivity and secretion in offspring. *Diabetes Care* 31(9):1872–1876. <https://doi.org/10.2337/dc08-0432>
77. Boyle KE, Patinkin ZW, Shapiro ALB et al (2017) Maternal obesity alters fatty acid oxidation, AMPK activity, and associated DNA methylation in mesenchymal stem cells from human infants. *Mol Metab* 6(11):1503–1516. <https://doi.org/10.1016/j.molmet.2017.08.012>
78. Boyle KE, Patinkin ZW, Shapiro AL, Baker PR 2nd, Dabelea D, Friedman JE (2016) Mesenchymal stem cells from infants born to obese mothers exhibit greater potential for adipogenesis: The Healthy Start BabyBUMP project. *Diabetes* 65(3):647–659. <https://doi.org/10.2337/db15-0849>
79. Baker PR 2nd, Patinkin ZW, Shapiro ALB et al (2017) Altered gene expression and metabolism in fetal umbilical cord mesenchymal stem cells correspond with differences in 5-month-old infant adiposity gain. *Sci Rep* 7(1):18095. <https://doi.org/10.1038/s41598-017-17588-4>
80. Baker PR, 2nd, Patinkin Z, Shapiro AL, et al. (2017) Maternal obesity and increased neonatal adiposity correspond with altered infant mesenchymal stem cell metabolism. *JCI Insight* 2(21):pii: 94200. <https://doi.org/10.1172/jci.insight.94200>
81. Shapiro AL, Boyle KE, Dabelea D et al (2016) Nicotinamide promotes adipogenesis in umbilical cord-derived mesenchymal stem cells and is associated with neonatal adiposity: The Healthy Start BabyBUMP Project. *PLoS One* 11(7):e0159575. <https://doi.org/10.1371/journal.pone.0159575>
82. Renault KM, Carlsen EM, Norgaard K et al (2015) Intake of carbohydrates during pregnancy in obese women is associated with fat mass in the newborn offspring. *Am J Clin Nutr* 102(6):1475–1481. <https://doi.org/10.3945/ajcn.115.110551>
83. Switkowski KM, Jacques PF, Must A, Kleinman KP, Gillman MW, Oken E (2016) Maternal protein intake during pregnancy and linear growth in the offspring. *Am J Clin Nutr* 104(4):1128–1136. <https://doi.org/10.3945/ajcn.115.128421>
84. Chen LW, Tint MT, Fortier MV et al (2016) Maternal macronutrient intake during pregnancy is associated with neonatal abdominal adiposity: The Growing Up in Singapore Towards healthy Outcomes (GUSTO) Study. *J Nutr* 146(8):1571–1579. <https://doi.org/10.3945/jn.116.230730>
85. Maslova E, Hansen S, Grunnet LG et al (2017) Maternal protein intake in pregnancy and offspring metabolic health at age 9–16 y: results from a Danish cohort of gestational diabetes mellitus pregnancies and controls. *Am J Clin Nutr* 106(2):623–636. <https://doi.org/10.3945/ajcn.115.128637>
86. Crume TL, Brinton JT, Shapiro A et al (2016) Maternal dietary intake during pregnancy and offspring body composition: The Healthy Start Study. *Am J Obstet Gynecol* 215(5):609.e601–609.e608. <https://doi.org/10.1016/j.ajog.2016.06.035>
87. Hu FB (2002) Dietary pattern analysis: a new direction in nutritional epidemiology. *Curr Opin Lipidol* 13(1):3–9. <https://doi.org/10.1097/00041433-200202000-00002>
88. Shapiro AL, Kaar JL, Crume TL et al (2016) Maternal diet quality in pregnancy and neonatal adiposity: the Healthy Start Study. *Int J Obes* (2005) 40(7):1056–1062. <https://doi.org/10.1038/ijo.2016.79>
89. Chatzi L, Rifas-Shiman SL, Georgiou V et al (2017) Adherence to the Mediterranean diet during pregnancy and offspring adiposity and cardiometabolic traits in childhood. *Pediatric obesity* 12(Suppl 1):47–56. <https://doi.org/10.1111/ijpo.12191>
90. O'Brien CM, Louise J, Deussen A, Dodd JM (2018) In Overweight or obese pregnant women, maternal dietary factors are not associated with fetal growth and adiposity. *Nutrients* 10(7):870. <https://doi.org/10.3390/nu10070870>
91. Tobias D, Bao W (2014) Diet during pregnancy and gestational weight gain. *Curr Nutri Rep* 3(3):289–297. <https://doi.org/10.1007/s13668-014-0092-4>
92. Hrolfsdottir L, Schalkwijk CG, Birgisdottir BE et al (2016) Maternal diet, gestational weight gain, and inflammatory markers during pregnancy. *Obesity* (Silver Spring, Md) 24(10):2133–2139. <https://doi.org/10.1002/oby.21617>
93. Shin D, Bianchi L, Chung H, Weatherspoon L, Song WO (2014) Is gestational weight gain associated with diet quality during pregnancy? *Matern Child Health J* 18(6):1433–1443. <https://doi.org/10.1007/s10995-013-1383-x>
94. Institute of Medicine National Research Council Committee to Reexamine IOM Pregnancy Weight Guidelines (2009) In: Rasmussen KM, Yaktine AL (eds) *Weight gain during pregnancy: reexamining the guidelines*. Washington, National Academies Press (US)
95. Deierlein AL, Siega-Riz AM, Adair LS, Herring AH (2011) Effects of pre-pregnancy body mass index and gestational weight gain on infant anthropometric outcomes. *J Pediatr* 158(2):221–226. <https://doi.org/10.1016/j.jpeds.2010.08.008>
96. Blackwell SC, Landon MB, Mele L et al (2016) Relationship between excessive gestational weight gain and neonatal adiposity in women with mild gestational diabetes mellitus. *Obstet Gynecol* 128(6):1325–1332. <https://doi.org/10.1097/AOG.0000000000001773>
97. Oken E, Taveras EM, Kleinman KP, Rich-Edwards JW, Gillman MW (2007) Gestational weight gain and child adiposity at age 3 years. *Am J Obstet Gynecol* 196(4):322.e321–322.e328. <https://doi.org/10.1016/j.ajog.2006.11.027>
98. Oken E, Rifas-Shiman SL, Field AE, Frazier AL, Gillman MW (2008) Maternal gestational weight gain and offspring weight in adolescence. *Obstet Gynecol* 112(5):999–1006. <https://doi.org/10.1097/AOG.0b013e31818a5d50>
99. Dello Russo M, Ahrens W, De Vriendt T et al (2013) Gestational weight gain and adiposity, fat distribution, metabolic profile, and blood pressure in offspring: the IDEFICS project. *Int J Obes* (2005) 37(7):914–919. <https://doi.org/10.1038/ijo.2013.35>
100. Fraser A, Tilling K, Macdonald-Wallis C et al (2011) Associations of gestational weight gain with maternal body mass index, waist circumference, and blood pressure measured 16 y after pregnancy: the Avon Longitudinal Study of Parents and Children (ALSPAC). *Am J Clin Nutr* 93(6):1285–1292. <https://doi.org/10.3945/ajcn.110.008326>
101. Margerison-Zilko CE, Shrimali BP, Eskenazi B, Lahiff M, Lindquist AR, Abrams BF (2012) Trimester of maternal gestational weight gain and offspring body weight at birth and age five. *Matern Child Health J* 16(6):1215–1223. <https://doi.org/10.1007/s10995-011-0846-1>
102. Hivert MF, Rifas-Shiman SL, Gillman MW, Oken E (2016) Greater early and mid-pregnancy gestational weight gains are associated with excess adiposity in mid-childhood. *Obesity* (Silver Spring, Md) 24(7):1546–1553. <https://doi.org/10.1002/oby.21511>
103. Sridhar SB, Xu F, Hedderson MM (2016) Trimester-Specific gestational weight gain and infant size for gestational age. *PLoS One* 11(7):e0159500–e0159500. <https://doi.org/10.1371/journal.pone.0159500>
104. Voerman E, Santos S, Patro Golab B et al (2019) Maternal body mass index, gestational weight gain, and the risk of overweight and obesity across childhood: an individual participant data meta-analysis. *PLoS Med* 16(2):e1002744. <https://doi.org/10.1371/journal.pmed.1002744>

105. Gillman MW (2012) Gestational weight gain: now and the future. *Circulation* 125(11):1339–1340. <https://doi.org/10.1161/circulationaha.112.091751>
106. Kramer MS, Kakuma R (2012) Optimal duration of exclusive breastfeeding. *Cochrane Database Syst Rev*, Issue 8, Art. no.: CD003517. <https://doi.org/10.1002/14651858.CD003517.pub2>
107. World Health Organization (2014) Exclusive breastfeeding to reduce the risk of childhood overweight and obesity. Available from [www.who.int/elena/titles/bbc/breastfeeding\\_childhood\\_obesity/en/](http://www.who.int/elena/titles/bbc/breastfeeding_childhood_obesity/en/). Accessed 7 April 2019
108. Oken E, Fields DA, Lovelady CA, Redman LM (2017) TOS Scientific Position Statement: Breastfeeding and Obesity. *Obesity* (Silver Spring, Md) 25(11):1864–1866. <https://doi.org/10.1002/oby.22024>
109. Jovanovic-Peterson L, Fuhrmann K, Hedden K, Walker L, Peterson CM (1989) Maternal milk and plasma glucose and insulin levels: studies in normal and diabetic subjects. *J Am Coll Nutr* 8(2):125–131. <https://doi.org/10.1080/07315724.1989.10720287>
110. Pettitt DJ, Knowler WC (1998) Long-term effects of the intrauterine environment, birth weight, and breast-feeding in Pima Indians. *Diabetes Care* 21(Suppl 2):B138–B141
111. Mayer-Davis EJ, Rifas-Shiman SL, Zhou L, Hu FB, Colditz GA, Gillman MW (2006) Breast-feeding and risk for childhood obesity: does maternal diabetes or obesity status matter? *Diabetes Care* 29(10):2231–2237. <https://doi.org/10.2337/dc06-0974>
112. Crume TL, Ogden L, Maligie M et al (2011) Long-term impact of neonatal breastfeeding on childhood adiposity and fat distribution among children exposed to diabetes in utero. *Diabetes Care* 34(3):641–645. <https://doi.org/10.2337/dc10-1716>
113. Kuh D, Ben-Shlomo Y, Lynch J, Hallqvist J, Power C (2003) Life course epidemiology. *J Epidemiol Community Health* 57(10):778–783. <https://doi.org/10.1136/jech.57.10.778>
114. Gaillard R, Steegers EA, Duijts L et al (2014) Childhood cardiometabolic outcomes of maternal obesity during pregnancy: the Generation R Study. *Hypertension* 63(4):683–691. <https://doi.org/10.1161/hypertensionaha.113.02671>
115. Branum AM, Parker JD, Keim SA, Schempf AH (2011) Prepregnancy body mass index and gestational weight gain in relation to child body mass index among siblings. *Am J Epidemiol* 174(10):1159–1165. <https://doi.org/10.1093/aje/kwr250>
116. Villamor E, Cnattingius S (2006) Interpregnancy weight change and risk of adverse pregnancy outcomes: a population-based study. *Lancet* 368(9542):1164–1170. [https://doi.org/10.1016/S0140-6736\(06\)69473-7](https://doi.org/10.1016/S0140-6736(06)69473-7)
117. Young JG, Hernan MA, Picciotto S, Robins JM (2010) Relation between three classes of structural models for the effect of a time-varying exposure on survival. *Lifetime Data Anal* 16(1):71–84. <https://doi.org/10.1007/s10985-009-9135-3>
118. Snowden JM, Rose S, Mortimer KM (2011) Implementation of G-computation on a simulated data set: demonstration of a causal inference technique. *Am J Epidemiol* 173(7):731–738. <https://doi.org/10.1093/aje/kwq472>
119. Dodd JM, Grivell RM, Crowther CA, Robinson JS (2010) Antenatal interventions for overweight or obese pregnant women: a systematic review of randomised trials. *BJOG* 117(11):1316–1326. <https://doi.org/10.1111/j.1471-0528.2010.02540.x>
120. Dodd JM, Deussen AR, Mohamad I et al (2016) The effect of antenatal lifestyle advice for women who are overweight or obese on secondary measures of neonatal body composition: the LIMIT randomised trial. *BJOG* 123(2):244–253. <https://doi.org/10.1111/1471-0528.13796>
121. Finer LB, Zolna MR (2016) Declines in unintended pregnancy in the United States, 2008–2011. *N Engl J Med* 374(9):843–852. <https://doi.org/10.1056/NEJMsal506575>
122. Ratner RE, Christophi CA, Metzger BE et al (2008) Prevention of diabetes in women with a history of gestational diabetes: effects of metformin and lifestyle interventions. *J Clin Endocrinol Metab* 93(12):4774–4779. <https://doi.org/10.1210/jc.2008-0772>
123. Hedderson MM, Gunderson EP, Ferrara A (2010) Gestational weight gain and risk of gestational diabetes mellitus. *Obstet Gynecol* 115(3):597–604. <https://doi.org/10.1097/AOG.0b013e3181cfce4f>

**Publisher's note** Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.