

 Open access • Journal Article • DOI:10.1136/BMJ.E281

## Diagnosis and management of the epilepsies in adults and children: summary of updated NICE guidance — [Source link](#)

[Vanessa Delgado Nunes](#), [L. Sawyer](#), [J Neilson](#), [Grammati Sarri](#) ...+1 more authors

**Institutions:** [Royal College of Physicians](#)

**Published on:** 26 Jan 2012 - [BMJ](#) (BMJ)

**Topics:** [Epilepsy](#), [Cost effectiveness](#) and [Guideline](#)

Related papers:

- [Definition of drug resistant epilepsy: consensus proposal by the ad hoc Task Force of the ILAE Commission on Therapeutic Strategies.](#)
- [Revised Terminology and Concepts for Organization of Seizures and Epilepsies: Report of the ILAE Commission on Classification and Terminology, 2005-2009](#)
- [Early Identification of Refractory Epilepsy](#)
- [ILAE Official Report: A practical clinical definition of epilepsy](#)
- [Updated ILAE evidence review of antiepileptic drug efficacy and effectiveness as initial monotherapy for epileptic seizures and syndromes](#)

Share this paper:    

View more about this paper here: <https://typeset.io/papers/diagnosis-and-management-of-the-epilepsies-in-adults-and-4r377fuq7j>

## CORRECTIONS

### Diagnosis and management of the epilepsies in adults and children: summary of updated NICE guidance

In this article by Vanessa Delgado Nunes and colleagues an editing error occurred in the section “Adjunctive treatment for generalised tonic-clonic seizures” (*BMJ* 2012;344:e281, doi:10.1136/bmj.e281). The second bullet point should read: “If there are absence seizures or myoclonic seizures [not “If myoclonic seizures are absent or if juvenile myoclonic epilepsy is

suspected”], do not offer carbamazepine, gabapentin, oxcarbazepine, phenytoin, pregabalin, tiagabine, or vigabatrin. (New recommendation.)”

Cite this as: *BMJ* 2012;344:e1173

© BMJ Publishing Group Ltd 2012