



Diaporthe: a genus of endophytic, saprobic and plant pathogenic fungi

R.R. Gomes¹, C. Glienke¹, S.I.R. Videira², L. Lombard², J.Z. Groenewald²,
P.W. Crous^{2,3,4}

Key words

Diaporthales
Diaporthe
Multi-Locus Sequence Typing (MLST)
Phomopsis
systematics

Abstract *Diaporthe* (*Phomopsis*) species have often been reported as plant pathogens, non-pathogenic endophytes or saprobes, commonly isolated from a wide range of hosts. The primary aim of the present study was to resolve the taxonomy and phylogeny of a large collection of *Diaporthe* species occurring on diverse hosts, either as pathogens, saprobes, or as harmless endophytes. In the present study we investigated 243 isolates using multi-locus DNA sequence data. Analyses of the rDNA internal transcribed spacer (ITS1, 5.8S, ITS2) region, and partial translation elongation factor 1-alpha (TEF1), beta-tubulin (TUB), histone H3 (HIS) and calmodulin (CAL) genes resolved 95 clades. Fifteen new species are described, namely *Diaporthe arengae*, *D. brasiliensis*, *D. endophytica*, *D. hongkongensis*, *D. inconspicua*, *D. infecunda*, *D. mayteni*, *D. neoarctii*, *D. oxe*, *D. paranensis*, *D. pseudomangiferae*, *D. pseudophoenicicola*, *D. raonikayaporum*, *D. schini* and *D. terebinthifoliae*. A further 14 new combinations are introduced in *Diaporthe*, and *D. anacardii* is epitypified. Although species of *Diaporthe* have in the past chiefly been distinguished based on host association, results of this study confirm several taxa to have wide host ranges, suggesting that they move freely among hosts, frequently co-colonising diseased or dead tissue. In contrast, some plant pathogenic and endophytic taxa appear to be strictly host specific. Given this diverse ecological behaviour among members of *Diaporthe*, future species descriptions lacking molecular data (at least ITS and HIS or TUB) should be strongly discouraged.

Article info Received: 27 November 2012; Accepted: 24 February 2013; Published: 28 March 2013.

INTRODUCTION

Species of *Diaporthe* and their *Phomopsis* asexual states have broad host ranges and are widely distributed, occurring as plant pathogens, endophytes or saprobes, but also as pathogens of humans and other mammals (Webber & Gibbs 1984, Carroll 1986, Boddy & Griffith 1989, Rehner & Uecker 1994, Garcia-Reyne et al. 2011, Udayanga et al. 2011). *Diaporthe* spp. are responsible for diseases on a wide range of plants hosts, some of which are economically important worldwide, causing root and fruit rots, dieback, cankers, leaf spots, blights, decay and wilt (Uecker 1988, Mostert et al. 2001a, van Rensburg et al. 2006, Santos et al. 2011, Thompson et al. 2011).

Currently, MycoBank (accessed Sept. 2012) lists more than 1 000 names in the genus *Phomopsis*, while *Diaporthe* contains more than 860 names. In the past species have chiefly been described under the assumption they are host-specific, leading to a proliferation of names based on the hosts from which they were isolated (Uecker 1988). However, subsequent studies have found that many species are able to colonise diverse hosts as opportunists, and that several different species could even co-occur on the same host or lesion (Brayford 1990, Rehner & Uecker 1994, Mostert et al. 2001a, Farr et al. 2002, Crous & Groenewald 2005). Curiously, some species of *Diaporthe* can be either pathogenic or harmless endophytes depending on the

host and its health. For example, *D. phaseolorum* is pathogenic to soybean (Santos et al. 2011), but endophytic in mangroves (*Laguncularia racemosa*) (Sebastiane et al. 2011). With the deletion of Art. 59 from the International Code of Nomenclature for algae, fungi, and plants (ICN), asexual and sexual names of fungi receive equal status (Hawksworth et al. 2011, Wingfield et al. 2012). Because the name *Diaporthe* (1870) predates *Phomopsis* (1905), *Diaporthe* is adopted in the present study for this group of fungi (Santos et al. 2010, 2011, Crous et al. 2011, Udayanga et al. 2012).

Diaporthe (incl. its *Phomopsis* state) has been reported as one of the most frequently encountered genera of endophytic fungi in several plant hosts (Murali et al. 2006, Botella & Diez 2011). The genus has also frequently been recognised as a producer of interesting enzymes and secondary metabolites (Isaka et al. 2001, Kobayashi et al. 2003, Dai et al. 2005, Elsaesser et al. 2005) with antibiotic (Bandre & Sasek 1977, Dettrakul et al. 2003, Lin et al. 2005) or anticancer (Kumaran & Hur 2009) activity. Furthermore, species of *Diaporthe* have in the past been noted to deter herbivory (Brayford 1990, Weber 2009, Vesterlund et al. 2011), have lignocellulolytic activities (Jordaan et al. 2006), or have been applied as bioherbicides (Ash et al. 2010).

The accurate application of accepted names of plant pathogenic fungi is essential for the development of effective biosecurity and trade policies (Crous & Groenewald 2005, Wingfield et al. 2012). The taxonomy of many groups of plant pathogenic fungi has in the past been based on host association (Crous et al. 2013, Groenewald et al. 2013). Although some species of *Diaporthe* are host specific, a great number have been noted to occur on more than one host (Brayford 1990, Rehner & Uecker 1994, Farr et al. 2002). Similar observations led Wehmeyer

¹ Department of Genetics, Universidade Federal do Paraná, Centro Politécnico, Box 19071, 81531-990, Curitiba, Brazil.

² CBS-KNAW Fungal Biodiversity Centre, Uppsalaalaan 8, 3584 CT, Utrecht, The Netherlands; corresponding author e-mail: p.crous@cbs.knaw.nl.

³ Microbiology, Department of Biology, Utrecht University, Padualaan 8, 3584 CH Utrecht, The Netherlands.

⁴ Wageningen University and Research Centre (WUR), Laboratory of Phytopathology, Droevedaalsesteeg 1, 6700 PB Wageningen, The Netherlands.

Table 1 Host/substrate, locality, collector and GenBank accession numbers of strains included in the study.

Species	Original name	Strain ¹	Isolation source	Host family	Locality	Collector	GenBank Accession numbers (ITS, CAL, HIS, TEF1, TUB) ²
<i>Diaporthe acutigena</i>	<i>D. acutigena</i>	CBS 129521; CPC 17622 (ex-type)	<i>Acacia retinodes</i> , leaves	<i>Mimosaceae</i>	Australia	P.W. Crous, I.G. Pascoe & J. Edwards	KC343247 KC343489 KC3433731 KC343973
<i>Diaporthe acerina</i>	<i>D. acerina</i>	CBS 137.27	<i>Acer saccharum</i>	<i>Aceraceae</i>	–	L.E. Wehnrey	KC343248 KC343490 KC343732 KC343974
<i>Diaporthe alleghaniensis</i>	<i>D. alleghaniensis</i>	CBS 495.72; ATCC 24097 (ex-type)	<i>Betula alleghaniensis</i> , branches	<i>Betulaceae</i>	Canada	R.H. Arnold	KC343733 KC343975
<i>Diaporthe alnea</i>	<i>D. alnea</i>	CBS 146.46	<i>Alnus sp.</i>	<i>Betulaceae</i>	–	S. Truter	KC343734 KC343976
<i>Diaporthe ambigua</i>	<i>D. ambigua</i>	CBS 159.47	<i>Alnus sp.</i>	<i>Betulaceae</i>	South Africa	S. Denman	KC343735 KC343977 KC343978
<i>Diaporthe ambigua</i>	<i>D. ambigua</i>	CBS 114015; STE-U 2657;	<i>Aspalathus linearis</i> , crown	<i>Fabaceae</i>	South Africa	J.C. Janse van Rensburg	KC343737 KC343979
<i>Diaporthe ambigua</i>	<i>D. ambigua</i>	CPC 2057 (ex-epitype)	<i>Foeniculum vulgare</i>	<i>Apiaceae</i>	Portugal	J.M. Santos	KC343254 KC343496 KC343738 KC343980
<i>Diaporthe ambigua</i>	<i>D. ambigua</i>	CBS 117167; STE-U 5414; CPC 5414	<i>Foeniculum vulgare</i>	<i>Apiaceae</i>	Portugal	J.M. Santos	KC343255 KC343497 KC343739 KC343981
<i>Diaporthe ambigua</i>	<i>D. ambigua</i>	CBS 123210; Di-C0029	<i>Platanus acerifolia</i>	<i>Platanaceae</i>	Italy	G. Granata	KC343256 KC343498 KC343740 KC343982
<i>Diaporthe ambigua</i>	<i>D. ambigua</i>	CBS 127746; IMI 398956	<i>Heilanthus annuus</i>	<i>Asteraceae</i>	Italy	A. Zazzarin	KC343257 KC343499 KC343741 KC343983
<i>Diaporthe ampelina</i> , comb. nov.	<i>D. ampelina</i>	CBS 137.87	<i>Vitis vinifera</i>	<i>Vitaceae</i>	USA: California	J.D. Cucuzzella	KC343258 KC343500 KC343742 KC343984
<i>Diaporthe ampelina</i> , comb. nov.	<i>P. viticola</i>	CPC 2073	<i>Vitis vinifera</i>	<i>Vitaceae</i>	France	P. Largnon	AF230751 AY745026 – AY745056 JX275452
<i>Diaporthe ampelina</i> , comb. nov.	<i>P. viticola</i>	PV F98-1 (ex-neotype)	<i>Vitis vinifera</i>	<i>Vitaceae</i>	Turkey	M. Erikan	KC343259 KC343501 KC343743 KC343985
<i>Diaporthe amygdali</i>	<i>P. viticola</i>	CBS 114867; STE-U 4708; CPC 4708	<i>Vitis vinifera</i>	<i>Vitaceae</i>	Italy	A. Zazzarin	KC343260 KC343502 KC343744 KC343986
<i>Diaporthe amygdali</i>	<i>P. amygdali</i>	CBS 267.80; STE-U 2671; CPC 2671	<i>Vitis vinifera</i>	<i>Vitaceae</i>	South Africa	L. Mostert	KC343261 KC343503 KC343745 KC343987
<i>Diaporthe amygdali</i>	<i>P. amygdali</i>	CBS 1181.1; STE-U 2632; CPC 2632	<i>Prunus persica</i> , cankers	<i>Rosaceae</i>	USA: Georgia	W. Uddin	KC343262 KC343504 KC343746 KC343988
<i>Diaporthe amygdali</i>	<i>P. amygdali</i>	CBS 15620; FAU 1005	<i>Prunus salicina</i> , wood	<i>Rosaceae</i>	South Africa	U. Damm	KC343263 KC343505 KC343747 KC343989
<i>Diaporthe amygdali</i>	<i>P. amygdali</i>	CBS 120840; STE-U 5833; CPC 5833	<i>Prunus dulcis</i>	<i>Rosaceae</i>	Portugal	E. Diogo	KC343264 KC343506 KC343748 KC343990
<i>Diaporthe amygdali</i>	<i>P. amygdali</i>	CBS 126679 (ex-epitype)	<i>Prunus dulcis</i>	<i>Rosaceae</i>	Portugal	E. Diogo	KC343265 KC343507 KC343749 KC343991
<i>Diaporthe anacardii</i>	<i>P. anacardii</i>	CBS 126680	<i>Anacardium occidentale</i>	<i>Anacardiaceae</i>	East Africa	M. Puccioni	KC343266 KC343508 KC343750 KC343992
<i>Diaporthe anacardii</i>	<i>P. foemiculi</i>	CBS 100871 (ex-epitype)	<i>Foeniculum vulgare</i> , dying twig	<i>Apiaceae</i>	Italy	L. Mugnai	KC343267 KC343509 KC343751 KC343993
<i>Diaporthe anacardii</i>	<i>P. foemiculi</i>	CBS 11591; AR 3724	<i>Heracleum sphondylium</i> , decaying stems	<i>Apiaceae</i>	Austria	A.Y. Rossman	KC343268 KC343510 KC343752 KC343994
<i>Diaporthe angelicae</i>	<i>D. angelicae</i>	CBS 11592; AR 3776 (ex-epitype)	<i>Heracleum sphondylium</i> , decaying stems	<i>Apiaceae</i>	Austria	A.Y. Rossman	KC343269 KC343511 KC343753 KC343995
<i>Diaporthe angelicae</i>	<i>D. angelicae</i>	CBS 123215; Ph-C133/1	<i>Foeniculum vulgare</i>	<i>Apiaceae</i>	Portugal	A.J.L. Phillips	KC343270 KC343512 KC343754 KC343996
<i>Diaporthe angelicae</i>	<i>D. angelicae</i>	CBS 344.86	<i>Eryngium marinum</i> , leaf spots	<i>Apiaceae</i>	France	H.A. van der Aa	KC343271 KC343513 KC343755 KC343997
<i>Diaporthe angelicae</i>	<i>D. angelicae</i>	CBS 501.90	<i>Heracleum sphondylium</i> , seeds	<i>Apiaceae</i>	France	H.A. van der Aa	KC343272 KC343514 KC343756 KC343998
<i>Diaporthe arctii</i>	<i>D. arctii</i>	CBS 136.25	<i>Arcium so.</i>	<i>Asteraceae</i>	–	A.W. Archer	KC343273 KC343515 KC343757 KC343999
<i>Diaporthe arecae</i> , comb. nov.	<i>D. arecae</i>	CBS 161.64 (ex-isotype)	<i>Arcea catenula</i> , fruit	<i>Arecaceae</i>	India	H.C. Srivastava	KC343274 KC343516 KC343758 KC343999
<i>Diaporthe arecae</i> , comb. nov.	<i>D. arecae</i>	CBS 161.64 (ex-isotype)	<i>Citrus sp.</i> , fruits	<i>Rutaceae</i>	I. Block	K.C. Hyde	KC343275 KC343517 KC343759 KC344001
<i>Diaporthe arengae</i> , sp. nov.	<i>P. pittospori</i>	CBS 535.75	<i>Arenga engleri</i>	<i>Arecaceae</i>	Hong Kong	J.C. Janse van Rensburg	KC343276 KC343518 KC343760 KC344002
<i>Diaporthe aspalathi</i>	<i>D. aspalathi</i>	CBS 14979; HKUCC 5527 (ex-type)	<i>Aspalathus linearis</i> , crown	<i>Fabaceae</i>	South Africa	J.C. Janse van Rensburg	KC343277 KC343519 KC343761 KC344003
<i>Diaporthe aspalathi</i>	<i>D. aspalathi</i>	CBS 177168; STE-U 5420; CPC 5420 (ex-type)	<i>Aspalathus linearis</i> , branch	<i>Fabaceae</i>	South Africa	K.C. Hyde	KC343278 KC343520 KC343762 KC344004
<i>Diaporthe australicana</i>	<i>D. australicana</i>	CBS 117500; STE-U 5408; CPC 5408 (ex-type)	<i>Aspalathus linearis</i>	<i>Fabaceae</i>	South Africa	S. Lamprecht	KC343037 KC343279 KC343521 KC344005
<i>Diaporthe australicana</i>	<i>D. australicana</i>	CBS 11886; STE-U 2676; CPC 2676 (ex-type)	<i>Vitis vinifera</i>	<i>Vitaceae</i>	Australia	R.W.A. Schepers	KC343038 KC343280 KC343522 KC343764
<i>Diaporthe batatas</i>	<i>D. batatas</i>	CBS 113497; STE-U 2655; CPC 2655	<i>Vitis vinifera</i>	<i>Vitaceae</i>	South Africa	L. Mostert	KC343281 KC343523 KC343765 KC344007
<i>Diaporthe beckhausii</i>	<i>D. beckhausii</i>	CBS 122.21	<i>Ipomoea batatas</i>	<i>Convolvulaceae</i>	USA	L.L. Harter	KC343282 KC343524 KC343766 KC344008
<i>Diaporthe brasiliensis</i> , sp. nov.	–	CBS 138.27	<i>Viburnum sp.</i>	<i>Caprifoliaceae</i>	–	L.E. Wehnrey	KC343283 KC343525 KC343767 KC344009
<i>Diaporthe brasiliensis</i> , sp. nov.	–	CBS 133.183; LGMF924; CPC 20300 (ex-type)	<i>Aspidospasma tomentosum</i> , endophytic in leaf	<i>Apocynaceae</i>	Brazil	K. Rodriguez	KC343284 KC343526 KC343768 KC344010
<i>Diaporthe brasiliensis</i> , sp. nov.	–	LGMF926; CPC 20302	<i>Aspidospasma tomentosum</i> , endophytic in leaf	<i>Apocynaceae</i>	Brazil	K. Rodriguez	KC343285 KC343527 KC343769 KC344011
<i>Diaporthe carpini</i>	<i>D. carpini</i>	CBS 114437; UPSC 2880	<i>Carpinus betulus</i>	<i>Corylaceae</i>	Sweden	K.C. Hyde	KC343286 KC343528 KC343770 KC344012
<i>Diaporthe carpini</i>	<i>D. carpini</i>	CBS 127298; Dpc1 (ex-neotype)	<i>Glycine max</i> , stem	<i>Fabaceae</i>	Croatia	K.C. Hyde	KC343287 KC343529 KC343771 KC344013
<i>Diaporthe carpini</i>	<i>D. carpini</i>	CBS 178.55; ATCC 12048; Afaria 243	<i>Glycine soja</i> , mature stem	<i>Fabaceae</i>	Canada	A.A. Hildebrand	KC343288 KC343530 KC343772 KC344014
<i>Diaporthe celastrina</i>	<i>D. celastrina</i>	CBS 139.27	<i>Celastrus scandens</i>	<i>Celastraceae</i>	–	L.E. Wehnrey	KC343289 KC343531 KC343773 KC344015
<i>Diaporthe chamaecyparis</i> , comb. nov.	<i>P. chamaecyparis</i>	CBS 454.81	<i>Chamaecyparis lawsoniana</i>	<i>Arecaceae</i>	Greece	H.A. van der Aa	KC343290 KC343532 KC343774 KC344016
<i>Diaporthe chamaecyparis</i> , comb. nov.	<i>P. chamaecyparis</i>	CBS 753.70	<i>Spartium junceum</i> , dead part of leaf	<i>Fabaceae</i>	Croatia	J.A. von Arx	KC343291 KC343533 KC343775 KC344017
<i>Diaporthe cinerascens</i>	<i>P. cinerascens</i>	CBS 719.96	<i>Ficus carica</i> , branch	<i>Moraceae</i>	Bulgaria	E. Ilieva	KC343292 KC343534 KC343776 KC344018
<i>Diaporthe citri</i>	<i>D. citri</i>	CBS 199.39	–	<i>Rutaceae</i>	Italy	G. Goldeinch	KC343293 KC343535 KC343777 KC344020
<i>Diaporthe citri</i>	<i>D. citri</i>	CBS 230.52	<i>Citrus sinensis</i> , decaying fruit	<i>Rutaceae</i>	Suriname	N.J. van Suchtelen	KC343294 KC343536 KC343778 KC344020

<i>Diaporthe convolvuli</i> , comb. nov.	Glycine max, seed	Fabaceae	Brazil	A. Almeida	KC344021
<i>P. convolvuli</i>	Convolvulus arvensis, leaves	Convolvulaceae	Turkey	D. Berner	KC344022
<i>D. crataegi</i>	Crataegus oxyacantha	Rosaceae	Sweden	K. & L. Holm	KC344023
<i>D. crotalariae</i>	Crotalaria spectabilis	Fabaceae	USA	G.F. Weber	KC344024
<i>P. cupataea</i>	<i>Aspalathus linearis</i>	Fabaceae	South Africa	J.C. Janse van Rensburg	KC344025
<i>Diaporthe cyanooids</i>	<i>Protea cynaroides</i> , leaf litter	Proteaceae	South Africa	S. Marincowitz	KC344026
<i>Diaporthe decedens</i>	<i>Corylus avellana</i>	Corylaceae	Austria	W. Jaklitsch	KC344027
<i>D. decedens</i>	<i>Corylus avellana</i>	Corylaceae	Sweden	K. & L. Holm	KC344028
<i>D. decessans</i>	<i>Berberis vulgaris</i>	Berberidaceae	Austria	A.Y. Rossman	KC344029
<i>D. decessans</i>	<i>Berberis vulgaris</i>	Berberidaceae	Sweden	K. & L. Holm	KC344030
<i>D. decessans</i>	<i>Berberis vulgaris</i>	Berberidaceae	–	L.E. Wehmeyer	KC344031
<i>P. elaeagni</i>	<i>Elaeagnus</i> sp., twig	Elaeagnaceae	Netherlands	J. Gremmen	KC344032
<i>Diaporthe endophytica</i> , sp. nov.	<i>Schinus terebinthifolius</i> , endophytic in leaf	Anacardiacae	Brazil	J. Lima	KC344033
<i>D. cyanoidis</i>	<i>Corylus avellana</i>	Anacardiacae	Brazil	J. Lima	KC344034
<i>D. cyanoidis</i>	<i>Corylus avellana</i>	Anacardiacae	Brazil	J. Lima	KC344035
<i>D. cyanoidis</i>	<i>Berberis vulgaris</i>	Berberidaceae	Brazil	R.R. Gomes	KC344036
<i>D. cyanoidis</i>	<i>Berberis vulgaris</i>	Berberidaceae	Brazil	R.R. Gomes	KC344037
<i>D. cyanoidis</i>	<i>Elaeagnus</i> sp., twig	Elaeagnaceae	Brazil	R.R. Gomes	KC344038
<i>D. cyanoidis</i>	<i>Schinus terebinthifolius</i> , endophytic in leaf	Anacardiacae	Brazil	R.R. Gomes	KC344039
<i>D. cyanoidis</i>	<i>Schinus terebinthifolius</i> , endophytic in leaf	Anacardiacae	Brazil	R.R. Gomes	KC344040
<i>D. cyanoidis</i>	<i>Maytenus ilicifolia</i> , endophytic in petiole	Celastraceae	Brazil	R.R. Gomes	KC344041
<i>D. cyanoidis</i>	<i>Maytenus ilicifolia</i> , endophytic in petiole	Celastraceae	Brazil	R.R. Gomes	KC344042
<i>D. cyanoidis</i>	<i>Maytenus ilicifolia</i> , endophytic in petiole	Celastraceae	Brazil	R.R. Gomes	KC344043
<i>D. cyanoidis</i>	<i>Maytenus ilicifolia</i> , endophytic in petiole	Celastraceae	Brazil	R.R. Gomes	KC344044
<i>D. cyanoidis</i>	<i>Maytenus ilicifolia</i> , endophytic in petiole	Celastraceae	Brazil	R.R. Gomes	KC344045
<i>D. cyanoidis</i>	<i>Glycine max</i> , seed	Fabaceae	Brazil	R.R. Gomes	KC344046
<i>D. cyanoidis</i>	<i>Fraxinus</i> sp., fallen fruit	Oleaceae	Netherlands	G.J.M. Verkley	KC344047
<i>D. cyanoidis</i>	<i>Juglans regia</i> , twig	Juglandaceae	Italy	M. Bisbach	KC344048
<i>D. cyanoidis</i>	<i>Acer campestre</i>	Aceraceae	Austria	W. Jaklitsch	KC344049
<i>D. cyanoidis</i>	<i>Arctium</i> sp., dead stems	Asteraceae	Netherlands	M. de Nooij	KC344050
<i>D. cyanoidis</i>	<i>Skimmia japonica</i> , dying twigs	Rutaceae	Netherlands	H.A. v. Kesteren	KC344051
<i>D. cyanoidis</i>	<i>Rhododendron</i> sp.	Ericaceae	Latvia	I. Apine	KC344052
<i>D. cyanoidis</i>	<i>Picea abies</i> , seedling	Pinaceae	UK	T.R. Peace	KC344053
<i>D. eres</i>	<i>CBS 101742</i>	Oleaceae	UK, Scotland	J.A. MacDonald	KC344054
<i>D. eres</i>	<i>CBS 102.81</i>	Oleaceae	Austria	W.G. Hutchinson	KC344055
<i>D. eres</i>	<i>CBS 109767</i> , AR 35388, WJ 1643	Oleaceae	Netherlands	I. de Boer	KC344056
<i>D. eres</i>	<i>CBS 110.85</i>	Oleaceae	Netherlands	H.A. van der Aa	KC344057
<i>D. eres</i>	<i>CBS 122.82</i>	Oleaceae	Netherlands	W.M. Loerakker	KC344058
<i>P. skimmiae</i>	<i>Phomopsis</i> sp. no. 23	CBS 129168	Netherlands	H.A. van der Aa	KC344059
<i>D. eres</i>	<i>CBS 186.37</i>	Oleaceae	Netherlands	H.A. van der Aa	KC344060
<i>D. eres</i>	<i>CBS 250.38</i>	Oleaceae	Netherlands	H.A. van der Aa	KC344061
<i>P. controversa</i>	<i>Fraxinus excelsior</i> , living and dead twigs	–	UK, Scotland	–	KC344062
<i>P. stictica</i>	<i>Laburnum × watereri</i> 'Vossii'	Fabaceae	Netherlands	W.G. Hutchinson	KC344063
<i>P. rufis</i>	<i>Allium giganteum</i> , dead stem	Alliaceae	Netherlands	I. de Boer	KC344064
<i>P. rangevicii</i>	<i>Sorbus aucuparia</i> , dead branch	Rosaceae	Netherlands	H.A. van der Aa	KC344065
<i>D. eres</i>	<i>CBS 287.74</i>	Oleaceae	Netherlands	W.M. Loerakker	KC344066
<i>P. osmanthi</i>	<i>Osmanthus aquifolium</i> , leaf tip	Cactaceae	Netherlands	H.A. van der Aa	KC344067
<i>P. cacti</i>	<i>Opuntia</i> sp., cladodes	Aquifoliaceae	Netherlands	H.A. van der Aa	KC344068
<i>P. crustosa</i>	<i>Ilex aquifolium</i> , dead leaf	Aquifoliaceae	Netherlands	H.A. van der Aa	KC344069
<i>D. perniciosa</i>	<i>Ilex aquifolium</i> , dead leaf	Rosaceae	–	Geily	KC344070
<i>P. phascoli</i>	<i>Malus sylvestris</i> , rotten fruit	Rosaceae	Netherlands	Goossens	KC344071
<i>P. cotoneasteri</i>	<i>Phaeoselus vulgaris</i>	Rosaceae	UK, Scotland	H. Buttin	KC344072
<i>P. cotoneasteri</i>	<i>Coloneaster</i> sp.	Rosaceae	Netherlands	G.H. Boerema	KC344073
<i>P. cruciferae</i>	(Isotype of <i>Phomopsis cotoneasteri</i>)	Brassicaceae	Netherlands	H.A. van der Aa	KC344074
<i>P. durandiana</i>	<i>Alliaria officinalis</i>	Polygonaceae	Netherlands	H.A. van der Aa	KC344075
<i>D. seposita</i>	<i>Rumex hydropathum</i> , dead stem	Fabaceae	Netherlands	H.A. van der Aa	KC344076
<i>P. abutilonis</i>	<i>Wistaria sinensis</i> , dead branch	Malvaceae	Netherlands	A. Apoot	KC344077
<i>P. crustosa</i>	<i>Abutilon</i> sp.	Aquifoliaceae	Netherlands	G.J.M. Verkley	KC344078
<i>P. magnolicaea</i>	<i>Ilex aquifolium</i> , twigs suffering from dieback	Magnoliaceae	Netherlands	H.A. van der Aa	KC344079
<i>P. trifolia</i>	<i>Magnolia × soulangeana</i> , withering leaf	Poaceae	Germany	M. Hostfeld	KC344080
<i>D. eugeniae</i>	<i>Hordeum</i> sp., leaf spot	Euphorbiaceae	West Sumatra	R. Kasim	KC344081
<i>D. fibrosa</i>	<i>Eugenia aromatica</i> , leaf	Rhamnaceae	Austria	A.Y. Rossman	KC344082
<i>D. foenulaeca</i>	<i>Rhamnus cathartica</i>	Rhamnaceae	Sweden	K. & L. Holm	KC344083
<i>D. foenulaeca</i>	<i>Fructicum vulgare</i> , base of senescent stem	Apiaceae	Spain	A.J.L. Phillips	KC344084
<i>D. foenulaeca</i>	<i>Fructicum vulgare</i> , base of senescent stem	Apiaceae	Portugal	A.J.L. Phillips	KC344085
<i>P. theicola</i>	<i>Pyrus pyrifolia</i>	Rosaceae	New Zealand	W. Kandula	KC344086
<i>D. neothecicola</i>	<i>Foenicum vulgare</i>	Apocynaceae	Portugal	A.J.L. Phillips	KC344087
<i>D. neothecicola</i>	<i>Foenicum vulgare</i>	Apocynaceae	Portugal	A.J.L. Phillips	KC344088
<i>P. malif sp. amygdali</i>	<i>Prunus amygdalus</i> , dried fruit	Rosaceae	Italy	A. Ciccarone	KC344089

P. malif sp. amygdali CBS 171.78
P. malif sp. amygdali CBS 123208; Di-C004/5 (ex-type of *D. neothecicola*)
P. malif sp. amygdali CBS 123209; Di-C004/4 (ex-type of *D. neothecicola*)

Species	Original name	Strain ¹	Isolation source	Host family	Locality	Collector	GenBank Accession numbers (ITS, CAL, HIS, TEF1, TUB) ²
<i>P. theicola</i>	CBS 187.27 (ex-type of <i>P. theicola</i>)		<i>Theaceae</i>	Italy	M. Curzi	KC343349	KC343591
<i>P. diospyri</i>	CBS 287.56		<i>Ebenaceae</i>	M. Ribaldi	KC343108	KC343592	KC344075
<i>D. seposita</i>	CBS 357.69		<i>Fabaceae</i>	Netherlands	H.A. van der Aa	KC343109	KC343534
<i>P. casuarinae</i>	CBS 400.48		–	India	S.R. Bose	KC343110	KC343593
<i>P. bougainvillae</i>	CBS 603.88		<i>Nyctaginaceae</i>	Portugal	H.A. van der Aa	KC343111	KC343536
<i>Diaporthella gariae</i> , comb. nov.	CBS 180.91; ILLS 43621 (ex-type)		<i>Cannabaceae</i>	USA: Illinois	J.M. McPartland	KC343354	KC343597
<i>Diaporthella gardeniae</i> comb. nov.	CBS 288.56		<i>Rubiaceae</i>	Italy	M. Ribaldi	KC343113	KC343598
<i>D. helianthi</i>	CBS 344.94		<i>Asteraceae</i>	–	–	KC343114	KC343598
<i>D. helianthi</i>	CBS 592.81 (ex-type)		<i>Asteraceae</i>	Serbia	M. Muntañola-Cvetkovic	KC343115	KC343599
<i>D. heveae</i>	CBS 852.97		<i>Euphorbiaceae</i>	Brazil	D.S. Attli	KC343116	KC343599
<i>P. heveae</i>	CBS 681.84		<i>Euphorbiaceae</i>	India	K. Jayaraman	KC343117	KC343558
<i>D. heveae</i>	CBS 145.26 (ex-type)		<i>Juglandaceae</i>	USA: Michigan	L.E. Wehmeyer	KC343118	KC343559
<i>D. hickoriae</i>	CBS 15448; HKUCC 9104;		<i>Hydrangeaceae</i>	Hong Kong	K.D. Hyde	KC343119	KC343560
<i>D. hickoriae</i>	AT 646 DF 24 (ex-type)		<i>Poaceae</i>	Norway	L. Sundheim	KC343120	KC343564
<i>D. hordei</i>	CBS 481.92		<i>Rosaceae</i>	Sweden	K. & L. Holm	KC343121	KC343565
<i>D. impulsa</i>	CBS 1144.34; UPSC 3052		–	–	L.E. Wehmeyer	KC343122	KC343566
<i>D. impulsa</i>	CBS 141.27		<i>Celastraceae</i>	Brazil	R.R. Gomes	KC343123	KC343567
<i>D. impulsa</i>	CBS 13381.27; LGMF930;		<i>Anacardiaceae</i>	Brazil	K. Rodriguez	KC343124	KC343568
<i>D. impulsa</i>	CPC 20306 (ex-type)		<i>Anacardiaceae</i>	Brazil	R.R. Gomes	KC343125	KC343569
<i>D. impulsa</i>	LGMF92; CPC 20288		<i>Celastraceae</i>	Brazil	J. Lima	KC343126	KC343568
<i>D. impulsa</i>	LGMF931; CPC 20307		<i>Anacardiaceae</i>	Brazil	K. Rodriguez	KC343124	KC343568
<i>D. impulsa</i>	CBS 13381.27; LGMF906;		<i>Schinus terebinthifolius</i> , endophytic in leaf	Brazil	R.R. Gomes	KC343125	KC343569
<i>D. impulsa</i>	CPC 20282 (ex-type)		<i>Schinus terebinthifolius</i> , endophytic in leaf	Brazil	J. Lima	KC343126	KC343569
<i>D. impulsa</i>	LGMF908; CPC 20284		<i>Anacardiaceae</i>	Brazil	K. Rodriguez	KC343124	KC343568
<i>D. impulsa</i>	LGMF912; CPC 20288		<i>Anacardiaceae</i>	Brazil	R.R. Gomes	KC343125	KC343569
<i>D. impulsa</i>	LGMF917; CPC 20283		<i>Anacardiaceae</i>	Brazil	J. Lima	KC343126	KC343569
<i>D. impulsa</i>	LGMF918; CPC 20294		<i>Anacardiaceae</i>	Brazil	J. Lima	KC343127	KC343569
<i>D. impulsa</i>	LGMF920; CPC 20296		<i>Anacardiaceae</i>	Brazil	J. Lima	KC343128	KC343569
<i>D. impulsa</i>	LGMF932; CPC 20309		<i>Anacardiaceae</i>	Brazil	K. Rodriguez	KC343129	KC343569
<i>D. juglandina</i>	LGMF940; CPC 20316		<i>Celastraceae</i>	Brazil	R.R. Gomes	KC343130	KC343569
<i>D. juglandina</i>	CBS 121004; DP 0659		<i>Juglandaceae</i>	USA: Tennessee	L. Vasileya	KC343131	KC343569
<i>D. juglandina</i>	CBS 194.36 (ex-type)		<i>Grossulariaceae</i>	Canada	L.E. Wehmeyer	KC343132	KC343569
<i>D. longispora</i>	CBS 123212; DiC00115 (ex-type)		<i>Apiaceae</i>	Portugal	J.M. Santos	KC343136	KC343578
<i>D. lusitanicae</i>	CBS 123213; DiC00113		<i>Apiaceae</i>	Portugal	J.M. Santos	KC343137	KC343580
<i>D. lusitanicae</i>	CBS 505.76		<i>Euphorbiaceae</i>	Rwanda	J. Senai	KC343138	KC343582
<i>D. lusitanicae</i>	CBS 133.185; LGMF938;		<i>Celastraceae</i>	Brazil	R.R. Gomes	KC343139	KC343581
<i>D. lusitanicae</i>	CPC 20314 (ex-type)		<i>Caprifoliaceae</i>	–	L.E. Wehmeyer	KC343140	KC343582
<i>D. lusitanicae</i>	CBS 435.87		<i>Fabaceae</i>	Indonesia	H. Vermeulen	KC343141	KC343583
<i>D. melonis</i>	CBS 507.78 (ex-isotype)		<i>Cucurbitaceae</i>	USA: Texas	L. Beraha & M.J. O'Brien	KC343142	KC343584
<i>D. melonis</i>	CBS 1295.19; CPC 17026 (ex-type)		<i>Musaceae</i>	Australia	P.W. Crous & R.G. Shivas	KC343143	KC343585
<i>D. musigena</i>	CBS 144.27		<i>Rosaceae</i>	–	L.E. Wehmeyer	KC343144	KC343586
<i>D. neiliae</i>	CBS 109490; GB 6421; AR 3450 (ex-type)		<i>Asteraceae</i>	USA:	G. Bills	KC343145	KC343587
<i>D. arctii</i>			<i>Fagaceae</i>	Korea	K.A. Seifert	KC343146	KC343588
<i>D. megalospora</i>	CBS 13470; DAOM 226800		<i>Rosaceae</i>	New Zealand	W. Kandula & L. Castlebury	KC343147	KC343589
<i>D. megalospora</i>	CBS 116953; NZ-26		<i>Castanea sativa</i> , chestnuts collected in grocery store			KC343148	KC343590
<i>D. megalospora</i>			<i>Pyrus pyrifolia</i>			KC343149	KC343591
<i>D. melonis</i>	CBS 16954; NZ-27		<i>Pyrus pyrifolia</i>	New Zealand	G.J. Samuels	KC343150	KC343592
<i>P. fukushii</i>	CBS 124030; GJS 77-49		<i>Malus pumila</i> , bark	New Zealand	I. Apine	KC343151	KC343593
<i>D. pennicosa</i>	CBS 129167		<i>Rhododendron</i> sp.	Latvia	Koithoff	KC343152	KC343594
<i>Phomopsis</i> sp. no. 22	CBS 200.39		<i>Laurus nobilis</i> , stem	Germany	M. Muntañola-Cvetkovic	KC343153	KC343595
<i>D. nobilis</i>	CBS 388.89		<i>Hedera helix</i>	Yugoslavia		KC343154	KC343596

<i>P. conorum</i>	CBS 587.79	<i>Pinus perniphylla</i> , bonsai imported from Japan into Netherlands	<i>Pinaceae</i>	Japan	G.H. Boerema	KC343153	KC343395	KC343637	KC34379	KC344121
<i>Diaporthe nomurai</i>	CBS 157.29	<i>Morus</i> sp.	<i>Moraceae</i>	Japan	K. Togashi	KC343154	KC343396	KC343638	KC343880	KC344122
<i>Diaporthe novem</i>	CBS 127289; 5-273-1	<i>Glycine</i> max, seed	<i>Fabaceae</i>	Croatia	T. Duvnjak	KC343155	KC343397	KC343639	KC343881	KC344123
<i>D. novem</i>	CBS 127270; 4-273-1 (ex-type)	<i>Glycine</i> max, seed	<i>Fabaceae</i>	Croatia	T. Duvnjak	KC343156	KC343398	KC343640	KC343882	KC344124
<i>D. novem</i>	CBS 354.71	<i>Polygonatum odoratum</i> , leaves	<i>Convallariaceae</i>	Romania	O. Constantinescu	KC343157	KC343399	KC343641	KC343883	KC344125
<i>D. pardalote</i> sp.	LGMF943; CPC 20319	<i>Maytenus ilicifolia</i> , endophytic in petiole	<i>Celastraceae</i>	Brazil	R.R. Gomes	KC343158	KC343400	KC343642	KC343884	KC344126
<i>D. oncostoma</i>	CBS 100454	<i>Robinia pseudoacacia</i> , leaf spot	<i>Fabaceae</i>	Germany	H. Buttin	KC343159	KC343401	KC343643	KC343885	KC344127
<i>D. oncostoma</i>	CBS 109741; AR 3445	<i>Robinia pseudoacacia</i>	<i>Fabaceae</i>	Russia	L. Vasilyeva	KC343160	KC343402	KC343644	KC343886	KC344128
<i>D. oncostoma</i>	CBS 588.78	<i>Robinia pseudoacacia</i> , dead branches	<i>Aquifoliaceae</i>	France	H.A. van der Aa	KC343161	KC343403	KC343645	KC343887	KC344129
<i>P. crustosa</i>	CBS 809.85	<i>Ilex aquifolium</i> , leaf	<i>Celastraceae</i>	Germany	M. Hesse	KC343162	KC343404	KC343646	KC343888	KC344130
<i>Diaporthe oxe</i> , sp. nov.	CBS 133186; LGMF942; CPC 20318 (ex-type)	<i>Maytenus ilicifolia</i> , endophytic in petiole	<i>Celastraceae</i>	Brazil	R.R. Gomes	KC343163	KC343405	KC343647	KC343889	KC344131
<i>D. decorticans</i>	CBS 114200; UPSC 2569	<i>Schinus terebinthifolius</i> , endophytic in leaf	<i>Anacardiacae</i>	Brazil	J. Lima	KC343164	KC343406	KC343648	KC343890	KC344132
<i>D. valisformis</i>	CBS 114649; LGMF945; CPC 20305 (ex-type)	<i>Maytenus ilicifolia</i> , endophytic in petiole	<i>Celastraceae</i>	Brazil	S.A.V. Pileggi	KC343167	KC343409	KC343649	KC343891	KC344133
<i>D. valisformis</i>	CBS 133184; LGMF92; CPC 109745; ARSEF 3461; AR 3461 (ex-epitype)	<i>Ulmus glabra</i>	<i>Ulmaceae</i>	Austria	R.Y. Rossman	KC343168	KC343410	KC343650	KC343892	KC344134
<i>Diaporthe paranensis</i> , sp. nov.	CBS 151.73	<i>Persea gratissima</i> , young fruit	<i>Lauraceae</i>	Netherlands	E. Laville	KC343169	KC343413	KC343651	KC343893	KC344135
<i>Diaporthe perfuncta</i>	<i>D. perfuncta</i>	<i>Olearia cf. rani</i>	<i>Asteraceae</i>	New Zealand	G.J.M. Verkley	KC343170	KC343416	KC343658	KC343900	KC344142
<i>Diaporthe perseae</i> comb. nov.	<i>P. perseae</i>	<i>Caperonia palustris</i>	<i>Euphorbiaceae</i>	USA: Mississippi	A. Mengistu	KC343171	KC343417	KC343659	KC343901	KC344143
<i>Diaporthe phaseolorum</i>	<i>P. oleariae</i>	<i>Aster exilis</i>	<i>Asteraceae</i>	USA: Mississippi	A. Mengistu	KC343172	KC343418	KC343660	KC343902	KC344144
<i>Diaporthe phaseolorum</i>	<i>D. phasedolorum</i>	<i>Actinidia chinensis</i> , rotting fruit	<i>Actinidiaceae</i>	New Zealand	S.R. Pennycook	KC343173	KC343419	KC343661	KC343903	KC344145
<i>Diaporthe phaseolorum</i>	<i>D. melonis</i>	—	—	—	L. Beraha	KC343178	KC343420	KC343662	KC343904	KC344146
<i>Diaporthe phaseolorum</i>	<i>D. melonis</i>	<i>Maytenus ilicifolia</i> , endophytic in petiole	<i>Celastraceae</i>	Brazil	R.R. Gomes	KC343179	KC343421	KC343663	KC343905	KC344147
<i>Diaporthe phaseolorum</i>	<i>D. mangiferae</i>	<i>Maytenus ilicifolia</i> , endophytic in petiole	<i>Anacardiacae</i>	Brazil	R.R. Gomes	KC343180	KC343422	KC343664	KC343906	KC344148
<i>Diaporthe phaseolorum</i>	<i>P. mangiferae</i>	<i>Mangifera indica</i>	<i>Anacardiacae</i>	Dominican Republic	P. de Leeuw	KC343181	KC343423	KC343665	KC343907	KC344149
<i>Diaporthe pseudophoenicicola</i> , sp. nov.	<i>P. pseudophoenicicola</i>	<i>Mangifera indica</i> , peel of fruit	<i>Anacardiacae</i>	Mexico	—	KC343182	KC343424	KC343666	KC343908	KC344150
<i>Diaporthe pseudophoenicicola</i> , sp. nov.	<i>P. pseudophoenicicola</i>	<i>Mangifera indica</i> , showing dieback	<i>Anacardiacae</i>	Iraq	M.S.A. Al-Momen	KC343183	KC343425	KC343667	KC343909	KC344151
<i>Diaporthe pseudophoenicicola</i> , sp. nov.	<i>P. pseudophoenicicola</i>	<i>Phoenix dactylifera</i> , dead tops of green leaves	<i>Arecaceae</i>	Spain	H.A. van der Aa	KC343184	KC343426	KC343668	KC343910	KC344152
<i>Diaporthe pustulata</i>	<i>D. pustulata</i>	<i>Acer pseudoplatanus</i>	<i>Aceraceae</i>	Austria	A.Y. Rossman	KC343185	KC343427	KC343669	KC343911	KC344153
<i>Diaporthe pustulata</i>	<i>D. pustulata</i>	<i>Prunus padus</i>	<i>Aceraceae</i>	Austria	A.Y. Rossman	KC343186	KC343428	KC343670	KC343912	KC344154
<i>Diaporthe pustulata</i>	<i>D. padi</i>	<i>Spondias mombin</i> , endophytic in leaf	<i>Rosaceae</i>	Brazil	K.C. Rodriguez	KC343187	KC343429	KC343671	KC343913	KC344155
<i>Diaporthe raonikayaporum</i> , sp. nov. —	<i>D. rhoina</i>	<i>Rhus toxicodendron</i>	<i>Anacardiacae</i>	—	L.E. Wehmeyer	KC343188	KC343431	KC343673	KC343915	KC344157
<i>Diaporthe raonikayaporum</i> , sp. nov. —	<i>P. mangiferae</i>	<i>Protea repens</i> , cankers	<i>Proteaceae</i>	South Africa	S. Denman	KC343189	KC343432	KC343674	KC343916	KC344158
<i>Diaporthe saccata</i> , comb. nov.	<i>P. saccata</i>	<i>Schinus terebinthifolius</i> , endophytic in leaf	<i>Anacardiacae</i>	Brazil	J. Lima	KC343190	KC343433	KC343675	KC343917	KC344159
<i>Diaporthe schimi</i> , sp. nov.	—	<i>Cucumis sativus</i> , root	<i>Cucurbitaceae</i>	Netherlands	H.A. van der Kesteren	KC343192	KC343434	KC343676	KC343918	KC344160
<i>Diaporthe sclerotoides</i>	<i>P. sclerotoides</i>	<i>Cucumis sativus</i> , root	<i>Cucurbitaceae</i>	Netherlands	—	KC343193	KC343435	KC343677	KC343919	KC344161
<i>Diaporthe sclerotoides</i>	<i>P. sclerotoides</i>	<i>Fraxinus excelsior</i> , living and dead twig	<i>Oleaceae</i>	UK: Scotland	J.A. MacDonald	KC343194	KC343436	KC343678	KC343920	KC344162
<i>Diaporthe sclerotoides</i>	<i>P. sclerotoides</i>	<i>Glycine soja</i>	<i>Euphorbiaceae</i>	Italy	P. Giunchi	KC343195	KC343437	KC343679	KC343921	KC344163
<i>Diaporthe sclerotoides</i>	<i>P. sclerotoides</i>	<i>Euphorbia nutans</i>	<i>Euphorbiaceae</i>	USA: Mississippi	A. Mengistu	KC343196	KC343438	KC343680	KC343922	KC344164
<i>Diaporthe sclerotoides</i>	<i>P. sclerotoides</i>	<i>Glycine max</i>	<i>Fabaceae</i>	Croatia	K. Vrandecic	KC343197	KC343439	KC343681	KC343923	KC344165
<i>Diaporthe sclerotoides</i>	<i>P. sclerotoides</i>	<i>Glycine max</i> , stem	<i>Fabaceae</i>	—	A.A. Hildebrand	KC343198	KC343440	KC343682	KC343924	KC344166
<i>Diaporthe sclerotoides</i>	<i>P. sclerotoides</i>	<i>Glycine soja</i> , mature stem	<i>Fabaceae</i>	USA	J. Marchinkowska	KC343199	KC343441	KC343683	KC343925	KC344167
<i>Diaporthe sclerotoides</i>	<i>P. sclerotoides</i>	<i>Glycine soja</i> , seedling	<i>Fabaceae</i>	sojæ	KC343200	KC343443	KC343685	KC343927	KC344169	KC344169

Species	Original name	Strain	Isolation source	Host family	Locality	Collector	GenBank Accession numbers (ITS, CAL, HIS, TEF1, TUB) ²
<i>Diaporthe</i> sp. 1	<i>D. phaseolorum</i>	CBS 119639; B-11861	Man. abscess	–	Germany	K. Piechulla	KC343928
	<i>Diaporthe</i> sp.	LGMF#47; CPC 20323	Glycine max, seed	Fabaceae	Brazil	A. Almeida	KC343929
	<i>Diaporthe</i> sp.	LGMF#32; CPC 20308	Maytenus ilicifolia, endophytic in petiole	Celastraceae	Brazil	R.R. Gomes	KC343930
	<i>P. conorum</i>	CBS 287.29	Pseudotisiga menziesii	Pinaceae	UK: Scotland	G.G. Hahn	KC343931
	<i>Diaporthe</i> sp.	LGMF#44; CPC 20320	Maytenus ilicifolia, endophytic in petiole	Celastraceae	Brazil	R.R. Gomes	KC343932
	<i>Acer opalus</i>	CBS 125575	Acer opalus	Aceraceae	Italy	W. Jaklitsch	KC343933
	<i>P. pittospori</i>	CBS 115584; HKUCC 7784; AT 7	Maesa perifolia, fruit	Myrsinaceae	Hong Kong	K.D. Hyde	KC343934
	<i>P. pittospori</i>	CBS 115595; HKUCC 10129	Maesa perifolia, fruit	Myrsinaceae	Hong Kong	K.D. Hyde	KC343935
	<i>P. anacardii</i>	CBS 488.78	Anacardium occidentale	Anacardaceae	India	H.C. Govindu	KC343936
	<i>Diaporthe</i> sp.	LGMF#25; CPC 20301	Aspidosperma tomentosum	Apocynaceae	Brazil	K. Rodriguez	KC343937
	<i>P. stictica</i>	CBS 370.54	Buxus sempervirens, dead twig	Buxaceae	Italy	M. Ribaldi	KC343938
	<i>Diaporthe</i> subordinaria	CBS 101711	Plantago lanceolata, blackened seed	Plantaginaceae	New Zealand	B. Alexander	KC343939
	<i>P. subordinaria</i>	CBS 484.90	Plantago lanceolata, stalk	Plantaginaceae	South Africa	R. Shivas	KC343940
	<i>P. tecomaie</i>	CBS 100547	Tabeaia sp., mycoecidium caused by Prosopodium tecomicola	Bignoniaceae	Brazil	A. Aptroot	KC343941
	<i>Diaporthe terebinthifoli</i> , sp. nov.	–	Schinus terebinthifolius, endophytic in leaf	Anacardiaceae	Brazil	J. Lima	KC343942
		CBS 133180; LGMF#14;	Schinus terebinthifolius, endophytic in leaf	Anacardiaceae	Brazil	J. Lima	KC343943
		CPC 20290 (ex-type)	Schinus terebinthifolius, endophytic in leaf	Anacardiaceae	Brazil	J. Lima	KC344185
		LGMF#07; CPC 20283	Schinus terebinthifolius, endophytic in leaf	Anacardiaceae	Brazil	J. Lima	KC344186
		LGMF#09; CPC 20285	Schinus terebinthifolius, endophytic in leaf	Anacardiaceae	Brazil	J. Lima	KC344187
		LGMF#13; CPC 20289	Schinus terebinthifolius, stem	Fabaceae	Western Australia J.B. Nunn	KC343946	KC344188
	<i>D. toxica</i>	CBS 534.93; ATCC 96741 (ex-type)	Lupinus angustifolius, stem	Fabaceae	Western Australia P.M. Williamson	KC343704	KC344189
	<i>D. toxica</i>	CBS 535.93	Lupinus sp.	Fabaceae	Western Australia P.M. Williamson	KC343705	KC344190
	<i>D. toxica</i>	CBS 546.93	Lupinus sp., stem	Ericaceae	USA: Michigan G.C. Adams	KC343948	KC344191
	<i>D. vaccinii</i>	CBS 118571; G.C.A. Dyack	Vaccinium corymbosum	Ericaceae	USA: New Jersey L. Carris	KC343949	KC344192
	<i>P. vaccinii</i>	CBS 122112; FAU 474	Vaccinium macrocarpon	Ericaceae	USA: Michigan D.C. Ramsdell	KC343950	KC344193
	<i>P. vaccinii</i>	CBS 122114; FAU 634	Vaccinium corymbosum	Ericaceae	USA: Michigan D.C. Ramsdell	KC343951	KC344194
	<i>P. vaccinii</i>	CBS 122115; FAU 580	Vaccinium corymbosum	Ericaceae	USA: North D.F. Farr	KC343952	KC344195
	<i>P. vaccinii</i>	CBS 122116; DF 5022	Vaccinium corymbosum	Ericaceae	Carolina	C.L. Shear	KC343953
	<i>D. vaccinii</i>	CBS 160.32; IFO 32646 (ex-type)	Oxycoccus macrocarpos	Ericaceae	USA: Massachusetts	C.L. Shear	KC343954
	<i>P. vexans</i>	CBS 127.14	Solanum melongena	Solanaceae	USA	L.L. Harter	KC343713
	<i>Diaporthe vitticola</i>	CBS 100170	Fraxinus excelsior, leaf spot	Oleaceae	Netherlands	H.A. van der Aa	KC343714
	<i>D. acutiae</i>	CBS 106.95	Aucuba japonica, branches and twigs	Aucubaceae	Netherlands	G.J.M. Verkley	KC343715
	<i>D. medusaea</i>	CBS 109492	Labium anagyroides	Fabaceae	Austria	A.Y. Rossman	KC343716
	<i>D. pardoiota</i>	CBS 109768; AR 3478	Ephedrum angustifolium	Onagraceae	Canada	M. Barr	KC343717
	<i>D. vitticola</i>	CBS 113201; STE-U 5683;	Vitis vinifera	Vitaceae	Portugal	A.J.L. Phillips	KC343718
	<i>D. vitticola</i>	CPC 5883 (ex-epitype)	Vitis vinifera	Vitaceae	Sweden	A.J.L. Phillips	KC343719
	<i>D. circumscripta</i>	CBS 114011; CPC 2677	Sambucus cf. racemosa	Caprifoliaceae	Portugal	KC343720	KC343955
	<i>D. medusaea</i>	CBS 114436; UPSC 2860	Rosa rugosa	Rosaceae	Netherlands	K. & L. Holm	KC343956
	<i>D. woodii</i>	CBS 266.85; PD 85/25	Lupinus arboreus, dead stem	Fabaceae	Netherlands	G.H. Boerema	KC343957
	<i>P. salicina</i>	CBS 446.62	Salix sp., twig	Salicaceae	Netherlands	H.A. van der Aa	KC343958
	<i>D. woodii</i>	CBS 449.82	Lupinus sp., dead stem	Fabaceae	Netherlands	H.A. van der Aa	KC343959
	<i>P. dipsaci</i>	CBS 502.85	Dipsacus fullonum, dead stem	Dipsacaceae	France	G.J.M. Verkley	KC343960
	<i>P. asphodelina</i>	CBS 759.95	Asphodelus albus, 1-yr-old stems	Asphodelaceae	UK	G.J.M. Verkley	KC343961
	<i>D. acutiae</i>	CBS 794.96	Aucuba japonica	Aucubaceae	Western Australia	P.M. Williamson	KC343962
	<i>D. woodii</i>	CBS 558.93	Lupinus sp., stem	Fabaceae	–	L.E. Wehmeyer	KC343963
	<i>D. woolworthii</i>	CBS 148.27	Ulmus americana	Ulmaceae	China: Fujian	L.N. Vasileva	KC343964
	<i>Diaporthe corylina</i>	–	Corylus sp., dying stems	Corylaceae	China: Fujian	L.N. Vasileva	KC343965
	<i>Diaporthe woolworthii</i>	CBS 12124; AR 4131	Convolvulus sp., dying stems	Convolvulaceae	Spain	KC343966	KC343967

¹ AR: Collection of A.Y. Rossman; ATCC: American type culture collection; CBS: CBS Fungal Biodiversity Centre, Utrecht, The Netherlands; CECT: Colección Española de Cultivos Típicos, Valencia, Spain; CPC: Collection Pedro Crous, housed at CBS; DAOM: Plant Research Institute, Department of Agriculture (Mycology), Ottawa, Canada; HKUCC: University of Hong Kong Culture Collection, Department of Ecology and Biodiversity, Hong Kong, China; IFCC: Institute for Fermentation, Osaka, Japan; IMI: International Mycological Institute, CABI-Bioscience, Egham, Berkshire, UK; LGMF: Culture collection of Laboratory of Genetics of Microorganisms, Federal University of Paraná, Curitiba, Brazil; MUCL: Université Catholique de Louvain, Louvain-la-Neuve, Belgium; NRRL: National Center for Agricultural Utilization Research, Peoria, Illinois, USA.; PD: Plant Protection Service, Wageningen, The Netherlands; SITE-U: Stellenbosch University culture collections, South Africa; UPSC: Fungal Culture Collection at the Botanical Museum, Uppsala University, Uppsala, Sweden

² TUB: partial beta-tubulin gene; CAL: partial calmodulin gene; HIS: partial histone H3 gene; ITS: internal transcribed spacer regions of the nrDNA and intervening 5.8S rRNA; TEF1: partial translation elongation factor 1-alpha gene.

(1933) to the conclusion that host-association was not informative enough in *Diaporthe*, thereby reducing the number of species from 650 to only 70 in the genus. However, this revision was based strictly on morphological characters of the *Diaporthe* sexual state, and connections to the *Phomopsis* asexual states (prior to molecular analyses) had been identified only in 20 % of the species (Wehmeyer 1933).

Although the classification of *Diaporthe* has been on-going, species are presently being redefined based on a combination of morphological, cultural, phytopathological, mating type and DNA sequence data (Rehner & Uecker 1994, Zhang et al. 1998, Mostert et al. 2001a, Farr et al. 2002, Santos et al. 2010).

However, even when using a combination of morphological and molecular data, the delimitation of species within the genus *Diaporthe* only proved satisfactory once multi-gene DNA sequence data were generated (Castlebury & Mengistu 2006, van Rensburg et al. 2006, Santos et al. 2010, Udayanga et al. 2012), since this adds valuable information in the resolution of complex evolutionary relationships. The aims of the present study were thus to: 1) provide a multi-gene phylogeny for the genus *Diaporthe* based on a large set of well-identified cultures deposited in the CBS culture collection; 2) to identify potential isolates for epitypification, thereby fixing the application of previously established names; 3) to link *Diaporthe* names to their *Phomopsis* asexual states; and 4) to identify a collection of mostly sterile endophytic *Diaporthe* strains isolated from several medicinal hosts in Brazil.

MATERIALS AND METHODS

Isolates

In the present study we analysed 243 *Diaporthe* isolates (Table 1), as well as the outgroup *Diaporthella corylina*. Isolates were obtained from several sources, including 40 endophytic strains isolated from medicinal plants in Brazil (LabGeM/UFPR collection, Curitiba, Brazil), and three isolates from the EMBRAPA-SOJA collection, Londrina, Brazil. A further 199 isolates were obtained from the CBS-KNAW Fungal Biodiversity Centre (CBS), or the working collection of P.W. Crous (CPC) housed at CBS.

DNA isolation, amplification and phylogenetic analysis

Colonies were cultivated on 2 % potato-dextrose agar (PDA), and genomic DNA extraction was undertaken using the Ultra-Clean™ Microbial DNA Kit (MO Bio, Carlsbad, CA, USA) according to manufacturer's instructions. Using 20 isolates, we screened nine loci, of which the five more informative loci were selected for multi-gene analyses.

The primers ITS5 and ITS4 (White et al. 1990) were used to amplify the internal transcribed spacer region (ITS) of the nuclear ribosomal RNA gene operon, including the 3' end of the 18S nrRNA, the first internal transcribed spacer region, the 5.8S nrRNA gene; the second internal transcribed spacer region and the 5' end of the 28S nrRNA gene. The primers EF1-728F and EF1-986R (Carbone & Kohn 1999) were used to amplify part of the translation elongation factor 1- α gene (TEF1) and the primers ACT-512F and ACT-783R (Carbone & Kohn 1999) were used to amplify part of the actin gene (ACT). The primers Gpd1-LM and Gpd2-LM (Mylllys et al. 2002) were used to amplify part of the glyceraldehyde-3-phosphate dehydrogenase (GPDH) gene, and part of the calmodulin (CAL) gene was sequenced using the primers CAL-228F and CAL-737R (Carbone & Kohn 1999). The primers CYLH3F (Crous et al. 2004b) and H3-1b (Glass & Donaldson 1995) were used to amplify part of the histone H3 (HIS) gene, and the primers T1 (O'Donnell & Cigelnik 1997) and Bt-2b (Glass & Donaldson 1995) to amplify

part of the β -tubulin gene (TUB). The primers NMS1 and NMS2 (Li et al. 1994) were used to amplify an internal region of the mitochondrial SSU (mtSSU). The partial large subunit nrDNA (LSU) was sequenced using the primers LSU1Fd (Crous et al. 2009a) and LR5 (Vilgalys & Hester 1990).

Amplification reactions had a total reaction volume of 12.5 μ L which was composed of 1× PCR buffer (Bioline GmbH, Luckenwalde, Germany), 5.6 % DMSO (v/v), 20 μ M dNTPs, 0.2 μ M of each forward and reverse primers, 0.25 U of BioTaq Taq DNA polymerase (Bioline GmbH, Luckenwalde, Germany), and 10 ng of genomic DNA. PCR conditions were the same for all loci, except for the MgCl₂ concentration: 2 mM MgCl₂ for the genes LSU and TEF1, 1.5 mM MgCl₂ for the genes ACT, GPDH, mtSSU, ITS and TUB, and 1 mM MgCl₂ for CAL and HIS genes. The PCR conditions were: start step of 2 min at 94 °C, followed by 40 cycles of 30 s at 94 °C, 1 min at adequate annealing temperature, and 1 min at 72 °C, followed by a finishing step of 3 min at 72 °C and a cool down step to 4 °C. The annealing temperature varied for each gene: 61 °C (ACT, GPDH, mtSSU); 58 °C (CAL, ITS, HIS); 55 °C (TEF1, TUB) and 48 °C (LSU).

However, some of these primer pairs failed to amplify with some isolates included in this study, and therefore additional combinations were used. The amplification reaction and cycle conditions were the same except the annealing temperature and MgCl₂ concentration. For the amplification of TEF1 with primers EF1-728F and EF2 (O'Donnell et al. 1998), 52 °C and 2 mM MgCl₂; TUB with primers T1 (O'Donnell & Cigelnik 1997) and CYLTUB1R (Crous et al. 2004b), 50 °C and 1 mM MgCl₂; CAL with primers CAL-228F and CAL2Rd (Quaedvlieg et al. 2011, Groenewald et al. 2013), 58 °C and 1 mM MgCl₂.

Amplicons were sequenced using both PCR primers with a BigDye Terminator Cycle Sequencing Kit v. 3.1 (Applied Biosystems, Foster City, CA, USA) according to the manufacturer's instructions, and sequences were analysed on an ABI Prism 3700 DNA Sequencer (Perkin-Elmer, Norwalk, Foster City, CA, USA). The consensus sequences were visually inspected using MEGA v. 5 software (Tamura et al. 2011). The alignment of obtained sequences was performed using the online MAFFT interface (Katoh & Toh 2008; <http://mafft.cbrc.jp/alignment/server>).

For the phylogenetic analyses based on Maximum Likelihood and Bayesian inference, we chose the best evolutionary models for each data partition using the software MrModelTest v. 2.3 (Nylander 2004). MrBayes v. 3.1.1 (Ronquist & Huelsenbeck 2003) was used to generate the phylogenetic trees under optimal criteria per data partition. The heating parameter was set at 0.3 and the Markov Chain Monte Carlo (MCMC) analysis of four chains was started in parallel from a random tree topology and lasted until the average standard deviation of split frequencies came below 0.01. Trees were saved each 10 000 generations and the resulting phylogenetic tree (Fig. 1) was printed with Geneious v. 5.5.4 (Drummond et al. 2011) and the layout of the tree was done in Adobe Illustrator v. CS5.1. *Diaporthella corylina* (CBS 121124) was used as outgroup in the phylogenetic analyses based on its position as sister family in *Diaporthales* (Vasilyeva et al. 2007). New sequences generated in this study were deposited in NCBI's GenBank nucleotide database (www.ncbi.nlm.nih.gov; Table 1) and the alignment and phylogenetic tree in TreeBASE (study S13943; www.treebase.org).

Locus resolution and SNP detection

Neighbour-joining analyses using the general time-reversible substitution model were applied to each data partition individually to check the stability and robustness of each species clade under each dataset using PAUP v. 4.0b10 (Swofford 2003) (TreeBASE study S13943). Alignment gaps were treated as missing data and all characters were unordered and of equal





Fig. 1 Consensus phylogram of 22 104 trees resulting from a Bayesian analysis of the combined 5-gene sequence alignment. Clades are numbered on the right of the boxes and *Diaporthe* species names in purple reflect new combinations and in red new species. Strain accession numbers are followed by the original species name (black, when applicable), the isolation source (green) and country of origin (blue). Accession numbers and names in bold represent strains known to be ex-type strains or are considered to be authentic for the species. Red dots indicate strains from medicinal plants and yellow dots from humans. Bayesian posterior probabilities are shown at the nodes and the scale bar represents the expected changes per site. The tree was rooted to *Diaporthe corylina* (strain CBS 121124).

weight. Any ties were broken randomly when encountered. The robustness of the trees obtained was evaluated by 1 000 bootstrap replications (Hillis & Bull 1993). In the present study, both the analysis of the combined alignment (Fig. 1) and of the individual loci were used to determine the species boundaries. For each clade in the combined analysis, the position of the members of that clade was determined in the phylogenetic tree obtained from each of the individual loci to check whether these members still represent a single clade in the individual gene tree. In this way the robustness of a given clade could

be evaluated together with the posterior probability value of that clade. A species was only counted if it was distinct from its closest relatives and the species clade contained all the associated strains.

Unique fixed nucleotide positions are used to characterise and describe several sterile species (see applicable species notes). For each sterile species that was described, the closest phylogenetic neighbour(s) were selected from Fig. 1 and this focused dataset was subjected to SNP analyses. These single nucleotide polymorphisms (SNPs) were determined for

Fig. 1 (cont.)




each aligned data partition using DnaSP v. 5.00.07 (Librado & Rozas 2009).

Taxonomy

All descriptions provided are based on colonies sporulating in culture, which for the most part only formed the asexual morph. Colonies were subcultured onto 2 % tap water agar supplemented with sterile pine needles (PNA; Smith et al. 1996), or

autoclaved leaf pieces of *Ilex aquifolium*, *Maytenus ilicifolia* or *Schinus terebinthifolius*, PDA, oatmeal agar (OA), and 2 % malt extract agar (MEA) (according to Crous et al. 2009b), and incubated at 20 °C under a 12 h near-ultraviolet light (400–315 nm) (Sylvania Blacklight-Blue, Osram Nederland B.V., Alphen aan den Rijn, The Netherlands), 12 h dark cycle to promote sporulation. Structures were mounted in clear lactic acid, and 50 measurements determined for conidia, and 30 for other

Fig. 1 (cont.)




structures. The 95 % confidence levels were determined, and the extremes given in parentheses. Colony diameters were determined at 25 °C in darkness on PDA, OA and MEA. Colony colours (surface and reverse) were described after 14 d using the colour charts of Rayner (1970). Nomenclatural novelties and descriptions were deposited in MycoBank (www.MycoBank.org; Crous et al. 2004a).

RESULTS

DNA sequencing and phylogenetic analysis

The most suitable genes for *Diaporthe* species delimitation in this study were found to be CAL, HIS, ITS, TEF1 and TUB. The amplified genomic regions of these genes were more informative, and the combined analysis provided a more robust species identification, from which phylogenetic relationships could be inferred.

Fig. 1 (cont.)




0.2

The manually adjusted, combined (ITS, TUB, CAL, TEF1 and HIS) alignment for the Bayesian analysis contained 243 isolates (including the outgroup sequences) and 2 435 characters were used in the phylogenetic analysis. The number of unique site patterns per data partition were 210, 616, 242, 281 and 183, respectively and were based on 466, 874, 355, 316 and 424 alignment positions, respectively. Based on the results of MrModeltest, the following priors were set in MrBayes for the different data partitions: all partitions had dirichlet base frequen-

cies and GTR+I+G models with inverse gamma-distributed rates were implemented for ITS and HIS, and HKY+I+G with inverse gamma-distributed rates for TUB, CAL and TEF1. The Bayesian analysis lasted 14 735 000 generations and the consensus trees and posterior probabilities were calculated from the 22 104 trees left after discarding 7 368 trees (the first 25 % of generations) for burn-in (Fig. 1). Ninety-five clades are recognised and discussed here.

Fig. 1 (cont.)



0.2

Locus resolution and SNP detection

The mtSSU and LSU regions had very few informative sites for the tested strains and were therefore not selected as good markers at species level. The ACT and GPDH regions were also discarded as suitable candidates for the multi-gene analyses because of their long branch lengths which made unambigu-

ous alignments impossible. These four loci were therefore not used for further amplification and sequencing on the complete dataset. The remaining five loci had varied success for species identification and some phylogenetic lineages were more prone to less variability than others. Fifty-eight of the 95 spe-

cies could be identified by all five loci. The loci are treated individually below:

CAL – The locus could distinguish 74 of the 95 species (78 % success). It had difficulty separating: *D. endophytica* and *D. phaseolorum* (clades 4, 5); *D. angelicae*, *D. arctii* and *D. subordinaria* (clades 17–19); *D. alleghaniensis*, *D. alnea*, *D. celastrina*, *D. eres*, *D. juglandina*, *D. neilliae* and *D. nobilis* (clades 60–62, 64–67); and *D. eugeniae*, *D. musigena*, *D. perseae*, *D. pseudomangiferae*, *D. pseudophoenicicola*, *Diaporthe* sp. 6 and *Diaporthe* sp. 7 (clades 82–86, 88, 89). A single strain each of *D. angelicae* (clade 17), *D. novem* (clade 22) and *D. terebinthifolii* (clade 9) clustered separate from the other strains of the species.

HIS – The locus could distinguish 84 of the 95 species (88 % success). It had difficulty separating: *D. australafricana* and *D. viticola* (clades 49, 50); *D. celastrina*, *D. eres* and *D. nobilis* (clades 66, 67, 62); *D. arecae* and *D. perseae* (clades 86, 87); and *D. pseudophoenicicola* and *Diaporthe* sp. 8 (clades 89, 90). A single strain each of *D. endophytica* (clade 5) and *D. terebinthifolii* (clade 9) clustered separate from the other strains of the species. This is the only locus that can distinguish *D. angelicae* (clade 17).

ITS – The locus could distinguish 75 of the 95 species (79 % success). It had difficulty separating: *D. angelicae*, *D. arctii* and *D. subordinaria* (clades 17–19); *D. cynaroides* and *D. viticola* (clades 48, 50); *D. alnea*, *D. neilliae* and *D. nobilis* (clades 60–62); *D. arengae*, *D. eugeniae* and *D. pseudomangiferae* (clades 81–83); *D. arecae* and *D. perseae* (clades 86, 87); and *D. aspalathi* and *D. woodii* (clades 93, 94). A single strain each of *D. arecae* (clade 87), *D. inconspicua* (clade 75), *D. novem* (clade 22) and *D. terebinthifolii* (clade 9), and two strains each of *D. impulsa* (clade 51) and *D. infecunda* (clade 23), clustered separate from the other strains of the species. This is the only locus that can distinguish *D. celastrina* (clade 17) and *D. eres* (clade 67).

TEF1 – The locus could distinguish 72 of the 95 species (76 % success). It had difficulty separating: *D. tecomaee*, *D. terebinthifolii* (clades 9, 10); *D. angelicae*, *D. arctii* and *D. subordinaria* (clades 17–19); *D. australafricana* and *D. viticola* (clades 49, 50); *D. celastrina* and *D. juglandina* (clades 65, 66); *D. eres* and *D. nobilis* (clades 67, 62); *D. chamaeropis*, *D. cinerascens* and *D. foeniculaceae* (clades 76–78); and *D. arengae*, *D. arecae*, *D. eugeniae*, *D. musigena*, *D. perseae*, *D. pseudomangiferae*, *D. pseudophoenicicola*, *Diaporthe* sp. 6 and *Diaporthe* sp. 8 (clades 81–87, 89, 90).

TUB – The locus could distinguish 84 of the 95 species (88 % success). It had difficulty separating: *D. endophytica* and *D. phaseolorum* (clades 4, 5); *D. alleghaniensis*, *D. celastrina*, *D. eres*, *D. juglandina*, *D. nobilis* and *D. vaccinii* (clades 62–67); and *D. aspalathi* and *D. woodii* (clades 93, 94). A single strain of *D. angelicae* (clade 17) clustered separate from the other strains of the species. This is the only locus that can distinguish *D. perseae* (clade 86).

Descriptions based on DNA characters are provided for three species in the Taxonomy section, namely *D. endophytica* (clade 5), *D. inconspicua* (clade 75) and *D. infecunda* (clade 23). *Diaporthe endophytica* (clade 5) was compared to *D. phaseolorum* (clade 4); *D. inconspicua* (clade 75) to *D. anacardii* (clade 69), *D. chamaeropis* (clade 77), *D. cinerascens* (clade 76), *D. elaeagni* (clade 73), *D. foeniculacea* (clade 78), *D. hickoriae* (clade 72), *D. oncostoma* (clade 70), *D. saccarata* (clade 71) and *D. stictica* (clade 74); and *D. infecunda* (clade 23) to *D. angelicae* (clade 17), *D. arctii* (clade 19), *D. cuppatea* (clade 20), *D. lusitanicae* (clade 21), *D. neoarctii* (clade 16), *D. novem* (clade 22) and *D. subordinaria* (clade 18).

Taxonomy

The multigene analyses resulted in 95 well-supported clades correlating to 243 isolates of *Diaporthe* (Table 1, Fig. 1). Fifteen new species are described, nine of which were isolated from medicinal plants (*Aspidosperma tomentosum*, *Maytenus ilicifolia*, *Schinus terebinthifolius*, *Spondias mombin*) in Brazil (clades 5, 9, 11, 23, 30, 31, 34, 35 and 36). Twenty-eight clades contain ex-type strains of presently known species, or strains accepted as authentic for the species name or which could be designated as epitypes in the present study, and were therefore well-resolved (7, 8, 12, 14, 17, 20–22, 24, 26–28, 40, 42, 43, 45, 48–50, 63, 64, 69, 71, 72, 84 and 91–93). The sexual-aseexual relationship was resolved for several taxa, and is reported below. New combinations in *Diaporthe* are introduced below for several *Phomopsis* names that represented well-resolved taxa. Several potential epitypes were identified during this study, which are discussed below.

Diaporthe acaciigena Crous, Pascoe & Jacq. Edwards, Persoonia 26: 123. 2011

Specimen examined. AUSTRALIA, Victoria, Otway Ranges, Anglesea, S38°23'21.7" E144°11'12.7", on leaves of *Acacia retinodes*, 16 Oct. 2009, P.W. Crous, I.G. Pascoe & J. Edwards (holotype CBS H-20581, ex-type culture CPC 17622 = CBS 129521).

Notes — Clade 43 contains the ex-type culture of *D. acaciigena* isolated from *Acacia retinodes* in Australia. This species is morphologically similar to *D. amygdali* (clade 42) (Crous et al. 2011), and closely related to *D. pustulata* (clade 44).

Diaporthe acerina (Peck) Sacc., Syll. Fung. (Abellini) 1: 611. 1882

Basionym. *Valsa acerina* Peck, Ann. Rep. N.Y. State Mus. Nat. Hist. 28: 73. 1876. 1874.

Specimen examined. UNKNOWN, from *Acer saccharum*, Sept. 1927, L.E. Wehmeyer (CBS 137.27).

Notes — Clade 39 is represented by *D. acerina*, isolated from *Acer saccharum*. This species is genetically similar to *D. perjuncta* (clade 40). It is known to occur in Europe and North America on dead limbs and trunks of *Acer pseudoplatanus*, *A. saccharinum*, *A. saccharum*, *A. spicatum*, and *Acer* sp. (Aceraceae) (Spielman 1985, Farr et al. 1989).

Diaporthe alleghaniensis R.H. Arnold, Canad. J. Bot. 45: 787. 1967

Specimen examined. CANADA, Ontario, on branches of *Betula alleghaniensis*, June 1972, R.H. Arnold (ex-type culture CBS 495.72 = ATCC 24097 = DAOM 45776).

Notes — Clade 64 contains the ex-type strain of *D. alleghaniensis*, isolated from *Betula alleghaniensis* in Canada. *Diaporthe alleghaniensis* causes canker and dieback of *B. alleghaniensis*, *B. lenta*, *B. papyrifera* and *B. pendula* in Canada (Arnold 1975), but has also been reported from Japan (Farr & Rossman 2012).

Diaporthe alnea Fuckel, Jahrb. Nassauischen Vereins Naturk. 23–24: 207. 1870 (1869–1870)

= *Phomopsis alnea* Höhn., Sber. Akad. Wiss. Wien, Math.-naturw. Kl., Abt. 1 115: 681 (33 of repr.). 1906.

Specimens examined. UNKNOWN, on *Alnus* sp., June 1946, S. Truter (CBS 146.46); on *Alnus* sp., Aug. 1947, S. Truter (CBS 159.47).

Notes — Clade 61 consists of two isolates from *Alnus* (presumably collected in the Netherlands). *Diaporthe alnea* causes dieback of *Alnus glutinosa* (alder) and *A. incana* (grey alder).

It has been reported from Europe, Russia and the USA (Munk 1957, Oak & Dorset 1983, Moricca 2002, Mel'nik et al. 2008, Farr & Rossman 2012).

Diaporthe ambigua Nitschke, Pyrenomycetes Germanici 2: 311. 1870

Specimens examined. ITALY, Sicily, Catania, on *Platanus acerifolia*, G. Granata (CBS 127746 = IMI 395956); Perugia, on *Helianthus annuus*, Mar. 1987, A. Zazzerini (CBS 187.87). – PORTUGAL, Vale Andeiro, on *Foeniculum vulgare*, J.M. Santos (CBS 123210 = Di-C003/10, CBS 123211 = Di-C002/9). – SOUTH AFRICA, Western Cape Province, from *Pyrus communis*, deposited 2002, S. Denman (ex-epitype culture CBS 114015 = CPC 2657); Western Cape Province, on crown of *Aspalathus linearis*, 15 May 1997, J.C. Janse van Rensburg (CBS 117167 = CPC 5414).

Notes — Clade 26 represents *D. ambigua*, which contains two isolates previously misidentified as *D. scabra* (CBS 127746) and *D. helianthi* (CBS 187.87), and four isolates of *D. ambigua*, including the ex-epitype culture. *Diaporthe ambigua* is an important pathogen of *Malus domestica*, *Prunus salicina* and *Pyrus communis* in South African fruit orchards. Infection by *D. ambigua* is associated with sunken lesions with longitudinal cracks on affected fruit trees. The fungus rapidly kills nursery rootstocks, but also kills mature rootstocks over a longer period of time (Smit et al. 1996). This species is also found as saprobe on wild fennel (Santos & Phillips 2009). It has been reported on *Aspalathus linearis* (van Rensburg et al. 2006), *Foeniculum vulgare*, *Malus domestica* (Smit et al. 1996, Santos & Phillips 2009), *Malus sylvestris* (Crous et al. 2000), *Prunus* spp. (Smit et al. 1996, Mostert et al. 2001a), *Pyrus communis* (Nitschke 1867), *Pyrus ussuriensis* (Tai 1979) and *Vitis vinifera* (van Niekerk et al. 2005). It is widely distributed, and is known from China, Cuba (Tai 1979), Germany (Nitschke 1867), South Africa (Smit et al. 1996), UK (Dennis 1986) and the USA (Washington) (Shaw 1973).

Diaporthe ampelina (Berk. & M.A. Curtis) R.R. Gomes, C. Glenke & Crous, comb. nov. — MycoBank MB802922

Basionym. *Phoma ampelina* Berk. & M.A. Curtis, Grevillea 2, 18: 81. 1873.
 ≡ *Phomopsis ampelina* (Berk. & M.A. Curt.) Grove, Bull. Misc. Inform. Kew 4: 184. 1919.
 = *Phoma viticola* Sacc., Michelia 2: 92. 1880.
 ≡ *Phomopsis viticola* (Sacc.) Sacc., Ann. Mycol. 13: 118. 1915.
 = *Fusicoccum viticolum* Reddick, Cornell Univ. Agric. Exp. Sta. Bull. 263: 331. 1909.
 ≡ *Phomopsis viticola* (Reddick) Goid., Atti Reale Accad. Naz. Lincei 26: 107. 1937.
 = *Phomopsis viticola* Sacc. var. *ampelopsis* Grove, Bull. Misc. Inform. Kew 4: 183. 1919.
 = *Diaporthe neoviticola* Udayanga, Crous & K.D. Hyde, Fung. Diversity 56: 166. 2012 (a nom. nov. based on *Phoma viticola* Sacc.).

Conidiomata pycnidial, eustromatic, subepidermal, brown to black, scattered or aggregated, globose, flask-like to conical, outer surface smooth, convoluted to unilocular, singly ostiolate, up to 430 µm wide and 190–300 µm tall, including short necks which rarely occur. Pycnidial wall consisting out of two regions of *textura angularis*; the outer region brown, 2–3 cells thick, 5–7 µm wide, inner region brown, 3–4 cells thick, 7–15 µm wide, with the outside cells compressed. *Conidial mass* globose or in cirrhi, white, pale-yellow to yellow, but predominantly pale-yellow. *Alpha conidiophores* cylindrical, some filiform, rarely septate and branched, 5–35 × 1–3 µm (av. = 25 × 2 µm). *Alpha conidiogenous cells* subcylindrical, tapering towards the apex, collarettes and periclinal thickening present, 3–19 × 1–2 µm (av. = 10 × 1.5 µm). *Alpha conidia* commonly found, fusoid-ellipsoidal, apex acutely rounded, base obtuse to subtruncate, multi-guttulate with guttules grouped at the polar ends, rarely biguttulate, (7–)9.5–10.5(–13) × (1.5–)2–3(–3.5) µm (av. = 10 × 2.5 µm). *Beta conidiophores* ampulliform to subcylindrical, rarely branched, 10–34 × 1–2 µm (av. = 26 × 1.5 µm). *Beta conidiogenous cells* subcylindrical, tapering towards the apex, collarette and periclinal thickening present, 7–14 × 1–2 µm (av. = 11–1.5 µm). *Beta conidia* less common than alpha conidia, straight, curved or hamate, 20–25 × 0.5–1 µm (av. = 23–1 µm). *Gamma conidia* rarely observed, fusoid to subcylindrical, apex acutely rounded, base subtruncate, multi-guttulate, 12–18 × 1.5–2 µm (av. = 15 × 2 µm). Description adapted from Mostert et al. (2001a).

Specimens examined. FRANCE, Bordeaux, Naujan-et-Postiac, on *Vitis vinifera* (Cabernet Sauvignon grapevine), May 1998, P. Larignon (PREM 56460 neotype, ex-neotype culture CBS 114016). – ITALY, Perugia, on *Vitis vinifera*, May 1980, A. Zazzerini (CBS 267.80 = CPC 2671). – TURKEY, from *Vitis vinifera*, 1 Dec. 2001, M. Erkan (CBS 114867 = CPC 4708). – USA, California, on *Vitis vinifera*, J.D. Cucuzza (CBS 111888 = ATCC 48153 = CPC 2673).

Notes — Grove (1919) distinguished *P. ampelina* (K 58408) from *P. viticola* by its external appearance on the host. However, Mostert et al. (2001a) re-examined the type specimen, and found alpha conidia to be ellipsoid-fusoid, 8–12 × 2.5–3.5 µm, within the range of *P. viticola* (Mostert et al. 2001a: f. 29), and thus considered them to be synonymous. Udayanga et al. (2012) proposed *D. neoviticola* as a nom. nov. for *P. viticola*, but this name is superfluous, as the older epithet 'ampelina' has precedence and should be adopted.

Diaporthe ampelina (clade 53) is a well-resolved species. It causes cane and leaf spot and infections of pruning wounds of *Vitis* and *Ampelopsis* spp. (Vitaceae). Several species of *Diaporthe* can infect the host and cause variable symptoms in different parts of the vine (canes, leaves and fruits) causing considerable confusion in the taxonomy of these species on grapevine (Phillips 1999, Scheper et al. 2000, Mostert et al.



Fig. 2 *Diaporthe anacardii* (CBS 720.97). a. Conidiomata sporulating on PNA; b. conidiomata sporulating on PDA; c, d. conidiogenous cells; e. beta conidia; f. alpha conidia. — Scale bars = 10 µm.

2001a). Merrin et al. (1995) studied the variation of *Diaporthe* in Australia using morphology. They identified two taxa (*Phomopsis* taxon 1 and taxon 2), which cause cane and leaf blight of *Vitis* spp.; and taxon 2 was identified as showing more resemblance to *P. viticola*. Mostert et al. (2001a) studied the species occurring on grapevines in South Africa using morphological, cultural, molecular and pathological characterisation and clarified the taxonomy of this complex. *Diaporthe ampelina* (= *Phomopsis viticola*, *D. neoviticola*, *Phomopsis* taxon 2 from Australia) was found to be the cause of cane and leaf spot disease, and was neotyped. Although the sexual morph has never been reported, Santos et al. (2010) found both MAT loci to be present in this species, and showed that it is heterothallic. However, the sexual morph could not be induced in culture by crossing opposing mating types.

***Diaporthe amygdali* (Delacr.) Udayanga, Crous & K.D. Hyde, Fung. Diversity 56: 166. 2012**

Basionym. *Fusicoccum amygdali* Delacr., Bull. Soc. Mycol. France 21: 280. 1905.

= *Phomopsis amygdali* (Delacr.) J.J. Tuset & M.T. Portilla, Canad. J. Bot. 67, 5: 1280. 1989.

Specimens examined. PORTUGAL, Mirandela, from *Prunus dulcis*, 2010, E. Diogo (ex-epitype culture CBS 126679); Tavira, on *Prunus dulcis*, 2010, E. Diogo (CBS 126680). – SOUTH AFRICA, Western Cape Province, on *Vitis vinifera*, 1 Mar. 1997, L. Mostert (CBS 111811 = CPC 2632); Western Cape Province, in wood on *Prunus salicina*, 2008, U. Damm (CBS 120840 = CPC 5833). – USA, Georgia, cankers on *Prunus persica*, Mar. 1994, W. Uddin (CBS 115620 = FAU 1005).

Notes — *Diaporthe amygdali* (clade 42) is the causal agent of twig canker and blight of almonds (*Prunus dulcis*) and peach (*P. persica*) wherever these hosts are grown (Diogo et al. 2010). It was first described as *Fusicoccum amygdali* causing cankers on almonds in France (Delacroix 1905). Tuset & Portilla (1989) re-examined the type specimen of *F. amygdali* and, based on morphology and symptomatology, they considered that it would be best accommodated in the genus *Phomopsis*. Clade 42 contains the ex-epitype strain (CBS 126679), five *Phomopsis amygdali* isolates from *Prunus dulcis* in Portugal, from *P. persica* in USA, *P. salicina* in South Africa, and from *Vitis vinifera* in South Africa.

***Diaporthe anacardii* (Early & Punith.) R.R. Gomes, C. Glienke & Crous, comb. nov. — MycoBank MB802923; Fig. 2**

Basionym. *Phomopsis anacardii* Early & Punith., Trans. Brit. Mycol. Soc. 59, 2: 345. 1972.

Conidiomata pycnidial, sporulating profusely on OA, globose, up to 600 µm diam, multilocular, black, erumpent; cream conidial droplets exuding from central ostioles; walls consisting of 3–6 layers of medium brown *textura angularis*. *Conidiophores* hyaline, smooth, 1–3-septate, branched, densely aggregated, cylindrical, straight to sinuous, 10–25 × 2–3 µm. *Conidiogenous cells* 9–16 × 1.5–2 µm, phialidic, cylindrical to cymbiform, terminal and lateral, with slight taper towards apex, 1–1.5 µm diam, with visible periclinal thickening; collarette slightly flared, up to 2 µm long when present. *Paraphyses* rarely present, hyaline, smooth, 1–3-septate, cylindrical with obtuse ends, extending above conidiophores. *Alpha conidia* aseptate, hyaline, smooth, guttulate, fusoid to ellipsoid, tapering towards both ends, straight, apex subobtuse, base bluntly rounded with flattened hilum, (6.5–)7–8(–9) × (2–)3(–3.5) µm. *Gamma conidia* not observed. *Beta conidia* spindle-shaped, aseptate, smooth, hyaline, apex subacutely rounded, base truncate, tapering from lower third towards apex, curved, (15–)20–25 × 1.5(–2) µm.

Culture characteristics — Colonies covering dish after 2 wk in the dark at 25 °C. On OA dirty white with moderate aerial mycelium and patches of iron-grey. On PDA having patches of dirty white and umber, reverse bay with patches of umber. On MEA having patches of dirty white and olivaceous-grey, reverse umber with patches of olivaceous-grey.

Specimens examined. EAST AFRICA, on *Anacardium occidentale*, Apr. 1997, M. Puccioni (epitype designated here CBS H-21101, culture ex-epitype CBS 720.97). – KENYA, on *Anacardium occidentale*, 4 Dec. 1969, M.P. Early (holotype IMI 144866).

Notes — *Phomopsis anacardii* (clade 69 as *D. anacardii*) was described from *Anacardi occidentalis* in Kenya, and also recorded from Nigeria, Guinea and Cuba (Early & Punithalingam 1972).




Fig. 3 *Diaporthe angelicae* (CBS 111591). a, b. Transverse section through conidiomata, showing conidiomatal wall; c, d. conidiogenous cells; e. alpha and beta conidia; f. conidiogenous cells giving rise to beta conidia; g. beta conidia. — Scale bars: a = 140 µm, all others = 10 µm.

Diaporthe angelicae (Berk.) D.F. Farr & Castl., Mycoscience 44: 204. 2003. — Fig. 3

Basionym. *Sphaeria angelicae* Berk., Mag. Zool. Bot.: 28. 1837.
 ≡ *Diaporthopsis angelicae* (Berk.) Wehm., The genus *Diaporthe* Nitschke: 228. 1933.
 ≡ *Mazzantia angelicae* (Berk.) Lar. N. Vassiljeva, Pyrenomycetes of the Russia Far East. I. Gnomoniaceae: 49. 1993.
 = *Leptosphaeria nigrella* (Auersw.), Mycol. Eur. Pyr. 5/6, pl. 12, f. 163. 1869.
 ≡ *Diaporthe nigrella* (Auersw.) Niessl, Beitr.: 51. 1872.
 ≡ *Diaporthopsis nigrella* (Auersw.) Fabre, Ann. Sci. Nat., Bot. 6 15: 35. 1883.

Conidiomata pycnidial, globose to ellipsoidal, aggregated or scattered, dark brown to black, immersed, ostiolate, 100–281 µm wide, 70–200 µm tall, lacking necks, with outer surface covered in hyphae; pycnidial wall consisting of brown, thick-walled cells of *textura angularis*; conidial mass globose or exuding in cirri, white to pale luteous or pale yellow. *Conidiophores* hyaline, subcylindrical, rarely branched, tapering towards the apex, aseptate, (12–)13–16(–18) × 3(–4) µm. *Conidiogenous cells* hyaline, subcylindrical, straight to curved, tapering towards the apex, collarette not flared, periclinal thickening inconspicuous, 8–10(–11) × 3(–3.5) µm. *Alpha conidia* hyaline, oblong to ellipsoid, apex bluntly rounded, base obtuse to subtruncate, bi- to multi-guttulate, (7–)8–10(–11) × 3(–4) µm. *Beta conidia* hyaline, smooth, spindle shaped, slightly curved, (19–)22–26(–28) × (1–)2 µm. *Gamma conidia* not observed (CBS 111592).

Culture characteristics — See Castlebury et al. (2003).

Specimens examined. AUSTRIA, Karnten, St. Margareten, decaying stems of *Heracleum sphondylium*, Aug. 2001, A.Y. Rossman (CBS 111591 = AR 3724); Niederösterreich, Ottenstein, decaying stems of *Heracleum sphondylium*, Aug. 2001, A.Y. Rossman (ex-epitype culture CBS 111592 = AR3776). — FRANCE, Bretagne, La Ville Borée, near Quessoy, on seeds of *Heracleum sphondylium*, 27 July 1990, H.A. van der Aa (CBS 501.90); sea dunes near Seignose le Penon, on *Eryngium maritimum*, leaf spots, 10 June 1986, H.A. van der Aa (CBS 344.86). — ITALY, San Casciano, Prov., Florence, twig blight of *Foeniculum vulgare*, July 1996, L. Mugnai (CBS 100871). — PORTUGAL, Malveira da Serra, Sintra, on *Foeniculum vulgare*, A.J.L. Phillips (CBS 123215 = Ph-C133/1).

Notes — *Diaporthe angelicae* (clade 17) is known to cause stem decay in several hosts including *Heracleum sphondylium* (Apiaceae) and *Foeniculum vulgare* (Apiaceae) in Europe and North America (Santos & Phillips 2009). Wehmeyer (1933) not only linked the conidial form of *Phomopsis asteriscus* to the sexual state *Diaporthopsis angelicae*, but also stated that *Diaporthe berkeleyi* was a synonym of *Diaporthopsis angelicae*. However, Castlebury et al. (2003) showed that *Diaporthopsis* is a synonym of *Diaporthe*, and also designated an epitype for *D. angelicae*.

Diaporthe arctii (Lasch) Nitschke, Pyrenomyces Germanici 2: 268. 1870

Basionym. *Sphaeria arctii* Lasch, in Rabenh., Klotzsch. Herb. Vivum Mycol.: no. 1046. 1846.
 ≡ *Phomopsis arctii* (Lasch) Traverso, Fl. Ital. Crypt., Pars 1: Fungi. Pyrenomycetidae. Xylariaceae, Valsaceae, Ceratostomataceae: 226. 1906.

Specimen examined. UNKNOWN, from *Arctium* sp., Sept. 1925, A.W. Archer (CBS 136.25).

Notes — There are several clades that contain isolates previously identified as *D. arctii* (clades 16, 19, part 2 and 67, part 4). We suspect that clade 19 may represent the real *D. arctii*, as it is basal to *D. subordinaria*, and Wehmeyer (1933) regarded the latter (from *Plantago lanceolata*) as synonym of *D. arctii* (from *Arctium*).

Diaporthe arecae (H.C. Srivast., Zakia & Govindar.) R.R.

Gomes, C. Glienke & Crous, comb. nov. — MycoBank MB802924

Basionym. *Subramanella arecae* H.C. Srivast., Zakia & Govindar., Mycologia 54, 1: 7. 1962.

Specimens examined. INDIA, on fruit of *Areca catechu*, Feb. 1964, H.C. Srivastava (isotype CBS H-7808, ex-isotype culture CBS 161.64). — SURINAME, on fruits of *Citrus* sp., Oct. 1975, I. Block (CBS 535.75).

Notes — The *Diaporthe* isolate from citrus (CBS 535.75) could well be distinct, but more strains are required to resolve this clade (clade 87).

Diaporthe arengae R.R. Gomes, C. Glienke & Crous, sp. nov.

— MycoBank MB802925; Fig. 4

Etymology. Named after the host genus from which it was collected, *Arenga*.

Pycnidia in culture on PNA sporulating poorly, subglobose, up to 250 µm diam, black, erumpent; cream conidial droplets exuding from central ostiole; walls consisting of 3–6 layers of medium brown *textura angularis*. *Conidiophores* hyaline in upper region, pale brown at base, smooth, 0–6-septate, branched, densely aggregated, cylindrical, straight to sinuous, 10–60 × 2.5–4 µm. *Conidiogenous cells* 8–15 × 1.5–2.5 µm, phialidic, cylindrical, terminal and lateral, with slight taper towards apex, 1–1.5 µm diam, with visible periclinal thickening; collarette not flared, up to 2 µm long when present. *Paraphyses* not observed. *Alpha conidia* aseptate, hyaline, guttulate, fusoid-ellipsoid, tapering towards both ends, apex subobtuse, base with flattened hilum, (5–)6–7(–9) × (2–)2.5(–3) µm. *Gamma conidia* not observed. *Beta conidia* rarely observed, subcylindrical, aseptate, smooth, hyaline, apex bluntly rounded, base truncate, tapering absent to very slight, curved, 20–25 × 1.5 µm.



Fig. 4 *Diaporthe arengae* (CBS 114979). a. Conidiomata sporulating on PNA; b, c. conidiogenous cells; d. beta conidia; e, f. alpha conidia. — Scale bars = 10 µm.

Culture characteristics — Colonies covering dish after 2 wk in the dark at 25 °C. On MEA surface with fluffy aerial mycelium, pale luteous, in reverse orange with patches of sienna. On OA umber with patches of sienna and saffron, in reverse umber with patches of saffron. On PDA surface with fluffy white aerial mycelium, umber with patches of saffron, in reverse umber with patches of pale luteous to luteous.

Specimen examined. HONG KONG, Victoria Peak, from *Arenga engleri*, 7 Oct. 1999, K.D. Hyde (holotype CBS H-21104, culture ex-type CBS 114979 = HKUCC 5527).

Notes — The *Diaporthe* species occurring on palms are summarised by Fröhlich et al. (1997). *Diaporthe arengae* (clade 81) is distinguished from known species based on a combination of its conidial morphology and host.

Diaporthe aspalathi E. Jansen, Castl. & Crous, Stud. Mycol. 55: 71. 2006

Basionym. *Diaporthe phaseolorum* var. *meridionalis* F.A. Fernández, Mycologia 88: 438. 1996 (non *D. meridionalis* Sacc., Syll. Fung. 1: 638. 1878).

Specimens examined. SOUTH AFRICA, Western Cape Province, Clanwilliam, Langebergpunt, in branch on *Aspalathus linearis*, J.C. Janse van Rensburg (ex-type culture CBS 117169 = CPC 5428); in crown on *Aspalathus linearis*, 17 Oct. 1997, J.C. Janse van Rensburg (CBS 117168 = CPC 5420); on *Aspalathus linearis*, 2 Dec. 1996, S. Lamprecht (CBS 117500 = CPC 5408).

Notes — *Diaporthe aspalathi* (clade 93) causes soybean stem canker in the South-eastern USA (Fernández & Hanlin 1996), and is not closely related to *D. phaseolorum* as might be expected. Although morphologically similar, this species clustered apart from the reference strain of *D. phaseolorum* (clade 4). *Diaporthe aspalathi* is also the main causal organism of canker and dieback of rooibos (*Aspalathus linearis*), and not *D. phaseolorum* as reported earlier (Smit & Knox-Davies 1989a, b, van Rensburg et al. 2006).

Diaporthe australafricana Crous & Van Niekerk, Australas. Pl. Pathol. 34: 33. 2005

Specimens examined. AUSTRALIA, on *Vitis vinifera*, 1 July 1995, R.W.A. Schepers (ex-type culture CBS 111886 = CPC 2676). — SOUTH AFRICA, on *V. vinifera*, 1 Nov. 1997, L. Mostert (CBS 113487 = CPC 2655).

Notes — Clade 49 contains two isolates of *D. australafricana*, one of them being the ex-type strain (CBS 111886), which is a sibling species of *D. viticola* in clade 50 (van Niekerk et al. 2005). Both species were described from *Vitis vinifera*, but *D. australafricana* is thus far only known from grapevines in Australia and South Africa.

Diaporthe batatas Harter & E.C. Field, Phytopathology 2: 121. 1912

Specimen examined. USA, on *Ipomoea batatas*, Feb. 1921, L.L. Harter (CBS 122.21).

Notes — Clade 8 consists of a single strain of *D. batatas* isolated from *Ipomoea batatas* in the USA. This species and *D. phaseolorum* have in the past been considered as varieties, namely *D. phaseolorum* var. *batatas* and *D. phaseolorum* var. *batatae*. However, the genetic data revealed no homology between the two species. Although it is not certain if CBS 122.21 (culture sterile) is an ex-type strain of *D. batatas*, it is regarded as authentic for the name.

Diaporthe beckhausii Nitschke, Pyrenomyces Germanici 2: 295. 1870

Specimen examined. UNKNOWN, from *Viburnum* sp., Sept. 1927, L.E. Wehmeyer (CBS 138.27).

Notes — Clade 47 is represented by *D. beckhausii*, which was isolated from *Viburnum* sp. (origin unknown, presumably North America, whereas the species was originally described from *Viburnum* collected in Germany). *Diaporthe beckhausii* is known from woody stems of *Betula* sp., *Cydonia japonica*, *Elaeagnus angustifolia*, *Halesia* sp., *Menispermum canadense*, *Menispermum* sp., *V. opulus*, *Viburnum* sp. and *V. tinus* in temperate North America and Europe (Farr & Rossman 2012).

Diaporthe brasiliensis R.R. Gomes, C. Glienke & Crous, sp. nov. — MycoBank MB802926; Fig. 5

Etymology. Named after the country where it was collected, Brazil.

Conidiomata pycnidial, globose to conical, immersed, scattered or aggregated, brown to black, ostiolate, 70–160 µm wide,



Fig. 5 *Diaporthe brasiliensis* (CBS 133183). a. Conidiomata sporulating on PDA; b, c. transverse section through conidiomata, showing conidiomatal wall; d, e. conidiogenous cells; f, g. alpha conidia. — Scale bars: b = 80 µm, all others = 10 µm.

60–140 µm tall, necks 60–130 µm tall, outer surface smooth; pycnidial wall consisting of brown, thick-walled cells of *textura angularis*; conidial mass globose, white to pale-luteous. Conidiophores hyaline, cylindrical, filiform, straight to curved, 1–3-septate, (17–)20–27(–30) × 2(–4) µm. Alpha conidiogenous cells hyaline, cylindrical, filiform, straight to curved, collarette flared, with slight periclinal thickening, (7–)8–12(–14) × 2(–3) µm. Alpha conidia hyaline, ellipsoid to irregular, apex bluntly rounded, base obtuse to subtruncate, bi- to multi-guttulate, 6–7(–8) × 2–3 µm. Beta and gamma conidia not observed.

Culture characteristics — Colonies on PDA flat, with an entire edge, surface mycelium dense and felty, buff, grey-olivaceous or olivaceous-grey; colonies covering dish after 2 wk at 25 °C in the dark; reverse olivaceous, dull green, olivaceous-buff. On OA raised, entire edge, surface mycelium dense felty, smoke-grey to grey-olivaceous; reverse purplish grey to pale purplish grey, grey olivaceous or olivaceous buff. On MEA raised, with an entire edge, buff, smoke-grey, with patches of olivaceous-grey and vinaceous-buff; reverse dark mouse-grey, buff.

Specimens examined. BRAZIL, Rio de Janeiro, endophytic species isolated from leaf of *Aspidosperma tomentosum* (popular name Peroba-do-campo), July 2007, K. Rodriguez (holotype CBS H-21100, ex-type culture CBS 133183 = LGMF 924 = CPC 20300); same collection details (LGMF 926 = CPC 20302).

Notes — Endophytic isolates (clade 36) from a medicinal plant in Brazil.

Diaporthe carpini (Pers.) Fuckel, Jahrb. Nassauischen Vereins Naturk. 23–24: 205. 1870 (1869–1870)

Basionym. *Sphaeria carpini* Pers., Syn. Meth. Fung. (Göttingen) 1: 39. 1801.

Specimen examined. SWEDEN, Skåne, S. Mellby par., Stenshuvud, on *Carpinus betulus*, 14 Apr. 1989, K. & L. Holm (CBS 114437 = UPSC 2980).

Notes — *Diaporthe carpini* (clade 55) is known from several European countries, where it occurs on *Carpinus* spp.

Diaporthe caulivora (Athow & Caldwell) J.M. Santos, Vranečić & A.J.L. Phillips, Persoonia 27: 13. 2011

Basionym. *Diaporthe phaseolorum* var. *caulivora* Athow & Caldwell, Phytopathology 44: 323. 1954.

Specimens examined. CANADA, Ontario, in mature stem on *Glycine soja*, Mar. 1955, A.A. Hildebrand (CBS 178.55 = ATCC 12048 = CECT 2023). — CROATIA, in stem on *Glycine max*, K. Vranečić (ex-neotype culture CBS 127268).

Notes — Clade 91 is represented by two isolates of *D. caulivora* on *Glycine soja* and *G. max*, respectively obtained from Canada (CBS 178.55) and Croatia (ex-neotype: CBS 127268). The soybean canker species complex was recently treated by Santos et al. (2011).

Diaporthe celastrina Ellis & Barthol., J. Mycol. 8, 4: 173. 1902

Specimen examined. UNKNOWN, on *Celastrus scandens*, Sept. 1927, L.E. Wehmeyer (CBS 139.27).

Notes — Strains from the USA are required to confirm the identity of this culture (clade 66).

Diaporthe chamaeropis (Cooke) R.R. Gomes, C. Glienke & Crous, comb. nov. — MycoBank MB802927; Fig. 6

Basionym. *Phoma chamaeropis* Cooke, Grevillea 13 (no. 68): 95. 1885.

≡ *Phomopsis chamaeropis* (Cooke) Petr., as '*Phomopsis chamaeropis*', Ann. Mycol. 17, 2/6: 83. 1920 (1919).

Conidiomata pycnidial in culture on PNA, globose, up to 400 µm diam (up to 600 µm diam on OA), black, erumpent; cream conidial droplets exuding from central ostioles; walls consisting of 3–6 layers of medium brown *textura angularis*. Conidiophores hyaline, smooth, 1–5-septate, branched, densely aggregated, cylindrical, straight to sinuous, 10–50 × 2–2.5 µm. Conidiogenous cells 10–20 × 1.5–2 µm, phialidic, cylindrical, terminal and lateral, with slight taper towards apex, 1–1.5 µm diam, with visible periclinal thickening; collarette not observed. Paraphyses not observed. Alpha conidia aseptate, hyaline, smooth, guttulate, fusoid to ellipsoid, tapering towards both ends, straight, apex subobtuse, base subtruncate, (5–)6–8(–9) × 2(–2.5) µm. Gamma conidia not observed. Beta conidia spindle-shaped, aseptate, smooth, hyaline, apex acutely rounded, base truncate, tapering from lower third towards apex, curved, (20–)22–27(–30) × 1.5(–2) µm.

Culture characteristics — Colonies covering dish after 2 wk in the dark at 25 °C. On OA with moderate aerial mycelium, surface dirty white with patches of pale olivaceous-grey, reverse with patches of dirty white and sienna. On MEA surface




Fig. 6 *Diaporthe chamaeropis* (CBS 454.81). a. Conidiomata sporulating on PDA; b. conidiomata sporulating on PNA; c–e. conidiogenous cells; f. alpha conidia; g. beta conidia. — Scale bars = 10 µm.

dirty white with patches of olivaceous-grey, reverse sienna, with patches of luteous. On PDA surface olivaceous-grey with patches of dirty white, reverse iron-grey.

Specimens examined. CROATIA, Rab, slope behind Hotel 'Imperial', on dead branch of *Spartium junceum*, July 1970, J.A. von Arx (CBS 753.70). – GREECE, Thessaloniki, dead part of leaf of *Chamaerops humilis*, Aug. 1981, H.A. van der Aa (CBS 454.81).

Notes — Conidial dimensions closely fit those provided in the original description (on *Chamaerops humilis* from Czechoslovakia; Uecker 1988), suggesting that these cultures (clade 77) could be authentic for the name.

***Diaporthe cinerascens* Sacc., Syll. Fung. (Abellini) 1: 679. 1882.**

— Fig. 7

= *Phoma cinerascens* Sacc., Michelia 1 (no. 5): 521. 1879.

≡ *Phomopsis cinerascens* (Sacc.) Traverso, Fl. Ital. Crypt. Pyrenomycetae 2, 1: 278. 1906.

Conidiomata pycnidial, sporulating poorly on MEA, globose, up to 300 µm diam, black, erumpent; creamy-luteous conidial droplets exuding from central ostioles; walls consisting of 3–6 layers of medium brown *textura angularis*. *Conidiophores* hyaline, smooth, 1–3-septate, branched, densely aggregated, cylindrical, straight to sinuous, 17–30 × 2–3 µm. *Conidiogenous cells* 8–18 × 2–3 µm, phialidic, cylindrical, terminal and lateral, with slight taper towards apex, 1.5–2 µm diam, with visible periclinal thickening; collarette mostly absent, slightly flared when present, up to 2 µm long. *Paraphyses* not observed. *Alpha conidia* aseptate, hyaline, smooth, guttulate, fusoid to ellipsoid, tapering towards both ends, straight, apex subobtuse, base subtruncate, 7–8(–9) × (2.5–)3 µm. *Gamma conidia* aseptate, hyaline, smooth, ellipsoid-fusoid, apex acutely rounded, base subtruncate, 8–12 × 3 µm. *Beta conidia* not observed.

Culture characteristics — Colonies covering dish after 2 wk in the dark at 25 °C. On MEA with profuse aerial mycelium, surface dirty white, reverse ochreous with patches of umber. On PDA with sparse aerial mycelium, surface olivaceous-grey, reverse iron-grey. On OA surface with moderate aerial mycelium, olivaceous-grey to pale olivaceous-grey.

Specimen examined. BULGARIA, Kostinbrod, Plant Protection Institute, on branch of *Ficus carica*, 1995, E. Ilieva (CBS 719.96).

Notes — *Diaporthe cinerascens* (clade 76) represents a European species occurring on *Ficus*, so the present culture could be authentic for the name, as the conidial dimensions match those provided in the original description. This species was originally associated with canker and dieback of *Ficus* spp. in Italy (Saccardo 1879), and the causal organism identified as *Phomopsis cinerascens* (sexual morph: *Diaporthe cinerascens*) by Grove (1935). *Diaporthe cinerascens* affects all commercial figs in California (Ogawa & English 1991), and is found in several geographical locations of the world (Hampson 1981,

Anderson & Hartman 1983, Benschop et al. 1984, Banihashemi & Javadi 2009). *Ficus* spp. are important exotic garden ornamentals across the USA and Canada as well as in the tropics.

***Diaporthe citri* F.A. Wolf, J. Agric. Res. 33, 7: 625. 1926**

= *Phomopsis citri* H.S. Fawc., Phytopathology 2, 3: 109. 1912.

Specimens examined. BRAZIL, on seed of *Glycine max*, A. Almeida EM-BRAPA/PR (LGMF 946 = CPC 20322). – ITALY, unknown host, June 1939, G. Goiánich (CBS 199.39). – SURINAME, Paramaribo, on decaying fruit of *Citrus sinensis*, Apr. 1932, N.J. van Suchtelen (CBS 230.52).

Notes — Clade 6 is represented by three isolates. One isolate (CBS 199.39) was previously identified as *D. conorum* from Italy, while another originates from soybean seed collected in Brazil (LGMF 946), and the third isolate is from *Citrus sinensis* in Suriname (CBS 230.52). Because *D. conorum* is regarded as synonym of *D. eres* (clade 67), we tentatively refer to this clade as *D. citri*, awaiting more isolates from *Citrus*. *Diaporthe citri* is a serious pathogen that is widely distributed, and associated with melanosis and stem-end rot of citrus fruits (Punithalingam & Holliday 1973, McKenzie 1992, Mondal et al. 2007, Farr & Rossman 2012).

***Diaporthe convolvuli* (Ormeno-Nuñez, Reeleder & A.K. Watson) R.R. Gomes, C. Glienke & Crous, comb. nov. — MycoBank MB802928**

Basionym. *Phomopsis convolvuli* Ormeno-Nuñez, Reeleder & A.K. Watson, Canad. J. Bot. 66, 11: 2232. 1988.

Specimen examined. TURKEY, isolated from leaves with anthracnose on *Convolvulus arvensis*, D. Berner (CBS 124654 = DP 0727).

Notes — *Phomopsis convolvuli* (clade 3) was originally described from diseased leaves of *Convolvulus arvensis* in Québec (Ormeno-Nuñez et al. 1988). The isolate of *Phomopsis convolvuli* studied here (CBS 124654), was found causing anthracnose on field bindweed (*Convolvulus arvensis*), a troublesome perennial weed to many important agricultural crops in the world, and was considered potentially useful as biological control agent (Kuleci et al. 2009).

***Diaporthe crataegi* (Carr.) Fuckel, Jahrb. Nassauischen Vereins Naturk. 23–24: 204. 1870**

Basionym. *Valsa crataegi* Carr., Trans. Linn. Soc. London 22: 278. 1858.

Specimen examined. SWEDEN, Skåne, Trolle-Ljungby par., Tosteberga, on *Crataegus oxyacantha*, 15 Apr. 1989, K. & L. Holm (CBS 114435 = UPSC 2938).

Notes — Clade 41 is represented by *D. crataegi* isolated from *Crataegus oxyacantha* in Sweden. The species is common on *C. chrysocarpa*, *C. laevigata* and *C. oxyacantha* in Canada and Europe (Farr & Rossman 2012).



Fig. 7 *Diaporthe cinerascens* (CBS 719.96). a. Conidiomata sporulating on PDA; b, c. conidiogenous cells; d. alpha conidia. — Scale bars = 10 µm.

Diaporthe crotalariae G.F. Weber, Phytopathology 23: 602.

1933

= *Phomopsis crotalariae* G.F. Weber, Phytopathology 23: 602. 1933.

Specimen examined. USA, on *Crotalaria spectabilis*, Oct. 1933, G.F. Weber (ex-type culture CBS 162.33).

Notes — Clade 92 contains the ex-type strain (CBS 162.33) of *D. crotalariae* isolated from *Crotalaria spectabilis* in the USA.

Diaporthe cuppatea (E. Jansen, Lampr. & Crous) Udayanga, Crous & K.D. Hyde, Fung. Diversity 56: 166. 2012

Basionym. *Phomopsis cuppatea* E. Jansen, Lampr. & Crous, Stud. Mycol. 55: 72. 2006.

Specimen examined. SOUTH AFRICA, Western Cape Province, on *Aspalathus linearis*, 2006, J. Janse van Rensburg (holotype CBS H-19687, ex-type culture CBS 117499 = STE-U 5431 = CPC 5431).

Notes — *Diaporthe cuppatea* (clade 20) is known only from the original collection made from dying branches of *Aspalathus linearis* in South Africa (van Rensburg et al. 2006).

Diaporthe cynaroidis Marinc., M.J. Wingf. & Crous, CBS Biodiversity Ser. (Utrecht) 7: 39. 2008

Specimen examined. SOUTH AFRICA, Western Cape Province, on leaf litter of *Protea cynaroides*, 26 June 2000, S. Marincowitz (ex-type culture CBS 122676 = CMW 22190 = CPC 13180).

Notes — Clade 48 contains the ex-type culture of *D. cynaroidis* (CBS 122676), which was isolated from *Protea cynaroides* in South Africa (Marincowitz et al. 2008). This species is closely related to *D. australaficana* and *D. viticola* (clades 49 and 50, respectively).

Diaporthe decedens (Pers.) Fuckel, Jahrb. Nassauischen Vereins Naturk. 23–24: 30. 1871

Basionym. *Sphaeria tessella* var. *decedens* Pers., Syn. Meth. Fung. (Göttingen) 1: 48. 1801.

Specimens examined. AUSTRIA, on *Corylus avellana*, Oct. 2001, W. Jakitsch (CBS 109772 = AR 3459). — SWEDEN, Öland, Kastlösa par., on *Corylus avellana*, 7 June 1989, K. & L. Holm (CBS 114281 = UPSC 2957).

Notes — *Diaporthe decedens* represents a European species on *Corylus*. Clade 68 consists of two isolates obtained on *Corylus avellana* from Austria and Sweden.

Diaporthe detrusa (Fr.) Fuckel, Jahrb. Nassauischen Vereins Naturk. 23–24: 205. 1869–1870

Basionym. *Sphaeria detrusa* Fr., in Kunze & Schmidt, Mykologische Hefte (Leipzig) 2: 43. 1823.

= *Phoma detrusa* Sacc., Michelia 2: 96. 1880.

≡ *Phomopsis detrusa* (Sacc.) Traverso, Fl. Ital. Crypt. Pars 1: Fungi. Pyrenomyctaceae. Xylariaceae, Valsaceae, Ceratostomataceae 1, 1: 195. 1906.

Specimens examined. AUSTRIA, on *Berberis vulgaris*, Oct. 2001, A.Y. Rossman (CBS 109770 = AR 3424). — SWEDEN, Uppland, Hällnäs par., on *Berberis vulgaris*, 14 May 1991, K. & L. Holm (CBS 114652 = UPSC 3371). — UNKNOWN, on *Berberis vulgaris*, Sept. 1927, L.E. Wehmeyer (CBS 140.27).

Notes — Clade 54 contains three isolates of *D. detrusa* obtained from *Berberis vulgaris* in Austria, Sweden and one of them with an unknown origin (presumably North America). This European species is known to also occur in the USA (Farr & Rossman 2012).

Diaporthe elaeagni Rehm, Syll. Fung. 14: 546. 1899. — Fig. 8

?= *Phoma elaeagni* Sacc., Michelia 1, 3: 354. 1878.

≡ *Phomopsis elaeagni* (Sacc.) Petr., Ann. Mycol. 19, 1–2: 48. 1921.

Specimen examined. NETHERLANDS, Maassluis, on twig of *Elaeagnus* sp., May 1972, J. Gremmen (CBS 504.72).

Notes — In culture CBS 504.72 (clade 73) primarily produces beta conidia (spindle shaped, 16–22 × 2 µm, thus wider than seen on average in most other species); alpha conidia rarely observed, fusoid-ellipsoidal, 7–10 × 2–3 µm, thus correlating with dimensions of *Phomopsis elaeagni* (Sacc.) Petr., which is a homonym of *P. elaeagni* Sacc. Furthermore, conidial dimensions of the asexual state of *D. elaeagni* are not known. Additional collections and type studies are thus required to resolve the complex occurring on *Elaeagnus*.

Diaporthe endophytica R.R. Gomes, C. Glienke & Crous, sp. nov. — MycoBank MB802929

Etymology. Named after its endophytic growth habit.

Cultures sterile. *Diaporthe endophytica* (clade 5) differs from its closest phylogenetic neighbour, *D. phaseolorum* (clade 4), by unique fixed alleles in five loci based on alignments of the separate loci deposited in TreeBase as study S13943: ITS positions 357 (C), 359 (G), 360 (T), 368 (A), 369 (A), 371 (A), 372 (G) and 373 (G); TUB positions 135 (C) and 592 (T); CAL position 145 (G); TEF1 positions 18 (G), 26 (T), 40 (T), 42 (A), 63 (A), 124 (A), 175 (A) and 343 (A); HIS position 369 (C).

Culture characteristics — Colonies with sparse aerial mycelium, covering the dish after 2 wk at 25 °C. On PDA buff, honey to isabelline; reverse smoke-grey. On OA smoke-grey to olivaceous-grey. On MEA buff with umber patches; reverse dark mouse-grey, with patches of isabelline.

Specimens examined. BRAZIL, endophytic in leaf on *Schinus terebinthifolius*, July 2007, J. Lima (LGMF 911 = CPC 20287, LGMF 919 = CPC 20295), (holotype CBS H-21107, culture ex-type LGMF 916 = CPC 20292




Fig. 8 *Diaporthe elaeagni* (CBS 504.72). a. Conidiomata sporulating on PNA; b. conidiomata sporulating on PDA; c, d. conidiogenous cells; e. beta conidia.
— Scale bars = 10 µm.

= CBS 133811); endophytic in petiole on *Maytenus ilicifolia*, July 2007, R.R. Gomes (LGMF 928 = CPC 20304, LGMF 934 = CPC 20310, LGMF 935 = CPC 20311, LGMF 937 = CPC 20313); in seed on *Glycine max*, A. Almeida EMBRAPA/PR (LGMF 948 = CPC 20324).

Notes — Clade 5 represents a distinct lineage, containing eight sterile isolates originating from Brazil. Four of them were isolated from *Maytenus ilicifolia*, three from *S. terebinthifolius* and one from soybean seeds. Isolates could not be induced to sporulate on any of the media defined in this study, nor on sterilised plant host tissue placed on WA.

Diaporthe eres Nitschke, Pyrenomycetes Germanici 2: 245. 1870

= *Phomopsis cotoneastri* Punith., Trans. Brit. Mycol. Soc. 60, 1: 157. 1973.
= *Phoma oblonga* Desm., Ann. Nat. Sci. Bot. 20: 218. 1853.
≡ *Phomopsis oblonga* (Desm.) Traverso, Fl. Ital. Crypt. Pars 1: Fungi. Pyrenomycetae. Xylariaceae, Valsaceae, Ceratostomataceae: 248. 1906.

Specimens examined. AUSTRIA, on *Acer campestre*, Oct. 2001, W. Jaklitsch (CBS 109767 = AR 3538 = WJ 1643). — GERMANY, Monheim, on leaf spot of *Hordeum* sp., 5 Aug. 1984, M. Hossfeld (CBS 841.84). — ITALY, Milano, on twig of *Juglans regia*, Dec. 1980, M. Bischiach (CBS 102.81). — LATVIA, on *Rhododendron* sp., I. Apine (CBS 129168). — NETHERLANDS, Oostvoorne, on dead stems of *Arctium* sp., 13 Dec. 1984, M. de Nooij (CBS 110.85); Soest, Dalweg, on fallen fruit of *Fraxinus* sp., 21 Feb. 1999, G. Verkley (CBS 101742); Veldhoven, on dead branch of *Sorbus aucuparia*, Nov. 1973, W.M. Loerakker (CBS 287.74); Baarn, garden Chopinlaan, on dead branch of *Wisteria sinensis*, 6 June 1983, H.A. van der Aa (CBS 528.83); Soest, inside house, on *Abutilon* sp., 26 Mar. 1997, A. Aptroot (CBS 688.97); Baarn, potted plant, on cladodes of *Opuntia* sp., 23 Sept. 1996, H.A. van der Aa (CBS 365.97); on *Alliaria officinalis*, Feb. 1962, G.H. Boerema (CBS 445.62); Baarn, on dead leaf of *Ilex aquifolium*, 11 June 1967, H.A. van der Aa (CBS 370.67 = MUCL 9931); Prov. Zuid-Holland, Huize Oud-Poelgeest, Oegstgeest, dieback of *Ilex aquifolium*, 21 Nov. 1994, G.J.M. Verkley (CBS 694.94); Baarn, garden Eemnesserweg 90, on dead stem of *Rumex hydrolapathum*, 19 Mar. 1996, H.A. van der Aa (CBS 485.96); Baarn, Cantonspark, on withering leaf of *Magnolia × soulangeana*, 23 Oct. 1968, H.A. van der Aa (CBS 791.68); Zuid-Holland, Ridderkerk, Huys ten Donck, on leaf tip of *Osmanthus aquifolium*, 7 May 1977, H.A. van der Aa (CBS 297.77); on *Phaseolus vulgaris*, Sept. 1950, Goossens (CBS 422.50); Baarn, on dead stem of *Allium giganteum*, May 1985, H.A. van der Aa (CBS 283.85); from *Laburnum × watereri 'Vossii'*, Apr. 1935, I. de Boer (CBS 267.55); Boskoop, nursery, dying twigs of *Skimmia japonica*, Nov. 1981, H.A. v. Kesteren (CBS 122.82). — UK, Scotland, on living and dead twig of *Fraxinus excelsior*, Feb. 1938, J.A. MacDonald

(CBS 250.38); on *Cotoneaster* sp., 1971, H. Butin (ex-type culture of *P. crotoneaster* CBS 439.82 = BBA P-407 = IMI 162181a); Oxford, on *Picea abies* seedling, Nov. 1937, T.R. Peace (CBS 186.37). — UNKNOWN, on rotten fruit of *Malus sylvestris*, May 1961, Geigy (CBS 375.61); May 1932, W.G. Hutchinson (CBS 267.32).

Notes — *Diaporthe eres* (clade 67) is the type species of the genus *Diaporthe*, and is present in several hosts, though it is known to be morphologically highly variable (Castlebury et al. 2002). Wehmeyer (1933) described this species on more than 60 hosts, and listed several synonymies based on morphological data. A detailed morphological study is required to designate a suitable epitype strain for *D. eres*, and to resolve the status of all its purported synonyms.

Diaporthe eugeniae (Punith.) R.R. Gomes, C. Glienke & Crous, comb. nov. — MycoBank MB802930

Basionym. *Phomopsis eugeniae* Punith., Trans. Brit. Mycol. Soc. 63, 2: 232. 1974.

Specimens examined. WEST SUMATRA, on *Eugenia aromatica*, May 1973, J. Waller (holotype IMI 177560); Lampung, on leaf of *Eugenia aromatica*, July 1982, R. Kasim (CBS 444.82).

Notes — *Diaporthe eugeniae* (clade 83) was originally described on *Eugenia aromatica* from West Sumatra. Although the present isolate could be authentic for the name, it unfortunately proved to be sterile.

Diaporthe fibrosa (Pers.) Fuckel, Jahrb. Nassauischen Vereins Naturk. 23–24: 204. 1870 (1869–1870)

Basionym. *Sphaeria fibrosa* Pers., Syn. Meth. Fung. (Göttingen) 1: 40. 1801.

Specimens examined. AUSTRIA, Vienna, on *Rhamnus cathartica*, Oct. 2001, A.Y. Rossman (CBS 109751 = AR 3425). — SWEDEN, Uppland, Dalby par., Hässleborg, on *Rhamnus cathartica*, 10 Mar. 1987, K. & L. Holm (CBS 113830 = UPSC 2117).

Notes — Clade 52 consists of two isolates from *Rhamnus cathartica* collected in Sweden and Austria. *Diaporthe fibrosa* was originally described from Europe on *Rhamnus*, so these cultures may well prove to be authentic for the name.




Fig. 9 *Diaporthe foeniculacea* (CBS 111554). a. Conidiomata sporulating on PDA; b, c. transverse section through conidiomata, showing conidiomatal wall; d–f. conidiogenous cells; g. beta conidia; h. alpha conidia. — Scale bars: b = 250 µm, all others = 10 µm.

Diaporthe foeniculacea Niessl, in von Thümen, Contr. Ad. Fl. Myc. Lusit. 2: 30. 1880. — Fig. 9

= *Phoma foeniculina* Sacc., Syll. Fung. 3: 125. 1884.
 ≡ *Phomopsis foeniculina* (Sacc.) Câmara, Agron. Lusit. 9: 104. 1947.
 = *Phomopsis theicola* Curzi, Atti Ist. Bot. Univ. Pavia, 3 sér., 3: 65. 1927.
 = *Diaporthe neotheicola* A.J.L. Phillips & J.M. Santos, Fung. Diversity 34: 120. 2009.

Conidiomata pycnidial, eustromatic, multilocular, immersed, ostiolate, dark brown, scattered or aggregated, 350–890 µm wide, 160–320 µm tall, necks absent, outer surface covered with hyphae; pycnidial wall consisting of brown, thick-walled cells of *textura angularis*; conidial mass globose to conical and exuding in cirri, yellow to reddish brown. *Conidiophores* hyaline, subcylindrical and cylindrical, filiform, branched above the septa, tapering towards the apex, 1–3-septate, (19–)20–28(–32) × 2(–3) µm. *Conidiogenous cells* hyaline, subcylindrical and filiform, straight, slightly tapering towards the apex, collarette not flared, prominent periclinal thickening, (10–)11–15(–17) × 2(–3) µm. *Alpha conidia* hyaline, oblong to ellipsoidal, apex bluntly rounded, base obtuse to subtruncate, bi- to multi-guttulate (6–)7–9 × 2(–3) µm. *Beta conidia* hyaline, smooth, slightly curved, (26–)28–32(–34) × 1(–2) µm. *Gamma conidia* not observed (based on isolate CBS 111554).

Specimens examined. INDIA, Calcutta, unknown host, Feb. 1948, S.R. Bose (CBS 400.48). — ITALY, on leaves and branches of *Camellia sinensis*, Oct. 1927, M. Curzi (ex-type culture of *P. theicola* CBS 187.27); Perugia, on *Diospyros kaki*, June 1956, M. Ribaldi (CBS 287.56); Apulia, near Bari, on *Prunus amygdalus*, winter 1974/75, A. Ciccarone (CBS 171.78). — NETHERLANDS, Baarn, 'Madoera', back frond, on *Wisteria sinensis*, 24 Apr. 1969, H.A. van der Aa (CBS 357.69). — NEW ZEALAND, Waikato region, on *Pyrus pyrifolia*, 2001, W. Kandula (CBS 116957). — PORTUGAL, near Lisbon, São Marcos, base of senescent stem of *Foeniculum vulgare*, Apr. 2002, A.J.L. Phillips (CBS 111554); Évora, *Foeniculum vulgare*, 1 Nov. 2007, A.J.L. Phillips (ex-type cultures of *Diaporthe neotheicola* CBS 123209, CBS 123208); Pedras del Rei, near Tavira, on *Bougainvillea spectabilis*, 15 June 1988, H.A. van der Aa (CBS 603.88); Madeira, Serra da Agua, base of senescent stem of *Foeniculum vulgare*, Aug. 2001, A.J.L. Phillips (CBS 111553).

Notes — *Diaporthe foeniculacea* (clade 78) was originally described from *Foeniculum vulgare* in Portugal, and represents an older name for *D. theicola* and *D. neotheicola*. There are many described species that occur on *Foeniculum vulgare* (wild fennel). Among them, *P. theicola* and its teleomorph *D. neotheicola* (Santos & Phillips 2009), and *D. foeniculacea*, the causal agent of stem necrosis of fennel. Phillips (2003) redescribed *D. foeniculacea*, and established the sexual-aseexual connection between *D. foeniculacea* and *Phomopsis foeniculina*. The synonymy of *D. neotheicola* under *D. foeniculacea* is based on the fact that the cultures matching the original descriptions are in fact genetically identical. However, as there are no ex-type strains of *D. foeniculacea*, this synonymy strongly relies on the earlier opinion of Phillips (2003). Either way, this matter can only be resolved once an epitype has been designated for *D. foeniculacea*, fixing the application of the name. We recommend that additional collections linked to stem necrosis of fennel in Portugal are obtained, before this decision is made.

Diaporthe ganjae (McPartl.) R.R. Gomes, C. Glienke & Crous, comb. nov. — MycoBank MB802932

Basionym. *Phomopsis ganjae* McPartl., Mycotaxon 18, 2: 527. 1983.

Specimen examined. USA, Illinois, Hannah City, dead leaf of *Cannabis sativa*, deposited Mar. 1991, J.M. McPartland (holotype HA 10987, ex-type culture ILLS 43621 = CBS 180.91).

Notes — *Diaporthe ganjae* (clade 24) is known only from the original collection taken from wilted, dead leaves of *Cannabis sativa* in Illinois, USA (McPartland 1983). Phylogenetically *D. ganjae* is closely related to an isolate identified as *D. mani-*

hotia (CBS 505.76), isolated from *Manihot utilissima* in Rwanda (clade 25).

Diaporthe gardeniae (Buddin & Wakef.) R.R. Gomes, C. Glienke & Crous, comb. nov. — MycoBank MB802933

Basionym. *Phomopsis gardeniae* Buddin & Wakef., Gard. Chron., ser. 3 103: 45. 1938.

= *Phomopsis gardeniae* H.N. Hansen & Barrett, Mycologia 30, 1: 18. 1938 (homonym).

Specimen examined. ITALY, on stem of *Gardenia florida*, June 1956, M. Ribaldi (CBS 288.56).

Notes — *Diaporthe gardeniae* (clade 59) causes gardenia canker in *Gardenia jasminoides*, *G. lucida* and *Gardenia* sp. (Farr & Rossman 2012). This disease is considered as serious (Tilford 1934, Huber 1936, Miller 1961). It was originally observed in 1894 in England (Cooke 1894), and has since been reported from the USA (Preston 1945) and India (Mathur 1979). All parts of the plant are susceptible to infection, including roots, stems and leaves (McKenzie et al. 1940), although cankered stems are the most diagnostic symptoms for this disease.

Diaporthe helianthi Munt.-Cvetk., Mihaljč. & M. Petrov, Nova Hedwigia 34: 433. 1981

= *Phomopsis helianthi* Munt.-Cvetk., Mihaljč. & M. Petrov, Nova Hedwigia 34: 433. 1981.

Specimens examined. SERBIA, Vojvodina, overwintering stem on *Helianthus annuus*, 1980, M. Muntañola-Cvetkovic (ex-type culture CBS 592.81 = CBS H-1540). — UNKNOWN, on seed of *H. annuus*, June 1994, Vanderhave Res., Rilland, Netherlands (CBS 344.94).

Notes — *Diaporthe helianthi* (clade 14) is associated worldwide with stem canker and grey spot disease of sunflower (*Helianthus annuus*) (Muntañola-Cvetkovic' et al. 1981). Yield reductions of up to 40 % have been recorded in Europe (Masicrevic & Gulya 1992) including the former Yugoslavia as well as France where it was considered a major pathogen of sunflower (Battilani et al. 2003, Debaeke et al. 2003). *Diaporthe helianthi* is also widespread in the sunflower growing regions of the USA (Gulya et al. 1997). The wide geographic distribution, and high genetic variability of the pathogen lead to the evolution of new strains that could be more aggressive, causing large yield losses and a decline in disease control (Pecchia et al. 2004, Rekab et al. 2004).

Diaporthe cf. heveae 1

Specimen examined. BRAZIL, São Paulo, from *Hevea brasiliensis*, Apr. 1997, D.S. Attili (CBS 852.97) (originally identified as *Phomopsis heveae*).

Notes — *Diaporthe heveae* and *Phomopsis heveae* were both described from *Hevea* in Sri Lanka, and could represent the same species. Two isolates deposited in CBS under this name, CBS 852.97 (from *Hevea brasiliensis* in Brazil) and CBS 681.84 (from *Hevea brasiliensis* in India) were shown to represent two distinct species (clades 46 and 80, respectively). However, as both were found to be sterile, their taxonomy could not be resolved.

Diaporthe cf. heveae 2

Specimen examined. INDIA, Kerala, Kottayam, in leaf on *Hevea brasiliensis*, Sept. 1984, K. Jayarathnam (CBS 681.84).

Notes — Isolate CBS 681.84 (clade 80, *P. heveae* from *Hevea brasiliensis* in India) is sterile, and thus its taxonomy could not be resolved. *Diaporthe heveae* has been reported from Brazil, China, India, Indonesia, Malaysia, Sri Lanka and Thailand (Holiday 1980, Zhuang 2001, Udayanga et al. 2011).

Diaporthe hickoriae Wehm., Monogr. Gen. Diaporthe Nitschke & Segreg., Univ. Michigan Stud., Sci. Ser. 9: 149. 1933

Specimen examined. USA, Michigan, on *Carya glabra*, June 1926, L.E. Wehmeyer (ex-type culture CBS 145.26).

Notes — *Diaporthe hickoriae* (clade 72) occurs on the bark of *Carya glabra* in the USA (Wehmeyer 1933).

Diaporthe hongkongensis R.R. Gomes, C. Glienke & Crous, sp. nov. — MycoBank MB802934; Fig. 10

Etymology. Named after the location where it was collected, Hong Kong.

Conidiomata pycnidial, superficial to embedded on PDA, solitary to aggregated, globose with central ostiole, exuding a creamy conidial cirrus; pycnidial up to 200 µm diam; wall of 3–6 layers of brown *textura angularis*. *Conidiophores* lining the inner cavity, reduced to conidiogenous cells. *Conidiogenous cells* phialidic, hyaline, smooth, ampulliform to subcylindrical with prominent apical taper, 5–12 × 2–4 µm; apex with periclinal thickening and minute collarette, 1 µm long. *Paraphyses* intermingled among conidiophores, hyaline, smooth, frequently branched below, up to 4-septate, with clavate terminal cell, up to 80 µm long, apex 2–8 µm diam. *Alpha conidia* hyaline, smooth, granular to guttulate, aseptate, fusiform, tapering towards both ends, mostly straight, apex acutely rounded, base truncate, (5–)6–7(–8) × (2–)2.5(–3) µm. *Gamma conidia* aseptate, hyaline, smooth, ellipsoid-fusoid, apex subobtuse, base truncate, 10–13 × 2 µm. *Beta conidia* aseptate, hyaline, smooth, spindle-shaped, apex acutely rounded, base truncate, widest in mid region, mostly curved in upper part, 18–22 × 1.5–2 µm.

Culture characteristics — Colonies covering dish after 2 wk in the dark at 25 °C, with moderate aerial mycelium. On OA surface dirty white with patches of pale olivaceous-grey, reverse dirty white with patches of olivaceous-grey and iron-grey. On PDA surface iron-grey, with patches of dirty white, reverse iron-grey. On MEA surface dirty white with patches of olivaceous-grey, reverse iron-grey with patches of dirty white.

Specimen examined. HONG KONG, Tai Po Kau, on fruit of *Dichroa febrifuga*, 20 Feb. 2002, K.D. Hyde (holotype CBS H-21103, culture ex-type CBS 115448 = HKUCC 9104).

Notes — Isolate CBS 115448 (clade 79; reported as *Phomopsis pittospori* on *Dichroa febrifuga* from Hong Kong) is morphologically distinct from *P. pittospori* (from *Pittosporum* twigs in California; alpha conidia 6–8 × 1.5 µm, beta conidia 18–20 × 1 µm), with wider alpha and beta conidia.

Diaporthe hordei (Punith.) R.R. Gomes, C. Glienke & Crous, comb. nov. — MycoBank MB802935

Basionym. *Phomopsis hordei* Punith., Trans. Brit. Mycol. Soc. 64, 3: 428. 1975.

Specimen examined. NORWAY, Fellesbygget, As, on root of *Hordeum vulgare*, Oct. 1992, L. Sundheim (CBS 481.92).

Notes — *Diaporthe hordei* (clade 13) was described from *Hordeum vulgare* in the UK. Although the present culture could be authentic (from *Hordeum* collected in Norway), it proved to be sterile, so its morphology could not be confirmed.

Diaporthe impulsa (Cooke & Peck) Sacc., Syll. Fung. (Abelini) 1: 618. 1882

Basionym. *Valsa impulsa* Cooke & Peck, Ann. Rep. N.Y. State Mus. Nat. Hist. 27: 109. 1875 (1874).

Specimens examined. SWEDEN, Uppland, Dalby par., Jerusalem, on *Sorbus aucuparia*, 24 Oct. 1989, K. & L. Holm (CBS 114434 = UPSC 3052). — UNKNOWN, on *Sorbus americana*, Sept. 1927, L.E. Wehmeyer (CBS 141.27).

Notes — Clade 51 is represented by two isolates of *D. impulsa* occurring on *Sorbus* spp. *Diaporthe impulsa* is a known pathogen of *Sorbus* spp., and has a wide geographic distribution (Farr & Rossman 2012). It was originally described from *Sorbus* in the USA, thus CBS 141.27 may well prove to be a good reference strain for the species.

Diaporthe inconspicua R.R. Gomes, C. Glienke & Crous, sp. nov. — MycoBank MB802936

Etymology. Referring to its inconspicuous nature, growing as endophyte in host tissue.

Cultures sterile. *Diaporthe inconspicua* (clade 75) differs from its closest phylogenetic neighbours, clade 68–74 and 76–78, by unique fixed alleles in four loci based on alignments of the separate loci deposited in TreeBase as study S13943: TUB positions 33 (A), 102–104 and 106–111 (indels), 127 (G), 149 (C), 151 (A), 195 (C), 204 (T), 357 (G), 446 (G), 449 (C), 465 (T), 484 (T), 559 (A), 592 (A), 629 (T), 653 (T), 708 (C), 732 (C), 754 (A), 763 (C), 784 (A) and 787 (G); CAL positions 28 (C), 102 (G), 114 (T), 148 (T), 152 (T), 153 (A), 157 (C), 170 (G), 199 (C) and 281 (C); TEF1 positions 9 (T), 16 (A), 22 (A), 29 (G), 30 (G), 81 (C), 86 (C), 87 (A), 88 (A), 89 (T), 131 (A), 275 (A), 298 (C) and 315 (T); HIS positions 139 (T), 211 (T), 244 (T) and 408 (T).

Culture characteristics — Colonies covering the dish after 2 wk in the dark at 25 °C. On OA spreading, flat with sparse aerial mycelium, surface cream in centre, umber in outer region.



Fig. 10 *Diaporthe hongkongensis* (CBS 115448). a, b. Conidiomata sporulating on PDA; c, d. conidiogenous cells; e. beta conidia; f. alpha conidia. — Scale bars = 10 µm.

On PDA surface and reverse cream to dirty white with sparse aerial mycelium. On MEA with sparse aerial mycelium, surface becoming folded, dirty white in centre, sienna in outer region, and luteous in reverse.

Specimens examined. BRAZIL, on petiole of *Maytenus ilicifolia*, July 2007, R.R. Gomes (holotype CBS H-21102, ex-type culture LGMF 930 = CPC 20306 = CBS 133813); same collection details (LGMF 931 = CPC 20307); on *Spondias mombin*, 2007, K. Rodriguez (LGMF 922 = CPC 20298).

Notes — Sterile endophytic isolates (clade 75) from medicinal plants in Brazil.

Diaporthe infecunda R.R. Gomes, C. Glienke & Crous, sp. nov.
— MycoBank MB802937

Etymology. Named after its sterile growth in culture.

Cultures sterile. *Diaporthe infecunda* (clade 23) differs from its closest phylogenetic neighbours, clade 16–22, by unique fixed alleles in five loci based on alignments of the separate loci deposited in TreeBase as study S13943: ITS positions 108 (T), 279 (C), 292 (G), 359 (C) and 360 (G); TUB positions 11 (indel), 106 (G), 138 (T), 140 (A), 153 (G), 155 (T), 184 (A), 197 (G), 202 (C), 302 (A), 354 (A), 369–374 (indels), 398 (G), 407 (indel), 414 (C), 422 (T), 424 (G), 425 (C), 432 (G), 452 (C), 454 (C), 458 (C), 461 (G), 479 (T), 482 (T), 486 (C), 540 (T), 572 (C), 622 (A), 694 (T), 696 (T), 697 (G), 716 (C), 728 (C), 776 (G), 778 (G) and 796 (C); CAL positions 64 (T), 83 (T), 104 (G), 146 (C), 151 (C), 155 (G), 159 (C), 172 (C), 176 (T), 179 (A), 184 (G), 197 (T), 206 (T), 212 (C) and 221 (T); TEF1 positions 6 (A), 9 (G), 13 (G), 16 (C), 21 (A), 30 (G), 32 (indel), 39 (A), 40 (A), 41 (G), 42 (T), 43 (A), 79 (G), 83 (T), 90 (T), 92 (T), 96 (A), 97 (C), 106 (C), 116 (A), 120 (C), 123 (A), 127 (A), 132 (A), 135 (G), 173 (G), 255 (T), 284 (A), 294 (C), 299 (C) and 309 (A); HIS positions 173 (T), 196 (T), 197 (G), 199 (C/T), 221 (C), 222 (C), 230 (G), 263 (C), 264 (T), 268 (C), 273 (T) and 279 (C).

Culture characteristics — Colonies covering the dish after 2 wk in the dark at 25 °C. On PDA surface umber with patches of white, reverse chestnut. On MEA surface dirty white, reverse umber. On OA surface with patches of dirty white and umber.

Specimens examined. BRAZIL, on leaf of *Schinus terebinthifolius*, July 2007, J. Lima (holotype CBS H-21095, ex-type culture LGMF 906 = CPC 20282 = CBS 133812); additional isolates with same collection details (LGMF 908 = CPC 20284, LGMF 912 = CPC 20288, LGMF 917 = CPC 20293, LGMF 918 = CPC 20294, LGMF 920 = CPC 20296); in petiole of *Maytenus ilicifolia*, July 2007, R.R. Gomes (LGMF 933 = CPC 20309, LGMF 940 = CPC 20316).

Notes — Clade 23 represents endophytic isolates from leaves of medicinal plants growing in Brazil. It consists of eight isolates, two from *Maytenus ilicifolia*, and six from *Schinus terebinthifolius*.

Diaporthe juglandina (Fuckel) Nitschke, Pyrenomycetes Germanici 2: 281. 1870

Basionym. *Aglaospora juglandina* Fuckel, Fungi Rhenani Exsicc., suppl. 7 (no. 2101–2200): no. 2159. 1868.

Specimen examined. USA, Tennessee, Great Smoky Mts National Park, dead wood of *Juglans* sp., L. Vasilyeva (CBS 121004).

Notes — *Diaporthe juglandina* (clade 65) represents a European taxon described from *Juglans*. European collections are required to confirm whether this name can be applied to the clade.

Diaporthe longispora (Wehm.) R.R. Gomes, C. Glienke & Crous, comb. nov. — MycoBank MB802938

Basionym. *Diaporthe strumella* var. *longispora* Wehm., Mycologia 28, 1: 46. 1936.

Specimen examined. CANADA, Ontario, Toronto, on *Ribes* sp., May 1936, L.E. Wehmeyer (ex-type culture CBS 194.36).

Notes — Clade 27 comprises the ex-type culture of *D. strumella* var. *longispora* isolated from *Ribes* sp., and forms a sister clade with *D. sclerotoides* (clade 28). *Diaporthe strumella* is found on woody limbs, especially of *Ribes* spp. in temperate North America and Europe (Farr & Rossman 2012). As *D. strumella* var. *longispora* is morphologically clearly a distinct species, we elevate this variety to species status.

Diaporthe lusitanicae A.J.L. Phillips & J.M. Santos, Fung. Diversity 34: 118. 2009

Specimen examined. PORTUGAL, Lisbon, Oeiras, Estação Agronómica Nacional, stem of *Foeniculum vulgare*, 14 Aug. 2007, J.M. Santos (ex-type cultures CBS 123212 = Di-C001/5, CBS 123213 = Di-C001/3).

Notes — This species (clade 21) was described in 2009 on senescent stems of *Foeniculum vulgare* (wild fennel) in Portugal by Santos & Phillips (2009).

Diaporthe manihotia Punith., Kavaka 3: 29. 1976 (1975)

= *Phomopsis manihotis* Swarup, L.S. Chauhan & Tripathi, Mycopathol. Mycol. Appl. 28, 4: 345. 1966.

Specimen examined. RWANDA, on leaves of *Manihot utilissima*, 9 July 1976, J. Semal (CBS 505.76).

Notes — *Phomopsis manihotis* (clade 25 as *D. manihotia*) causes leaf spot of cassava (*Manihot esculenta*), though the disease is also referred to as *Phomopsis* blight of tapioca. Severe infection leads to defoliation and stem lesions. Affected areas become shrivelled with numerous pycnidia embedded in the tissue. On severely infected stems the bark starts to gradually peel off, leading to partial or total girdling. The disease is known from Africa (Ethiopia, Nigeria), Asia (India), Central America and West Indies (S.E. Dominica), and South America (Colombia) (Sarbhoy et al. 1971, Mathur 1979, Farr & Rossman 2012).

Diaporthe mayteni R.R. Gomes, C. Glienke & Crous, sp. nov.
— MycoBank MB802939; Fig. 11

Etymology. Named after the host genus from which it was collected, *Maytenus*.

Conidiomata pycnidial, globose, immersed, scattered and aggregated, brown to black, ostiolate, 70–230 µm wide, 40–150 µm tall, with short necks, 40–140 µm; outer surface smooth or covered in hyphae; pycnidial wall consisting of brown, thick-walled cells of *textura angularis*; conidial mass globose or exuding in cirri; predominantly yellow, pale luteous to cream. *Conidiophores* hyaline, subcylindrical to cylindrical, rarely branched above the septa, tapering towards the apex, 1–3-septate, (10–)13–27(–36) × (2–)3(–4) µm. *Conidiogenous cells* hyaline, subcylindrical, rarely tapering towards the apex, collarette present and not flared, with prominent periclinal thickening, (5–)6–10(–13) × 2(–3) µm. *Alpha conidia* hyaline, oblong to ellipsoid, apex bluntly rounded, base obtuse; biguttulate, (5–)6(–7) × (2–)3 µm. *Beta* and *gamma conidia* absent.

Culture characteristics — Colonies on PDA flat, with entire edge, cottony, olivaceous buff, with primrose aerial mycelium in concentric rings, with olivaceous patches; colonies reaching 66 mm diam after 2 wk at 25 °C; reverse olivaceous buff and greenish olivaceous. On OA flat, with entire edge, cottony




Fig. 11 *Diaporthe mayteni* (CBS 133185). a. Conidiomata sporulating on PNA; b, c. transverse section through conidiomata, showing conidiomatal wall; d, e. conidiogenous cells; f. beta conidia; g. alpha conidia. — Scale bars: b = 85 µm, all others = 10 µm.

appressed, buff, white, the center of the colony pale olivaceous-grey, patches isabelline and luteous; colonies reaching 56 mm diam; reverse buff and pale olivaceous grey. On MEA flat, with entire edge, aerial mycelium cottony, white to pale olivaceous grey or olivaceous buff; colonies reaching 37 mm diam; reverse hazel, ochreous, with patches greenish black and olivaceous black.

Specimen examined. BRAZIL, Paraná, Colombo, endophytic species isolated from petiole of *Maytenus ilicifolia* (popular name Espinheira Santa), July 2007, R.R. Gomes (holotype CBS H-21096, ex-type culture CBS 133185 = LGMF 938 = CPC 20314).

Notes — *Diaporthe mayteni* (clade 30) grows endophytically in *Maytenus ilicifolia* in Brazil.

Diaporthe megalospora Ellis & Everh., Proc. Acad. Nat. Sci. Philadelphia 42: 235. 1890

Specimen examined. UNKNOWN, from *Sambucus canadensis*, Sept. 1927, L.E. Wehmeyer (CBS 143.27).

Notes — *Diaporthe megalospora* (clade 15) is known on *Sambucus canadensis* from North America (Wehmeyer 1933, Hanlin 1963, Farr & Rossman 2012). Fresh collections are required to designate an epitype, and fix the genetic application of the name.

Diaporthe melonis Beraha & M.J. O'Brien, Phytopathol. Z. 94, 3: 205. 1979

= *Phomopsis cucurbitae* McKeen, Canad. J. Bot. 35: 46. 1957.

Specimens examined. INDONESIA, Java, Muneng, Exp. Station, on *Glycine soja*, Sept. 1987, H. Vermeulen (CBS 435.87). – USA, Texas, Rio Grande Valley, on *Cucumis melo*, 1978, L. Beraha & M.J. O'Brien (ex-isotype culture CBS 507.78, specimen derived from culture CBS H-891).

Notes — Clade 7 represents *D. melonis* (Beraha & O'Brien 1979), and contains the ex-isotype culture, and one isolate previously identified as *D. phaseolororum* var. *sojae* (though the two isolates are not identical). *Diaporthe melonis* is frequently reported on soybean (Santos et al. 2011). *Phomopsis cucurbitae* (treated here as synonym) is reported to have a cosmopolitan distribution, and to cause black rot disease of greenhouse

cucumbers (McKeen 1957, Punithalingam & Holliday 1975, Ohsawa & Kobayashi 1989).

Diaporthe musigena Crous & R.G. Shivas, Persoonia 26: 119. 2011

Specimen examined. AUSTRALIA, Queensland, Brisbane Botanical Garden, on leaves of *Musa* sp., 14 July 2009, P.W. Crous & R.G. Shivas (ex-type culture CBS 129519 = CPC 17026).

Notes — Clade 84 represents *D. musigena*, isolated from *Musa* sp. in Australia (Crous et al. 2011).

Diaporthe neilliae Peck, Ann. Rep. N.Y. State Mus. Nat. Hist. 39: 52. 1887 (1886)

Specimen examined. UNKNOWN, on *Spiraea* sp., Sep. 1927, L.E. Wehmeyer (CBS 144.27).

Notes — *Diaporthe neilliae* (clade 60) was originally described from *Spiraea* sp. from North America. The origin of the present isolate, however, remains unclear (presumably North America).

Diaporthe neoarctii R.R. Gomes, C. Glienke & Crous, sp. nov.
— MycoBank MB802940, Fig. 12

Etymology. Named after its superficial resemblance to *Diaporthe arctii*.

Conidiomata pycnidial, ampulliform to finger-like, aggregated, dark brown to black, immersed, ostiolate, 300–450 µm wide, 200–670 µm tall, with prominent necks 240–560 µm long, outer surface covered with hyphae; pycnidial wall consisting of brown, thick-walled cells of *textura angularis*; conidial mass globose, pale yellow. *Conidiophores* hyaline, ampulliform to subcylindrical, filiform, branched above the septa, tapering towards the apex, rarely septate, (12–)13–17(–18) × (2–)3 µm. *Conidiogenous cells* hyaline, subcylindrical, filiform, straight, tapering towards the apex, collarette flared, periclinal thickening prominent, (10–)11–13(–14) × (1.5–)2(–3) µm. *Alpha conidia* hyaline, fusoid, apex acute, base obtusely rounded to subtruncate, bi- to multi-guttulate, (9–)11–13(–14) × 3(–4) µm. *Beta and gamma conidia* not observed.




Fig. 12 *Diaporthe neoarctii* (CBS 109490). a. Conidiomata sporulating on PDA; b, c. transverse section through conidiomata, showing conidiomatal wall; d–f. conidiogenous cells; g. alpha conidia. — Scale bars: b = 225 µm, all others = 10 µm.

Culture characteristics — Colonies with sparse aerial mycelium, covering the dish after 2 wk in the dark at 25 °C. On MEA umber with patches of greyish sepia, umber in reverse. On PDA fuscous-black on surface and in reverse.

Specimen examined. USA, New Jersey, isolated from *Ambrosia trifida*, May 2001, G. Bills (holotype CBS H-21094, ex-type culture CBS 109490 = GB 6421 = AR 3450).

Notes — Isolates originally identified as *D. arctii* cluster in clades 19 and 67 (Fig. 1). *Diaporthe neoarctii* (clade 16) was isolated from *Ambrosia trifida* in New Jersey, USA, and differs morphologically from the ex-type culture of *D. arctii* (alpha conidia 7 × 3–3.5 µm) (clade 19). Based on these differences *D. neoarctii* is described as a novel species.

Diaporthe nobilis complex

Specimens examined. GERMANY, Münster, on stem of *Laurus nobilis*, Feb. 1939, Kotthoff (CBS 200.39). — JAPAN, isolate from *Pinus pentaphylla* bonsai plant imported from Japan into the Netherlands, May 1979, G.H. Boerema (CBS H-16732, culture CBS 587.79). — KOREA, on imported chestnuts (*Castanea sativa*), collected in grocery store in Sydney, Australia, 5 July 1999, K.A. Seifert (CBS 113470 = DAOM 226800). — LATVIA, *Rhododendron* sp., *I. Apine* (CBS 129167). — NEW ZEALAND, on bark of *Malus pumila*, G.J. Samuels (CBS 124030 = GJS 77-49); Waikato region, on *Pyrus pyrifolia*, 2001, isol. W. Kandula, det. L. Castlebury (CBS 116953 = NZ-26, CBS 116954 = NZ-27). — YUGOSLAVIA, on *Hedera helix*, July 1989, M. Muntañola-Cvetkovic (CBS 338.89).

Notes — Clade 62 is poorly resolved in this dataset, but has some internal structure, suggesting that it contains several potentially distinct species. More isolates would be required to resolve their taxonomy. Isolates in this clade were originally identified as *Phomopsis fukushii* (on *Pyrus pyrifolia*, New Zealand), *P. conorum* (on *Pinus pentaphylla*, the Netherlands), *P. castanea* (on *Castanea sativa*, UK), *Diaporthe perniciosa* (*Malus pumila*, New Zealand), *D. pulla* (on *Hedera helix*, Yugoslavia) and *D. nobilis* (on *Laurus nobilis*, Germany).

Diaporthe nomurai Hara, in Hara, Diseases of cultivated plants: 140. 1925. — Fig. 13

Conidiomata in culture on OA sporulating poorly, globose, up to 300 µm diam, black, erumpent; cream conidial droplets exuding from central ostioles; walls consisting of 3–6 layers of medium brown *textura angularis*. *Conidiophores* hyaline, smooth, 0–1-septate, rarely branched, densely aggregated, cylindrical, straight to sinuous, 10–20 × 2–3 µm. *Conidiogenous cells* 6–10 × 1.5–3 µm, phialidic, cylindrical, terminal, with slight taper towards apex, 1–1.5 µm diam, with visible periclinal thickening; collarette not flared, minute. *Paraphyses* not observed. *Alpha conidia* aseptate, hyaline, smooth, guttulate, fusoid-ellipsoid to clavate, straight to variously curved, tapering towards both ends, straight, apex subobtuse, base truncate, (7–)9–11(–13) × (2.5–)3 µm. *Gamma conidia* not observed. *Beta conidia* spindle-




Fig. 13 *Diaporthe nomurai* (CBS 157.29). a. Conidiomata sporulating on PDA; b–e. conidiogenous cells; f. alpha conidia; g. beta conidia. — Scale bars = 10 µm.

shaped, aseptate, smooth, hyaline, apex acutely rounded, base truncate, tapering from lower third towards apex, gently curved, (20–)25–27(–30) × 1.5(–2) µm.

Culture characteristics — Colonies reaching up to 8 cm diam after 2 wk in the dark at 25 °C. On MEA surface isabelline, reverse sepia. On OA surface pale mouse grey with concentric rings of mouse grey; reverse mouse grey. On PDA surface and reverse fuscous-black, with sparse aerial mycelium.

Specimen examined. JAPAN, on *Morus* sp., Dec. 1929, K. Togashi (CBS 157.29).

Notes — Clade 58 represents *D. nomurai* from *Morus* sp. in Japan. *Diaporthe nomurai* is known from hosts such as *Morus alba*, *M. bombycina*, *M. latifolia* and *Morus* sp. (Farr & Rossman 2012).

Diaporthe novem J.M. Santos, Vrandečić & A.J.L. Phillips, Persoonia 27: 14. 2011

= *Phomopsis* sp. 9 van Rensburg et al., Stud. Mycol. 55: 65. 2006.

Specimens examined. BRAZIL, endophytic in petiole on *Maytenus ilicifolia*, July 2007, R.R. Gomes (LGMF 943 = CPC 20319). — CROATIA, Slavonija, in seed on *Glycine max*, Sept. 2008, T. Duvnjak (holotype CBS H-20462, ex-type cultures CBS 127270 = 4-27/3-1, CBS 127271 = 5/27/3-3, CBS 127269 = 5-27/3-1). — ROMANIA, Calugareni, Distr. Mizil, living leaves on *Polygonatum odoratum*, 31 July 1970, O. Constantinescu (CBS 354.71).

Notes — Clade 22 represents *D. novem* (Santos et al. 2011), and contains an endophytic isolate (LGMF 43) from *Maytenus ilicifolia*, one isolate previously identified as *Diaporthe pardalota* on *Polygonatum odoratum* from Romania, and three isolates of *D. novem* which includes the ex-type isolate. Isolate LGMF 943 represents higher genetic variation than the other isolates, and appears to represent a different species. Since this isolate did not sporulate, further morphological characterisation was not possible and we refrain from excluding it from the species pending collection of more strains to clarify its status.

Diaporthe novem was reported as pathogen of *Aspalathus linearis* (van Rensburg et al. 2006) as *Phomopsis* sp. 9. It was recently described as pathogen of *Glycine max* (Santos et al. 2011). This species was also reported on *Hydrangea macrophylla* (Santos et al. 2010), *Helianthus annuus* and *Vitis vinifera*

(Santos et al. 2011). It is known to occur in Brazil, Romania, Croatia, Italy (Rekab et al. 2004), Portugal (Santos et al. 2010) and South Africa (van Niekerk et al. 2005, van Rensburg et al. 2006).

Diaporthe oncostoma (Duby) Fuckel, Jahrb. Nassauischen Vereins Naturk. 23–24: 205. 1870. (1869–1870). — Fig. 14

Basionym. *Sphaeria oncostoma* Duby, in Rabenh., Klotzsch. Herb. Vivum Mycol.: no. 253. 1854.

Conidiomata pycnidial, globose to ellipsoidal, aggregated as well as scattered, dark brown to black, immersed, ostiolate, 430–1170 µm wide, 370–790 µm tall, lacking necks, with outer surface covered in brown hyphae; pycnidial wall consisting of brown, thick-walled cells of *textura angularis*; conidial mass globose or exuding in cirri, white to pale luteous or pale yellow. *Conidiophores* hyaline, subcylindrical, branched above the septa, tapering towards the apex, 1–2-septate, (10–)11–19(–22) × 3(–4) µm. *Conidiogenous cells* hyaline, subcylindrical, straight or curved, tapering towards the apex, collarette not flared, periclinal thickening prominent, (6–)7–9(–10) × (2–)3 µm. *Alpha conidia* hyaline, fusoid to ellipsoidal, straight to slightly curved, acute at apex, subobtuse at base, bi- or multi-guttulate, (7.5–)9–11(–12) × (2–)3(–4) µm. *Gamma conidia* hyaline, smooth, ellipsoid-fusoid, apex acutely rounded, and tapering towards truncate base, (11–)12–16 × 3(–3.5) µm. *Beta conidia* and sexual morph not observed in culture.

Culture characteristics — Colonies covering dish after 2 wk in the dark at 25 °C. On MEA surface dirty white with profuse aerial mycelium, reverse umber. On OA surface dirty white with patches of umber, same in reverse. On PDA surface and reverse sienna, with sparse aerial mycelium.

Specimens examined. FRANCE, Hte Savoie, Aigueblanche-Bellecombe, outlet of river Morel in Isère, on dead branches of *Robinia pseudoacacia*, 17 July 1978, H.A. van der Aa (CBS 589.78). — GERMANY, Wolfenbüttel, on leaf spot of *Robinia pseudoacacia*, 15 Nov. 1996, H. Butin (CBS 100454); Berlin, on leaf of *Ilex aquifolium*, Nov. 1985, M. Hesse (CBS 809.85). — RUSSIA, on *Robinia pseudoacacia*, June 2000, L. Vasilyeva (CBS 109741 = AR 3445).

Notes — *Diaporthe oncostoma* (clade 70) has been considered to be a saprobic, or low virulence pathogen, which plays some role in natural pruning and self-thinning of black locust




Fig. 14 *Diaporthe oncostoma* (CBS 100454). a. Conidiomata sporulating on OA; b, c. transverse section through conidiomata, showing conidiomatal wall; d–f. conidiogenous cells; g. alpha conidia. — Scale bars: b = 225 µm, all others = 10 µm.

forests (*Robinia pseudoacacia*) (Vajna 2002). However, this fungus has been reported as a causal agent of canker and severe dieback disease of black locust in Russia (Scerbin-Parfenenko 1953) and in Greece (Michalopoulos-Skarmoutsos & Skarmoutsos 1999).

Although isolate CBS 809.85 was obtained from *Ilex aquifolium* in Germany, we treat it as belonging to *D. oncostoma*, as it matches the other strains phylogenetically as well as morphologically.

***Diaporthe oxe* R.R. Gomes, C. Glienke & Crous, sp. nov. —**
Mycobank MB802941; Fig. 15

Etymology. The word 'oxe' is an expression used in northeastern Brazil that means amazement or surprise, in relation to the number of novel species isolated as endophytes from medicinal plants in Brazil.

Conidiomata pycnidial ampulliform to finger-like, eustromatic, convoluted to unilocular, semi-immersed, scattered, dark brown to black, ostiolate, 60–170 µm wide, 60–220 µm tall; necks variable in length, 20–150 µm, outer surface covered with hyphae; pycnidial wall consisting of brown, thick-walled cells of *textura angularis*; conidial mass globose or exuding in cirri, pale-luteous to cream or pale-yellow. *Conidiophores* hyaline, ampulliform to subcylindrical, branched above the septa, tapering towards the apex, 1–2-septate, (14–)17–25(–27) × (2–)3 µm. *Conidiogenous cells* hyaline, subcylindrical, filiform, straight to curved, tapering towards the apex, collarette flared, periclinal thickening prominent, (5–)6–10(–12) × 2(–3) µm. *Alpha conidia* hyaline, oblong to ellipsoid, apex bluntly rounded, base obtuse to subtruncate, bi- to multi-guttulate, (5–)6–7(–8) × (2–)3 µm. *Beta conidia* hyaline, smooth, curved or hamate, (17–)22–30(–33) × 2–3 µm. *Gamma conidia* not observed.

Culture characteristics — Colonies on PDA flat, with an entire edge, surface mycelium dense and felty, ochreous to fulvous, dark brick, honey, buff, exudates rarely present as colourless drops; colonies reaching 49 mm diam after 2 wk at 25 °C; reverse umber, ochreous to fulvous. On OA flat, with an entire edge, surface mycelium dense and felty, rosy buff, pale olivaceous-grey, iron-grey, with patches olivaceous buff, exudates in colourless and pale luteous drops; colonies reaching 40 mm diam; reverse dark brick, olivaceous. On MEA raised, with an entire edge, surface mycelium dense and felty, buff,

rosy-buff, with chestnut coloured exudates in the centre of the colony, and pale luteous at the periphery; colonies reaching 49 mm diam; reverse chestnut and bay.

Specimens examined. BRAZIL, on petiole of *Maytenus ilicifolia*, July 2007, R.R. Gomes (holotype CBS H-21098, ex-type culture CBS 133186 = LGMF 942 = CPC 20318); same collection details (CBS 133187 = LGMF 936 = CPC 20312); on leaf of *Schinus terebinthifolius*, July 2007, J. Lima (LGMF 915 = CPC 20291); on petiole of *M. ilicifolia*, S.A.V. Pileggi (LGMF 939 = CPC 20315); on petiole of *M. ilicifolia*, July 2007, R.R. Gomes (LGMF 945 = CPC 20321).

Notes — Endophytic isolates (clade 34) from medicinal plants in Brazil.

***Diaporthe padi* var. *padi* G.H. Otth, Mitt. Naturf. Ges. Bern: 99. 1871 (1870)**

Specimens examined. SWEDEN, Uppland, Dalby par., Tuna, on *Prunus padus*, 17 Apr. 1988, K. & L. Holm (CBS 114200 = UPSC 2569); Dalarna, Folkärna par., Sonnbo, on *Alnus glutinosa*, Dec. 1992, K. & L. Holm (CBS 114649 = UPSC 3496).

Notes — *Diaporthe padi* var. *padi* (clade 56) represents a European taxon occurring on *Prunus*. We chose the name *D. padi* over *D. decorticans*, as the basionym of the latter, *Sphaeria decorticans*, is an illegitimate homonym.

***Diaporthe paranensis* R.R. Gomes, C. Glienke & Crous, sp. nov. —**
Mycobank MB802942, Fig. 16

Etymology. Named after Paraná, the state in Brazil from where it was collected.

Conidiomata pycnidial, ampulliform, semi-immersed, scattered, brown to black, ostiolate, 130–220 µm wide, 60–130 µm tall; prominent necks 50–210 µm long, outer surface smooth or covered in hyphae; pycnidial wall consisting of brown, thick-walled cells of *textura angularis*; conidial mass globose, predominantly pale-luteous to yellow and some cases green-olivaceous. *Conidiophores* hyaline, subcylindrical to cylindrical, filiform, branched above the septa on a globose cell, not tapering towards the apex, 2–3-septate, (14–)15–22(–26) × (2–)3(–4) µm. *Conidiogenous cells* hyaline, subcylindrical, filiform, rarely tapering towards the apex, collarette present and flared, slight periclinal thickening, (5–)8–14(–15) × 2(–3) µm.




Fig. 15 *Diaporthe oxe* (CBS 133186). a. Conidiomata sporulating on PDA; b, c. transverse section through conidiomata, showing conidiomatal wall; d, e. conidiogenous cells; f. beta conidia; g. alpha conidia. — Scale bars: b = 100 µm, all others = 10 µm.




Fig. 16 *Diaporthe paranensis* (CBS 133184). a. Conidiomata sporulating on PDA; b, c. transverse section through conidiomata, showing conidiomatal wall; d, e. conidiogenous cells; f. alpha and beta conidia. — Scale bars: b = 100 µm, all others = 10 µm.

Alpha conidia hyaline, fusoid-ellipsoidal, apex bluntly rounded, base obtuse to subtruncate, bi- to multi-guttulate, (6–)7–8(–9) × (2–)3 µm. *Beta conidia* hyaline, smooth, curved or hamate and slightly curved, (16–)17–21(–23) × (1–)2 µm. *Gamma conidia* not observed.

Culture characteristics — Colonies on PDA flat, with an entire edge, mycelium growing in concentric rings, cottony texture, white to smoke-grey; colonies reaching up to 64 mm diam after 2 wk at 25 °C; reverse buff and isabelline. On OA flat, with an entire edge, aerial mycelium in concentric rings, ranging in colour from smoke-grey to grey-olivaceous and white in the centre; colonies reaching 44 mm diam; reverse iron-grey, grey-olivaceous to olivaceous-buff. On MEA flat, with an entire edge, aerial mycelium growing in concentric rings, with cottony texture, pale olivaceous-grey to grey-olivaceous and buff; colonies reaching 56 mm diam; reverse umber, fulvous with patches of greenish black.

Specimen examined. BRAZIL, Paraná, Colombo, endophytic species isolated from petiole of *Maytenus ilicifolia* (popular name Espinheira Santa), July 2007, R.R. Gomes (holotype CBS H-21099, ex-type culture CBS 133184 = LGMF 929 = CPC 20305).

Notes — Endophytic isolate (clade 35) from medicinal plant in Brazil.

Diaporthe perjuncta Niessl, Hedwigia 15: 153. 1876

Specimen examined. AUSTRIA, from *Ulmus glabra*, Oct. 2001, A.Y. Rossman (ex-epitype culture CBS 109745 = ARSEF 3461 = AR 3461).

Notes — *Diaporthe perjuncta* (clade 40) is associated with fallen branches of *Ulmus campestris* and *U. glabra* (*Ulmaceae*). This species is found in Austria, Germany and Portugal. *Diaporthe perjuncta* is distinguished from *D. viticola* and *D. australafricana* based on morphology and DNA sequence data (van Niekerk et al. 2005). Pathogenicity studies and endophytic isolation of '*D. perjuncta*' from grapevines in Australia and South Africa in fact represent isolates of *D. australafricana* (Mostert et al. 2001a, Rawnsley et al. 2004, van Niekerk et al. 2005).

Diaporthe perseae (Zerova) R.R. Gomes, C. Glienke & Crous, comb. nov. — MycoBank MB802944; Fig. 17

Basionym. *Phomopsis perseae* Zerova, J. Bot. Acad. Sci. RSS Ukraine 1, 1–2: 307. 1940.

Conidiomata pycnidial in culture on MEA, globose, up to 400 µm diam, black, erumpent; cream conidial droplets exuding from central ostioles; walls consisting of 3–6 layers of medium brown *textura angularis*. Conidiophores hyaline, smooth, 1–3-septate, branched, densely aggregated, cylindrical, straight to sinuous, 15–35 × 3–4 µm. Conidiogenous cells 8–17 × 1.5–2.5 µm,




Fig. 17 *Diaporthe perseae* (CBS 151.73). a. Conidiomata sporulating on PDA; b–d. conidiogenous cells; e. alpha and beta conidia. — Scale bars = 10 µm.

phialidic, cylindrical, terminal and lateral, with slight taper towards apex, 1–1.5 µm diam, with visible periclinal thickening; collarette prominent, up to 5 µm long. Paraphyses hyaline, smooth, subcylindrical with obtuse ends, 2–4-septate, up to 60 µm long, 3 µm diam. Alpha conidia aseptate, hyaline, smooth, guttulate, fusoid to ellipsoid, tapering towards both ends, straight, apex subobtuse, base subtruncate, (6–)7–8(–9) × 2(–2.5) µm. Gamma conidia aseptate, hyaline, smooth, ellipsoid-fusoid, apex acutely rounded, base subtruncate, 9–14 × 1.5–2 µm. Beta conidia spindle-shaped, aseptate, smooth, hyaline, apex acutely rounded, base truncate, tapering from lower third towards apex, curved, (15–)22–25(–28) × 1.5(–2) µm.

Culture characteristics — Colonies covering dish after 2 wk in the dark at 25 °C, with moderate aerial mycelium. On OA surface ochreous, with patches of dirty white and iron-grey. On PDA surface dirty white with patches of sienna, reverse sienna with patches of umber. On MEA surface sienna, with patches of umber, reverse umber with patches of sienna.

Specimen examined. NETHERLANDS ANTILLES, Martinique, on young fruit of *Persea gratissima*, 10 July 1972, E. Laville (CBS 151.73).

Notes — *Diaporthe perseae* (clade 86) was originally described from branches of dying *Persea gratissima* trees in Russia. Based on the morphology (alpha conidia 7–10.2 × 2.3–2.5 µm; Uecker 1988), this strain could be authentic for the name.

Diaporthe phaseolorum (Cooke & Ellis) Sacc., Syll. Fung. 1: 692. 1882

Basionym. *Sphaeria phaseolorum* Cooke & Ellis, Grevillea 6, 39: 93. 1878.

Specimens examined. BRAZIL, endophytic in petiole on *Maytenus ilicifolia*, July 2007, R.R. Gomes (LGMF 927 = CPC 20303, LGMF 941 = CPC 20317). — NEW ZEALAND, from *Olearia cf. rani*, 22 Jan. 2003, G.J.M. Verkley (CBS 113425); *Actinidia chinensis*, rotting fruit, kiwifruit orchard, S.R. Pennycook (CBS 127465 = GJS 83-379). — UNKNOWN, Apr. 1980, L. Beraha (CBS 257.80). — USA, Mississippi, from *Caperonia palustris*, Oct. 2003, A. Mengistu (CBS 116019); Mississippi, from *Aster exilis*, Oct. 2003, A. Mengistu (CBS 116020).

Notes — Clade 4 represents isolates of *D. phaseolorum*. It includes two endophytic isolates from *Maytenus ilicifolia* collected in Brazil, one isolate previously misidentified as *D. melonis* (CBS 257.80), two isolates respectively from *Caperonia palustris* and *Aster exilis* in the USA (Mengistu et al. 2007), one isolate from *Olearia cf. rami*, and one from *Actinidia chinensis*. The ITS and TEF1 sequences of this clade are similar to sequences (GenBank U11323, U11373 and EU222020, respectively) of a well-characterised isolate of *D. phaseolorum* (ATCC 64802 = FAU458). By accepting this clade as authentic for

D. phaseolorum, we follow the precedent set by van Rensburg et al. (2006), Mengistu et al. (2007) and Santos et al. (2011).

Diaporthe pseudomangiferae R.R. Gomes, C. Glienke & Crous, sp. nov. — MycoBank MB802945; Fig. 18

Etymology. Named after its morphological similarity to *Phomopsis mangiferae*.

Conidiomata pycnidial, erumpent to superficial on PDA, globose, up to 300 µm diam with elongated necks with central ostioles that exude yellow-orange to cream conidial droplets; walls of 6–8 layers of brown *textura angularis*. Conidiophores hyaline, smooth, 1–3-septate, branched, densely aggregated, cylindrical, straight to sinuous, 20–30 × 2–2.5 µm. Conidiogenous cells phialidic, cylindrical, terminal and lateral with slight apical taper, 10–15 × 2–3 µm; collarette flared, up to 3 µm long. Paraphyses hyaline, smooth, cylindrical, septate, extending above conidiophores, straight to flexuous, unbranched or branched below, up to 80 µm long, 2–3 µm wide at base. Alpha conidia aseptate, hyaline, smooth, guttulate to granular, fusiform, tapering towards both ends, apex acutely rounded, base truncate, (6–)7–9(–10) × (2–)2.5(–3) µm. Beta and gamma conidia not seen (description based on CBS 101339).

Culture characteristics — Colonies covering dish after 2 wk in the dark at 25 °C, with moderate aerial mycelium. On OA surface and reverse dirty white with patches of iron-grey. On PDA surface dirty white to ochreous, reverse umber. On MEA surface greyish sepia with patches of iron-grey, reverse greyish sepia with patches of iron-grey.

Specimens examined. DOMINICAN REPUBLIC, from *Mangifera indica*, P. de Leeuw, ATO-DLO, Wageningen (holotype CBS H-21105, culture ex-type CBS 101339). — MEXICO, on fruit peel of *Mangifera indica* (CBS 388.89).

Notes — Although these isolates (clade 82) were originally described as representative of *P. mangiferae* (dead leaves of *Mangifera indica*, Pakistan), they differ in having larger conidiomata, longer conidiophores and larger alpha conidia.

Diaporthe pseudophoenicicola R.R. Gomes, C. Glienke & Crous, sp. nov. — MycoBank MB803839; Fig. 19

Etymology. Named after its morphological similarity to *Diaporthe phoenicicola*.

Conidiomata pycnidial on MEA, up to 400 µm diam, erumpent, globose with neck; ostiole exuding yellow-orange conidial droplets; walls consisting of 3–6 layers of medium brown *textura angularis*. Conidiophores hyaline, smooth, densely aggregated, 1–3-septate, branched, cylindrical, straight to curved, 12–45 × 1.5–3 µm. Conidiogenous cells phialidic, cylindrical, terminal




Fig. 18 *Diaporthe pseudomangiferae* (CBS 101339). a. Conidiomata sporulating on PNA; b. conidiomata sporulating on PDA; c, d. conidiogenous cells; e. beta conidia; f. alpha conidia. — Scale bars = 10 µm.



Fig. 19 *Diaporthe pseudophoenicicola* (CBS 462.69). a, b. Conidiomata sporulating on PDA; c, d. conidiogenous cells; e. alpha conidia. — Scale bars = 10 µm.

and lateral with slight apical taper, 12–20 × 1.5–2 µm, with visible periclinal thickening; collarette flared, 2–5 µm long. Paraphyses hyaline, smooth, cylindrical, 1–3-septate, extending above conidiophores, straight to flexuous, unbranched or branched, up to 100 µm long, and 3 µm wide at base. Alpha conidia aseptate, hyaline, granular, smooth, fusiform, tapering towards both ends, straight, acutely rounded apex, and truncate base, (6–)7–8(–9) × (2–)2.5(–3) µm. Beta and gamma conidia not seen (description based on CBS 462.69).

Culture characteristics — Colonies covering dish after 2 wk in the dark at 25 °C, with sparse aerial mycelium. On MEA surface dirty white with patches of sienna, reverse umber with patches of sienna. On OA surface dirty white with patches of sienna. On PDA surface ochreous with patches of olivaceous-grey, reverse iron-grey with patches of ochreous.

Specimens examined. IRAQ, Prov. Basrah, Shalt El Arab, showing dieback on *Mangifera indica*, 1976, M.S.A. Al-Momen (CBS 176.77). — SPAIN, Mallorca, Can Pastilla, dead tops of green leaves on *Phoenix dactylifera*, 27 May 1969, H.A. van der Aa (holotype CBS H-21106, culture ex-type CBS 462.69).

Notes — *Diaporthe pseudophoenicicola* (clade 89) is distinct from *D. phoenicicola* (conidia 8–12 × 2–2.5 µm; Uecker 1988) by having shorter, and wider alpha conidia. A similar strain was isolated from *Mangifera indica* in Iraq (CBS 176.77), suggesting that this species has a wider host range.

Diaporthe pustulata Sacc., Syll. Fung. (Abellini) 1: 610. 1882

Specimens examined. AUSTRIA, on *Acer pseudoplatanus*, Oct. 2001, A.Y. Rossman (CBS 109742 = AR 3430 and CBS 109760 = AR 3535); Raab, Au Wald, on *Prunus padus*, Oct. 2001, A. Y. Rossman (CBS 109784 = AR 3419).

Notes — Clade 44 contains one isolate from *Prunus padus* and two isolates from *Acer pseudoplatanus*, all isolated from Austria. Clade 56 contains another isolate on *Prunus padus* from Sweden. Clearly there are two different species from *Prunus*, one isolated in Austria and another in Sweden. Because *D. pustulata* was originally described on *Acer pseudoplatanus*, we tentatively apply this name to isolates in clade 44. To clarify the status of isolates in clades 44 and 56, however, additional isolates and a comparison with type materials would be required.

Diaporthe raonikayaporum R.R. Gomes, C. Glienke & Crous, sp. nov. — MycoBank MB802947; Fig. 20

Etymology. Raoni + Kayapo = after the name of a leader (Raoni) of the indigenous Kayapo ethnic tribe in Brazil. The Kayapos are inhabitants of the Amazon region in Brazil. They use the medicinal plant *Spondias mombin*, from which this species was isolated, as adornment or ornament, and for its medicinal properties.

Conidiomata pycnidial, globose to conical or ampulliform, eustromatic and convoluted or unilocular, scattered, dark brown to black, immersed, ostiolate, 110–200 µm wide, 50–130 µm tall,




Fig. 20 *Diaporthe raonikayaporum* (CBS 133182). a. Conidiomata sporulating on PNA; b, c. transverse section through conidiomata, showing conidiomatal wall; d. conidiogenous cells; e. beta with a few alpha conidia; f. alpha conidia. — Scale bars: b = 100 µm, all others = 10 µm.

with prominent necks 40–140 µm long, outer surface smooth or covered in hyphae; pycnidial wall consisting of brown, thick-walled cells of *textura angularis*; conidial mass globose or exuding in cirrhi, white to pale-luteous. *Conidiophores* hyaline, ampulliform to subcylindrical, filiform, branched above the septa, tapering towards the apex, 1–3-septate, (16–)17–22(–26) × (2–)3 µm. *Conidiogenous cells* hyaline, subcylindrical, filiform, straight to curved, tapering towards the apex, collarette not flared, periclinal thickening prominent, (5–)7–9(–10) × (2–)3 µm. *Alpha* and *gamma* conidia are formed in the same conidiogenous cells. *Alpha* conidia hyaline, oblong to ellipsoid, apex bluntly rounded, base obtuse to subtruncate, bi- to multi-guttulate, (6–)7(–8) × (2–)3 µm. *Beta* conidia not observed. *Gamma* conidia hyaline, fusoid to subcylindrical, slightly curved, apex bluntly rounded, base obtuse to subtruncate, bi- to multi-guttulate, or eguttulate, (7–)9–11(–13) × (1–)2 µm.

Culture characteristics — Colonies on PDA flat, with an entire edge, aerial mycelium forming concentric rings with cottony texture, olivaceous-buff, isabelline to honey on surface; colonies reaching 63 mm diam after 2 wk at 25 °C; reverse pale purplish grey to smoke-grey. On OA flat, with an entire edge, aerial mycelium forming concentric rings, white, olivaceous on surface, colonies reaching 31 mm diam; reverse buff and greenish olivaceous. On MEA flat, with a lobate edge, aerial mycelium forming woolly concentric rings, olivaceous-grey, greenish olivaceous and patches of amber on surface, colonies reaching 51 mm diam; reverse brown-vinaceous.

Specimen examined. BRAZIL, Pará, Redenção, endophytic species isolated from leaf of *Spondias mombin* (popular name Cajazeira and Taperebá), July 2007, K. Rodriguez (holotype CBS H-21097, ex-type culture CBS 133182 = LGMF 923 = CPC 20299).

Notes — Endophytic isolate (clade 31) from medicinal plant in Brazil.

Diaporthe rhoina Feltgen, Vorstud. Pilzfl. Luxemb., Nachtr. III: 145. 1903

Specimen examined. UNKNOWN, on *Rhus toxicodendron*, Sept. 1927, L.E. Wehmeyer (CBS 146.27).

Notes — This species (clade 95) was originally described on *Rhus typhina* from Luxembourg. European isolates of this

pathogen will need to be collected to confirm the identity of CBS 146.27, which is presumably of North American origin.

Diaporthe saccarata (J.C. Kang, L. Mostert & Crous) Crous, comb. nov. — MB802948

Basionym. *Phomopsis saccarata* J.C. Kang, L. Mostert & Crous, Sydowia 53, 2: 230. 2001.

Specimen examined. SOUTH AFRICA, Western Cape Province, Jonkershoek Mountains, Stellenbosch, on cankers of *Protea repens*, Mar. 1999, S. Denman (ex-type culture CBS 116311 = CPC 3743).

Note — *Diaporthe saccarata* (clade 71) is known to cause a canker disease on shoots of *Protea repens* in South Africa (Mostert et al. 2001b).

Diaporthe schini R.R. Gomes, C. Glienke & Crous, sp. nov. — MycoBank MB802949; Fig. 21

Etymology. Named after the host genus from which it was isolated, *Schinus*.

Conidiomata pycnidial, eustromatic, multilocular, immersed to erumpent, ostiolate, dark brown to black, scattered or aggregated, 80–270 µm wide, 70–240 µm tall, prominent necks 70–220 µm long, outer surface covered with hyphae; pycnidial wall consisting of brown, thick-walled cells of *textura angularis*; conidial mass globose, pale-luteous to cream. *Conidiophores* hyaline, subcylindrical, filiform, rarely branched, tapering towards the apex, 0–1-septate, (11–)12–17(–20) × (2–)3(–4) µm. *Conidiogenous cells* hyaline, subcylindrical and filiform, straight, tapering towards the apex, collarette not observed, with prominent periclinal thickening 5–6(–7) × (1–)2 µm. *Beta* conidia hyaline, smooth, curved or hamate (14–)22–28(–30) × (1–)2 µm. *Alpha* and *gamma* conidia not observed.

Culture characteristics — Colonies on PDA flat, with a lobate margin, surface mycelium sparse, felty and appressed, buff, honey to isabelline; colonies reaching 30 mm diam after 2 wk at 25 °C; reverse greyish sepia, smoke-grey. On OA with a lobate margin, surface mycelium flat, sparse, felty and appressed, smoke-grey, olivaceous-grey, or olivaceous buff; colonies reaching 21 mm diam; reverse pale mouse-grey to



Fig. 21 *Diaporthe schini* (CBS 133181). a. Conidiomata sporulating on PDA; b, c. transverse section through conidiomata, showing conidiomatal wall; d, e. conidiogenous cells; f. beta conidia. — Scale bars: b = 135 µm, all others = 10 µm.

olivaceous-grey or buff. On MEA with a lobate margin, surface mycelium flat, dense, felty and appressed, buff with umber patches; colonies reaching 30 mm diam; reverse dark mouse-grey, umber, with patches of isabelline or luteous.

Specimen examined. BRAZIL, Paraná, Curitiba, endophytic species isolated from leaf of *Schinus terebinthifolius* (popular name Aroeira), July 2007, J. Lima (holotype CBS H-21093, culture ex-type CBS 133181 = LGMF 921 = CPC 20297); same collection details (LGMF 910).

Notes — Other than *D. schini* (clade 11), additional endophytic isolates were also obtained from *Schinus terebinthifolius* in Brazil, but these are morphologically different and cluster in clades 5 and 9 (*D. endophytica* and *D. terebinthifolii*).

***Diaporthe sclerotiooides* (Kesteren) Udayanga, Crous & K.D. Hyde, Fung. Diversity 56: 166. 2012**

Basionym. *Phomopsis sclerotiooides* Kesteren, Neth. Jl. Pl. Path. 73: 115. 1967.

Specimens examined. NETHERLANDS, Maarssen, on root of *Cucumis sativus*, June 1967, H.A. van der Kesteren (ex-type culture CBS 296.67 = ATCC 18585 = IMI 151828 = PD 68/690); Roermond, on root of *C. sativus*, Dec. 1976 (CBS 710.76 = PD 76/674).

Notes — *Diaporthe sclerotiooides* (clade 28) was originally described from roots of *Cucumis sativus* in the Netherlands. This species has subsequently been reported to cause black root rot of *Citrullus lanatus*, *Cucurbita sativus*, *C. ficifolia*, *C. maxima* and *C. moschata* in various countries in the world (Udayanga et al. 2011).

***Diaporthe scobina* Nitschke, Pyrenomycetes Germanici 2: 293. 1870**

= *Phomopsis scobina* Höhn., Sber. Akad. Wiss. Wien, Math.-naturw. Kl., Abt. 1 115: 681 (33 of repr.). 1906.

Specimen examined. SCOTLAND, living and dead twig of *Fraxinus excelsior*, Feb. 1938, J.A. MacDonald (CBS 251.38).

Notes — Clade 38 is represented by *D. scobina* isolated from *Fraxinus excelsior* in Scotland. The fungus is known on this host from Scotland and Poland (Mulenko et al. 2008, Farr & Rossman 2012).

***Diaporthe sojae* Lehman, Ann. Missouri Bot. Gard. 10: 128. 1923**

≡ *Diaporthe phaseolorum* var. *sojae* (Lehman) Wehm., The genus Diaporthe Nitschke and its segregates 47: 1933.

= *Phomopsis longicolla* Hobbs, Mycologia 77: 542. 1985.

≡ *Diaporthe longicolla* (Hobbs) J.M. Santos, Vrandečić & A.J.L. Phillips, Persoonia 27: 13. 2011.

Specimens examined. CROATIA, on *Glycine max* stem, Sept. 2005, K. Vrandečić (specimen CBS H-20460, culture CBS 127267). – ITALY, Bologna, from *Glycine soja*, 1986, P. Giunchi (specimen CBS H-16776, culture CBS 100.87). – UNKNOWN, on *Glycine soja* ('Blackhawk') mature stem, A.A. Hildebrand (CBS 180.55 = ATCC 12050 = CECT 2024). – USA, Mississippi, from *Euphorbia nutans*, A. Mengistu (CBS 116017 = DP 0508) and from *Glycine max*, Oct. 2003, A. Mengistu (CBS 116023); on *Glycine soja* seedling, J. Marcinkowska (CBS 659.78 = NRRL 13656).

Notes — Isolates of *D. phaseolorum* var. *sojae* clustered in two distinct clades (clade 1, *D. sojae*; clade 7, *D. melonis*). *Diaporthe sojae* causes pod and stem blight of soybean, while *P. longicolla* is known to cause seed decay (Santos et al. 2011). Several authors have found it difficult to distinguish them based on disease symptoms alone, and usually report them together (Almeida & Seixas 2010). Hobbs et al. (1985) described *P. longicolla* as a different species to *D. sojae* (*Diaporthe phaseolorum* var. *sojae*) based on morphological characters. Both symptom types, however, have also been linked to the same species

(Kulik 1984, Morgan-Jones 1989, Kulik & Sinclair 1999). Considering their genetic similarity based on the five genes studied here, disease etiology and common host, it appears that these isolates belong to the same species, which is distinct from *D. phaseolorum* (clade 4). *Diaporthe sojae* (clade 1) is an older name than *D. longicolla*, and is therefore applied to this clade.

***Diaporthe* sp. 1**

Specimens examined. BRAZIL, EMBRAPA/PR, on *Glycine max* seed, A. Almeida (LGMF 947 = CPC 20323). – GERMANY, Bielefeld, human abscess, K. Plechulla (CBS 119639 = B 11861).

Notes — Isolates from clade 2 appear to represent a novel species, *Diaporthe* sp. 1 (sterile). It is represented by CBS 119639, isolated from an abscess of a male patient in Germany, and isolate LGMF 947, obtained from soybean seeds in Brazil. Isolates from this clade share a low genetic homology to isolates of the clade 4 (*D. phaseolorum*; Fig. 1, part 1).

Diaporthe species commonly described from soybean were also reported as opportunistic human pathogens. In 1999, a species of *Phomopsis* was reported as etiological agent of a subcutaneous infection on the finger of an immunosuppressed farmer and this genus was added to the list of fungi capable to cause human disease (Sutton et al. 1999). In 2011, *D. sojae* (as *Phomopsis longicolla*), a known pathogen of soybean, was identified as causing skin infection in an immunocompromised patient after kidney transplantation. The authors believed that this patient acquired the fungus at least 5 yr before, when he had contact with seeds or soybean plants in Equatorial Guinea (Garcia-Reyne et al. 2011).

Another phytopathogenic species also described in soybean, *Diaporthe phaseolorum*, was reported causing osteomyelitis in patients with positive serology for human lymphotropic virus type 1 (HTLV-1), disturbing the immune response. The patient was a farmer and inoculation occurred possibly through injury with *Amaranthus spinosus* thorns (Iriart et al. 2011).

***Diaporthe* sp. 2**

Specimen examined. BRAZIL, on petiole of *Maytenus ilicifolia*, July 2007, R.R. Gomes (LGMF 932 = CPC 20308).

Notes — Sterile, endophytic isolate from medicinal plant in Brazil, which appears to represent a novel species (clade 29).

***Diaporthe* sp. 3**

Specimen examined. SCOTLAND, on *Pseudotsuga menziesii*, Mar. 1929, G.G. Hahn (CBS 287.29).

Notes — Clade 32 was tentatively named *Diaporthe* sp. 3, and is represented by a single isolate previously identified as *Phomopsis conorum*, and obtained from *Pseudotsuga menziesii* in Scotland. This clade was not resolved, because there are at least eight different conifer species without any ex-type cultures (Udayanga et al. 2011).

***Diaporthe* sp. 4**

Specimen examined. BRAZIL, endophytic in petiole on *Maytenus ilicifolia*, July 2007, R.R. Gomes (LGMF 944 = CPC 20320).

Notes — Sterile endophytic isolate (clade 33) from a medicinal plant in Brazil, appearing to represent an undescribed species.

Diaporthe sp. 5

Specimen examined. ITALY, from *Acer opalus*, W. Jaklitsch (CBS 125575).

Notes — This isolate (clade 37) represents a novel species occurring on *Acer*, which will be treated separately as part of another study (W. Jaklitsch, pers comm.).

Diaporthe sp. 6

Specimens examined. HONG KONG, University Drive, on fruit of *Maesa perlarius*, 18 Dec. 2000, K.D. Hyde (CBS 115595 = HKUCC 10129, CBS 115584 = HKUCC 7784).

Notes — The two strains (clade 85) studied here were originally identified as *P. pittospori* (described from *Pittosporum* twigs, USA, California), which seems highly unlikely, as they were isolated from fruit of *Maesa perlarius* in Hong Kong. Unfortunately both strains proved to be sterile, so their identity could not be confirmed.

Diaporthe sp. 7

Specimen examined. INDIA, Bangalore, on *Anacardium occidentale*, Aug. 1978, H.C. Govindu (CBS 458.78).

Notes — The identity of the present isolate (identified as *Phomopsis anacardii*) could not be confirmed, as the culture proved to be sterile. However, phylogenetically (clade 88) it represents a distinct taxon from *D. anacardii* (clade 69), and when recollected, should be described as new.

Diaporthe sp. 8

Specimen examined. BRAZIL, from *Aspidosperma tomentosum*, K. Rodriguez (LGMF 925 = CPC 20301).

Culture characteristics — Colonies covering dish after 2 wk in the dark at 25 °C, with moderate aerial mycelium. On PDA surface ochreous, reverse pale luteous. On OA surface and reverse luteous. On MEA surface pale luteous, reverse orange to apricot.

Notes — Although this isolate (clade 90) appears to represent an undescribed species based on phylogenetic data, it proved to be sterile. As we presently only have a single strain of this taxon, its treatment will have to await further collections.

Diaporthe stictica (Berk. & Broome) R.R. Gomes, C. Glienke & Crous, comb. nov. — MycoBank MB802950

Basionym. *Phoma stictica* Berk. & Broome, Ann. Mag. Nat. Hist., ser. II 5: 370. 1850.

≡ *Phomopsis stictica* (Berk. & Broome) Traverso, Fl. Ital. Crypt. 2, 1: 276. 1906.

Specimen examined. ITALY, Perugia, on dead twig of *Buxus sempervirens*, Dec. 1954, M. Ribaldi (CBS 370.54).

Notes — *Diaporthe stictica* (clade 74) represents a European species occurring on *Buxus sempervirens* (Italy, Germany). Although the present isolate could be authentic for the name, this could not be confirmed based on morphology, as the isolate proved to be sterile.

Diaporthe subordinaria (Desm.) R.R. Gomes, C. Glienke & Crous, comb. nov. — MycoBank MB802951

Basionym. *Phoma subordinaria* Desm., Ann. Sci. Nat., Bot. ser. 3, 9: 284. 1849.

≡ *Phomopsis subordinaria* (Desm.) Traverso, Fl. Ital. Crypt. Pars 1: Fungi. Pyrenomycetidae. Xylariaceae, Valsaceae, Ceratostomataceae: 232. 1906.

Specimens examined. NEW ZEALAND, blackened seed of *Plantago lanceolata*, Apr. 1999, B. Alexander (CBS 101711). — SOUTH AFRICA, Eastern Cape Province, Grahamstown, on stalks of *Plantago lanceolata*, 2 Dec. 1989, R. Shivas (CBS 464.90).

Notes — *Diaporthe subordinaria* (clade 18) has a global distribution on *Plantago lanceolata*, on which it causes a stalk disease (de Nooit & van der Aa 1987). It is possible that the disease relates to several different species occurring on *Plantago*, but this matter can only be resolved following further collections and correlation with type material.

Diaporthe tecomaie Sacc. & P. Syd., Syll. Fung. 14: 550. 1899 (nom. nov. for *D. interrupta* Niessl). — Fig. 22

?= *Phoma tecomaie* Sacc., Nuovo Giorn. Bot. Ital. 8: 201. 1876.

≡ *Phomopsis tecomaie* (Sacc.) Traverso & Spessa, Bol. Soc. Brot. Coimbra, sér. 1, 25: 124. 1910.

Conidiomata pycnidial, sporulating poorly on OA, globose, up to 1 mm diam, black, erumpent, multilocular; cream conidial droplets exuding from central ostioles; walls consisting of 3–6 layers of medium brown *textura angularis*. *Conidiophores* hyaline in upper region, pale brown at base, smooth, 1–3-septate, branched, densely aggregated, cylindrical, straight to sinuous, 20–30 × 2–3 µm. *Conidiogenous cells* 8–15 × 1.5–3 µm, phialidic, cylindrical, terminal and lateral, with slight taper towards apex, 1 µm diam, with visible periclinal thickening; collarette not flared, minute. *Paraphyses* not observed. *Beta conidia* spindle-shaped, aseptate, smooth, hyaline, apex acutely rounded, base truncate, tapering from lower third towards apex, apex strongly curved, (17–)22–24(–26) × 1.5(–2) µm. *Alpha* and *gamma* conidia not observed.

Culture characteristics — Colonies covering dish after 2 wk in the dark at 25 °C. On OA fluffy, dirty white with patches of grey olivaceous. On PDA dirty white with patches of olivaceous grey and isabelline, reverse with patches of dirty white, brown vinaceous and dark brick. On MEA dirty white with patches of isabelline and olivaceous grey, reverse brown vinaceous with patches of dark brick.




Fig. 22 *Diaporthe tecomaie* (CBS 100547). a. Conidiomata forming on PNA; b, c. conidiogenous cells; d. beta conidia. — Scale bars = 10 µm.

Specimen examined. BRAZIL, São Paulo, Serra da Mantiqueira, mycocecidium caused by *Prosopodium tecomicola* on living young branch of *Tabebuia* sp., 27 Sept. 1997, coll. A. Aptroot, isol. H.A. van der Aa (specimen CBS H-16834, culture CBS 100547).

Notes — *Diaporthe tecomaiae* was a new name proposed for *D. interrupta* Niessl (on *Tecoma radicans*, Portugal), as the epithet was already occupied. The link between the *Diaporthe* and *Phomopsis* state remains to be proven. The asexual morph was originally described as *Phoma tecomaiae* (from Italy on *Tecoma radicans*, conidiophores 20 × 1 µm, conidia 8 × 3 µm; Saccardo 1878), and is probably distinct from the fungus represented by CBS 100547, which occurs on *Tabebuia* sp. in Brazil. However, as no ex-type strains are available of *D. tecomaiae* (clade 10), and no alpha conidia were observed in culture, this could not be confirmed, and is pending fresh collections.

Diaporthe terebinthifoliae R.R. Gomes, C. Glienke & Crous, sp. nov. — MycoBank MB802952; Fig. 23

Etymology. Named after the host species from which it was isolated, *Schinus terebinthifolius*.

Conidiomata pycnidial, globose to conical, immersed, ostiolate, brown to black, scattered or aggregated, 95–110 µm wide, 140–160 µm tall, rarely forms necks, but when present, they are short and covered with hyphae; pycnidial wall consisting of brown, thick-walled cells of *textura angularis*; conidial mass globose, white or pale-luteous to cream. **Conidiophores** hyaline, subcylindrical, filiform, branched above septa, tapering towards the apex, 1–2-septate, (13–)15–21(–22) × 2(–3) µm. **Beta conidiogenous cells** hyaline, ampulliform to subcylindrical and filiform, tapering towards the apex, collarette present and not flared, slight periclinal thickening, (3–)6–10(–14) × 2(–3) µm. **Beta conidia** hyaline, smooth, curved or hamate, (18–)20–24(–26) × 1(–2) µm. **Alfa and gamma conidia** not observed.

Culture characteristics — Colonies on PDA flat, with an entire edge, aerial mycelium cottony, greyish white, colonies reaching 64 mm diam after 2 wk in the dark at 25 °C; reverse buff. On OA flat, entire edge, aerial mycelium cottony, with concentric rings, pale olivaceous-grey, smoke-grey and greyish white, colonies

reaching 48 mm diam; reverse olivaceous-grey and olivaceous buff. On MEA flat, with an entire edge; aerial mycelium cottony, smoke-grey, colonies reaching 60 mm diam; reverse umber with patches of fuscous-black.

Specimens examined. BRAZIL, Paraná, Curitiba, endophytic species isolated from leaf of *Schinus terebinthifolius* (popular name Aroeira), July 2007, J. Lima (holotype CBS H-21097, ex-type culture CBS 133180 = LGMF 914 = CPC 20290); same collection details (LGMF 909 = CPC 20285, LGMF 907 = CPC 20283, LGMF 913 = CPC 20289).

Notes — The multigene analysis of isolates in clade 9 exhibited insignificant homology to sequences found in GenBank. An isolate previously identified as *Phomopsis tecomaiae* (CBS 100547) also resides in this clade, but is morphologically distinct. No morphologically similar isolates are known from *S. terebinthifolius*, and thus we designate these isolates as representative of a new taxon.

Diaporthe toxica P.M. Will., Hight, W. Gams & Sivasith., Mycol. Res. 98: 1367. 1994

Specimens examined. WESTERN AUSTRALIA, Morawa, on stem of *Lupinus angustifolius*, 6 May 1991, J.B. Nunn (ex-type culture CBS 534.93 = ATCC 96741); Serpentine, on *Lupinus* sp., 8 June 1993, P.M. Williamson (CBS 535.93); Medina, on *Lupinus* sp., 8 June 1993, P.M. Williamson (CBS 546.93).

Notes — Clade 45 contains three isolates of *D. toxica*, including the ex-type culture (CBS 534.93), isolated from *Lupinus angustifolius* in Western Australia. Two varieties of *Phomopsis* (*P. leptostromiformis* var. *leptostromiformis* and *P. leptostromiformis* var. *occidentalis*) were identified as causing disease in *Lupinus* sp. *Diaporthe woodii* was later recognised as the sexual state of *P. leptostromiformis* var. *occidentalis* (Punithalingam 1974), while Williamson et al. (1994) designated the name *D. toxica* for the sexual state of the toxicogenic variety, *P. leptostromiformis* var. *leptostromiformis*.

Lupins (*Lupinus* spp.) are grown in many parts of the world as a grain legume crop. The seeds are used for animal feed and increasingly as flour for human consumption. The plants increase soil nitrogen and are grown in rotation with other crops.




Fig. 23 *Diaporthe terebinthifoliae* (CBS 133180). a. Conidiomata sporulating on PNA; b, c. transverse section through conidiomata, showing conidiomatal wall; d, e. conidiogenous cells; f. beta conidia. — Scale bars: b = 100 µm, c = 25 µm, all others = 10 µm.

In Australia the stubble left after harvesting aids soil conservation and is a valuable summer feed for livestock. *Diaporthe toxica* is considered to be an important limiting factor to more extensive sowing of lupins. This organism has been reported to cause stem blight in young lupins (*Lupinus luteus*) (Ostazeski & Wells 1960) and to produce phomopsins (Culvenor et al. 1977). These mycotoxins cause the animal liver disease known as lupinosis (Gardiner 1975, Allen & Wood 1979).

Diaporthe vaccinii Shear, U.S. Dept. Agric. Tech. Bull. 258: 7. 1931

= *Phomopsis vaccinii* Shear, U.S. Dept. Agric. Tech. Bull. 258: 7. 1931.

Specimens examined. USA, Massachusetts, on *Oxycoccus macrocarpos*, Mar. 1932, C.L. Shear (ex-type culture CBS 160.32 = IFO 32646); Michigan, on *Vaccinium corymbosum*, G.C. Adams (CBS 118571); New Jersey, on *V. macrocarpon*, 1988, L. Carris (CBS 122112 = FAU 474); Michigan, on *V. corymbosum*, 1992, D.C. Ramsdell (CBS 122114 = FAU 634, CBS 122115 = FAU 590); North Carolina, from *V. corymbosum*, pre-1999, D.F. Farr (CBS 122116 = DF 5022).

Notes — Clade 63 consists of six isolates of *D. vaccinii*, including the ex-type strain (CBS 160.32) isolated on *Vaccinium corymbosum* from the USA. *Diaporthe vaccinii* causes fruit rot and twig blight and leaf spots of *Vaccinium* spp. (blueberries) in the USA (Alfieri et al. 1984, Farr et al. 2002, Farr & Rossman 2012). The principal hosts are American and European cranberries (*Vaccinium macrocarpon*, *V. oxycoccus*, *V. oxycoccus* var. *intermedium*), highbush blueberry (*V. corymbosum*) and rabbiteye blueberry (*V. Ashei*). *Diaporthe vaccinii* is restricted to cultivated *Vaccinium* species. The wild European species, *V. oxycoccus*, which usually occurs in mountain bogs, could be a potential reservoir for the pest. In the EPPO region it has been reported from Romania (found in experimental plots of introduced American cultivars, but did not establish (Teodorescu et al. 1985)), UK (found in plants originally imported from the Netherlands and USA, but did not establish (Wilcox & Falconer 1961, Baker 1972)).

Symptoms in susceptible blueberry cultivars include blighting of 1-yr-old woody stems with flower buds. Infected succulent, current-year shoots wilt in 4 d and become covered with minute lesions. The fungus continues to travel downward through the stem, killing major branches, and often entire plants (Wilcox 1939, Daykin & Milholland 1990). Infected fruits turn reddish-brown, soft, mushy, often splitting and causing leakage of juice (Milholland & Daykin 1983).

Diaporthe vexans (Sacc. & P. Syd.) Gratz, Phytopathology 32: 542. 1942

Basionym. *Phoma vexans* Sacc. & P. Syd., Syll. Fung. (Abellini) 14, 2: 889. 1899.

≡ *Phomopsis vexans* (Sacc. & P. Syd.) Harter, J. Agric. Res. 2, 5: 338. 1914.

Specimen examined. USA, from *Solanum melongena*, Dec. 1914, L.L. Harter (CBS 127.14).

Notes — *Diaporthe vexans* (clade 12) causes fruit rot, leaf spot, stem and tip blight disease of eggplants (*Solanum melongena* and *S. wendlandii*) and other solanaceous species, *Acacia* sp. (Fabaceae), *Prunus* sp. (Rosaceae) and *Sorghum bicolor* (Poaceae), *Capsicum annuum* and *Lycopersicon esculentum* (Solanaceae). The disease is widespread in North America, the West Indies, and Eastern and Central Asia, also in Africa (Senegal, Tanzania, Zambia) and Mauritius (Punithalingam & Holliday 1972). Additional records include Brunei, Haiti, Iran, Iraq and Romania (Harter 1914, Farr & Rossman 2012).

Diaporthe viticola Nitschke, Pyrenomyctes Germanici 2: 264. 1870

Specimens examined. AUSTRIA, Vienna, Risenbergbach-Weg, on *Laburnum anagyroides*, May 2001, A.R. Rossman (CBS 109492). — CANADA, British Columbia, Sidney, on *Epilobium angustifolium*, Oct. 2001, M. Barr (CBS 109768 = AR 3478). — FRANCE, Dordogne, near Sarlat la Canéda, 1-yr-old stems on *Asphodelus albus*, 20 May 1995, G. Verkley (CBS 759.95). — NETHERLANDS, Utrecht, Baarn, in branches and twigs of *Aucuba japonica*, Jan. 1995, G. Verkley (CBS 106.95); on *Rosa rugosa*, 18 Mar. 1985, G.H. Boerema (CBS 266.85 = PD 85/25); Lelystad, in dead stem on *Lupinus* sp., May 1982, H.A. van der Aa (CBS 449.82); Wieringermeer, Robbenoordbos, in dead stem on *Lupinus arboreus*, 12 Mar. 1991, H.A. van der Aa & F. Meurs (CBS 312.91); Flevoland, trees in front of Info Centre Lepelaarsplassen, in leaf spot on *Fraxinus excelsior*, 31 Aug. 1997, H.A. van der Aa (CBS 100170); Baarn, in dead stem on *Dipsacus fullonum*, 14 June 1985, H.A. van der Aa (CBS 502.85); on twig on *Salix* sp., Apr. 1962, G.H. Boerema (CBS 446.62). — PORTUGAL, on *Vitis vinifera* (Galego durado), 1 Jan. 1998, A.J.L. Phillips (CBS 114011 = CPC 2677); Burgaes, Santo Tirso, on *Vitis vinifera*, 16 Feb. 1998, A.J.L. Phillips (ex-type culture CBS 113201 = CPC 5683). — SWEDEN, Skåne, Maglehem par., on *Sambucus* cf. *racemosa*, 14 Apr. 1989, K. Holm & L. Holm (CBS 114436 = UPSC 2960). — UK, Sheffield, on *A. japonica*, July 1996, G. Verkley (CBS 794.96).

Notes — *Diaporthe viticola* (clade 50) is known from several hosts, but especially from grapevines, on which it causes a cane spot disease in Europe (Portugal, Germany). Merrin et al. (1995) referred to several Australian isolates from grapevines as *Phomopsis* taxon 1. The same species was reported by Phillips (1999) as *D. perjuncta* and by Schepers et al. (2000) as *D. viticola*. In a subsequent study, Mostert et al. (2001a) chose to follow Phillips (1999) and applied the name *D. perjuncta* to taxon 1. However, they also noted that minor morphological differences existed in perithecia and ascospores between the European and Southern Hemisphere material, which led to the description of a novel taxon, *D. australafricana*, for isolates from Australia and South Africa (van Niekerk et al. 2005), and the epitypification of *D. viticola* based on European material. Based on the results obtained here, *D. viticola* (clade 50) is closely related to *D. australafricana* (clade 49), and is clearly distinguishable from *D. perjuncta* (clade 40).

Diaporthe woodii Punith., Mycol. Pap. 136: 51. 1974

= *Phomopsis leptostromiformis* var. *occidentalis*, R.G. Shivas, J.G. Allen & P.M. Will., Mycol. Res. 95: 322. 1991.

Specimen examined. WESTERN AUSTRALIA, Medina, stems of *Lupinus* sp., 8 July 1993, P.M. Williamson (CBS H-5319, culture CBS 558.93).

Notes — Clade 94 represents *Diaporthe woodii* (CBS 558.93), which was characterised by Williamson et al. (1994), based on the ex-type strain (IMI 166508). *Diaporthe crotalariae* (clade 92), *D. aspalathi* (clade 93) and *D. woodii* are closely related species. *Diaporthe woodii* causes stem rot, stem cankers, leaf infections and seed decay of *Lupinus angustifolius* and *L. cosenfinii*, and blight and seed discoloration of *L. albus*, *L. angustifolius*, *L. cosentinii*, *L. luteus*, *L. pilosus* and *Trifolium subterraneum* (subterranean clover). The fungus is known to occur in Brazil, South Africa, USA (Florida), and Western Australia (Williamson et al. 1994).

Diaporthe woolworthii (Peck) Sacc., Syll. Fung. (Abellini) 1: 615. 1882

Basionym. *Valsa woolworthii* Peck, Ann. Rep. N.Y. State Mus. Nat. Hist. 28: 73. 1876 (1875).

Specimen examined. UNKNOWN, on *Ulmus americana*, Sept. 1927, L.E. Wehmeyer (CBS 148.27).

Notes — Clade 57 contains a single isolate of *D. woolworthii* from *Ulmus americana*. This taxon represents an American species occurring on *Ulmus*, so this culture (presumably from North America), could prove to be authentic for the name.

DISCUSSION

A major aim of the present study was to resolve the taxonomy of *Diaporthe* species occurring on diverse hosts, either as pathogens, saprobes, or as harmless endophytes. To delimitate these taxa, nine genes were screened, from which the best five were selected to conduct a multi-gene phylogenetic analysis (ITS, TEF1, ACT, HIS and CAL). *Diaporthe* represents a highly complex genus containing numerous cryptic species, several of which are newly described in the present study, while others remain unclear, awaiting fresh collections and type studies. Many *Diaporthe* species that are morphologically similar proved to be genetically distinct, and several isolates that were formerly identified based on their host, were shown to represent different taxa.

Although the genera *Diaporthe* and *Phomopsis* have received much taxonomic attention, few phylogenetic studies have thus far been conducted, and hence the taxonomy of this group is still problematic. Due to the lack of reference strains, and the fact that few gene loci other than ITS have in the past been used for DNA analysis, most of the conclusions reached thus far have been incorrect, meaning that published literature will have to be interpreted with care.

In this study we studied 15 endophytic *Diaporthe* species from Brazil. Three were not identified to species level, two were identified as *D. novem* and *D. phaseolorum*, while a further 10 were described as new. High genetic diversity was found amongst the analysed isolates from medicinal plants. Species of *Diaporthe* are commonly isolated as endophytes from several hosts in temperate and tropical regions (Bussaban et al. 2001, Murali et al. 2006, Rossman et al. 2007, Botella & Diez 2011, González & Tello 2011). Skaltsas et al. (2011) isolated 108 *Diaporthe* isolates from asymptomatic leaves and bark of three different hosts (*Hevea brasiliensis*, *H. guianensis* and *Micandra* spp.) from Cameroon, Mexico and Peru. Using a multigene approach, the authors found more than 40 phylogenetic species, of which several appeared to represent novel taxa (Skaltsas et al. 2011).

Despite members of *Diaporthe* commonly being described as phytopathogenic, an increasing number of reports link this genus to endophytic studies, focusing on its potential as a producer of enzymes and novel secondary metabolites, with antibiotic, fungicide and anticancer activity (Dai et al. 2005, Elsaesser et al. 2005, Lin et al. 2005, Silva et al. 2005, Wu et al. 2008, Kumaran & Hur 2009, Weber 2009, Vesterlund et al. 2011).

The ecology of species of *Diaporthe* remains poorly understood, as some endophytes isolated from the medicinal plant *Maytenus ilicifolia* were identified as *D. phaseolorum* (clade 4) and *D. novem* (clade 22), respectively known as pathogen of soybean (Santos et al. 2011) and *Aspalathus linearis* (van Rensburg et al. 2006). *Diaporthe novem* is also reported from hosts such as *Hydrangea macrophylla* (Santos et al. 2010), *Helianthus annuus* and *Vitis vinifera* (Santos et al. 2011). These reports agree with the pogo stick hypothesis, postulating that host-specific fungal plant pathogens frequently exhibit the ability to colonise non-host tissue, enabling them to disperse further, in an attempt to find the host on which they are pathogenic (Crous & Groenewald 2005).

The taxonomy of *Diaporthe* (incl. *Phomopsis*) has traditionally been based on host association, with species being described on the assumption that they are host-specific. In the present study the taxonomy of all *Diaporthe* isolates deposited in the CBS culture collection over time were reviewed, based on this assumption. The employment of this criterion, has led to an exponential growth in the number of taxa described in *Diaporthe* thus far (Uecker 1988). However, in spite of the

apparent synonymies outlined in this study, there was evidence for a huge proliferation of cryptic taxa that were formerly overlooked based on a morphological approach in the absence of molecular data. Species delimitation in *Diaporthe* based on morphological characters is challenging, as most taxa in culture do not produce all spore states of the asexual (alpha, beta and gamma conidia) or the sexual morph. The description of novel taxa in *Diaporthe* in the absence of molecular data (at least ITS and HIS or TUB; see discussion in next section) should thus be strongly discouraged in the future.

In conclusion thus, it seems that in spite of the fact that these taxa readily colonise or co-colonise non-hosts (see also Rehner & Uecker 1994, Mostert et al. 2001a, Farr et al. 2002, Diogo et al. 2010), there is still a multitude of undescribed taxa awaiting further study in this complex. It is thus hoped that the phylogenetic backbone generated here provides a stable platform to enable future studies by others interested in the biology of *Diaporthe*.

Phylogenetic species recognition by genealogical concordance

Taylor et al. (2000) developed the Genealogical Concordance Phylogenetic Species Recognition (GCPSR) concept to define the limits of sexual species, using the phylogenetic concordance of multiple unlinked genes. This concept has proved greatly useful in fungi, because it is more finely discriminating than other species concepts, as several species are unable to be crossed, or cannot be recognised due to the lack of distinguishing morphological characters or sterility (Reynolds 1993, Taylor et al. 2000, Cai et al. 2011). The adoption of genealogical concordance for species recognition in *Diaporthe* enabled us to distinguish species that were otherwise not possible to identify due to either sterility, or the loss of specific character states. For instance, *D. viticola* and *D. australafricana* are two closely related species (clades 50 and 49, respectively) associated with grapevines. They are morphologically similar, but occur on different continents (van Niekerk et al. 2005). These species have probably accumulated genetic differences due to their geographical isolation. Several cryptic species were recently described in other genera using the GCPSR criterion, some of which are consistent with allopatric divergence, because these species occupy non-overlapping areas separated by geographic barriers, e.g. in *Cladosporium* (Bensch et al. 2012), *Colletotrichum* (Damm et al. 2012a, b), *Harknessia* (Crous et al. 2012), *Ilyonectria* (Cabral et al. 2012a, b) and *Phyllosticta* (Glienke et al. 2011), to name but a few. Using the GCPSR concept it is possible to define the genetic variation observed in some species, but still insufficient to establish them as distinct species, since genetic flow still occurs between them. For example, isolates of clades 79–90 clustered differently based on analyses of the different genes, probably because of recent gene flow among them.

We have compared the location and monophyly of the strains in each clade in the phylogenetic tree of the combined alignment (Fig. 1) to those phylogenetic trees obtained from the individual loci to determine the species boundaries and species resolution. The five loci selected for the Bayesian phylogeny have a similar resolution for species discrimination, ranging from TEF1 resolving 72 out of the 95 species, to HIS and TUB resolving 84 of the 95 species.

The ITS region, which is often considered to be less than optimal for closely related species, was not much better or worse (resolving 75 of the 95 species) than the other included loci. However, given the recent acceptance of the ITS region as official fungal barcode (Schoch et al. 2012) and its intermediate resolving power in the present study, this locus should not

be discarded from future studies. Also, TEF1, which has in the past been used as additional locus for phylogenetic studies of *Diaporthe*, performed the worst in this study (resolving 72 of the 95 species), although this was not much worse than ITS and CAL (resolving 75 and 74 of the 95 species, respectively). The HIS and TUB regions appear to have the best resolution for species discrimination in the present study and therefore are good candidates as secondary markers to the commonly used ITS region. Similar results were also reported for ITS, CAL, TEF1 and TUB by Udayanga et al. (2012), who suggested that TUB be considered as secondary phylogenetic marker for *Diaporthe*.

The importance of epitypification in *Diaporthe*

The best option to supplement poor type material is via epitypification (Cannon et al. 2012). To employ the GCPSR concept in fungi, DNA is mostly extracted from poorly preserved, ancient herbarium specimens with difficulty, and in many cases it only results in short sequences of the ITS region (Quaedvlieg et al. 2011, Cheewangkoon et al. 2012). Therefore, epitypification of living material, and its preservation and deposit in publically available collections and databases, are important steps to provide a stable platform to enable others to test future hypotheses. Although it is not a prerequisite, it is strongly recommended that the chosen epitype should originate from the same geographical region and host, and have morphological, cultural and pathological characteristics similar to those described in the original publication (see Damm et al. 2012a, b, Cannon et al. 2012, Weir et al. 2012).

Despite the fact that close to 2 000 species of *Diaporthe* (incl. *Phomopsis*) have been described in literature, hardly any ex-type strains are available today, the majority of which were included in the present study. Due to the lack of ex-type strains, the taxonomy of several species continue to be unresolved, some of which are important plant pathogens. A serious effort will thus be called for to recollect and redescribe all these old names. An alternative approach would be to simply start over, ensuring that all newly described names are based not only on morphology, but also supplemented by DNA barcodes. However, as long as fungal nomenclature is governed by the ICN, this seems unobtainable. Eventually though, all mycologists will realise that a stable fungal nomenclature must incorporate DNA data, and that this is only achievable if mycology follows a code of nomenclature that incorporates this requirement.

Acknowledgements We thank the technical staff, Arien van Iperen (cultures), Marjan Vermaas (photographic plates) and Mieke Starink-Willems (DNA isolation, amplification and sequencing) for their invaluable assistance. We are grateful to the Brazilian agency CAPES for financial support to Renata R. Gomes.

REFERENCES

- Alfieri JR, Langdon KR, Wehlburg C, Kimbrough JW. 1984. Index of plant diseases in Florida (Revised). Florida Department of Agriculture and Consumer Services, Division of Plant Industry Bulletin 11: 1–389.
- Allen JG, Wood PM. 1979. The prevention of lupinosis by making lupin hay. Australian Veterinary Journal 55: 38–39.
- Almeida AMR, Seixas CDS (eds). 2010. Soyabeans: root and stem diseases and interrelationships with soil management and culture. Embrapa Soja, Londrina.
- Anderson RG, Hartman JR. 1983. Phomopsis twig blight on weeping figs indoors: a case study. Foliage Digest 6: 38–58.
- Arnold RH. 1975. *Diaporthe alleghaniensis*. Fungi Canadenses 70: 1–2.
- Ash GJ, Stodart B, Sakuanrungsirikul S, Anschaw E, Crump N, et al. 2010. Genetic characterization of a novel *Phomopsis* sp., a putative biocontrol agent for *Carthamus lanatus*. Mycologia 102: 54–61.
- Baker JJ. 1972. Report on diseases of cultivated plants in England and Wales for the years 1957–1968. Ministry of Agriculture, Fisheries and Food Technical Bulletin 25: 148.
- Bandre TR, Sasek V. 1977. Antibiotic activity of pyrenomyctes under submerged conditions. Folia Microbiologica 22: 269–274.
- Banishehmi Z, Javadi AR. 2009. Further investigations on the biology of *Phomopsis cinerascens*, the cause of fig canker in Iran. Phytopathologia Mediterranea 48: 454–460.
- Battilani P, Rossi V, Girometta B, Delos M, Rouzet J, et al. 2003. Estimating the potential development of *Diaporthe helianthi* epidemics in Italy. OEPP/EPPO Bulletin 33: 427–431.
- Bensch K, Braun U, Groenewald JZ, Crous PW. 2012. The genus Cladosporium. Studies in Mycology 72: 1–401.
- Benschop K, Tewari JP, Toop EW. 1984. Phomopsis twig die-back of some woody interior ornamentals in Alberta. Canadian Plant Disease Survey 64: 29–31.
- Beraha LR, O'Brien MJ. 1979. *Diaporthe melonis* sp. nov., a new soft rot of market cantaloupes. Phytopathologische Zeitschrift 94: 199–207.
- Boddy L, Griffith GS. 1989. Role of endophytes and latent invasion in the development of decay communities in sapwood of angiospermous trees. Sydowia 41: 41–73.
- Botella L, Diez JJ. 2011. Phylogenetic diversity of fungal endophytes in Spanish stands of *Pinus halepensis*. Fungal Diversity 47: 9–18.
- Brayford D. 1990. Variation in *Phomopsis* isolates from *Ulmus* species in the British Isles and Italy. Mycology Research 94: 691–697.
- Bussaban B, Lumyong L, Lumyong P, McKenzie EHC, Hyde KD. 2001. Endophytic fungi from *Amomum siamense*. Canadian Journal of Microbiology 47: 943–948.
- Cabral A, Groenewald JZ, Rego C, Oliveira H, Crous PW. 2012a. Cylindrocarpon root rot: multi-gene analysis reveals novel species within the *Ilyonectria radicicola* species complex. Mycological Progress 11: 655–688.
- Cabral A, Rego C, Nascimento T, Oliveira H, Groenewald JZ, Crous PW. 2012b. Multi-gene analysis and morphology reveal novel *Ilyonectria* species associated with black foot disease of grapevines. Fungal Biology 116: 62–80.
- Cai L, Giraud T, Zhang N, Begerow D, Cai G, Shivas RG. 2011. The evolution of species concepts and species recognition criteria in plant pathogenic fungi. Fungal Diversity 50: 121–133.
- Cannon PF, Damm U, Johnston PR, Weir BS. 2012. Colletotrichum – current status and future directions. Studies in Mycology 73: 181–213.
- Carbone I, Kohn LM. 1999. A method for designing primer sets for speciation studies in filamentous ascomycetes. Mycologia 91: 553–556.
- Carroll GC. 1986. The biology of endophytism in plants with particular reference to woody perennials. In: Fokkema NJ, Heuvel J van den (eds), Microbiology of the phyllosphere: 205–222. Cambridge University Press, Cambridge.
- Castlebury LA, Farr DF, Rossman AY, Jaklitsch WJ. 2003. *Diaporthe angelicae* comb. nov., a modern description and placement of *Diaporthopsis* in *Diaporthe*. Mycoscience 44: 203–208.
- Castlebury LA, Mengistu A. 2006. Phylogenetic distinction of *Diaporthe/ Phomopsis* isolates from soybeans. Systematic Mycology and Microbiology 57: 1–13.
- Castlebury LA, Rossman AY, Jaklitsch WJ, Vasilyeva LN. 2002. A preliminary overview of the Diaporthales based on large subunit nuclear ribosomal DNA sequences. Mycologia 94: 1017–1031.
- Cheewangkoon R, Groenewald JZ, Hyde KD, To-anun C, Crous PW. 2012. Chocolate spot of Eucalyptus. Mycological Progress 11: 61–69.
- Cooke MC. 1894. Answers to correspondents. Gardeners' Chronicle 15: 605.
- Crous PW, Braun U, Hunter GC, Wingfield MJ, Verkley GJM, et al. 2013. Phylogenetic lineages in *Pseudocercospora*. Studies in Mycology 75: 37–114.
- Crous PW, Gams W, Stalpers JA, Robert V, Stegehuis G. 2004a. MycoBank: an online initiative to launch mycology into the 21st century. Studies in Mycology 50: 19–22.
- Crous PW, Groenewald JZ. 2005. Hosts, species and genotypes: opinions versus data. Australasian Plant Pathology 34: 463–470.
- Crous PW, Groenewald JZ, Risede J-M, Hywel-Jones NL. 2004b. *Calonectria* species and their *Cylindrocladium* anamorphs: species with sphaeroendoculate vesicles. Studies in Mycology 50: 415–430.
- Crous PW, Groenewald JZ, Shivas RG, Edwards J, Seifert KA, et al. 2011. Fungal Planet description sheets: 69–91. Persoonia 26: 108–156.
- Crous PW, Phillips AJL, Baxter AP. 2000. Phytopathogenic fungi from South Africa. University of Stellenbosch, Department of Plant Pathology Press.
- Crous PW, Schoch CL, Hyde KD, Wood AR, Gueidan C, et al. 2009a. Phylogenetic lineages in the Capnodiales. Studies in Mycology 64: 17–47.
- Crous PW, Summerell BA, Shivas RG, Carnegie AJ, Groenewald JZ. 2012. A re-appraisal of *Harknessia* (Diaporthales), and the introduction of *Harknessiaceae* fam. nov. Persoonia 28: 49–65.
- Crous PW, Verkley GJM, Groenewald JZ, Samson RA (eds). 2009b. Fungal Biodiversity. CBS Laboratory Manual Series 1: 1–269. Centraalbureau voor Schimmelcultures, Utrecht, Netherlands.

- Culvenor CCJ, Beck AB, Clarke M, Cockrum PA, Edgar JA, et al. 1977. Isolation of toxic metabolites of *Phomopsis leptostromiformis* responsible for lupinosis. *Australian Journal of Biological Sciences* 30: 269–277.
- Dai J, Krohn K, Floerke U, Gehle D, Aust HJ, et al. 2005. Novel highly substituted diaryl ethers, phomopsines D-G, isolated from endophytic fungus *Phomopsis* sp. from *Adenocarpus foliolosus*. *European Journal of Organic Chemistry* 23: 5100–5105.
- Damm U, Cannon PF, Woudenberg JHC, Baroncelli R, Crous PW. 2012a. The *Colletotrichum acutatum* species complex. *Studies in Mycology* 73: 1–36.
- Damm U, Cannon PF, Woudenberg JHC, Johnston PR, Weir BS, et al. 2012b. The *Colletotrichum boninense* species complex. *Studies in Mycology* 73: 1–36.
- Daykin ME, Milholland RD. 1990. Histopathology of blueberry twig blight caused by *Phomopsis vaccinii*. *Phytopathology* 80: 736–740.
- Debaeke P, Estragnat A, Reau R. 2003. Influence of crop management on sunflower stem canker (*Diaporthe helianthi*). *Agronomie* 23: 581–592.
- Delacroix G. 1905. Sur une maladie des Amendiers en Provence. *Bulletin de la Société Mycologique de France* 21: 180–185.
- Dennis RWG. 1986. Fungi of the Hebrides. Royal Botanic Gardens, Kew.
- Dettrakul S, Kittakoop P, Isaka M, Nopichai S, Suyarnsestakorn C, et al. 2003. Antimycobacterial pimarane diterpenes from the fungus *Diaporthe* sp. *Bioorganic & Medicinal Chemistry Letters* 7: 1253–1255.
- Diogo ELF, Santos JM, Phillips AJL. 2010. Phylogeny, morphology and pathogenicity of *Diaporthe* and *Phomopsis* species on almond in Portugal. *Fungal Diversity* 44: 107–115.
- Drummond AJ, Ashton B, Buxton S, Cheung M, Cooper A, et al. 2011. Geneious v5.4, Available from <http://www.geneious.com/>.
- Early MP, Punithalingam E. 1972. *Phomopsis anacardii* sp. nov. on *Anacardium occidentale*. *Transactions of British Mycological Society* 59: 345–347.
- Elsaesser B, Krohn K, Floerke U, Root N, Aust HJ, et al. 2005. X-ray structure determination absolute configuration and biological activity of phomoxanthone. *European Journal of Organic Chemistry* 21: 4563–4570.
- Farr DF, Bills GF, Chamuris GP, Rossman AY. 1989. Fungi on plants and plant products in the United States. American Phytopathological Society, St. Paul, Minnesota.
- Farr DF, Castlebury LA, Rossman AY, Putnam ML. 2002. A new species of *Phomopsis* causing twig dieback of *Vaccinium vitis-idaea* (lingonberry). *Mycological Research* 106: 745–752.
- Farr DF, Rossman AY. 2012. Fungal databases, systematic mycology and microbiology laboratory, ARS, USDA. Retrieved December, 2012, from <http://nt.ars-grin.gov/fungal databases/>.
- Fernández FA, Hanlin RT. 1996. Morphological and RAPD analyses of *Diaporthe phaseolorum* from soybean. *Mycologia* 88: 425–440.
- Fröhlich J, Hyde KD, Guest DI. 1997. Fungi associated with leaf spots of palms in north Queensland, Australia. *Mycological Research* 101: 721–732.
- Garcia-Reyne A, López-Medrano F, Morales JM, Esteban CG, Martín I, et al. 2011. Cutaneous infection by *Phomopsis longicolla* in a renal transplant recipient from Guinea: first report of human infection by this fungus. *Transplant Infectious Disease* 13: 204–207.
- Gardiner MR. 1975. Lupinosis. *Journal of Agriculture* 16: 26–30.
- Glass NL, Donaldson G. 1995. Development of primer sets designed for use with PCR to amplify conserved genes from filamentous ascomycetes. *Applied and Environmental Microbiology* 61: 1323–1330.
- Glienke C, Pereira OL, Stringari D, Fabris J, Kava-Cordeiro V, et al. 2011. Endophytic and pathogenic *Phyllosticta* species, with reference to those associated with Citrus Black Spot. *Persoonia* 26: 47–56.
- González V, Tello ML. 2011. The endophytic mycota associated with *Vitis vinifera* in central Spain. *Fungal Diversity* 47: 29–42.
- Groenewald JZ, Nakashima C, Nishikawa J, Shin HD, Park JH, et al. 2013. Species concepts in Cercospora: spotting the weeds among the roses. *Studies in Mycology* 75: 115–170.
- Grove WB. 1919. Species placed by Saccardo in the genus *Phoma*. Part 1. *Bulletin of Miscellaneous Information of the Royal Botanical Gardens Kew* 4: 177–201.
- Grove WB. 1935. British stem- and leaf-fungi (Coelomycetes). Cambridge University Press, UK.
- Gulya TJ, Rashid KY, Masirevic SM. 1997. Sunflower diseases. In: Schneiter AA (ed), *Sunflower technology and production*: 313–319. American Society of Agronomy, Madison USA.
- Hampson MC. 1981. *Phomopsis* canker on weeping fig in Newfoundland. *Canadian Plant Disease Survey* 61, 1: 3–5.
- Hanlin RT. 1963. A revision of the Ascomycetes of Georgia. Mimeo series, Georgia Agricultural Experiment Station 175: 1–65.
- Harter LL. 1914. Fruit-rot, leaf-spot and stem-blight of the eggplant caused by *Phomopsis vexans*. *Journal of Agricultural Research* 2: 331–338.
- Hawksworth DL, Crous PW, Redhead SA, Reynolds DR, Samson RA, et al. 2011. The Amsterdam Declaration on Fungal Nomenclature. *IMA Fungus* 2: 105–112.
- Hillis DM, Bull JJ. 1993. An empirical test of bootstrapping as a method for assessing confidence in phylogenetic analysis. *Systematic Biology* 42: 182–192.
- Hobbs TW, Schmittbrenner AF, Kuter GA. 1985. A new *Phomopsis* species from soybean. *Mycologia* 77: 535–544.
- Holiday P. 1980. Fungus diseases of tropical crops. Cambridge University Press, Cambridge, UK.
- Huber GA. 1936. A *Phomopsis* canker and gall disease of gardenia. *Florist Exchange and Horticultural Trade World* 86, 10: 11.
- Iriart X, Binois R, Fior A, Blanchet D, Berry A, et al. 2011. Eumycetoma caused by *Diaporthe phaseolorum* (*Phomopsis phaseoli*): a case report and a mini-review of *Diaporthe/Phomopsis* spp. invasive infections in humans. *Clinical Microbiology and Infection* 17: 1492–1494.
- Isaka M, Jaturapat A, Rukseree K, Danwisetkanjana K, Tanticharoen M, Thebtaranonth Y. 2001. Phomoxanthones A and B, novel xanthone dimers from the endophytic fungus *Phomopsis* species. *Journal Natural Products* 64: 1015–1018.
- Jordaan A, Taylor JE, Rossenkhahn R. 2006. Occurrence and possible role of endophytic fungi associated with seed pods of *Colophospermum mopane* (Fabaceae) in Botswana. *South African Journal of Botany* 72: 245–255.
- Katoh K, Toh H. 2008. Recent developments in the MAFFT multiple sequence alignment program. *Briefings in Bioinformatics* 9: 286–298.
- Kobayashi H, Meguro S, Yoshimoto T, Namikoshi M. 2003. Absolute structure, biosynthesis, and anti-microtubule activity of phomopsidin, isolated from a marine derived fungus *Phomopsis* sp. *Tetrahedron* 59: 455–459.
- Kuleci E, Tunali B, Berner DK, Cavin CA, Castlebury LA. 2009. First report of leaf anthracnose caused by *Phomopsis convolvuli* on field bindweed in Turkey. *Plant Disease* 93: 847.
- Kulik MM. 1984. Failure of *Phomopsis phaseoli* to produce mature pycnidia in senescent soybean stems at the end of the growing season. *Mycologia* 76: 863–867.
- Kulik MM, Sinclair JB. 1999. *Phomopsis* seed decay. In: Hartman GL, Sinclair JB, Rupe JC (eds), *Compendium of soybean diseases*: 31–32. American Phytopathological Society, St. Paul, USA.
- Kumaran RS, Hur B. 2009. Screening of species of the endophytic fungus *Phomopsis* for the production of the anticancer drug taxol. *Biotechnology and Applied Biochemistry* 54: 21–30.
- Li KN, Rouse D, German TL. 1994. PCR primers that allow intergeneric differentiation of ascomycetes and their application to *Verticillium* spp. *Applied and Environmental Microbiology* 60: 4324–4331.
- Librado P, Rozas J. 2009. DnaSP v. 5: a software for comprehensive analysis of DNA polymorphism data. *Bioinformatics* 25: 1451–1452.
- Lin X, Huang Y, Fang M, Wang J, Zheng Z, Su W. 2005. Cytotoxic and antimicrobial metabolites from marine lignicolous fungi, *Diaporthe* sp. *FEMS Microbiology Letters* 251: 53–58.
- Marincowitz S, Crous PW, Groenewald JZ, Wingfield MJ. 2008. Microfungi occurring on the Proteaceae in the fynbos. *CBS Biodiversity Series* 7: 1–166. CBS Fungal Biodiversity Centre, Utrecht, Netherlands.
- Masirevic S, Gulya TJ. 1992. Sclerotinia and *Phomopsis* – two devastating sunflower pathogens. *Field Crops Research* 30: 271–300.
- Mathur RS. 1979. The Coelomycetes of India. Delhi, India: Bishen Singh Mahendra Pal Singh, India.
- McKeen CD. 1957. *Phomopsis* black rot of cucurbits. *Canadian Journal of Botany* 35: 43–50.
- McKenzie EHC. 1992. Fungi of the Kermadec Islands. *Mycotaxon* 45: 149–170.
- McKenzie MA, Jones LH, Gilgut CJ. 1940. *Phomopsis gardeniae* in relation to gardenia culture. *Plant Disease Reporter* 24: 58–62.
- McPartland J. 1983. *Phomopsis ganjae* sp. nov. on *Cannabis sativa*. *Mycotaxon* 18: 527–530.
- Mel'nik VA, Shabunin DA, Popov ES. 2008. Contributions to the studies of mycobiota in Novgorod and Pskov regions. II. Coelomycetes. *Mikrobiologiya i Fitopatologiya* 42: 43–52.
- Mengistu A, Castlebury LA, Smith JR, Rossman AY, Reddy KN. 2007. Isolates of *Diaporthe-Phomopsis* from weeds and their effect on soybean. *Canadian Journal of Plant Pathology* 29: 283–289.
- Merrin SJ, Nair NG, Tarjan J. 1995. Variation in *Phomopsis* recorded on grapevine in Australia and its taxonomic and biological implications. *Australasian Plant Pathology* 24: 44–56.
- Michalopoulos-Skarmoutsos HG, Skarmoutsos G. 1999. Pathogenicity of fungi affecting black locust (*Robinia pseudoacacia*) in Greece. *Phytoparasitica* 27: 233–234.
- Milholland RD, Daykin ME. 1983. Blueberry fruit rot caused by *Phomopsis vaccinii*. *Plant Disease* 67: 325–326.

- Miller HN. 1961. Annual report of the agricultural experiment stations, Florida, for the year ending June 30, 1961. *Revue of Applied Mycology* 42: 236.
- Mondal SN, Vicent A, Reis RF, Timmer LW. 2007. Saprophytic colonisation of citrus twigs by *Diaporthe citri* and factors affecting pycnidia production and conidial survival. *Plant Disease* 91: 387–392.
- Morgan-Jones G. 1989. The *Diaporthe/Phomopsis* complex: taxonomical considerations. In: Pascale AJ (ed), *World Soybean Research Conference IV Proceedings: 1699–1706*. Orientación Gráfica Editora, Buenos Aires, Argentina.
- Moricca S. 2002. *Phomopsis alnea*, the cause of dieback of black alder in Italy. *Plant Pathology* 51: 755–764.
- Mostert L, Crous PW, Kang JC, Phillips AJL. 2001a. Species of *Phomopsis* and a *Libertia* sp. occurring on grapevines with specific reference to South Africa: morphological, cultural, molecular and pathological characterization. *Mycologia* 93: 146–167.
- Mostert L, Kang JC, Crous PW, Denman S. 2001b. *Phomopsis saccharata* sp. nov., causing a canker and die-back disease of *Protea repens* in South Africa. *Sydowia* 53: 227–235.
- Mułenko W, Majewski T, Ruszkiewicz-Michalska M (eds). 2008. A preliminary checklist of micromycetes in Poland. In: *Biodiversity of Poland* 9: 752. W. Szafer Institute of Botany, Polish Academy of Sciences, Kraków.
- Munk A. 1957. Danish pyrenomyces. A preliminary flora. *Dansk Botanisk Arkiv Udgivet af Dansk Botanisk Forening*, Copenhagen 17: 1–491.
- Muntañola-Cvetkovic M, Mihaljevic M, Petrov M. 1981. On the identity of the causative agent of a serious *Phomopsis-Diaporthe* disease in sunflower plants. *Nova Hedwigia* 34: 417–435.
- Murali TS, Suryanarayanan TS, Geeta R. 2006. Endophytic *Phomopsis* species: host range and implications for diversity estimates. *Canadian Journal of Microbiology* 52: 673–680.
- Mylllys L, Stenroos S, Thell A. 2002. New genes for phylogenetic studies of lichenized fungi: glyceraldehyde-3-phosphate dehydrogenase and beta-tubulin genes. *Lichenologist* 34: 237–246.
- Niekerk JM van, Groenewald JZ, Farr DF, Fourie PH, Halleen F, Crous PW. 2005. Reassessment of *Phomopsis* species on grapevines. *Australasian Plant Pathology* 34: 27–39.
- Nitschke T. 1867. Pyrenomycetes Germanici. Die Kernpilze Deutschlands, Breslau 1.
- Nooit MP de, Aa HA van der. 1987. *Phomopsis subordinaria* and associated stalk disease in natural populations of *Plantago lanceolata*. *Canadian Journal of Botany* 65: 2318–2325.
- Nylander JAA. 2004. MrModeltest v2. Program distributed by the author. Evolutionary Biology Centre, Uppsala University, Sweden.
- O'Donnell K, Cigelnik E. 1997. Two divergent intragenomic rDNA ITS2 types within a monophyletic lineage of the fungus *Fusarium* are nonorthologous. *Molecular Phylogenetics and Evolution* 7: 103–116.
- O'Donnell K, Cigelnik E, Nirenberg HI. 1998. Molecular systematics and phylogeography of the *Gibberella fujikuroi* species complex. *Mycologia* 90: 465–493.
- Oak SW, Dorset RD. 1983. *Phomopsis* canker of European black alder found in Kentucky seed-production areas. *Plant Disease* 67: 691–693.
- Ogawa JM, English H. 1991. Disease of temperate zone tree fruit and nut crops. University of California, Division of Agriculture and Natural Resources, Oakland, California. Publication 3345.
- Ohsawa T, Kobayashi T. 1989. Concave rot of melon fruit caused by two *Phomopsis* fungi. *Annals of the Phytopathological Society of Japan* 55: 410–419.
- Ormeno-Núñez ZR, Eeleder D, Watson K. 1988. A new species of *Phomopsis* recovered from field bindweed (*Convolvulus arvensis*). *Canadian Journal of Botany* 66: 2228–2233.
- Ostazeski SA, Wells HD. 1960. A *phomopsis* stem blight of yellow lupins (*Lupinus luteus* L.). *Plant Disease Reporter* 44: 66–67.
- Pecchia S, Mercatelli E, Vannacci G. 2004. Intraspecific diversity within *Diaporthe helianthi*: evidence from rDNA intergenic spacer (IGS) sequence analysis. *Mycopathologia* 157: 317–326.
- Phillips AJL. 1999. The relationship between *Diaporthe perjuncta* and *Phomopsis viticola* on grapevines. *Mycologia* 91: 1001–1007.
- Phillips AJL. 2003. Morphological characterization of *Diaporthe foeniculacea* and its *Phomopsis* anamorph on *Foeniculum vulgare*. *Sydowia* 55: 274–285.
- Preston DA. 1945. Host index of Oklahoma plant diseases. Oklahoma Agricultural College, Agricultural Experiment Station Technical Bulletin 21: 1–168.
- Punithalingam E. 1973. Two new species of *Phomopsis*. *Transactions of the British Mycological Society* 60: 157–160.
- Punithalingam E. 1974. Studies on *Sphaeropsidales* in culture. II. *Mycological Papers* 136: 1–63.
- Punithalingam E, Holliday P. 1972. *Phomopsis vexans*. CMI Descriptions of Pathogenic Fungi and Bacteria No. 338.
- Punithalingam E, Holliday P. 1973. *Diaporthe citri*. CMI Descriptions of Pathogenic Fungi and Bacteria No. 396.
- Punithalingam E, Holliday P. 1975. *Phomopsis cucurbitae*. CMI Descriptions of Pathogenic Fungi and Bacteria No. 469.
- Quaedvlieg W, Kema GHJ, Groenewald JZ, Verleyen GJM, Seifbarghi S, et al. 2011. *Zymoseptoria* gen. nov.: a new genus to accommodate *Septoria*-like species occurring on graminicolous hosts. *Persoonia* 26: 57–69.
- Rawnsley B, Wicks TJ, Scott ES, Stummer BE. 2004. *Diaporthe perjuncta* does not cause *Phomopsis* cane and leaf spot disease of grapevine in Australia. *Plant Disease* 88: 1005–1010.
- Rayner RW. 1970. A mycological colour chart. Commonwealth Mycological Institute, Kew, UK.
- Rehner SA, Uecker FA. 1994. Nuclear ribosomal internal transcribed spacer phylogeny and host diversity in the coelomycete *Phomopsis*. *Canadian Journal of Botany* 72: 1666–1674.
- Rekab D, Sorbo G del, Reggio C, Zoina A, Firrao G. 2004. Polymorphisms in nuclear rDNA and mtDNA reveal the polyphyletic nature of isolates of *Phomopsis* pathogenic to sunflower and a tight monophyletic clade of defined geographic origin. *Mycological Research* 108: 393–402.
- Rensburg JCJ van, Lampecht SC, Groenewald JZ, Castlebury LA, Crous PW. 2006. Characterization of *Phomopsis* spp. associated with die-back of rooibos (*Aspalathus linearis*) in South Africa. *Studies in Mycology* 55: 65–74.
- Reynolds DR. 1993. The fungal holomorph: an overview. In: Reynolds DR, Taylor JW (eds), *The fungal holomorph: mitotic, meiotic and pleomorphic speciation in fungal systematics*: 15–25. CAB International, Wallingford, UK.
- Ronquist F, Huelsenbeck JP. 2003. MrBayes 3: Bayesian phylogenetic inference under mixed models. *Bioinformatics* 19: 1572–1574.
- Rossman AY, Farr DF, Castlebury LA. 2007. A review of the phylogeny and biology of the Diaporthales. *Mycoscience* 48: 135–144.
- Saccardo PA. 1878. *Fungi Veneti novi vel critici auctore. Series VIII. Appendicula*. *Michelia* 1: 351–355.
- Saccardo PA. 1879. *Fungi Galicii lecti a cl. viris P. Brunaud, C.C. Gillet et Abb. Letendre*. *Michelia* 1: 500–538.
- Santos JM, Correia VG, Phillips AJL. 2010. Primers for mating-type diagnosis in *Diaporthe* and *Phomopsis*: their use in teleomorph induction in vitro and biological species definition. *Fungal Biology* 114: 255–270.
- Santos JM, Phillips AJL. 2009. Resolving the complex of *Diaporthe* (*Phomopsis*) species occurring on *Foeniculum vulgare* in Portugal. *Fungal Diversity* 34: 111–125.
- Santos JM, Vrandečić K, Čosić J, Duvnjak T, Phillips AJL. 2011. Resolving the *Diaporthe* species occurring on soybean in Croatia. *Persoonia* 27: 9–19.
- Sarbhoj AK, Lal G, Varshney JL. 1971. *Fungi of India*. Navyug Traders, New Delhi, India.
- Scerbin-Parfenenko AL. 1953. *Rakovye i sosudistye bolezni listvennyh porod*. Goslesbumisdat, Moskva-Leningrad.
- Scheper RWA, Crane DC, Whisson DL, Scott ES. 2000. The *Diaporthe* teleomorph of *Phomopsis* Taxon 1 on grapevine. *Mycological Research* 104: 226–231.
- Schoch CL, Seifert KA, Huhndorf S, Robert V, Spouge JL, et al. 2012. Nuclear ribosomal internal transcribed spacer (ITS) region as a universal DNA barcode marker for Fungi. *Proceedings of the National Academy of Sciences of the United States of America* 109: 6241–6246.
- Sebastiane FLS, Lacava PT, Fávaro LCL, Rodrigues MBC, Araújo WL, et al. 2011. Genetic transformation of *Diaporthe phaseolorum*, an endophytic fungus found in mangrove forests, mediated by *Agrobacterium tumefaciens*. *Current Genetics* 58: 21–33.
- Shaw CG. 1973. Host fungus index for the Pacific Northwest - I. Hosts. Washington State University Agricultural Experiment Station Bulletin 765: 1–121.
- Silva GH, Teles HL, Trevisan HC, Bolzani VS, Young MCM, et al. 2005. New bioactive metabolites produced by *Phomopsis cassiae*, an endophytic fungus in *Cassia spectabilis*. *Journal of Brazilian Chemical Society* 16: 1463–1466.
- Skaltsas D, Castlebury L, Chaverri P. 2011. Delimitation of tropical endophytic *Phomopsis* species from three euphorbiaceous hosts: *Hevea brasiliensis*, *H. guianensis*, and *Micandra* sp. *Inoculum* 62, 3: 41.
- Smit WA, Knox-Davies PS. 1989a. Die-back of rooibos tea caused by *Diaporthe phaseolorum*. *Phytophylactica* 21: 183–188.
- Smit WA, Knox-Davies PS. 1989b. Comparison of *Diaporthe phaseolorum* isolates from rooibos tea, *Aspalathus linearis*. *Phytophylactica* 21: 301–306.
- Smit WA, Wingfield MJ, Wingfield BD. 1996. A new canker disease of apple, pear, and plum rootstocks caused by *Diaporthe ambigua* in South Africa. *Plant Disease* 80: 1331–1335.
- Smith H, Wingfield MJ, Crous PW, Coutinho TA. 1996. *Sphaeropsis sapinea* and *Botryosphaeria dothidea* endophytic in *Pinus* spp. and *Eucalyptus* spp. in South Africa. *South African Journal of Botany* 62: 86–88.

- Spielman LJ. 1985. A monograph of *Valsa* on hardwoods in North America. Canadian Journal of Botany 63: 1355–1378.
- Sutton DA, Timm WD, Morgan-Jones G, Rinaldi MG. 1999. Human phaeohyphomycotic osteomyelitis caused by the coelomycete *Phomopsis saccardo 1905*: criteria for identification, case history, and therapy. Journal of Clinical Microbiology 37: 807–811.
- Swofford DL. 2003. PAUP*: phylogenetic analysis using parsimony (*and other methods), version 4. Sinauer Associates, Sunderland, Massachusetts.
- Tai FL. 1979. *Sylloge Fungorum Sinicorum*. Science Press, Academica Sinica, Peking.
- Tamura K, Peterson D, Peterson N, Stecher G, Nei M, Kumar S. 2011. MEGA5: Molecular evolutionary genetics analysis using maximum likelihood, evolutionary distance, and maximum parsimony methods. Molecular Biology and Evolution 28: 2731–2739.
- Taylor JW, Jacobson DJ, Kroken S, Kasuga T, Geiser DM, et al. 2000. Phylogenetic species recognition and species concepts in fungi. Fungal Genetics and Biology 31: 21–32.
- Teodorescu G, Copăescu V, Florea S. 1985. The behaviour of some blueberry cultivars to the main mycoses in Romania. Acta Horticulturae 165: 159–165.
- Thompson SM, Tan YP, Young AJ, Neate SM, Aitken EA, Shivas RG. 2011. Stem cankers on sunflower (*Helianthus annuus*) in Australia reveal a complex of pathogenic Diaporthe (*Phomopsis*) species. Persoonia 27: 80–89.
- Tilford PE. 1934. Stem canker disease of gardenia. Bulletin of the Ohio Agriculture Experiment Station 19 (168): 116–117.
- Tuset JJ, Portilla MT. 1989. Taxonomic status of *Fusicoccum amygdali* and *Phomopsis amygdalina*. Canadian Journal of Botany 67: 1275–1280.
- Udayanga D, Xingzhong L, Crous PW, McKenzie EHC, Chukeatirote E, Hyde KD. 2012. A multi-locus phylogenetic evaluation of Diaporthe (*Phomopsis*). Fungal Diversity 56: 157–171.
- Udayanga D, Xingzhong L, McKenzie EHC, Chukeatirote E, Bahkali AHA, Hyde KD. 2011. The genus *Phomopsis*: biology, applications, species concepts and names of common pathogens. Fungal Diversity 50: 189–225.
- Uecker FA. 1988. A world list of *Phomopsis* names with notes on nomenclature, morphology and biology. Mycological Memoirs 13: 1–231.
- Vajna L. 2002. Diaporthe oncostoma causing stem canker of black locust in Hungary. Plant Pathology 51: 393.
- Vasiliyeva LN, Rossman AY, Farr DF. 2007. New species of the Diaporthales from eastern Asia and eastern North America. Mycologia 99: 916–923.
- Vesterlund SR, Helander M, Faeth SH, Hyvönen T, Saikkonen K. 2011. Environmental conditions and host plant origin override endophyte effects on invertebrate communities. Fungal Diversity 47: 109–118.
- Vilgalys R, Hester M. 1990. Rapid genetic identification and mapping of enzymatically amplified ribosomal DNA from several *Cryptococcus* species. Journal of Bacteriology 172: 4238–4246.
- Webber J, Gibbs JN. 1984. Colonization of elm bark by *Phomopsis oblonga*. Transactions of the British Mycological Society 82: 348–352.
- Weber D. 2009. Endophytic fungi, occurrence and metabolites. In: Anke T, Weber D (eds), *Physiology and genetics. The mycota* 15: 153–195. Springer-Verlag, Berlin.
- Weber GF. 1933. Stem canker of *Crotalaria spectabilis* caused by *Diaporthe crotalariae* n. sp. Phytopathology 23: 602.
- Wehmeyer LE. 1933. The genus *Diaporthe* Nitschke and its segregates. University of Michigan Studies Scientific Series 9: 1–349.
- Weir B, Johnston PR, Damm U. 2012. The *Colletotrichum gloeosporioides* species complex. Studies in Mycology 73: 115–180.
- White TJ, Bruns T, Lee J, Taylor J. 1990. Amplification and direct sequencing of fungal ribosomal RNA genes for phylogenetics. In: Innis MA, Gelfand DH, Sninsky JJ, White TJ (eds), *PCR protocols: a guide to methods and applications*: 315–322. Academic Press, San Diego, California, USA.
- Wilcox HJ, Falconer MA. 1961. New or uncommon plant pests. Plant Pathology 10: 123–124.
- Wilcox MS. 1939. *Phomopsis* twig blight of blueberry. Phytopathology 29: 136–142.
- Williamson PM, Higher AS, Gams W, Sivasithamparam K, Cowling WA. 1994. *Diaporthe toxica* sp. nov. the cause of lupinosis in sheep. Mycology Research 98: 1364–1368.
- Wingfield MJ, Beer ZW de, Slippers B, Wingfield BD, Groenewald JZ, et al. 2012. One fungus one name promotes progressive plant pathology. Molecular Plant Pathology 13: 604–613.
- Wu SH, Chen YW, Shao SC, Wang LD, Li ZY, et al. 2008. Ten-membered lactones from *Phomopsis* sp., an endophytic fungus of *Azadirachta indica*. Journal of Natural Products 71: 731–734.
- Zhang AW, Riccioni L, Pedersen WL, Kollipara KP, Hartman GL. 1998. Molecular identification and phylogenetic grouping of *Diaporthe phaseolorum* and *Phomopsis longicolla* isolates from soybean. Phytopathology 88: 1306–1314.
- Zhuang WY. 2001. Higher fungi of tropical China. Mycotaxon, Ithaca, USA.