

 Open access • Journal Article • DOI:10.1002/IJC.26295

Dickkopf-3 is regulated by the MYCN-induced miR-17-92 cluster in neuroblastoma

— [Source link](#) 

Sara De Brouwer, Pieter Mestdagh, Irina Lambertz, Filip Pattyn ...+8 more authors

Institutions: [Ghent University Hospital](#), [German Cancer Research Center](#), [Boston Children's Hospital](#)

Published on: 01 Jun 2012 - [International Journal of Cancer](#) (Wiley Subscription Services, Inc., A Wiley Company)

Topics: [N-Myc](#) and [Neuroblastoma](#)

Related papers:

- [MYCN regulates oncogenic MicroRNAs in neuroblastoma.](#)
- [MYCN-regulated miRNA-92 inhibits secretion of the tumor suppressor DICKKOPF-3 \(DKK3\) in neuroblastoma](#)
- [The miR-17-92 MicroRNA Cluster Regulates Multiple Components of the TGF- \$\beta\$ Pathway in Neuroblastoma](#)
- [MYCN/c-MYC-induced microRNAs repress coding gene networks associated with poor outcome in MYCN/c-MYC-activated tumors](#)
- [Neuroblastoma: biological insights into a clinical enigma.](#)

Share this paper:    

View more about this paper here: <https://typeset.io/papers/dickkopf-3-is-regulated-by-the-mycn-induced-mir-17-92-538bmwaa0t>

Dickkopf-3 is regulated by the MYCN-induced miR-17-92 cluster in neuroblastoma

Sara De Brouwer¹, Pieter Mestdagh¹, Irina Lambertz¹, Filip Pattyn¹, Anne De Paepe¹, Frank Westermann², Christina Schroeder², Johannes H. Schulte³, Alexander Schramm³, Katleen De Preter¹, Jo Vandesompele¹ and Frank Speleman¹

¹ Center for Medical Genetics, Ghent University Hospital, Ghent, Belgium

² Department of Tumor Genetics, German Cancer Research Center, Heidelberg, Germany

³ Division of Hematology and Oncology, University of Children's Hospital, Essen, Germany

Neuroblastoma (NB) is a paediatric tumour with a remarkable diverse clinical behaviour. Approximately half of the high stage aggressive tumours are characterized by *MYCN* gene amplification but our understanding of the role of *MYCN* in NB oncogenesis is incomplete. Previous studies have shown that *MYCN* expression is inversely correlated with expression of *Dickkopf-3* (*DKK3*), a gene encoding an extracellular protein with presumed tumour suppressor activity, but direct *MYCN* regulation of *DKK3* was excluded leaving the mechanism of regulation unexplained. Given the recently established role of *MYCN*-regulated miRNAs in downregulation of protein-coding genes and predicted seeds for miR-17-92 cluster members within the *DKK3* 3'UTR, we hypothesized that this mechanism would act in *MYCN* regulation of *DKK3*. To investigate this, we used a validated miR-17-92-inducible cellular system and could demonstrate robust downregulation of *DKK3* mRNA and protein levels upon miR-17-92 overexpression. Next, two of the three predicted miRNAs, miR-19b and miR-92a, were shown to lower *DKK3* protein levels, in addition to measurable *DKK3* mRNA knock-down by miR-92a. Direct interaction between miR-19b or miR-92a and the 3'UTR of *DKK3* was validated using luciferase reporter assays. In conclusion, this study demonstrates that the *MYCN*-induced downregulation of *DKK3* results from direct upregulation of miR-17-92 components effecting both *DKK3* mRNA stability and translation which further contributes to the pleiotropic oncogenic effect of elevated *MYCN* levels. The strict *MYCN*-mediated regulation of *DKK3* is suggestive for an important downstream function of the *MYCN* protein and thus warrants further investigations to unravel the role of *DKK3* in NB.

Neuroblastoma (NB) is a paediatric tumour with a worldwide incidence of 10 cases per million in children younger than 15 years old.¹ NB originates from sympathetic neuronal progenitors and is characterized by a remarkable diverse clinical behaviour ranging from spontaneously regressing to highly aggressive metastasized tumours. Amplification of *MYCN* occurs with a high incidence in aggressive tumour phenotype and, furthermore, *MYCN* status can be used as an independent predictor of poor prognosis. *MYCN* is member of the

MYC family of transcription factors which contain a basic helix-loop-helix leucine zipper (HLH-zip) domain that is important for protein dimerization, sequence specific DNA binding and regulation of transcription.^{2,3} Evidence for a direct role for *MYCN* in NB development was obtained through the generation of a transgenic mouse model where a tyrosine hydroxylase-driven overexpression of *MYCN* in sympathetic neural crest progenitor cells resulted in the development of NB tumours.⁴ Subsequently, further studies aimed at identifying the transcriptional targets of *MYCN* in NB cells by using *MYCN*-regulable constructs,⁵⁻⁷ retinoic acid treatment^{8,9} and siRNA-mediated *MYCN* knock-down.¹⁰⁻¹² Remarkably, despite these in-depth investigations, only a limited number of *bona fide* upregulated target genes could be identified, including *ODC1*, *MDM2*, *MCM7* and *hTERT*.^{5,13-15} Transcriptional upregulation results from binding of the *MYCN*/*MAX* heterodimer to E-box sequences in the promoter region of the target genes as well as through binding to non-canonical sequences.¹⁶ In addition to activation of transcription, *MYCN* can also function as a transcriptional repressor of genes such as *TRKA* and *p75NTR*. Here, *MYCN* associates with transcription factors *MIZ1*/*SP1* followed by a recruitment of *HDAC1*, thereby inducing a repressed chromatin state.¹⁷ These and other studies indicate that *MYCN* controls a broad regulatory network implicated

Key words: neuroblastoma, dickkopf-3, miR-17-92 cluster

Additional Supporting Information may be found in the online version of this article.

Grant sponsors: Flemish League against Cancer, Belgian Program of Interuniversity Poles of Attraction, initiated by the Belgian State, Prime Minister's Office, Science Policy Programming, Fund for Scientific Research; **Grant numbers:** G.O198.08, 01G01910

DOI: 10.1002/ijc.26295

History: Received 3 Mar 2011; Accepted 24 Jun 2011; Online 27 Jul 2011

Correspondence to: Frank Speleman, Center for Medical Genetics, Ghent University Hospital, De Pintelaan 185, B-9000 Ghent, Belgium, Tel.: +32-9-332-2451, Fax: +32-9-332-6549, E-mail: franki.speleman@UGent.be

in cell cycle, DNA damage response, differentiation and apoptosis. Unravelling the downstream pathways controlled by MYCN is essential for understanding how this transcription factor contributes to the process of tumour development, particularly that of NB, which will further be important in the choice of appropriate therapeutic targets.

In addition to the above-mentioned direct MYCN targets, the *Dickkopf-3* (*DKK3*) gene has also been studied as it appears to be consistently regulated in several systems upon modulation of MYCN expression.¹⁸ *DKK3* is a member of the DKK family of proteins which play important roles in normal development and in the negative modulation of the Wnt/ β -catenin signaling pathway.^{19–22} High expression levels of this putative tumour suppressor gene was shown to be positively correlated with good prognosis for NB patients,^{23,24} whereas reduced *DKK3* expression has been observed in multiple cell lines and tumours of endothelial and epithelial origin. In certain tumour entities, decreased *DKK3* expression levels were found in association with promoter-hypermethylation.^{23,25–28} Furthermore, overexpression of *DKK3* was shown to inhibit proliferation and motility both *in vitro* and *in vivo*.^{25,29–36} Taken together, these data are in keeping with a tumour suppressor function for *DKK3*.

Koppen and colleagues observed that *DKK3* mRNA expression in NB cell lines was negatively correlated with MYCN expression levels and further demonstrated MYCN-mediated downregulation of *DKK3* using a MYCN-inducible cellular system. Chromatin immunoprecipitation assays, however, failed to identify direct interaction between MYCN and the promoter region of the *DKK3* gene, implying a mechanism of regulation different from direct transcriptional inhibition.²³

Recently, several studies have demonstrated that an indirect MYCN-mediated downregulation of gene expression results from direct activation of specific microRNAs (miRNAs).^{6,37} MiRNAs are small non-coding RNAs that negatively regulate the expression of target mRNAs at post-transcriptional level and they are known important regulators of oncogenes and tumour suppressor genes. In a previous study, we explored the role of the MYCN-driven oncogenic miR-17-92 cluster in NB cells. MiR-17-92 activation was shown to result in a profound downregulation of TGF- β pathway activity, accompanied by increased cell proliferation *in vitro* and *in vivo*. Moreover, high expression of the miR-17-92 cluster in NB tumours was significantly correlated to poor patient survival.³⁸ Since *DKK3* is indirectly downregulated by MYCN, we hypothesized that *DKK3* translational repression and/or degradation could be indirectly controlled through the MYCN-activated miR-17-92 cluster. In this paper, we confirmed a direct regulation of both *DKK3* mRNA and *DKK3* protein levels by miR-17-92 components, thus providing a mechanistic basis for the tight downregulation of *DKK3* levels through MYCN in NB cells.

Material and Methods

Samples and cell lines

A representative cohort of 101 primary untreated NB tumours of all clinical stages was used for analysis. Written

informed consent was obtained from the patients' relatives and an ethical approval from the Ghent University Hospital Ethical Committee (EC2008/159) was also obtained.

The SHEP-Tet21N cell line is derived from the SHEP NB cell line and contains a MYCN tet-off system where, more specifically, the expression of a MYCN construct is repressed by tetracycline in the culture medium. The SHEP-TR-miR-17-92 NB cell line contains a tet-on system to induce the oncogenic miR-17-92 cluster with the addition of 2 μ g/ml of tetracycline to the culture medium. In the SHEP MYCN-ER cell line, the cDNA of MYCN is fused to a mutated estrogen responsive domain (ER) which can bind 4-hydroxy-tamoxifen (4-OHT) but is unable to bind with natural estrogen.⁶ Addition of 200 nM 4-OHT to the culture medium activates the MYCN-ER expression.

Data from the NCI-60 cell line panel was downloaded from the National Cancer Institute's Developmental Therapeutic Program Web site (www.sanger.ac.uk/genetics/CGP/NCI60/).

Cell culture and transfection

All NB cell lines were cultured in RPMI1640 medium (Invitrogen) supplemented with 10% FBS, 1% PEN/STREP, 1% kanamycin and 1% glutamine at 37°C in a 5% CO₂ atmosphere. DLD1Dicer^{hypo} cells were grown in DMEM medium (Invitrogen) using the same supplements as the RPMI1640 medium.

SHEP cells were seeded at 150,000 cells per 6-well, 24 hr prior to transfection. Transfection was performed using the XtremeGene lipid (Roche) mixed with OPTI-MEM (Invitrogen) and 100 nM pre-miR (-19a, 19b, -92a or non-targeting control). Conditioned medium was harvested for ELISA and cells were pelleted for RNA isolation (miRNeasy mini kit, Qiagen) and for subsequent *DKK3* expression analysis 24 hr, 48 and 72 hr post-transfection.

SHEP MYCN-ER cells were seeded at 150,000 cells per 6-well, 24 hr prior to 4-OHT induction. Transfection was performed 24 hr post 4-OHT induction using the XtremeGene lipid (Roche) mixed with OPTI-MEM (Invitrogen) and 100 nM anti-miR (-92a or non-targeting control). Cells were pelleted for RNA isolation 48 hr post-transfection.

Quantitative polymerase reaction (qPCR)

cDNA synthesis was performed on 2 μ g total RNA (isolated using the miRNeasy mini-kit, Qiagen according to the manufacturer's protocol) with 4 μ l of iScript reaction mix and 1 μ l of iScript reverse transcriptase (iScript, Bio-Rad) in a final volume of 20 μ l. Subsequently, this mix was incubated for 5' at 25°C, 30' at 42°C and 5' at 85°C using the iCycler (Bio-Rad). The expression levels were analyzed using 5 ng cDNA, 2.5 μ l of SYBR Green I mastermix (Eurogentec) and 5 μ M primers in a total volume of 5 μ l. Expression levels were normalized against two of four reference gene (*GAPDH*, *HPRT1*, *YWHAZ*, *SDHA*) and analyzed using the qBasePlus software (<http://www.biogazelle.com>). All used qPCR assays are




Figure 1. Inverse correlation of *DKK3* and *MYCN* in a series of NB cell lines. In cell lines with *MYCN* amplification (SK-MYC2, TR14, LAN-5, NGP, IMR-32, SMS-KAN, SK-N-BE), the expression of *DKK3* is low while *DKK3* expression is high in cell lines that are *MYCN* single copy (SK-N-AS, SK-N-SH, CLB-GA, SK-N-FI, NBLS). The y-axis is shown in logarithmic scale.

available in RTPprimerDB with IDs 3 (*GAPDH*), 5 (*HPRT1*), 9 (*YWHAZ*), 7 (*SDHA*), 8229 (*DKK3*) and 180 (*MYCN*) (<http://www.rtpprimerdb.org>).³⁹

In silico miRNA target predictions

MiRNAs putatively targeting the 3'UTR of *DKK3* were investigated using the online prediction databases TargetScan v5.1 (<http://www.targetscan.org>) and DIANA microT v3.0 (<http://diana.cslab.ece.ntua.gr/microT/>).

Luciferase assay

The 3'UTR of the human *DKK3* gene was cloned in a vector containing the firefly luciferase open reading frame (pSGG_3'UTR, SwitchGear Genomics). Seed regions were mutated using the QuickChange II mutagenesis kit (Stratagene). Specifically, the miR-19 complementary site, gTtacCA, was changed into tTgcaCA and the miR-92a complementary site, gTgcaATA, was changed into tTtacATA (mutated sites in lower case). Prior to seeding of the cells, plates were coated with fibronectin (Tebu-Bio) for a better attachment of the DLD1Dicer^{hypo} cells. Cells were plated 24 hr prior to transfection at 10,000 cells per 96-well in DMEM medium containing 10% FBS. Transfection mixes contained RPMI1640, 50 nM pre-miR (Applied Biosystems), 0.4% DharmaFect Duo, 100 ng 3'UTR vector and 20 ng pRL-TK vector, containing the *Renilla* luciferase (Promega). The ratios between firefly and *Renilla* luciferase were measured 48 hr post-transfection using the dual-glo luciferase kit (Promega) according to the manufacturer's protocol.

Enzyme-linked immunosorbent assay (ELISA)

Cell culture supernatants were collected at various time points and cleared of cellular debris by microcentrifugation. The concentration of secreted *DKK3* was measured using the human Dkk-3 DuoSet ELISA kit (R&D Systems, DY1118)




Figure 2. Correlation of *DKK3* and *MYCN* expression in a series of 101 primary untreated NB tumours. Expression of *DKK3* (x-axis) and *MYCN* (y-axis) are significantly inversely correlated in 101 NB tumours (Spearman correlation $r = -0.30$, $p < 0.01$).

according to the manufacturer's instructions. Optical density was measured using a Multiskan RC plate reader (LabSystems) and values were plotted against a 4-PL standard curve (Genesis software). Finally, *DKK3* concentration values were normalized against total micrograms of protein in the cell culture supernatants, determined by the DC protein assay (Biorad, 500-0116).

Statistical analysis

Correlation analysis of gene expressions in 101 tumour samples was evaluated with the Spearman correlation coefficient using the R statistical package. Protein or mRNA expression levels between different conditions or samples are assessed with the unpaired two-sided Student *t*-test and a significance level of 0.05 as cut-off. Calculation of standard errors of repeated experiments was performed using the error propagation rules.

Results

MYCN/c-MYC mRNA levels are inversely correlated with *DKK3* expression. Expression of *DKK3* has previously been correlated to *MYCN* and *c-MYC* expression levels (here referred to as *MYCN/c-MYC*).^{23,40,41} We confirmed the previously reported²³ inverse relationship between *DKK3* expression levels and *MYCN* in a set of 12 NB cell lines which were either *MYCN* amplified or *MYCN* single-copy (Fig. 1) and through analysis of their expression levels in an independent cohort of 101 primary untreated NB tumours ($r = -0.30$, $p < 0.01$) (Fig. 2). Analysis of the SHEP-Tet21N cell line with a tetracycline-regulable *MYCN* construct^{5,16} further confirmed this negative correlation, as shown by the decreased levels of *DKK3* after *MYCN* induction (Fig. 3).




Figure 3. Inverse correlation between *DKK3* and *MYCN* using the SHEP-TET21N model system. Expression of *DKK3* and *MYCN* are shown after *MYCN*-induction at different timepoints in the SHEP-TET21N model system (shown on x-axis). The y-axis is shown in a logarithmic scale.

Additionally, we evaluated the expression levels of *DKK3* and *MYCN/c-MYC* in the NCI-60 cell line panel which contains cell lines of several tumour entities other than NB with elevated expression of *MYCN* or *c-MYC*. This analysis also yielded a significant negative correlation between *DKK3* and *MYCN/c-MYC* (Spearman correlation $r = -0.51$, $p < 0.01$). Altogether, these results confirm the tight relationship between *MYCN* and *DKK3* expression levels.

***DKK3* expression levels are inversely correlated with the expression of members of the miR-17-92 cluster**

Using the SHEP-Tet21N model system, *DKK3* downregulation only became apparent 24 hr after *MYCN*-induction suggesting indirect regulation of *DKK3* (Fig. 3). In view of the recently established role of the miR-17-92 cluster as an effector for indirect *MYCN*-induced downregulation of protein-coding genes, we hypothesized that this cluster might also regulate *DKK3*. To investigate this possibility we first verified *DKK3* mRNA expression following conditional upregulation of the miR-17-92 cluster in the SHEP-TR-miR-17-92 NB cell line. This indeed showed a significant downregulation of *DKK3* transcripts (Fig. 4a; $p = 0.02$ after 72 hr) which was subsequently confirmed at the protein level (Fig. 4b; $p < 0.01$ after 72 hr), in keeping with the possibility that *DKK3* is a direct target of one or more components from the miR-17-92 cluster.

Next, we performed an *in silico* search for putative miRNA target sites within the 3'UTR of *DKK3* using the TargetScan and DIANA microT prediction databases. Several members of the *MYCN*-activated miR-17-92 cluster, more specifically miR-19a, miR-19b and miR-92a, were predicted to target the 3'UTR of *DKK3* (Fig. 5a). We further looked for correlation between *DKK3* expression and each of these three predicted miRNAs in a set of 101 primary untreated




Figure 4. *DKK3* expression levels using the SHEP-TR-miR-17-92 cell line. (a) mRNA expression levels of *DKK3* after miR-17-92 induction in the SHEP-TR-miR-17-92 cell line. (b) Protein expression levels of *DKK3* after miR-17-92 induction in the SHEP-TR-miR-17-92 cell line, measured using ELISA. Ctr = SHEP-TR-miR-17-92 cells treated with ethanol as control, Tetr = SHEP-TR-miR-17-92 cells treated with tetracycline for induction of the miR-17-92 cluster.

NB tumours and all three miRNAs showed a significant inverse correlation with *DKK3* (Supporting Information Table 1) which we could further confirm in the NCI-60 cell line collection (Supporting Information Table 2). These results suggest that the miR-17-92 cluster is the missing link between *MYCN* and *DKK3*.

The *DKK3* 3'UTR is directly targeted by members of the miR-17-92 cluster

To experimentally establish the direct role of the miR-17-92 cluster in *DKK3* regulation, we tested the three miRNAs from the cluster with predicted putative binding sites in the 3'UTR of *DKK3*, i.e., miR-19a, miR-19b and miR-92a. Luciferase assays were performed in a cell line derived from colon carcinoma containing a hypomorphic mutation in Dicer (*DLD1Dicer^{hypo}*). This cell line has a reduced amount of mature miRNA levels which makes the effect of pre-miR transfection higher than in non-mutated cell lines. Co-




Figure 5. miRNA interaction with the 3'UTR of *DKK3*. (a) Localisation of putative binding sites of miR-19 and miR-92a in the 3'UTR of *DKK3*. The positions of mutations of the miRNA seed sequences (miR-19 or miR-92a) in the 3'UTR of *DKK3* are shown in bold. (b) Relative luciferase activity for transfection of a wild type or mutated 3'UTR *DKK3* vector in combination with pre-miR-19a or pre-miR-92a. WT = wild type; MUT = mutated vector.

transfection of a selected pre-miR (pre-miR-19a, pre-miR-19b or pre-miR-92a) and a vector containing the 3'UTR of *DKK3* showed significant reduction for both pre-miR-19b ($p < 0.01$) and pre-miR-92a ($p < 0.01$) indicating a direct interaction between these two miRNAs and the 3'UTR of *DKK3* (Fig. 5b). Introduction of mutations in the miRNA seeds sites of miR-19b and miR-92a rescued the luciferase output which suggests that the effect of luciferase reduction depend on the presence of the miRNA seed sequences in the 3'UTR (Fig. 5b). No reduction in relative luciferase output could be observed upon repeated experiments for pre-miR-19a, thus excluding this miRNA as an effector of *DKK3* expression (data not shown).

miR-19b and miR-92a directly regulate *DKK3* expression

Since both miR-19b and miR-92a interact with the 3'UTR of *DKK3*, we overexpressed each individual miRNA in the NB SHEP cell line using pre-miR mimics and evaluated the effect on both mRNA and protein *DKK3* levels. As expected, for both miRNAs, *DKK3* protein levels were shown to be significantly reduced after 72 hr of pre-miR-19b or pre-miR-92a overexpression, respectively, 12% reduction ($p = 0.02$) and 58% reduction ($p < 0.01$) (Fig. 6a). Interestingly, only with pre-miR-92a transfections, reduced *DKK3* mRNA levels were




Figure 6. *DKK3* levels after pre-miR-19b or pre-miR-92a transfection relative to control transfection. (a) Protein expression levels of *DKK3* after pre-miR-19b or pre-miR-92a transfection of NB SHEP cells at 24, 48, and 72 hr of transfection. (b) mRNA expression levels of *DKK3* after transfection of NB SHEP cells significantly reduced after pre-miR-92a overexpression. Pre-miR-19b overexpression did not result in reduction of *DKK3* mRNA levels.

observed (Fig. 6b; $p = 0.01$) indicating that each of these two components of the miR-17-92 cluster has a distinct effect on *DKK3* regulation. To the best of our knowledge this is the first example of combined repressive effect on protein expression through targeting of mRNA stability and translation by different members of one miRNA cluster.

Anti-miR-92a transfection restores *DKK3* expression

MYCN is induced in the SHEP MYCN-ER *in vitro* cell line model system by adding 4-OHT to the medium. As expected, transfection of this cell line with an anti-miR-neg ctr before and after MYCN-induction resulted in a significant ($p < 0.01$) decrease in *DKK3* levels upon MYCN-induction (Supporting Information Fig. 1). Anti-miR-92a transfection after MYCN-induction was, however, able to restore the *DKK3* levels. The increase of *DKK3* levels compared to the non-

induced cells line is possibly explained by the fact that anti-miR-92a transfection not only reduces the MYCN-induced miR-92a levels but additionally reduces the endogenous miR-92a levels.

In line with our hypothesis, these results exclude a significant contribution in DKK3 regulation through other MYCN regulated genes.

Discussion

The MYCN transcription factor is known to regulate a broad range of genes affecting various cellular functions implicated in normal development and cancer.² Studies of MYCN-regulated genes have mainly focussed on direct upregulated genes. A few studies also demonstrate a role for MYCN in downregulation of genes but these mechanisms are less well understood. At least for some target genes, MYCN is recruited to the promoter by the DNA binding protein MIZ1 followed by epigenetic silencing, in part due to concomitant action of DNMT3a.¹⁷ Recently, miRNA-mediated silencing has emerged as an alternative mechanism for MYCN-driven downregulation of gene expression. Several MYCN-upregulated miRNAs were identified and have provided evidence that miRNA activation contributes to widespread MYCN-induced mRNA repression.^{6,38} Using *in silico* analyses based on predicted seeds in the 3'UTR of downregulated mRNAs and predicted MYCN downregulated genes from the MYCNot database (<http://medgen.ugent.be/MYCNot>), we provided strong evidence that activated miRNA expression could serve as a mechanism for such a MYCN-mediated mRNA repression. In this study, we present experimental evidence for indirect downregulation of DKK3 through direct MYCN-induced transcriptional activation of the miR-17-92 cluster.

DKK3 is a member of the Dickkopf family together with DKK1, 2, 4 and Soggy. While DKK1, 2 and 4 have a well-established role as WNT-inhibitors, the functions of DKK3 and its homologue Soggy are less firmly established, although recent studies on DKK3 also provide evidence for a role in the canonical WNT pathway.^{35,41} In the study of Tudzarova and colleagues, DKK3 is shown to interact with the betaTrCP which is a negative regulator of beta-catenin, a key component in the WNT signaling pathway. DKK3 has been proposed as a tumour suppressor given the observed reduced expression levels in several tumour entities and tumour suppressing effects upon overexpression in certain cancer cell types. These observations are in line with reduced expression following MYCN overexpression but a relation between DKK3 and WNT signaling could not be established in NB.

Here, we elucidate the mechanism of downregulation of DKK3 as a result of post-transcriptional regulation through the miR-17-92 cluster. Evidence for such an interaction was obtained for two of the six members of the cluster, miR-19b and miR-92a. Although pre-miR-19a was also predicted to target the 3'UTR of DKK3, we could not observe a reduced luciferase output nor reduced DKK3 mRNA or protein levels

after transient transfection (data not shown). For miR-19b and miR-92a, reduced DKK3 protein levels were detected upon a 72-hr overexpression in the MYCN single-copy SHEP NB cell line. Interestingly, decrease of the mRNA levels were only noted after overexpression of miR-92a which could subsequently be restored upon anti-miR-92a transfection. These findings suggest that different members of the same miRNA cluster regulate target gene expression on distinctive levels by effecting mRNA stability or protein translation.

In this study we focussed on miR-17-92 and its components given its direct regulation through MYCN. It should be noted however that other miRNAs, such as members of the miR-17-92 paralog clusters, miR-106b-25 and miR-106a-363, are also predicted to target DKK3.

In addition to the present report, our recent investigation of putative miR-17-92 targets also demonstrated regulation of several components of the TGF- β pathway.³⁸ Together with other previously described target genes, this most probably only represents a small proportion of all *bona fide* target genes which are regulated by this important oncogenic miRNA cluster. It will be of interest to investigate which of these miRNAs and subsequent target genes play a crucial role in the tumour phenotype as a prelude to design new therapies. Also, given the tight regulation of DKK3 and its presumed role as tumour suppressor gene, further investigations towards unravelling the function of this protein are warranted. One possible line of further investigation is the role of DKK3 in relation to control of the so-called origin activation checkpoint, as described in a recent paper by Tudzarova et al.⁴¹ In this study it is shown that DKK3 is a direct target of TP53 which is activated as a consequence of disruption of the origin activation checkpoint leading to G1/S arrest. Reduction of DKK3 levels through MYCN could very well fit with one of the major functions of MYCN as driver of proliferation and progression through the cell cycle.

In conclusion, this study shows a direct regulation of DKK3 mRNA and protein levels by members of the highly oncogenic miR-17-92 cluster. As MYCN has been shown to induce this cluster and because MYCN amplification is a marker for poor prognosis, these results suggest a critical role for DKK3 as tumour suppressor in NB development. Further investigation of the biological function of DKK3 and its role as tumour suppressor gene in NB pathogenesis are therefore of great interest in the light of development of novel therapeutic strategies.

Acknowledgements

The authors thank Fanny De Vloed for her excellent practical work. S.D.B. is supported by an "Emmanuel van der Schueren" grant from the "Vlaamse Liga tegen Kanker", P.M. by the Ghent University Research Fund (BOF 01D31406), F.P.T. and K.D.P. are postdoctoral fellows of the Research Foundation Flanders (F.W.O.) and F.W. was funded by the Bundesministerium für Bildung und Forschung (BMBF, NGFNPlus). The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

References

- Maris JM. Recent advances in neuroblastoma. *N Engl J Med* 2010;362:2202–11.
- Bell E, Chen L, Liu T, Marshall GM, Lunec J, Tweddle DA. MYCN oncprotein targets and their therapeutic potential. *Cancer Lett* 2010;293:144–57.
- Luscher B. Function and regulation of the transcription factors of the Myc/Max/Mad network. *Gene* 2001;277:1–14.
- Weiss WA, Aldape K, Mohapatra G, Feuerstein BG, Bishop JM. Targeted expression of MYCN causes neuroblastoma in transgenic mice. *EMBO J* 1997;16:2985–95.
- Lutz W, Stohr M, Schurmann J, Wenzel A, Lohr A, Schwab M. Conditional expression of N-myc in human neuroblastoma cells increases expression of alpha-prothymosin and ornithine decarboxylase and accelerates progression into S-phase early after mitogenic stimulation of quiescent cells. *Oncogene* 1996;13:803–12.
- Schulte JH, Horn S, Otto T, Samans B, Heukamp LC, Eilers UC, Krause M, Astrahantseff K, Klein-Hitpass L, Buettner R, Schramm A, Christiansen H, et al. MYCN regulates oncogenic MicroRNAs in neuroblastoma. *Int J Cancer* 2008;122:699–704.
- Tang XX, Zhao H, Kung B, Kim DY, Hicks SL, Cohn SL, Cheung NK, Seeger RC, Evans AE, Ikegaki N. The MYCN enigma: significance of MYCN expression in neuroblastoma. *Cancer Res* 2006;66:2826–33.
- Cetinkaya C, Hultquist A, Su Y, Wu S, Bahram F, Pahlman S, Guzhova I, Larsson LG. Combined IFN-gamma and retinoic acid treatment targets the N-Myc/Max/Mad1 network resulting in repression of N-Myc target genes in MYCN-amplified neuroblastoma cells. *Mol Cancer Ther* 2007;6:2634–41.
- Nakamura M, Matsuo T, Stauffer J, Neckers L, Thiele CJ. Retinoic acid decreases targeting of p27 for degradation via an N-myc-dependent decrease in p27 phosphorylation and an N-myc-independent decrease in Skp2. *Cell Death Differ* 2003;10:230–9.
- Kang JH, Rychahou PG, Ishola TA, Qiao J, Evers BM, Chung DH. MYCN silencing induces differentiation and apoptosis in human neuroblastoma cells. *Biochem Biophys Res Commun* 2006;351:192–7.
- Nara K, Kusafuka T, Yoneda A, Oue T, Sangkhathat S, Fukuzawa M. Silencing of MYCN by RNA interference induces growth inhibition, apoptotic activity and cell differentiation in a neuroblastoma cell line with MYCN amplification. *Int J Oncol* 2007;30:1189–96.
- Woo CW, Tan F, Cassano H, Lee J, Lee KC, Thiele CJ. Use of RNA interference to elucidate the effect of MYCN on cell cycle in neuroblastoma. *Pediatr Blood Cancer* 2008;50:208–12.
- Mac SM, D’Cunha CA, Farnham PJ. Direct recruitment of N-myc to target gene promoters. *Mol Carcinog* 2000;29:76–86.
- Shohet JM, Hicks MJ, Plon SE, Burlingame SM, Stuart S, Chen SY, Brenner MK, Nuchtern JG. Minichromosome maintenance protein MCM7 is a direct target of the MYCN transcription factor in neuroblastoma. *Cancer Res* 2002;62:1123–8.
- Slack A, Chen Z, Tonelli R, Pule M, Hunt L, Pession A, Shohet JM. The p53 regulatory gene MDM2 is a direct transcriptional target of MYCN in neuroblastoma. *Proc Natl Acad Sci USA* 2005;102:731–6.
- Westermann F, Muth D, Benner A, Bauer T, Henrich KO, Oberthuer A, Brors B, Beissbarth T, Vandesompele J, Pattyn F, Hero B, Konig R, et al. Distinct transcriptional MYCN/c-MYC activities are associated with spontaneous regression or malignant progression in neuroblastomas. *Genome Biol* 2008;9:R150.
- Iraci N, Diolaiti D, Papa A, Porro A, Valli E, Gherardi S, Herold S, Eilers M, Bernardoni R, Della Valle G, Perini G. A SPI1/MIZ1/MYCN repression complex recruits HDAC1 at the TRKA and p75NTR promoters and affects neuroblastoma malignancy by inhibiting the cell response to NGF. *Cancer Res* 2011;71:404–12.
- Bell E, Lunec J, Tweddle DA. Cell cycle regulation targets of MYCN identified by gene expression microarrays. *Cell Cycle* 2007;6:1249–56.
- Glinka A, Wu W, Delius H, Monaghan AP, Blumenstock C, Niehrs C. Dickkopf-1 is a member of a new family of secreted proteins and functions in head induction. *Nature* 1998;391:357–62.
- Kawano Y, Kypta R. Secreted antagonists of the Wnt signalling pathway. *J Cell Sci* 2003;116:2627–34.
- Mao B, Niehrs C. Kremen2 modulates Dickkopf2 activity during Wnt/LRP6 signaling. *Gene* 2003;302:179–83.
- Wu W, Glinka A, Delius H, Niehrs C. Mutual antagonism between dickkopf1 and dickkopf2 regulates Wnt/beta-catenin signalling. *Curr Biol* 2000;10:1611–14.
- Koppen A, Ait-Aissa R, Koster J, Ora I, Bras J, van Sluis PG, Caron H, Versteeg R, Valentijn LJ. Dickkopf-3 expression is a marker for neuroblastic tumor maturation and is down-regulated by MYCN. *Int J Cancer* 2008;122:1455–64.
- Revet I, Huizenga G, Koster J, Volckmann R, van Sluis P, Versteeg R, Geerts D. MSX1 induces the Wnt pathway antagonist genes DKK1, DKK2, DKK3, and SFRP1 in neuroblastoma cells, but does not block Wnt3 and Wnt5A signalling to DVL3. *Cancer Lett* 2010;289:195–207.
- Kobayashi K, Ouchida M, Tsuji T, Hanafusa H, Miyazaki M, Namba M, Shimizu N, Shimizu K. Reduced expression of the REIC/Dkk-3 gene by promoter-hypermethylation in human tumor cells. *Gene* 2002;282:151–8.
- Lodygin D, Epanchintsev A, Menssen A, Diebold J, Hermeking H. Functional epigenomics identifies genes frequently silenced in prostate cancer. *Cancer Res* 2005;65:4218–27.
- Roman-Gomez J, Jimenez-Velasco A, Agirre X, Castillejo JA, Navarro G, Barrios M, Andreu EJ, Prosper F, Heiniger A, Torres A. Transcriptional silencing of the Dickkopf-3 (Dkk-3) gene by CpG hypermethylation in acute lymphoblastic leukaemia. *Br J Cancer* 2004;91:707–13.
- Tsuji T, Nozaki I, Miyazaki M, Sakaguchi M, Pu H, Hamazaki Y, Iijima O, Namba M. Antiproliferative activity of REIC/Dkk-3 and its significant down-regulation in non-small-cell lung carcinomas. *Biochem Biophys Res Commun* 2001;289:257–63.
- Abarzua F, Sakaguchi M, Takaishi M, Nasu Y, Kurose K, Ebara S, Miyazaki M, Namba M, Kumon H, Huh NH. Adenovirus-mediated overexpression of REIC/Dkk-3 selectively induces apoptosis in human prostate cancer cells through activation of c-Jun-NH2-kinase. *Cancer Res* 2005;65:9617–22.
- Hoang BH, Kubo T, Healey JH, Yang R, Nathan SS, Kolb EA, Mazza B, Meyers PA, Gorlick R. Dickkopf-3 inhibits invasion and motility of Saos-2 osteosarcoma cells by modulating the Wnt-beta-catenin pathway. *Cancer Res* 2004;64:2734–9.
- Hsieh SY, Hsieh PS, Chiu CT, Chen WY. Dickkopf-3/REIC functions as a suppressor gene of tumor growth. *Oncogene* 2004;23:9183–9.
- Kawano Y, Kitaoka M, Hamada Y, Walker MM, Waxman J, Kypta RM. Regulation of prostate cell growth and morphogenesis by Dickkopf-3. *Oncogene* 2006;25:6528–37.
- Kuphal S, Lodermeier S, Bataille F, Schuierer M, Hoang BH, Bosserhoff AK. Expression of Dickkopf genes is strongly reduced in malignant melanoma. *Oncogene* 2006;25:5027–36.
- Kurose K, Sakaguchi M, Nasu Y, Ebara S, Kaku H, Kariyama R, Arai Y, Miyazaki M, Tsushima T, Namba M, Kumon H, Huh NH. Decreased expression of REIC/Dkk-3

- in human renal clear cell carcinoma. *J Urol* 2004;171:1314–8.
35. Lee EJ, Jo M, Rho SB, Park K, Yoo YN, Park J, Chae M, Zhang W, Lee JH. Dkk3, downregulated in cervical cancer, functions as a negative regulator of beta-catenin. *Int J Cancer* 2009;124:287–97.
 36. Tsuji T, Miyazaki M, Sakaguchi M, Inoue Y, Namba M. A REIC gene shows down-regulation in human immortalized cells and human tumor-derived cell lines. *Biochem Biophys Res Commun* 2000;268:20–4.
 37. Mestdagh P, Fredlund E, Pattyn F, Schulte JH, Muth D, Vermeulen J, Kumps C, Schlierf S, De Preter K, Van Roy N, Noguera R, Laureys G, et al. MYCN/c-MYC-induced microRNAs repress coding gene networks associated with poor outcome in MYCN/c-MYC-activated tumors. *Oncogene* 2010;29:1394–404.
 38. Mestdagh P, Bostrom AK, Impens F, Fredlund E, Van Peer G, De Antonellis P, von Stedingk K, Ghesquiere B, Schulte S, Dews M, Thomas-Tikhonenko A, Schulte JH, et al. The miR-17-92 microRNA cluster regulates multiple components of the TGF-beta pathway in neuroblastoma. *Mol Cell* 2010;40:762–73.
 39. Lefever S, Vandesompele J, Speleman F, Pattyn F. RTPrimerDB: the portal for real-time PCR primers and probes. *Nucleic Acids Res* 2009;37:D942–5.
 40. Clevers H. Wnt/beta-catenin signaling in development and disease. *Cell* 2006;127:469–80.
 41. Tudzarova S, Trotter MW, Wollenschlaeger A, Mulvey C, Godovac-Zimmermann J, Williams GH, Stoeber K. Molecular architecture of the DNA replication origin activation checkpoint. *EMBO J* 2010;29:3381–94.