

**Abstract**

This paper seeks to catalyse a reappraisal of the nature, fate and biological significance in humans of phenols, polyphenols and tannins (PPT) consumed in normal diets, and in particular questions the primacy of PPT radical-scavenging mechanisms for the supposed health benefits of diets rich in fruits and vegetables. PPT are classified by structure and function. Arguments are presented to show that cinnamates and derived polyphenols make significantly larger contributions to the total PPT intake than the flavonols and flavones upon which the vast majority of attention has been focussed previously. Daily intakes of total PPT may range from less than 100 mg to in excess of 2 g, and the critical importance of coffee and black tea as the major dietary sources is shown. Only some 5% of the dietary PPT is absorbed in the duodenum, and of this only some 5%, mainly flavanols, reaches the plasma unchanged, the balance being mammalian conjugates. Over 95% of the intake passes to the colon and is fermented by the gut microflora. A fraction of the resulting microbial metabolites is absorbed and appears in the plasma primarily as mammalian conjugates. Even following high intakes of PPT, the plasma metabolites collectively make a very small (less than 5%) and transient contribution to the total concentration of redox active substances in plasma. This explains the failure of

most studies that sought to detect an increase in plasma antioxidant power after consuming a PPT-rich meal or supplement. The powerfully antioxidant PPT aglycones, much used in *in vitro* studies, do not reach the plasma. The redox potential of those unchanged PPT and PPT metabolites that reach the plasma enables them to scavenge damaging radicals, but the endogenous plasma antioxidants, especially ascorbate, are required for disposal of the resultant phenoxyl radicals. Black tea and coffee, the major sources of PPT, are poor sources of ascorbate. It is suggested that if diets rich in fruits and vegetables are health-promoting, and if these effects are due to PPT, then alternatives to radical-scavenging mechanisms must be sought. Evidence is presented to show that some mammalian metabolites of PPT may indeed be able to protect the vascular endothelium and that diets rich in PPT may in humans at normal dietary levels have the ability to protect against Type II diabetes and the metabolic syndrome through effects on glucose absorption and associated hormones. Such effects are recommended for further investigation.

**Key words**

Absorption · antioxidants · black tea · chlorogenic acids · cinnamates · coffee · derived polyphenols · diabetes · flavonoids · fruit · GIP · GLP1 · gut microflora · hydroxycinnamates · insulin · metabolism · phenols · polyphenols · tannins · review · vegetables

**Affiliation**

Centre for Nutrition & Food Safety, School of Biomedical & Molecular Sciences, University of Surrey, Guildford, UK

**Correspondence**

M. N. Clifford · Centre for Nutrition & Food Safety · School of Biomedical & Molecular Sciences · University of Surrey · Guildford · Surrey GU2 7XH · UK · Fax: +44-1483-300374 · E-mail: M.Clifford@Surrey.ac.uk

Received March 5, 2004 · Accepted June 4, 2004

**Bibliography**

Planta Med 2004; 70: 1103–1114 · © Georg Thieme Verlag KG Stuttgart · New York  
DOI 10.1055/s-2004-835835  
ISSN 0032-0943

## Introduction

Simple phenols, polyphenols and tannins (PPT) have been of great interest for many years, in part because of their impact on the colour, odour and flavour of foods and beverages [1], but more recently because of the possibility that these substances may have health-protecting properties [2], [3], [4], [5], [6], [7], [8], [9], [10], [11], [12], [13]. PPT may be classified in several ways, for example, by biosynthetic origin, occurrence, function or effect, or structure [1], [12], [14]. A classification based on structure and function will be used in this paper [15]. Simple phenols are substances containing only one aromatic ring and bearing at least one phenolic hydroxy group and possibly other substituents, whereas polyphenols contain more than one such aromatic ring. Phenols and polyphenols may occur as unconjugated aglycones or as conjugates, frequently with sugars, organic acids, amino acids, lipids, etc. [16].

## The Diversity of Dietary Phenols, Polyphenols and Tannins

The commonest simple phenols are cinnamates that have a C<sub>6</sub>-C<sub>3</sub> structure [17], [18] accompanied by C<sub>6</sub>-C<sub>2</sub> and C<sub>6</sub>-C<sub>1</sub> compounds, and a few unsubstituted phenols [19], [20], [21]. In general these occur as conjugates. Flavonoids are the most extensively studied polyphenols, all characterised by a C<sub>6</sub>-C<sub>3</sub>-C<sub>6</sub> structure, subdivided by the nature of the C<sub>3</sub> element into anthocyanins, chalcones, dihydrochalcones, flavanols, flavanones, flavones, flavonols, isoflavones and proanthocyanidins. The flavanols and proanthocyanidins generally occur unconjugated but the others normally occur as glycosides. Since the seminal paper of Hertog et al. [22] there has been a tendency to think of dietary PPT as encompassing only the flavonoids, and the flavonoids *per se* to consist only of the three flavonols and two flavones that featured in that study, but this is misleading and was never intended. It is not possible to say with precision just how many individual PPT occur regularly in human diets, but on present evidence a figure in excess of 200 seems reasonable [16].

The term "tannin" refers historically to crude plant preparations that are capable of converting hides to leather [23] and such preparations are not consumed as human food. However, the functional polyphenols contained therein at high concentrations may also occur in certain foods and beverages but at comparatively low concentrations that would render them totally ineffective for producing leather. These polyphenols may be subdivided into the flavonoid-derived proanthocyanidins (condensed tannins) [24] and the gallic acid-derived and ellagic acid-derived hydrolysable tannins, this latter subgroup being of more restricted occurrence in human food (but commoner in some animal feeds) [25]. The phloroglucinol-derived phlorotannins, while never used for preparing leather, also have a limited occurrence in human food [19]. The more recent term "phytoestrogen" refers to substances with oestrogenic and/or anti-androgenic activity at least *in vitro*, and encompasses some isoflavones, some stilbenes, some lignans and some coumarins [26]. The lignans are not oestrogen-active until transformed by the gut microflora [26], [27]. "Antioxidants" is a third function-based term much used to describe PPT, but individual compounds differ markedly in their ability to scavenge reactive oxygen species and reactive nitrogen

species, and inhibit oxidative enzymes. Mammalian metabolites of PPT do not necessarily retain fully the antioxidant ability of the PPT found in plants and especially not that of their aglycones as commonly tested *in vitro* [28], [29].

The PPT discussed above are substances found in healthy and intact plant tissues, and in the main are of known structure. However, many traditional foods and beverages as consumed have been produced by more or less extensive processing of such plant tissues, resulting in biochemical or chemical transformations of the naturally-occurring PPT. In some cases, black tea, matured red wines, and coffee beverage, for example, these transformations may be substantial, generating large quantities of substances not found in the original plant material. Despite significant advances in the last decade [30], [31], [32], [33], [34], [35], [36], [37], [38], [39], [40], [41], the structures of the majority of these novel compounds have yet to be elucidated. Although often described as tannins, these substances are not functional tanning agents, and should be referred to collectively as derived polyphenols until such time as their full structural characterisation permits a more precise nomenclature.

## The Consumption of PPT

There have been several attempts to estimate the quantities of PPT consumed, either by using diet diaries or food frequency questionnaires and data on the typical composition of individual commodities [3], [8], [42], [43], [44], [45], [46], [47], [48], [49], [50], or by diet analysis [6], [22], [51]. In comparison with the comprehensive databases providing the content in the diet of the established micro- and macro-nutrients, data for the contents of PPT are much more limited. Those data available for PPT content have been obtained by many different methods of analysis, rarely take account of the effects of agricultural practice, season, cooking or commercial processing, are not necessarily just for the edible portion, and may be for varieties of fruit and vegetable different from those consumed in a particular diet under investigation [45], [46], [52].

These are potentially serious limitations since quantitatively cultivars may differ substantially in composition, and the non-edible parts of fruits and vegetables may differ greatly both quantitatively and qualitatively, compared with the flesh or juice [53]. In addition, domestic cooking and commercial processing may in some cases cause extensive leaching and destruction [54], [55], [56], [57], [58], [59], [60].

Data based upon analysis of particular diets avoid these limitations but are usually restricted to a few PPT because of the difficulties and cost associated with quantifying so many individual compounds of known structure, to say nothing of the serious difficulties associated with quantifying the uncharacterised derived polyphenols [61]. When such data are available, they are usually for PPT as aglycones released by hydrolysis (to simplify the analysis still further) and generally for the flavonols and flavones first studied by Hertog et al. [22] since these are amongst the easiest to determine [44], [47], [48], [51], [52], [62], [63], [64], [65], [66], [67], [68]. There are more limited data for flavanones [44], [64] and isoflavones after hydrolysis

[44], [52], and flavanols, and proanthocyanidins [6], [69] (which occur as aglycones).

In an attempt at modest cost to shed more light on the nature and quantity of PPT consumed by various populations we developed a computerised spreadsheet database beginning with the several hundred papers identified in the NEODIET reviews [16] with continued updating as more information has been published or found [17], [52], [70], [71], [72], [73], [74], [75]. The database covers 80 commodities, including five alcoholic beverages, six fruit juices and three other non-alcoholic beverages, and 14 PPT subgroups including derived polyphenols, with every entry labelled to show the paper(s) from which the information was taken. Data sources have been restricted to papers using specific methods of analysis: data for “total phenols by Folin-Ciocalteu” or “total antioxidant power as gallic acid equivalents” and similar, have been excluded. In order to reflect the variability in composition, data have always been entered as “high” and “low” values (mean  $\pm 2\sigma$  wherever possible) and an overall mean for each commodity used in determining the amounts of PPT consumed.

While recognising the limitations (discussed above) of such an approach to estimating diet composition and the intake of PPT, using this database in conjunction with diet diaries available from our other studies [76], [77], [78] has produced interesting data (Table 1) and insights.

From Table 1 it is clear that PPT intakes may vary substantially, and that the flavones and flavonols, upon which most emphasis has so far been placed [22], [44], [47], [48], [51], [52], [62], [63], [64], [65], [66], [67], [68], form a comparatively small part of the

total intake for the populations studied. The relatively low consumption of chalcones and dihydrochalcones, isoflavones, anthocyanins, and stilbenes reflects the comparatively low consumption of apples and ciders, soya products, dark berries and red wines by these populations. The significant contributions made by the hydroxycinnamates (in these populations primarily reflecting coffee consumption [17], [18]) and derived polyphenols (in these populations primarily reflecting black tea consumption [79], [80], [81]) are striking. In this context “black tea” refers to the beverage prepared from the fermented leaf (as distinct from green tea) and not to the addition or otherwise of milk to the beverage prior to consumption. This domination by PPT from black tea and coffee indicates the importance also of considering the hydroxycinnamates and derived polyphenols whenever assessing the dietary significance of PPT, and clearly shows the limitations of looking only at flavonols and flavones after hydrolysis no matter how precise *per se* the data for these aglycones might be.

It is important to stress that data for the composition of black tea and coffee beverage reflect exactly what is consumed (with the exception of the dregs left in the cup) since all transformations associated with processing and domestic preparation have already taken place. Moreover, the NEODIET database is replete with analytical data from numerous sources for the composition of these beverages (thus better avoiding extreme values associated with any peculiarity of the material analysed or method of analysis) compared with data for many fruits and vegetables. Accordingly, the estimated consumption figures obtained using this database are likely to be more accurate than would have been the case if solid foods were the major sources of PPT, and data were for raw rather than after cooking/processing. This ar-

Table 1 Mean dietary intakes of 14 classes of PPT as determined from diet-diaries and a food composition data base

PPT	103 UK females aged 20–30 years <sup>a</sup>		50 UK males aged 27–57 years <sup>b</sup>	
	Estimated as conjugates	Estimated as aglycones <sup>c</sup>	Estimated as conjugates	Estimated as aglycones <sup>c</sup>
Total, range	100–2300		30–2200	
Total, mean	780	451	1058	598
Hydroxybenzoates	15		23	
Hydroxycinnamates	353	176	670	335
Total flavonoids	210	105	205	103
Anthocyanins	5		9	
Chalcones and dihydrochalcones	0.7			
Flavanols	64		58	
Flavanones	22		89	
Flavones	72		17	
Flavonols	35		26	
Isoflavones	9		0.13	
Proanthocyanidins	7		6	
Ellagitannins	23			
Derived polyphenols	170	170	160	160
Stilbenes	9			
Lignans	0.04			

<sup>a</sup> Ref. [49].

<sup>b</sup> Ref. [50].

<sup>c</sup> Aglycones are estimated approximately by taking rutin as a representative flavonoid and 5-caffeoylquinic acid as a representative hydroxycinnamate and adjusting for the relevant molecular masses.

gument applies also to the data for PPT delivered by wines and juices. Using this approach has led us to estimate typical mean intakes of PPT for the two populations so far studied to be in the range 450 to 600 mg as aglycones.

### Absorption and Metabolism of PPT

Extensive studies in humans and animals have indicated that some PPT can be absorbed in the small intestine, for example, certain cinnamate conjugates [82], [83], flavanols [84] (that occur naturally as aglycones), and certain flavonoid glucosides [85], [86] (but not the corresponding flavonoid rutosides [87]). The mechanisms of absorption have not been completely elucidated but involve *inter alia* interaction of certain glucosides with the active sugar transporter (SGIT1) and luminal lactase-phloridzin hydrolase, passive diffusion of the more hydrophobic aglycones, and interaction with cytosolic  $\beta$ -glucosidase. Although varying with PPT subclass and matrix, when expressed relative to the total intake of PPT, only some 5 to 10% of the amount consumed is absorbed at this site. The major part of that absorbed (90 to 95% for every substance so far studied) enters the circulation as mammalian conjugates produced by a combination of methylation, sulphate conjugation, glucuronide conjugation and, in the case of some phenolic acids, also by glycine conjugation [29]. Only a very small amount of the total PPT consumed, maximally 5 to 10%, enters the plasma unchanged.

The 90 to 95% of the total PPT ingested, plus any mammalian glucuronides excreted through the bile, pass to the colon where they are metabolised by the gut microflora. Transformations may be extensive, and include the removal of sugars, removal of phenolic hydroxyl groups, fission of aromatic rings, and metabolism to carbon dioxide, possibly via oxaloacetate [88]. A substantial range of microbial metabolites has been identified, including phenols and aromatic/phenolic acids/lactones possessing 0, 1 or 2 phenolic hydroxyl groups and up to five carbons in the side chain [89], [90], [91], [92], [93], [94], [95], [96], [97], [98], [99], [100], [101], [102], [103], [104], [105], [106], [107], [108], [109], [110], [111]. *Eubacterium* is of particular interest since this species not only metabolises dietary (poly)phenols [99], [100], [105], [106], [112], [113], [114], [115], [116], but also produces butyrate [117], a preferred energy source for colonic epithelial cells thought to play an important role in maintaining colon health in humans. The yield of phenolic/aromatic acids is variable (up to  $\times 10$ ) between individuals, but can be substantial (up to 50%) relative to the intake of PPT substrates [95], [103], [104], [108], [109], [110].

There is evidence from cell culture studies that some of the aromatic/phenolic acids, e.g., benzoic, salicylic, *p*-coumaric and ferulic acids, are transported actively by the monocarboxylate transporter MCT1 [118], [119], [120], [121], [122]. A comparatively small percentage of these microbial metabolites may eventually appear unchanged in plasma or urine but the majority is subject to mammalian conjugation as described for intact PPT.

Table 2 summarises in a semi-quantitative manner so far as current knowledge allows the fate of a "typical" daily consumption of some 450 to 600 mg of PPT (as aglycones) previously defined in Table 1.

### Plasma Pseudo-Pharmacokinetics

Since for the majority of dietary PPT, neither the conjugates consumed, nor their free aglycones, are detectable in plasma, it is rarely possible to perform true pharmacokinetic analyses. Most so-called pharmacokinetic data that have been published relate to the concentrations of aglycones released after hydrolysis of mammalian conjugates in plasma or urine with commercial  $\beta$ -glucuronidase and/or sulphatase, and the data so obtained are better referred to as pseudo-pharmacokinetics. Published data are summarised in Table 3. Although the maximum concentration achieved transiently varies to some extent with PPT subclass and matrix in which consumed, it is unlikely that plasma metabolite concentrations will routinely exceed 10  $\mu$ M in total, and approximately 1  $\mu$ M for total aglycones. The reported  $T_{\max}$  values range from 1 to 2.5 hours for substances absorbed in the duodenum [85], [86], [87], [123], [124], [125], [126], [127], [128], [129], [130], up to 5 to 12 hours when microbial metabolism is a prerequisite [87], [104], [131]. Elimination half-lives are very variable, ranging from as low as 1 hour [132], [133] to values in excess of 20 hours [85], [6], [87]. The very low values may be artefacts of observation periods being less than the true half-life, whereas the very high values may be exaggerated because of a biphasic elimination reflecting significant entero-hepatic circulation of glucuronides. When studied explicitly, repeat dosing has failed to provide evidence of accumulation in plasma suggesting that, in general, significant elimination occurs in a time period shorter than the interval between repeat doses [123].

Table 4 summarises the concentrations of a range of endogenous (i.e., non-dietary) simple phenols, including  $\alpha$ -tocopherol, and ascorbate in plasma from healthy individuals. The total simple phenol and ascorbate concentration is between 159 and

Table 2 Fate of ingested PPT

	Aglycones (mg)
Estimated mean daily consumption (from Table 1)	450–600
~ 5–10% of intake absorbed in duodenum and excreted in urine. Of this	22–60
5–10% unchanged plant (poly)phenols, and	< 6
90–95% mammalian conjugates	20–55
~90–95% fermented in colon (unabsorbed PPT+ enteric and entero-hepatic cycling of glucuronides, etc.)	400–570
Poorly-defined and very variable portion (5 to 50%) absorbed depending on individual's flora and substrates.	
Mainly mammalian conjugates of microbial metabolites	20–285

Table 3 Plasma pseudo-pharmacokinetics after consumption of normal portions of rich sources

PPT Subclass	$C_{max}$ (nM) unchanged <sup>a</sup>	$C_{max}$ (nM) mammalian conjugates	% urine excretion
Anthocyanins	10–150	traces	N.D. - 0.1 <sup>b</sup>
Flavanols, low fat	40–140	1000–2000	0.5–4.0
Flavanols, high fat	150–220	up to 6200	25–30
Flavonol glycosides	Minute traces	N.D. <sup>b</sup>	0.5–2.5
Flavonol aglycones	Minute traces	350–1100	
Flavanone glycosides	Minute traces	120–1500	4–10
Isoflavone glycosides	Minute traces	900–4000	20–55
Isoflavone aglycones	10–150	500–6500	
Cinnamates & chlorogenic acids	up to 120	up to 500	1–2
Phenolic gut flora metabolites		20–60	Up to 50
<b>Hypothetical total if all consumed in one meal</b>	<b>250–780</b>	<b>2890–21660</b>	

<sup>a</sup>  $C_{max}$  = maximum concentration achieved transiently in plasma.

<sup>b</sup> N.D. = not detected.

Table 4 Plasma concentrations ( $\mu$ M) of endogenous (non-dietary) phenols and other plasma antioxidants

	Plasma concentration, healthy individuals
Homogentisic acid	0.014–0.070 <sup>a</sup>
<i>p</i> -Hydroxyphenyl lactate	40–90 <sup>b</sup>
<i>p</i> -Hydroxyphenyl pyruvate	14–60 <sup>b</sup>
Tyrosine	60–130 <sup>b,c</sup>
Ascorbate	40–70 <sup>d,e</sup>
$\alpha$ -Tocopherol	5–30 <sup>f</sup>
<b>Total endogenous phenols &amp; antioxidants</b>	<b>159–380</b>
Hypothetical total diet-derived phenols	3.1–22.4 <sup>g</sup>
Averaged over three meals gives a transient increase of between 0.3 and 5%.	$\approx$ 1–7.5 <sup>g</sup>
Many people consume much less	

<sup>a</sup> Ref. [201], <sup>b</sup> Ref. [202], <sup>c</sup> Ref. [203], <sup>d</sup> Ref. [204], <sup>e</sup> Ref. [205], <sup>f</sup> Ref. [206], <sup>g</sup> From Table 3

380  $\mu$ M. The maximum additional concentration that is likely to be achieved from dietary sources, 3 to 22  $\mu$ M, is marginal by comparison adding only between 0.3 and 5% if it is assumed, quite reasonably, that the “typical” mean intake is taken over three equal meals. Many people consume a much smaller quantity of dietary PPT and even those consuming double the average amount (450 to 600 mg aglycones) adopted in this paper will only achieve a transient 5 to 10% increase in total plasma antioxidant content.

Many investigators have attempted to demonstrate increases in plasma antioxidant capability following the consumption of foods, beverages or supplements rich in PPT. Table 5 summarises the outcomes of 34 such studies [127], [130], [131], [132], [133], [134], [135], [136], [137], [138], [139], [140], [141], [142], [143], [144], [145], [146], [147], [148], [149], [150], [151], [152], [153], [154], [155], [156], [157], [158], [159]. The test substances included a range of fruit and vegetable products, including juices, alcoholic beverages, tea, and chocolate. In view of the calculations presented in Tables 3 and 4, it is perhaps not surprising that increases in plasma antioxidant capacity were often undetectable, and at best, small and transient. Moreover, in four studies that

produced increases in plasma antioxidant capability it could be attributed, at least in part, to increased plasma ascorbate [149], [156], [158].

In view of these observations, it is instructive also to consider the redox potentials of PPT-derived mammalian metabolites that are known to reach plasma, and to compare these with the corresponding values for the endogenous plasma antioxidants. The polyphenols with the lowest redox potentials are flavonoids with vicinal hydroxyl groups in the B-ring, and conjugation extending to the A-ring, e.g., quercetin aglycone (330 mV at pH 7) [160]. If the conjugation does not extend beyond the B-ring, then the redox potential is significantly higher even for (–)-epigallocatechin gallate (480 mV at pH 7) [161] with three vicinal hydroxyl groups. The value rises again when there are only two vicinal hydroxyl groups {e.g., (+)-catechin 570 mV [162] or caffeic acid 540 mV [162]}, a single *para* hydroxyl group (e.g., hesperidin 720 mV [162]) or isolated (*meta*) hydroxyl groups (e.g., resorcinol 810 mV [163]). These comparisons are extended to the endogenous (non-dietary) plasma antioxidants in Table 6. Fig. 1 illustrates the marked effects of mammalian and microbial metabolism on the redox potential of PPT aglycones that are frequently examined in *in vitro* systems designed to demonstrate their potent antioxidant properties. When the aglycones of such powerful antioxidants are given intravenously to humans [164] or intraperitoneally to animals [165], [166], thus circumventing the protection

Table 5 The outcome of 34 studies<sup>a</sup> in which volunteers were given foods, beverages or supplements rich in PPT and plasma was analysed for total antioxidant activity


34 studies, (poly)phenol-rich diet compared with control
– 13 studies (3 high & 3 very high doses) showed no change in plasma antioxidant status <i>ex vivo</i>
– 21 studies (8 low & 7 moderate doses) showed small and transient increases in plasma antioxidant status <i>ex vivo</i>
– 1 showed reduction in plasma Vitamin E
– 1 showed reduction in plasma ascorbate and glutathione

<sup>a</sup> Refs: [127], [130], [131], [132], [133], [134], [135], [136], [137], [138], [139], [140], [141], [142], [143], [144], [145], [146], [147], [148], [149], [150], [151], [152], [153], [154], [155], [156], [157], [158], [159].

**Table 6** A summary of published data for transient maximal plasma concentrations of diet-derived (poly)phenols, typical plasma concentrations of endogenous phenols and antioxidants, and associated redox potentials (pH 7)

Mammalian metabolite hydroxylation pattern	Maximal transient concentration ( $\mu\text{M}$ )	Redox potential (mV) at pH 7
1,2,3- <i>vic</i>	0.14 <sup>a</sup>	400–600 <sup>d,e,f</sup>
1,2- <i>vic</i>	0.8 <sup>a</sup>	500–650 <sup>d,g,h</sup>
Single <i>para</i> or isolated <i>meta</i> hydroxy groups	10 <sup>a</sup>	700–1050 <sup>d,e,g,i,j,k</sup>
Blocked/inactive	?	Inactive
Damaging radicals		Redox potential (mV) at pH 7
Hydroxyl radical		2310 <sup>h</sup>
Superoxide radical anion		1800 <sup>h</sup>
Alkoxy radical		1600 <sup>l</sup>
Alkyl-peroxy radical		1000 $\pm$ 60 <sup>h,l,m</sup>
PUFA (bis-allylic) radical		600 <sup>h</sup>
Endogenous phenols and antioxidants in plasma		Redox potential (mV) at pH 7
Endogenous <i>p</i> -phenols		$\approx$ 700 <sup>d,e,g,i,j,k</sup>
$\alpha$ -Tocopherol		$\approx$ 500 <sup>d,h</sup>
Ascorbate (depleted)		$\approx$ 280 <sup>e,h</sup>
Glutathione		– 276 <sup>n</sup>

<sup>a</sup> From Table 3, <sup>b</sup> From Table 4, <sup>c</sup> Ref. [170], <sup>d</sup> Ref. [161], <sup>e</sup> Ref. [163], <sup>f</sup> Ref. [207], <sup>g</sup> Ref. [162], <sup>h</sup> Ref. [208], <sup>i</sup> Ref. [209], <sup>j</sup> Ref. [210], <sup>k</sup> Ref. [211], <sup>l</sup> Ref. [212], <sup>m</sup> Ref. [213], <sup>n</sup> Ref. [214]



**Fig. 1** Illustration of the effects of mammalian metabolism and microbial metabolism on the redox potential of (poly)phenols found in plasma compared with their precursors in the diet and the aglycones commonly used in *in vitro* studies.

offered by the gastric mucosa and Phase II conjugations, redox cycling causes serious and possibly fatal liver and kidney damage. One may conclude that it is better to avoid high plasma concentrations of the more potent PPT antioxidants (such as unconjugated quercetin), and that it might be ill-advised grossly to supplement normal diets with capsules and concentrates of such potent antioxidants.

Table 6 indicates that the diet-derived PPT metabolites are able thermodynamically to scavenge some or all of the damaging radicals should they come into contact. However, these metabolites are so hydrophilic {e.g., quercetin 3-glucuronide ( $K = 0.008$ )

[167], [168] compared with quercetin ( $K = 66$ ) [167], [168] and  $\alpha$ -tocopherol ( $K = 550$ ) [169]] that it is unlikely they will encounter lipid-derived radicals. However, any phenoxyl radicals generated will have to be removed either by transfer of the unpaired electron to an endogenous scavenger such as  $\alpha$ -tocopherol, ascorbate, glutathione or serum albumin, or by dismutation or disproportionation although these latter mechanisms seem somewhat unlikely *in vivo* because of the relatively low phenoxyl radical concentrations. The implied demand for  $\alpha$ -tocopherol and ascorbate is particularly interesting, since two of the supplementation studies (Table 5) produced reductions in plasma ascorbate and  $\alpha$ -tocopherol [150], and the major sources of dietary PPT identified from the NEODIET database (coffee and black tea) supply neither. Moreover, it is known that for approximately 14% of the over-65 population subgroup in the UK the mean plasma ascorbate value is below 11  $\mu\text{M}$  [170], indicating biochemical depletion [171], suggesting that for heavy consumers of black tea or coffee within this population subgroup the transient concentration of PPT metabolites may approach or even exceed plasma ascorbate.

From the data assembled, it is difficult to envisage how diet-derived PPT metabolites can make a major contribution to radical scavenging in plasma compared with the contribution to be expected from the endogenous antioxidants in healthy individuals replete in ascorbate. It follows that if diets rich in fruits and vegetables are advantageous, at least in part, by virtue of their content of PPT then mechanisms other than radical scavenging are implicated.

### Protective Mechanisms other than Radical Scavenging

Although classically, mammalian conjugates of drugs are viewed as biologically significantly less active than the parent drug, this is not inevitably the case when PPT are considered. Some quercetin conjugates are able to inhibit lipoxygenase and xanthine oxidase. Quercetin 3-glucuronide, one of the three major human conjugates of dietary quercetin glycosides,

has been shown *in vitro* to protect the vascular endothelium [172], [173] and suppresses peroxy-nitrite-induced consumption of lipophilic antioxidants in human LDL [174]. Another human metabolite, quercetin 4'-glucuronide, inhibits xanthine oxidase *in vitro* at a concentration in plasma that on normal diets can realistically be approached ( $K_i = 0.25 \mu\text{M}$ ) [175]. As these observations are more widely appreciated, and mammalian metabolites become more readily available, it is quite possible that other biologically interesting properties will be identified for mammalian conjugates of PPT.

Effects occurring in the gastro-intestinal tract prior to absorption also deserve greater consideration. For example, there is a growing body of evidence suggesting that diets rich in PPT may influence the absorption and metabolism of glucose, resulting in a lower glycaemic index [176] than would otherwise be expected. Red wine [177], coffee [178] and apple juice [179] have all been shown in controlled volunteer studies to slow glucose absorption and reduce the post-prandial surge in plasma glucose, an event known to be an independent risk factor for CHD [180]. Studies in which volunteers consumed normal portions of PPT-rich foods [178] have also produced reductions in the post-prandial concentrations of plasma insulin and glucose-dependent insulinotropic polypeptide and elevation in the concentration of glucagon-like polypeptide-1. A prospective study of 17,000 people suggested that the mean relative risk of developing Type II diabetes was only 0.5 (0.35–0.72) in those individuals habitually consuming six or more cups of coffee per day compared with those consuming two or less ( $p = 0.0002$ ) [181], and a polyphenol-enriched diet has been reported to reduce the incidence and severity of nephropathy in Type II diabetics [182].

The reduced glycaemic index has been attributed to PPT-mediated inhibition of  $\alpha$ -amylase [183], [184], maltase [185] or  $\alpha$ -glucosidase (sucrase) [184], [186], but this mechanism would not operate with preformed glucose as observed in our studies [178], [179]. In studies using bolus doses of glucose, the observation is more conveniently explained by an effect on the active glucose transporter (SGLT1) in the duodenum. Phloridzin, a dihydrochalcone glucoside characteristic of apples and apple products [53], but now known to be more widely distributed [187], competes for the active site both *in vitro* and *in vivo* when given intraperitoneally [188], [189], [190], [191]. Other dietary PPT [(–)-epigallocatechin gallate, (–)-epigallocatechin and 5-caffeoylquinic acid] have been shown *in vitro* to dissipate the sodium gradient essential to the operation of SGLT1 [192], [193], and several quercetin glucosides have been shown to interact with it and thus to have the potential to interfere in glucose transport [194], [195], [196], [197], [198], [199], [200]. While these effects on glucose absorption and the associated hormones are modest, they have been achieved in volunteers consuming sensible quantities of common dietary components (as distinct from effects seen only *in vitro* with high levels of PPT aglycones never seen in the diet). Such effects repeated daily, or even several times daily for say 30 years, might in part explain the reduced incidence of chronic disease, especially Type II diabetes and the metabolic syndrome, in later life associated with diets rich in fruits and vegetables.

## Conclusions

Evidence has been presented to indicate that very little of the dietary PPT consumed reaches human plasma, and that this transient fraction contains only weak antioxidants able to make little contribution to the total antioxidant activity of the plasma. This suggests that radical scavenging by PPT is unlikely to be the mechanism by which diets rich in fruits and vegetables protect against chronic diseases. Instead, it is proposed that modulation of glucose absorption in the duodenum prior to the absorption of PPT, with protection by PPT metabolites through mechanisms other than radical scavenging, might over a lifetime offer modest protection against chronic diseases, especially Type II diabetes and the metabolic syndrome.

## References

- Shahidi F, Naczk M. Food Phenolics: Sources; Chemistry; Effects; Applications. Lancaster, Pennsylvania: Technomic Publishing Inc., 1995: pp 1–321
- Higdon JV, Frei B. Tea catechins and polyphenols: health effects, metabolism, and antioxidant functions. *CRC Crit Rev Food Sci Nutr* 2003; 43: 89–143
- Sesso HD, Gaziano JM, Liu S, Buring JE. Flavonoid intake and the risk of cardiovascular disease in women. *Am J Clin Nutr* 2003; 77: 1400–8
- Stewart JR, Artime MC, O'Brian CA. Resveratrol: a candidate nutritional substance for prostate cancer prevention. *J Nutr* 2003; 133: S2440–3
- Kris-Etherton PM, Keen CL. Evidence that the antioxidant flavonoids in tea and cocoa are beneficial for cardiovascular health. *Curr Opin Lipidol* 2002; 13: 41–9
- Arts IC, Jacobs DR, Gross M, Harnack LJ, Folsom AR. Dietary catechins and cancer incidence among postmenopausal women: the Iowa Women's Health Study (United States). *Cancer Causes Control* 2002; 13: 373–82
- Reed J. Cranberry flavonoids, atherosclerosis and cardiovascular health. *CRC Crit Rev Food Sci Nutr* 2002; 42: 301–16
- Sampson L, Rimm E, Hollman PC, de Vries JH, Katan MB. Flavonol and flavone intakes in US health professionals. *J Am Diet Assoc* 2002; 102: 1414–20
- Youdim KA, Spencer JP, Schroeter H, Rice-Evans C. Dietary flavonoids as potential neuroprotectants. *Biol Chem* 2002; 383: 503–19
- Dillard CJ, German JB. Phytochemicals: nutraceuticals and human health. *J Sci Food Agric* 2000; 80: 1744–56
- Riemersma RA, Rice-Evans CA, Tyrrell RM, Clifford MN, Lean MEJ. Tea flavonoids and cardiovascular health. *Quart J Med* 2001; 94: 277–82
- Parr AJ, Bolwell GP. Phenols in the plant and in man. The potential for possible nutritional enhancement of the diet by modifying phenolic content or composition. *J Sci Food Agric* 2000; 80: 985–1012
- Department of Health. Nutritional Aspects of the Development of Cancer. London: Her Majesty's Stationery Office, 1998: pp 1–274
- Ribereau-Gayon P. Plant phenolics. New York: Hafner Publishing Company, 1972
- Clifford MN. The health effects of tea and tea components. Appendix 1. A nomenclature for phenols with special reference to tea. *CRC Crit Rev Food Sci Nutr* 2001; 41: 393–7
- Lindsay DG, Clifford MN. Special issue devoted to critical reviews produced within the EU Concerted Action "Nutritional Enhancement of Plant-based Food in European Trade" (NEODIET). *J Sci Food Agric* 2000; 80: 793–1137
- Clifford MN. Chlorogenic acids and other cinnamates – nature, occurrence and dietary burden. *J Sci Food Agric* 1999; 79: 362–72
- Clifford MN. Chlorogenic acids and other cinnamates – nature, occurrence, dietary burden, absorption and metabolism. *J Sci Food Agric* 2000; 80: 1033–42
- Clifford MN. Miscellaneous phenols in foods and beverages – nature, occurrence and dietary burden. *J Sci Food Agric* 2000; 80: 1126–37
- Soler-Rivas C, Espin JC, Wichers HJ. Oleuropein and related compounds. *J Sci Food Agric* 2000; 80: 1013–23

- 21 Tomás-Barberán FA, Clifford MN. Dietary hydroxybenzoic acid derivatives – nature, occurrence and dietary burden. *J Sci Food Agric* 2000; 80: 1024–32
- 22 Hertog MGL, Hollman PCH, Katan MB. Content of potentially anticarcinogenic flavonoids of 28 vegetables and 9 fruits consumed in The Netherlands. *J Agr Food Chem* 1992; 40: 2379–83
- 23 Haslam E. Practical Polyphenolics. From structure to molecular recognition and physiological action. Cambridge: Cambridge University Press, 1998: pp 1–438
- 24 Santos-Buelga C, Scalbert A. Proanthocyanidins and tannin-like compounds – nature, occurrence, dietary intake and effects on nutrition and health. *J Sci Food Agric* 2000; 80: 1094–1117
- 25 Clifford MN, Scalbert A. Ellagitannins – nature, occurrence and dietary burden. *J Sci Food Agric* 2000; 80: 1118–25
- 26 Cassidy A, Hanley B, Lamuela-Raventós RM. Isoflavones, lignans and stilbenes – origins, metabolism and potential importance to human health. *J Sci Food Agric* 2000; 80: 1044–62
- 27 Setchell KDR, Lawson AM, Borriello SP, Harkness R, Gordon H, Morgan DML, Kirk DN, Adlercreutz H, Anderson LC. Lignan formation in man – microbial involvement and possible roles in relation to cancer. *Lancet* 1981; ii: 4–7
- 28 Diplock AT. Defence against reactive oxygen species. *Free Radic Res* 1999; 29: 4637
- 29 Kroon PA, Clifford MN, Crozier A, Day AJ, Donovan JL, Manach C, Williamson G. COMMENTARY: How should we assess the effects of exposure to dietary polyphenols *in vitro*. *Am J Clin Nutr* 2004; 80: 15–21
- 30 Cameira-dos-Santos P-J, Brillouet J-M, Cheynier V, Moutounet M. Detection and partial characterisation of new anthocyanin-derived pigments in wine. *J Sci Food Agric* 1996; 70: 204–8
- 31 Fulcrand H, Cameira-dos-Santos P-J, Sarni-Manchado P, Cheynier V, Favre-Bonvin J. Structure of new anthocyanin-derived wine pigments. *J Chem Soc Perkin Trans* 1996; 1: 735–9
- 32 Bakker J, Bridle P, Honda P, Kuwano H, Saito N, Terahara N, Timberlake CF. Identification of an anthocyanin occurring in some red wines. *Phytochemistry* 1997; 44: 1375–82
- 33 Bakker J, Timberlake CF. Isolation, identification and characterization of new color-stable anthocyanins occurring in some red wines. *J Agr Food Chem* 1997; 45: 35–43
- 34 Davis AL, Lewis JR, Cai Y, Powell C, Davies AP, Wilkins JPG, Pudney P, Clifford MN. A polyphenolic pigment from black tea. *Phytochemistry* 1997; 46: 1397–402
- 35 Fulcrand H, Benabdelljalil C, Rigaud J, Cheynier V, Moutounet M. A new class of wine pigments yielded by reaction between pyruvic acid and the grape anthocyanins. *Phytochemistry* 1998; 47: 1401–7
- 36 Davies AP, Goodsall C, Cai Y, Davis AL, Lewis JR, Wilkins J, Wan X, Clifford MN, Powell C, Thiru A, Safford R, Nursten HE. Black tea dimeric and oligomeric pigments – structures and formation. In: Gross GG, Hemingway RW and Yoshida T, editors. *Plant Polyphenols 2*. Chemistry, Biology, Pharmacology, Ecology Indian bend, Oregon, USA: Kluwer Academic, 1999: pp 697–724
- 37 Vivar-Quintana AM, Santos-Buelga C, Francia-Aricha E, Rivas-Gonzalo JC. Formation of anthocyanin-derived pigments in experimental red wines. *Food Science and Technology International* 1999; 5: 347–52
- 38 Tanaka T, Betsumiya Y, Mine C, Kouno I. Theanaphthoquinone, a novel pigment oxidatively derived from theaflavin during tea fermentation. *Chem Commun*, 2000: 1365–6
- 39 Mateus N, Silva AM, Santos-Buelga C, Rivas-Gonzalo JC, de F V. Identification of anthocyanin-flavanol pigments in red wines by NMR and mass spectrometry. *J Agr Food Chem* 2002; 50: 2110–6
- 40 Mateus N, de Pascual-Teresa S, Rivas-Gonzalo JC, Santos-Buelga C, de Freitas V. Structural diversity of anthocyanin-derived pigments in port wines. *Food Chem* 2002; 76: 335–42
- 41 Sang S, Tian W, Meng X, Stark RE, Rosen RT, Yang CS, Ho C-T. Theadibenzotropolone A, a new type pigment from enzymatic oxidation of (–)-epicatechin and (–)-epigallocatechin gallate and characterized from black tea using LC/MS/MS. *Tetrahedron Lett* 2002; 43: 7129–33
- 42 Kühnau J. The flavonoids. A class of semi-essential food components: their role in human nutrition. *World Rev Nutr Diet* 1976; 24: 117–91
- 43 Knekt P, Jarvinen R, Reunanen A, Maatela J, Lunetta M, di Mauro M, Crimi S, Mughini L. Flavonoid intake and coronary mortality in Finland: a cohort study. No important differences in glycaemic responses to common fruits in type 2 diabetic patients. *Brit Med J* 1995; 12: 674–8
- 44 Kimira M, Arai Y, Shimoi K, Watanabe S. Japanese intake of flavonoids and isoflavonoids from foods. *J Epidemiol* 1998; 8: 168–5
- 45 Linseisen J, Radtke J, Wolfram G. Flavonoid intake of adults in a Bavarian subgroup of the National Food Consumption survey. *Z Ernährungswiss* 1997; 36: 403–12
- 46 Radtke J, Linseisen J, Wolfram G. Phenolic acid intake of adults in a Bavarian subgroup of the national food consumption survey. *Z Ernährungswiss* 1998; 37: 190–7
- 47 Hirvonen T, Pietinen P, Virtanen M, Ovaskainen ML, Hakkinen S, Albanes D, Virtamo J. Intake of flavonols and flavones and risk of coronary heart disease in male smokers. *Epidemiology* 2001; 12: 62–7
- 48 Hirvonen T, Virtamo J, Korhonen P, Albanes D, Pietinen P. Flavonol and flavone intake and the risk of cancer in male smokers (Finland). *Cancer Causes Control* 2001; 12: 789–96
- 49 Gosnay SL, Bishop JA, New SA, Catterick J, Clifford MN. Estimation of the mean intakes of fourteen classes of dietary phenolics in a population of young British women aged 20–30 years. *Proc Nutr Soc* 2002; 61: 125A
- 50 Woods E, Clifford MN, Gibbs M, Hampton S, Arendt J, Morgan L. Estimation of mean intakes of 14 classes of dietary phenols in a population of male shift workers. *Proc Nutr Soc* 2003; 62: 60A
- 51 Hertog MGL, Sweetman PM, Fehily AM, Elwood PC, Kromhout D. Antioxidant flavonols and ischaemic heart disease in a Welsh population of men: the Caerphilly Study. *Am J Clin Nutr* 1997; 65: 1489–94
- 52 Arai Y, Watanabe S, Kimira M, Shimoi K, Mochizuki R, Kinai N. Dietary intakes of flavonols, flavones and isoflavones by Japanese women and the inverse correlation between quercetin intake and plasma LDL cholesterol concentration. *J Nutr* 2000; 130: 2243–50
- 53 Tomás-Barberán FA, Clifford MN. Flavanones, chalcones and dihydrochalcones – nature, occurrence and dietary burden. *J Sci Food Agric* 2000; 80: 1073–80
- 54 Price KR, Bacon JR, Rhodes MJC. Effect of storage and domestic processing on the content and composition of flavonol glucosides in onion (*Allium cepa*). *J Agr Food Chem* 1997; 45: 938–42
- 55 Ewald C, Fjelkner-Modig S, Johanssen K, Sjöholm I, Åkesson B. Effect of processing on major flavonoids in processed onions, green beans, and peas. *Food Chem* 1999; 64: 231–5
- 56 Chuda Y, Suzuki M, Nagata T, Tsuchida T. Contents and cooking loss of three quinic acid derivatives from Garland (*Chrysanthemum coronarium* L.). *J Agr Food Chem* 1998; 46: 1437–9
- 57 Price KR, Casuscelli F, Colquhoun IJ, Rhodes MJC. Composition and content of flavonol glycosides in broccoli florets (*Brassica oleracea*) and their fate during cooking. *J Sci Food Agric* 1998; 77: 468–72
- 58 Ioku K, Aoyama Y, Tokuno A, Terao J, Nakatani N, Takei Y. Various cooking methods and the flavonoid content in onion. *J Nutr Sci Vitaminol (Tokyo)* 2001; 47: 78–83
- 59 Brenes M, Garcia A, Dobarganes MC, Velasco J, Romero C. Influence of thermal treatments simulating cooking processes on the polyphenol content in virgin olive oil. *J Agr Food Chem* 2002; 50: 5962–7
- 60 Valledo F, Tom FA, Garc C. Phenolic compound contents in edible parts of broccoli inflorescences after domestic cooking. *J Sci Food Agric* 2003; 83: 1511–6
- 61 Bond TA, Lewis JR, Davis A, Davies AP. Analysis and purification of catechins and their transformation products. In: Santos-Buelga C, Williamson G, editors. *Methods in Polyphenol Analysis* Cambridge: Royal Society of Chemistry, 2003: pp 229–66
- 62 Garcia-Closas R, Gonzalez CA, Agudo A, Riboli E. Intake of specific carotenoids and flavonoids and the risk of gastric cancer in Spain. *Cancer Causes Control* 1999; 10: 71–5
- 63 Justesen U, Knuthsen P. Composition of flavonoids in fresh herbs and calculation of flavonoid intake by use of herbs in traditional Danish dishes. *Food Chem* 2001; 73: 245–50
- 64 Justesen U, Knuthsen P, Andersen NL, Leth T. Estimation of daily intake distribution of flavonols and flavanones in Denmark. *Scand J Nutr* 2000; 44: 158–60
- 65 Keli SO, Hertog MG, Feskens EJ, Kromhout D. Dietary flavonoids, antioxidant vitamins, and incidence of stroke: the Zutphen study. *Arch Intern Med* 1996; 156: 637–42
- 66 Knekt P, Jarvinen R, Seppanen R, Hellewaara M, Teppo L, Pukkala E, Aromaa A. Dietary flavonoids and the risk of lung cancer and other malignant neoplasms. *Am J Epid* 1997; 146: 223–30
- 67 Knekt P, Jarvinen R, Reunanen A, Maatela J. Flavonoid intake and coronary mortality in Finland: a cohort study. *Brit Med J* 1996; 312: 478–81



- 68 Knekt P, Isotupa S, Rissanen H, Heliovaara M, Jarvinen R, Hakkinen S, Aromaa A, Reunanen A. Quercetin intake and the incidence of cerebrovascular disease. *Eur J Clin Nutr* 2000; 54: 415–7
- 69 Pascual-Teresa S, Santos-Buelga C, Rivas-Gonzalo JC. Quantitative analysis of flavan-3-ols in Spanish foodstuffs and beverages. *J Agric Food Chem* 2000; 48: 5331–7
- 70 Arts ICW, van de Putte B, Hollman PCH. Catechin contents of foods commonly consumed in the Netherlands. 1. Fruits, vegetables, staple foods, and processed foods. *J Agr Food Chem*, 2000: 1746–51
- 71 Arts ICW, Hollman PCH, Feskens EJM, de Mesquita BB, Kromhout D. Catechin intake is inversely associated with coronary heart disease mortality; Results from the Zutphen Elderly Study. *Circulation* 2000; 101: 4
- 72 Stewart AJ, Bozonnet S, Mullen W, Jenkins GI, Lean ME, Crozier A. Occurrence of flavonols in tomatoes and tomato-based products. *J Agr Food Chem* 2000; 48: 2663–9
- 73 Crozier A, Burns J, Aziz AA, Stewart AJ, Rabiasz HS, Jenkins G, Edwards CA, Lean MEJ. Antioxidant flavonols from fruits, vegetables and beverages: measurement and bioavailability. *Biol Res* 2000; 33: 79–88
- 74 Howard LR, Pandjaitan N, Morelock T, Gil MI. Antioxidant capacity and phenolic content of spinach as affected by genetics and growing season. *J Agr Food Chem* 2002; 50: 5891–6
- 75 Cacace JE, Mazza G. Extraction of anthocyanins and other phenolics from blackcurrants with sulfured water. *J Agr Food Chem* 2002; 50: 5939–46
- 76 Porteous L. Nutritional and dietary risk factors for osteoporosis: An investigation of association between diet and indices of bone health in young British women aged 25–30 years. B.Sc. Thesis. University of Surrey, 2001
- 77 Bennett G. Food choices, dietary patterns and circulation adaptation in shift workers: an investigation into the possible impact on coronary heart disease risk. B.Sc. Thesis. University of Surrey, 2001
- 78 Paul N. Do eating patterns and post-prandial hormonal and metabolic markers vary significantly according to shift type and circadian status in swing shift workers in the offshore oil industry. B.Sc. Thesis. University of Surrey, 2001
- 79 Balentine DA, Wiseman SA, Bouwens CM. The chemistry of tea flavonoids. *CRC Crit Rev Food Sci Nutr* 1997; 37: 693–704
- 80 Clifford MN. Tea. In: Ranken MD, Kill RC and Baker CJG, editors. *Food Industries Manual*; 24th edition London: Blackie, 1997: pp 379–95
- 81 Harbowy ME, Balentine DA. Tea Chemistry. *CRC Crit Rev Plant Sci* 1997; 16: 415–80
- 82 Olthof MR, Hollman PC, Buijsman MN, van Amelsvoort JM, Katan MB. Chlorogenic acid, quercetin-3-rutinoside and black tea phenols are extensively metabolized in humans. *J Nutr* 2003; 133: 1806–14
- 83 Nardini M, Cirillo E, Natella F, Scaccini C. Absorption of phenolic acids in humans after coffee consumption. *J Agr Food Chem* 2002; 50: 5735–41
- 84 Lee M-J, Wang Z-Y, Li H, Chen L, Sun Y, Gobbo S, Balentine DA, Yang CS. Analysis of plasma and urinary tea polyphenols in human subjects. *Cancer Epidemiol Biomarkers Prev* 1995; 4: 393–9
- 85 Hollman PC, Katan MB. Health effects and bioavailability of dietary flavonols. *Free Radic Res* 1999; 31 Suppl: S75–80
- 86 Olthof MR, Hollman PC, Vree TB, Katan MB. Bioavailabilities of quercetin-3-glucoside and quercetin-4-glucoside do not differ in humans. *J Nutr* 2000; 130: 1200–3
- 87 Hollman PCH, van Trijp JM, Buysman MN, van der Gaag MS, Mengelers MJ, de Vries JHM, Katan MB. Relative bioavailability of the antioxidant quercetin from various foods in man. *FEBS Lett* 1997; 418: 152–6
- 88 Walle T, Walle UK, Halushka PV. Carbon dioxide is the major metabolite of quercetin in humans. *J Nutr* 2001; 131: 2648–52
- 89 Krishnamurthy HG, Cheng KJ, Jones GA, Simpson FJ, Watkin JE. Identification of products produced by the anaerobic degradation of rutin and related flavonoids by *Butyrivibrio* sp. C3. *Can J Microbiol* 1970; 16: 759–67
- 90 Scheline RR. The metabolism of (+)-catechin to hydroxyphenylvaleric acids by the intestinal microflora. *Biochim Biophys Acta* 1970; 222: 228–30
- 91 Scheline RR. Mammalian Metabolism of Plant Xenobiotics. London: Academic Press, 1978: pp 1–502
- 92 Griffiths LA. Mammalian metabolism of flavonoids. In: Harborne JB, Mabry TJ, editors. *The Flavonoids: Advances in Research* London: Chapman and Hall, 1982: pp 681–718
- 93 Groenewoud G, Hundt HKL. The microbial metabolism of (+)-catechins to two novel diarylpropan-2-ol metabolites *in vitro*. *Xenobiotica* 1984; 14: 711–7
- 94 Bokkenheuser VD, Shackleton CH, Winter J. Hydrolysis of dietary flavonoid glycosides by strains of intestinal *Bacteroides* from humans. *Biochem J* 1987; 248: 953–6
- 95 Sawai Y, Kohsaka K, Nishiyama Y, Ando K. Serum concentrations of rutoside metabolites after oral administration of a rutoside formulation to humans. *Arzneimittel-Forsch* 1987; 37: 729–32
- 96 Bokkenheuser VD, Winter J. Hydrolysis of flavonoids by human intestinal bacteria. *Progress in Clinical & Biological Research* 1988; 280: 143–5
- 97 Winter J, Moore LH, Dowell VR, Bokkenheuser VD. C-ring cleavage of flavonoids by human intestinal bacteria. *Appl Environ Microbiol* 1989; 55: 1203–8
- 98 Coldham NG, King LJ, Macpherson DD, Sauer MJ. Biotransformation of genistein in the rat: elucidation of metabolite structure by product ion mass fragmentology. *J Steroid Biochem Mol Biol* 1999; 70: 169–84
- 99 Schneider H, Schwiertz A, Collins MD, Blaut M. Anaerobic transformation of quercetin-3-glucoside by bacteria from the human intestinal tract. *Arch Microbiol* 1999; 171: 81–091
- 100 Simmering R, Kleessen B, Blaut M. Quantification of the flavonoid-degrading bacterium *Eubacterium ramulus* in human fecal samples with a species-specific oligonucleotide hybridization probe. *Appl Environ Microbiol* 1999; 65: 3705–9
- 101 Deprez S, Brezillon C, Rabot S, Philippe C, Mila I, Lapierre C, Scalbert A. Polymeric proanthocyanidins are catabolized by human colonic microflora into low-molecular-weight phenolic acids. *J Nutr* 2000; 130: 2733–8
- 102 Justesen U, Arrigoni E, Larsen BR, Amado R. Degradation of flavonoid glycosides and aglycones during *in vitro* fermentation with human faecal flora. *Lebensm Wiss Technol* 2000; 33: 424–30
- 103 Clifford MN, Copeland EL, Bloxside JP, Mitchell LA. Hippuric acid is a major excretion product associated with black tea consumption. *Xenobiotica* 2000; 30: 317–26
- 104 Li C, Lee M-J, Sheng S, Meng X, Prabhu S, Winnik B, Huang B, Chung JY, Yan S, Ho C-T, Yang CS. Structural identification of two metabolites of catechins and their kinetics in human urine and blood after tea ingestion. *Chem Res Toxicol* 2000; 13: 177–84
- 105 Schneider H, Blaut M. Anaerobic degradation of flavonoids by *Eubacterium ramulus*. *Arch Microbiol* 2000; 173: 71–5
- 106 Schneider H, Simmering R, Hartmann L, Blaut M. Degradation of quercetin-3-glucoside in gnotobiotic rats associated with human intestinal bacteria. *J Appl Microbiol* 2000; 89: 1027–37
- 107 Jenkins DI, Wolever TM, Taylor RH. Glycaemic index of foods: a physiological basis for carbohydrate exchange. *Am J Clin Nutr* 1981; 34: 362–6
- 108 Couteau D, McCartney AL, Gibson GR, Williamson G, Faulds CB. Isolation and characterization of human colonic bacteria able to hydrolyse chlorogenic acid. *J Appl Microbiol* 2001; 90: 873–881
- 109 Olthof MR. Bioavailability of flavonoids and cinnamic acids and their effect on plasma homocysteine in humans. PhD Thesis. Wageningen University, 2001: pp 136
- 110 Olthof MR, Hollman PCH, Katan MB. Chlorogenic acid and caffeic acid are absorbed in humans. *J Nutr* 2001; 131: 66–71
- 111 Aura AM, O'Leary KA, Williamson G, Ojala M, Bailey M, Puupponen-Pimia R, Nuutila AM, Oksman-Caldentey KM, Poutanen K. Quercetin derivatives are deconjugated and converted to hydroxyphenylacetic acids but not methylated by human fecal flora *in vitro*. *J Agr Food Chem* 2002; 50: 1725–30
- 112 Schoefer L, Mohan R, Braune A, Birringer M, Blaut M. Anaerobic C-ring cleavage of genistein and daidzein by *Eubacterium ramulus*. *FEMS Microbiol Lett* 2002; 208: 197–202
- 113 Braune A, Gutschow M, Engst W, Blaut M. Degradation of quercetin and luteolin by *Eubacterium ramulus*. *Appl Environ Microbiol* 2001; 67: 5558–67
- 114 Wang LQ, Meselhy MR, Li Y, Nakamura N, Min BS, Qin GW, Hattori M. The heterocyclic ring fission and dehydroxylation of catechins and related compounds by *Eubacterium* sp. strain SDG-2, a human intestinal bacterium. *Chem Pharm Bull* 2001; 49: 1640–3
- 115 Hur H, Rafii F. Biotransformation of the isoflavonoids biochanin A, formononetin, and glycitein by *Eubacterium limosum*. *FEMS Microbiol Lett* 2000; 192: 21–5

- <sup>116</sup> Wang LQ, Meselhy MR, Li Y, Qin GW, Hattori M. Human intestinal bacteria capable of transforming secoisolariciresinol diglucoside to mammalian lignans, enterodiols and enterolactone. *Chem Pharm Bull* 2000; 48: 1606–10
- <sup>117</sup> Barcenilla A, Pryde SE, Martin JC, Duncan SH, Stewart CS, Henderson C, Flint HJ. Phylogenetic relationships of butyrate-producing bacteria from the human gut. *Appl Environ Microbiol* 2000; 66: 1654–61
- <sup>118</sup> Takanaga H, Tamai I, Tsuji A. pH-dependent and carrier-mediated transport of salicylic acid across Caco-2 cells. *J Pharm Pharmacol* 1994; 46: 567–70
- <sup>119</sup> Tsuji A, Takanaga H, Tamai I, Terasaki T. Transcellular transport of benzoic acid across Caco-2 cells by a pH-dependent and carrier-mediated transport mechanism. *Pharm Res* 1994; 11: 30–7
- <sup>120</sup> Tamai I, Takanaga H, Maeda H, Yabuuchi H, Sai Y, Suzuki Y, Tsuji A. Intestinal brush-border membrane transport of monocarboxylic acids mediated by proton-coupled transport and anion antiport mechanisms. *J Pharm Pharmacol* 1997; 49: 108–12
- <sup>121</sup> Konishi Y, Shimizu M. Transepithelial transport of ferulic acid by monocarboxylic acid transporter in Caco-2 cell monolayers. *Biosci Biotechnol Biochem* 2003; 67: 856–62
- <sup>122</sup> Konishi Y, Kobayashi S, Shimizu M. Transepithelial transport of *p*-coumaric acid and gallic acid in Caco-2 cell monolayers. *Biosci Biotechnol Biochem* 2003; 67: 2317–24
- <sup>123</sup> Baba S, Osakabe N, Yasuda A, Natsume M, Takizawa T, Nakamura T, Terao J. Bioavailability of (-)-epicatechin upon intake of chocolate and cocoa in human volunteers. *Free Radic Res* 2000; 33: 635–41
- <sup>124</sup> Bell JRC, Donovan JL, Wong R, Waterhouse AL, German JB, Walzem RL, Kasim-Karakas SE. (+)-Catechin in human plasma after ingestion of a single serving of reconstituted red wine. *Am J Clin Nutr* 2000; 71: 103–8
- <sup>125</sup> Hollman PCH, van der Gaag MS, Mengelers MJB, van Trijp JMP, de Vries JHM, Katan MB. Absorption and disposition kinetics of the dietary antioxidant quercetin in man. *Free Radic Biol Med* 1996; 21: 703–7
- <sup>126</sup> Holt RR, Lazarus SA, Sullards MC, Zhu QY, Schramm DD, Hammerstone JF, Fraga CG, Schmitz HH, Keen CL. Procyanidin dimer B2 [epicatechin-(4beta-8)-epicatechin] in human plasma after the consumption of a flavanol-rich cocoa. *Am J Clin Nutr* 2002; 76: 798–804
- <sup>127</sup> Kimura M, Umegaki K, Kasuya Y, Sugisawa A, Higuchi M. The relation between single/double or repeated tea catechin ingestions and plasma antioxidant activity in humans. *Eur J Clin Nutr* 2002; 56: 1186–93
- <sup>128</sup> McAnlis GT, McEneny J, Pearce J, Young IS. Absorption and antioxidant effects of quercetin from onions, in man. *Eur J Clin Nutr* 1999; 53: 92–6
- <sup>129</sup> van het Hof K, Wiseman SA, de Boer H, Weststrate NLJ, Tijburg L. Bioavailability and antioxidant activity of tea polyphenols in humans. In: Amadó R, Andersson H, Bardócs S and Serra F, editors. COST 916 Bioactive plant cell wall components in nutrition and health. Polyphenols in food Luxembourg: EC, 1998: 115
- <sup>130</sup> Wang JF, Schramm DD, Holt RR, Ensunsa JL, Fraga CG, Schmitz HH, Keen CL. A dose-response effect from chocolate consumption on plasma epicatechin and oxidative damage. *J Nutr* 2000; 130: 2115S–9
- <sup>131</sup> Manach C, Morand C, Gil-Izquierdo A, Bouteloup-Demange C, Remesy C. Bioavailability in humans of the flavanones hesperidin and narirutin after the ingestion of two doses of orange juice. *Eur J Clin Nutr* 2003; 57: 235–42
- <sup>132</sup> Yang CS, Lee MJ, Chen L. Human salivary tea catechin levels and catechin esterase activities: implication in human cancer prevention studies. *Cancer Epidemiology, Biomarkers & Prevention* 1999; 8: 83–9
- <sup>133</sup> Meng X, Lee MJ, Li C, Sheng S, Zhu N, Sang S, Ho CT, Yang CS. Formation and identification of 4'-O-methyl(-)-epigallocatechin in humans. *Drug Metab Dispos* 2001; 29: 789–93
- <sup>134</sup> Caccetta RA, Croft KD, Beilin LJ, Puddey IB. Ingestion of red wine significantly increases plasma phenolic acid concentrations but does not acutely affect *ex vivo* lipoprotein oxidizability. *Am J Clin Nutr* 2000; 71: 67–74
- <sup>135</sup> Carbonneau MA, Leger CL, Monnier L, Bonnet C, Michel F, Fouret G, Dedieu F, Descomps B. Supplementation with wine phenolic compounds increases the antioxidant capacity of plasma and vitamin E of low-density lipoprotein without changing the lipoprotein Cu(2+)-oxidizability: possible explanation by phenolic location. *Eur J Clin Nutr* 1997; 51: 682–90
- <sup>136</sup> Conquer JA, Maiani G, Azzini E, Raguzzini A, Holub BJ. Supplementation with quercetin markedly increases plasma quercetin concentration without effect on selected risk factors for heart disease in healthy subjects. *J Nutr* 1998; 128: 593–7
- <sup>137</sup> Eccleston C, Baoru Y, Tahvonen R, Kallio H, Rimbach GH, Minihane AM. Effects of an antioxidant-rich juice (sea buckthorn) on risk factors for coronary heart disease in humans. *J Nutr Biochem* 2002; 13: 346–54
- <sup>138</sup> Ghiselli A, Natella F, Guidi A, Montanari L, Fantozzi P, Scaccini C. Beer increases plasma antioxidant capacity in humans. *J Nutr Biochem* 2000; 11: 76–80
- <sup>139</sup> Hodgson JM, Puddey IB, Croft KD, Burke V, Mori TA, Caccetta RA, Beilin LJ. Acute effects of ingestion of black and green tea on lipoprotein oxidation. *Am J Clin Nutr* 2000; 71: 1103–7
- <sup>140</sup> Hodgson JM, Puddey IB, Croft KD, Mori TA, Rivera J, Beilin LJ. Isoflavonoids do not inhibit *in vivo* lipid peroxidation in subjects with high-normal blood pressure. *Atherosclerosis* 1999; 145: 167–72
- <sup>141</sup> Leenen R, Roodenburg AJ, Tijburg LB, Wiseman SA. A single dose of tea with or without milk increases plasma antioxidant activity in humans. *Eur J Clin Nutr* 2000; 54: 87–92
- <sup>142</sup> Mathur S, Devaraj S, Grundy SM, Jialal I. Cocoa products decrease low density lipoprotein oxidative susceptibility but do not affect biomarkers of inflammation in humans. *J Nutr* 2002; 132: 3663–7
- <sup>143</sup> Mazza G, Kay CD, Cottrell T, Holub BJ. Absorption of anthocyanins from blueberries and serum antioxidant status in human subjects. *J Agr Food Chem* 2002; 50: 7731–7
- <sup>144</sup> Mitchell JH, Collins AR. Effects of a soy milk supplement on plasma cholesterol levels and oxidative DNA damage in men – a pilot study. *Eur J Nutr* 1999; 38: 143–8
- <sup>145</sup> Nakagawa K, Ninomiya M, Okubo T, Aoi N, Juneja LR, Kim M, Yamanaoka K, Miyazawa T. Tea catechin supplementation increases antioxidant capacity and prevents phospholipid hydroperoxidation in plasma of humans. *J Agr Food Chem* 1999; 47: 3967–73
- <sup>146</sup> Natella F, Belelli F, Gentili V, Ursini F, Scaccini C. Grape seed proanthocyanidins prevent plasma postprandial oxidative stress in humans. *J Agr Food Chem* 2002; 50: 7720–5
- <sup>147</sup> Nigdikar SV, Williams NR, Griffin BA, Howard AN. Consumption of red wine polyphenols reduces the susceptibility of low-density lipoproteins to oxidation *in vivo*. *Am J Clin Nutr* 1998; 68: 258–65
- <sup>148</sup> O'Byrne DJ, Devaraj S, Grundy SM, Jialal I. Comparison of the antioxidant effects of Concord grape juice flavonoids alpha-tocopherol on markers of oxidative stress in healthy adults. *Am J Clin Nutr* 2002; 76: 1367–74
- <sup>149</sup> Pedersen CB, Kyle J, Jenkinson AM, Gardner PT, McPhail DB, Duthie GG. Effects of blueberry and cranberry juice consumption on the plasma antioxidant capacity of healthy female volunteers. *Eur J Clin Nutr* 2000; 54: 405–8
- <sup>150</sup> Princen HM, van Duyvenvoorde W, Buytenhek R, Blonk C, Tijburg LB, Langius JA, Meinders AE, Pijl H. No effect of consumption of green and black tea on plasma lipid and antioxidant levels and on LDL oxidation in smokers. *Arterioscler Thromb Vasc Biol* 1998; 18: 833–41
- <sup>151</sup> Rein D, Lotito S, Holt RR, Keen CL, Schmitz HH, Fraga CG. Epicatechin in human plasma: *In vivo* determination and effect of chocolate consumption on plasma oxidation status. *J Nutr* 2000; 130: 2109S–14
- <sup>152</sup> Serafini M, Maiani G, Ferro-Luzzi A. Alcohol-free red wine enhances plasma antioxidant capacity in humans. *J Nutr* 1998; 128: 1003–7
- <sup>153</sup> Simonetti P, Ciappellano S, Gardana C, Bramati L, Pietta P. Procyanidins from *Vitis vinifera* seeds: *in vivo* effects on oxidative stress. *J Agr Food Chem* 2002; 50: 6217–21
- <sup>154</sup> Stein JH, Keevil JG, Wiebe DA, Aeschlimann S, Folts JD. Purple grape juice improves endothelial function and reduces the susceptibility of LDL cholesterol to oxidation in patients with coronary artery disease. *Circulation* 1999; 100: 1050–5
- <sup>155</sup> Tikkanen MJ, Wahala K, Ojala S, Vihma V, Adlercreutz H. Effect of soybean phytoestrogen intake on low density lipoprotein oxidation resistance. *Proc Natl Acad Sci USA* 1998; 95: 3106–10
- <sup>156</sup> van den Berg R, van Vliet T, Broekmans WMR, Cnubben NHP, Vaes WHJ, Roza L, Haenen GRMM, Bast A, van den Berg H. A vegetable/fruit concentrate with high antioxidant capacity has no effect on biomarkers of antioxidant status in male smokers. *J Nutr* 2001; 131: 1714–22
- <sup>157</sup> Wan Y, Vinson JA, Etherton TD, Proch J, Lazarus SA, Kris-Etherton PM. Effects of cocoa powder and dark chocolate on LDL oxidative susceptibility and prostaglandin concentrations in humans. *Am J Clin Nutr* 2001; 74: 596–602

- <sup>158</sup> Young JF, Nielsen SE, Haraldsdottir J, Daneshvar B, Lauridsen ST, Knuthsen P, Crozier A, Sandstrom B, Dragsted LO. Effect of fruit juice intake on urinary quercetin excretion and biomarkers of antioxidative status. *Am J Clin Nutr* 1999; 69: 87–94
- <sup>159</sup> Lotito SB, Frei B. Relevance of apple polyphenols as antioxidants in human plasma: contrasting *in vitro* and *in vivo* effects. *Free Radic Biol Med* 2004; 36: 201–11
- <sup>160</sup> Jovanovic SV, Steenken S, Hara Y, Simic MG. Reduction potentials of flavonoid and model phenoxyl radicals. Which ring in flavonoids is responsible for antioxidant activity? *J Chem Soc Perkin Trans* 1996; 2: 2497–504
- <sup>161</sup> Jovanovic SV, Hara Y, Steenken S, Simic MG. Antioxidant potential of gallic catechins. A pulse radiolysis and laser photolysis study. *J Am Chem Soc* 1995; 117: 9881–8
- <sup>162</sup> Jovanovic SV, Steenken S, Tomic M, Marjanovic B, Simic MG. Flavonoids as antioxidants. *J Am Chem Soc* 1994; 116: 4846–51
- <sup>163</sup> Steenken S, Neta P. One-electron redox potentials of phenols. Hydroxy and amino-phenols and related compounds of biological interest. *J Phys Chem*, 1982: 3661–7
- <sup>164</sup> Ferry DR, Smith A, Malkhandi J, Fyfe DW, deTakats PG, Anderson D, Baker J, Kerr DJ. Phase I clinical trial of the flavonoid quercetin: pharmacokinetics and evidence for *in vivo* tyrosine kinase inhibition. *Clin Cancer Res* 1996; 2: 659–68
- <sup>165</sup> Ayrton AD, Lewis DFV, Walker R, Ioannides C. Antimutagenicity of ellagic acid toward the food mutagen IQ: investigations into the possible mechanisms of action. *Food Chem Toxicol* 1992; 30: 289–95
- <sup>166</sup> Hirose Y, Tanaka T, Kawamori T, Ohnishi M, Makita H, Mori H, Satoh K, Hara A. Chemoprevention of urinary-bladder carcinogenesis by the natural phenolic compound protocatechuic acid in rats. *Carcinogen* 1995; 16: 2337–42
- <sup>167</sup> Shirai M, Moon JH, Tsushida T, Terao J. Inhibitory effect of a quercetin metabolite, quercetin 3-O- $\beta$ -D-glucuronide, on lipid peroxidation in liposomal membranes. *J Agr Food Chem* 2001; 49: 5602–8
- <sup>168</sup> Terao J, Murota K, Moon J-H. Quercetin glucosides as dietary antioxidants in blood plasma: modulation of their function by metabolic conversion. In: Yoshikawa T, Toyokuni Y, Yamamoto Y and Naito Y, editors. *Free radicals in chemistry, biology and medicine* London: OIAC International, 2000: pp 462–75
- <sup>169</sup> Liao K, Yin M. Individual and combined antioxidant effects of seven phenolic agents in human erythrocyte membrane ghosts and phosphatidylcholine liposome systems: importance of the partition coefficient. *J Agr Food Chem* 2000; 48: 2266–70
- <sup>170</sup> Finch S, Doyle W, Lowe C, Bates CJ, Prentice A, Smithers G, Clarke PC. National Diet and Nutrition Survey. People aged 65 years and over. London: The Stationery Office, 1998
- <sup>171</sup> Sauberlich HE. Human requirements and needs. Vitamin C status: methods and findings. *Ann NY Acad Sci* 1975; 258: 438–50
- <sup>172</sup> Yoshizumi M, Tsuchiya K, Suzaki Y, Kirima K, Kyaw M, Moon JH, Terao J, Tamaki T. Quercetin glucuronide prevents VSMC hypertrophy by angiotensin II via the inhibition of JNK and AP-1 signaling pathway. *Biochem Biophys Res Commun* 2002; 293: 1458–65
- <sup>173</sup> Nees S, Weiss DR, Reichenbach-Klinke E, Rampp F, Heilmeier B, Kanbach J, Esperester A. Protective effects of flavonoids contained in the red vine leaf on venular endothelium against the attack of activated blood components *in vitro*. *Arzneimittel-Forsch* 2003; 53: 33–41
- <sup>174</sup> Terao J, Yamaguchi S, Shirai M, Miyoshi M, Moon JH, Oshima S, Inakuma T, Tsushida T, Kato Y. Protection by quercetin and quercetin 3-O- $\beta$ -D-glucuronide of peroxynitrite-induced antioxidant consumption in human plasma low-density lipoprotein. *Free Radic Res* 2001; 35: 925–31
- <sup>175</sup> Day AJ, Bao Y, Morgan MRA, Williamson G. Conjugation position of quercetin glucuronides and effect on biological activity. *Free Radic Biol Med* 2000; 29: 1234–43
- <sup>176</sup> Thompson LU, Yoon JH, Jenkins DJ, Wolever TM, Jenkins AL. Relationship between polyphenol intake and blood glucose response of normal and diabetic individuals. *Am J Clin Nutr* 1984; 39: 745–51
- <sup>177</sup> Gin H, Rigalleau V, Caubet O, Masquelier J, Aubertin J. Effects of red wine, tannic acid, or ethanol on glucose tolerance in non-insulin-dependent diabetic patients and on starch digestibility *in vitro*. *Metabolism* 1999; 48: 1179–83
- <sup>178</sup> Johnston KL, Clifford MN, Morgan LM. Coffee acutely modifies gastrointestinal hormone secretion and glucose tolerance in humans: glycemic effects of chlorogenic acid and caffeine. *Am J Clin Nutr* 2003; 78: 728–33
- <sup>179</sup> Johnston KL, Clifford MN, Morgan LM. Possible role for apple juice phenolic compounds in the acute modification of glucose tolerance and gastrointestinal hormone secretion in humans. *J Sci Food Agric* 2002; 82: 1800–5
- <sup>180</sup> Coutinho M, Gerstein HC, Wang Y, Yusef S. The relationship between glucose and incident cardiovascular events: a metaregression analysis of published data from 20 studies of 95,783 individuals followed for 12.4 years. *Diabetes Care* 1999; 22: 233–40
- <sup>181</sup> van Dam RM, Feskens EJ. Coffee consumption and risk of type 2 diabetes mellitus. *Lancet* 2002; 360: 1477–8
- <sup>182</sup> Facchini FS, Saylor KL. A low-iron-available, polyphenol-enriched, carbohydrate-restricted diet to slow progression of diabetic nephropathy. *Diabetes* 2003; 52: 1204–9
- <sup>183</sup> Hara Y, Honda M. The inhibition of  $\alpha$ -amylase by tea polyphenols. *Agric Biol Chem* 1990; 54: 1939–45
- <sup>184</sup> Matsumoto N, Ishigaki F, Ishigaki A, Iwashina H, Hara Y. Reduction of blood-glucose levels by tea catechin. *Biosci Biotechnol Biochem* 1993; 57: 525–7
- <sup>185</sup> Matsui T, Ebuchi S, Kobayashi M, Fukui K, Sugita K, Terahara N, Matsumoto K. Anti-hyperglycemic effect of diacylated anthocyanin derived from *Ipomoea batatas* cultivar Ayamurasaki can be achieved through the  $\alpha$ -glucosidase inhibitory action. *J Agr Food Chem* 2002; 50: 7244–8
- <sup>186</sup> Kawabata J, Mizuhata K, Sato E, Nishioka T, Aoyama Y, Kasai T. 6-hydroxyflavonoids as  $\alpha$ -glucosidase inhibitors from marjoram (*Origanum majorana*) leaves. *Biosci Biotechnol Biochem* 2003; 67: 445–7
- <sup>187</sup> Hilt P, Schieber A, Yildirim C, Arnold G, Klaiber I, Conrad J, Beifuss U, Carle R. Detection of phloridzin in strawberries (*Fragaria x ananassa* Duch.) by HPLC-PDA-MS/MS and NMR spectroscopy. *J Agr Food Chem* 2003; 51: 2896–9
- <sup>188</sup> Nakazawa F. Influence of phloridzin on intestinal absorption. *Tohoku J Exptl Med* 1922; 3: 288–94
- <sup>189</sup> Alvarado F. Hypothesis for the interaction of phlorizin and phloretin with membrane carriers for sugars. *Biochim Biophys Acta* 1967; 135: 483–95
- <sup>190</sup> Alvarado F, Crane RK. Phlorizin as a competitive inhibitor of the active transport of sugars by hamster small intestine. *Biochim Biophys Acta* 1962; 56: 170–2
- <sup>191</sup> Crane RK. Intestinal absorption of sugars. *Phys Rev* 1960; 40: 789–825
- <sup>192</sup> Welsch CA, Lachance PA, Wasserman BP. Dietary phenolic compounds: Inhibition of Na<sup>+</sup>-dependent d-glucose uptake in rat intestinal brush border membrane vesicles. *J Nutr* 1989; 119: 1698–1704
- <sup>193</sup> Kobayashi Y, Suzuki M, Satsu H, Arai S, Hara Y, Suzuki K, Miyamoto Y, Shimizu M. Green tea polyphenols inhibit the sodium-dependent glucose transporter of intestinal epithelial cells by a competitive mechanism. *J Agr Food Chem* 2000; 48: 5618–23
- <sup>194</sup> Noteborn HPJM, Jansen E, Benito S, Mengelers MJB. Oral absorption and metabolism of quercetin and sugar-conjugated derivatives in specific transport systems. *Cancer Lett* 1997; 114: 175–7
- <sup>195</sup> Walgren RA, Walle UK, Walle T. Transport of quercetin and its glucosides across human intestinal epithelial Caco-2 cells. *Biochem Pharmacol* 1998; 55: 1721–7
- <sup>196</sup> Gee JM, DuPont MS, Rhodes MJC, Johnson IT. Quercetin glucosides interact with the intestinal glucose transport pathway. *Free Radic Biol Med* 1998; 25: 19–25
- <sup>197</sup> Walgren RA, Lin JT, Kinne RK, Walle T. Cellular uptake of dietary flavonoid quercetin 4'- $\beta$ -glucoside by sodium-dependent glucose transporter SGLT1. *J Pharmacol Exp Ther* 2000; 294: 837–43
- <sup>198</sup> Walgren RA, Karnaky KJ Jr., Lindenmayer GE, Walle T. Efflux of dietary flavonoid quercetin 4'-beta-glucoside across human intestinal Caco-2 cell monolayers by apical multidrug resistance-associated protein-2. *J Pharmacol Exp Ther* 2000; 294: 830–6
- <sup>199</sup> Song J, Kwon O, Chen S, Daruwala R, Eck P, Park JB, Levine M. Flavonoid inhibition of sodium-dependent vitamin C transporter 1 (SVCT1) and glucose transporter isoform 2 (GLUT2), intestinal transporters for vitamin C and glucose. *J Biol Chem* 2002; 277: 15252–60
- <sup>200</sup> Wolfram S, Block M, Ader P. Quercetin-3-glucoside is transported by the glucose carrier SGLT1 across the brush border membrane of rat small intestine. *J Nutr* 2002; 132: 630–5
- <sup>201</sup> Deutsch JC, SanthoshKumar CR. Quantitation of homogentisic acid in normal human plasma. *J Chromatogr B Biomed Appl* 1996; 667: 147–51

- <sup>202</sup> Deutsch JC. Determination of *p*-hydroxyphenylpyruvate, *p*-hydroxyphenyllactate and tyrosine in normal human plasma by gas chromatography-mass spectrometry isotope-dilution assay. *J Chromatogr B* 1997; 690: 1–6
- <sup>203</sup> Heinecke JW. Tyrosyl radical production by myeloperoxidase: a phagocyte pathway for lipid peroxidation and dityrosine cross-linking of proteins. *Toxicology* 2002; 177: 11–22
- <sup>204</sup> Gregory JR, Collins DL, Davies PSW, Hughes JM, Clarke PC. National Diet and Nutrition Survey. Children aged one and a half years to four and a half years. London: HMSO, 1995
- <sup>205</sup> Gregory J, Lowe S, Bates CJ, Prentice A, Jackson LV, Smithers G, Wenlock R, Farron M. National Diet and Nutrition Survey. Young people aged 4 to 18 years. London: The Stationery Office, 2000
- <sup>206</sup> Halliwell B, Gutteridge JM. *Free Radicals in Biology and Medicine*. 3th edition. Oxford: Oxford University Press, 1999
- <sup>207</sup> Jovanovic SV, Hara Y, Steenken S, Simic MG. Antioxidant potential of theaflavins. a pulse radiolysis study. *J Am Chem Soc* 1997; 119: 5337–43
- <sup>208</sup> Buettner GR. The pecking order of free radicals and antioxidants: lipid peroxidation, alpha-tocopherol, and ascorbate. *Arch Biochem Biophys* 1993; 300: 535–43
- <sup>209</sup> Jovanovic SV, Steenken S, Boone CW, Simic MG. H-atom transfer is a preferred antioxidant mechanism of curcumin. *J Am Chem Soc* 1999; 121: 9677–81
- <sup>210</sup> Steenken S, Neta P. Electron transfer rates and equilibria between substituted phenoxide ions and phenoxyl radicals. *J Phys Chem* 1979; 83: 1134–7
- <sup>211</sup> Jovanovic SV, Tosic M, Simic MG. Use of Hammett correlation and  $\sigma^+$  for calculation of one-electron redox potentials of antioxidants. *J Phys Chem* 1991; 95: 10824–7
- <sup>212</sup> Koppenol WH. Oxyradical reactions: from bond-dissociation energies to reduction potentials. *FEBS Lett* 1990; 264: 165–7
- <sup>213</sup> Jovanovic SV, Jankovic I, Josimovic L. Electron transfer reactions of alkyl peroxy radicals. *J Am Chem Soc* 1992; 114: 9018–21
- <sup>214</sup> Jones DP, Carlson JL, Mody VC, Cai J, Lynn MJ, Sternberg P. Redox state of glutathione in human plasma. *Free Radic Biol Med* 2000; 28: 625–35