

Dietary Knowledge Among Female Diabetic Patients in Amman, Jordan

JAFAR M.F. EL-QUDAH

Department of Nutrition and Food Processing, Faculty of Agricultural Technology, Al-Balqa Applied University , Al-Salt 19117, Jordan.

<http://dx.doi.org/10.12944/CRNFSJ.4.2.04>

(Received: June 29, 2016; Accepted: July 28, 2016)

ABSTRACT

The aim of this study was to assess dietary knowledge among females with diabetes in Amman, Jordan. We performed a cross-sectional study of diabetic patients attending outpatient diabetes clinics, at several hospitals in Amman . A questionnaires and 24-hour dietary recall were completed by the participants. The study used a face to face interview during which a questionnaire was filled by the investigator about knowledge and attitudes of diabetic patients. The study sample was composed of 180 females. Most of the patients have diabetes for more than two years (78.3%) and had a positive family history of diabetes as a first degree relative. About 50% suffered from diabetes related health disease and about half of the patients were overweight and 32.2% were obese. More than half of patients (56.1%) gave wrong answers for meals recommended for diabetes/day and that should be used with hypoglycemia (44.5%), food rich in protein (47.6%), number of snacks (50.6%) and their role (43.1%), drinks permitted with unlimited amounts (61.4%) and suitable time to drink juice (78.1%). Most of the patients gave correct answers for food rich in fats (48.4%), cholesterol (47.8%), and controlling blood sugar (51.7%), right and number of snacks recommended (68.7%) and food which raises blood sugar (49.1%). According to the results, the dietary knowledge of diabetic patients is inadequate and need improvement. The study recommends that there is a need to set up a health education programme on diabetes mellitus which would also include a self-monitoring programme.

Keywords: Dietary knowledge , Diabetes mellitus , Food items , Jordan.

INTRODUCTION

Diabetes mellitus (DM) is a major health problem worldwide, that requires continuing medical care and ongoing patient self-management, education and support¹. According to WHO , the global average prevalence is around 10%². However the Arab countries have a higher prevalence of diabetes than the global average. Based on a study in 2010, in Jordan the prevalence of type 2 DM was 17.1%³. The prevalence rates of DM in the United Arab Emirates, Bahrain, Saudi Arabia and Kuwait were 20.1%, 20.1%, 30%, and 14.8% respectively^{4,5}.

Suitable dietary practices are basic and integral part treating DM and may reduce the development of disease complications⁶. In Jordan, around 45% of 195 diabetic patients previously diagnosed, had unsatisfactory glycemic control³. It is documented that poorly controlled diabetes leads to serious complication⁷⁻¹¹.

It was documented that diet control can improve glycemic control¹²⁻¹⁵. Adherence of diabetic patients to diabetic medications, to dietary advice and to physical exercise has been found to have a large effect on the degree of diabetic control^{3,16,17}. The aim of this study was to assess

dietary knowledge among female diabetic patients in Amman, Jordan.

METHODS

A convenience sampling technique was used to approach adult patients with DM, attending outpatient clinics from different hospitals in Amman, Jordan. One hundred and eighty diagnosed female diabetics were randomly selected with the help of the chief dietician at each hospital. Dietary knowledge was assessed by 14 multiple choice questions including food that raises and control blood sugar, food sources of nutrients, meals and snacks recommended. The response to each dietary knowledge question was scored from 0-2. With 2 for correct answer, 1 for incorrect and incomplete answer and 0 for wrong and don't answer. Total dietary score was ranged from 0 – 28 and was leveled as good when score is 20-28, fair 10-19 and poor 0-9. Perceived benefits items were measured as questions formatted as 3 points Likert scale with 3 for high perceived benefit. Total perceived benefits score was 12 – 36, which leveled into: 30 – 36 high perceived benefit, 21 – 29 moderate perceived benefit and 12– 20 low perceived benefits. Patients' files were reviewed to collect data on weight and height. The 24 hour recall was used to evaluate the food consumption pattern. The evaluation of the daily food intake was made by using ESHA Food Processor¹⁸, as well as Food Composition Tables for use in the Middle East¹⁹ and Food Composition Tables for Kingdom of Bahrain²⁰.

Statistical Methods

Data analysis was carried out by using the SPSS software package version 16. Absolute and relative frequencies and means were used as descriptive statistics. Multiple regression was used to identify the variables related to knowledge. The internal consistency of the questionnaire was checked by Cronbach's alpha and it was considered good (0.87 for total knowledge).

RESULTS

The study sample was composed of 180 females. As shown in Table (1), the majority of the patients were married (82.7%) in the age group 41-50 years (49.2%) and were housewives (62.9%). The

patients were uniformly distributed across different education levels and the majority with monthly family income between 300 and less than 800 Jordanian Dinar (JD) (1 US\$ = 0.71 JD). Most of the patients have diabetes for more than two years (78.3%) and had a positive family history of diabetes as a first degree relative. About 50% suffered from diabetes related health disease and about half of the patients were overweight and about 32% were obese. Table (2) shows dietary knowledge of diabetic patients. More than half of patients gave wrong answers for meals recommended for diabetes/day (56.1%) and that should be used with hypoglycemia (44.5%), food rich in protein (47.6%), number of snacks (50.6%) and their role (43.1%), drinks permitted with unlimited amounts (61.4%) and suitable time to drink juice (78.1%). Most of the patients gave correct answers for food rich in fats (48.4%), cholesterol (47.8%), and controlling blood sugar (51.7%), right and number of snacks recommended (68.7%) and food which raises blood sugar (49.1%).

Table 3 shows the perceived benefits of diabetic patients. Most of the patients perceived benefits of adherence of dieting regime (93.5%), regulating time and quantities of meals (65%), exercising (97.2%) and maintain or reducing weight (96.2%). A considerable percentage believed they should avoid starches food (51.3%) and honey will not affect blood sugar (57.4%). The benefit of herbal and traditional description in the management of diabetes is strong (48.9%). The assessment of daily nutrient intake of diabetic patients (Table 4) showed that the mean total energy intake was 1781.2 ± 671 kcal, the percentage of total calories from carbohydrate 56%, protein 14% and fat 30%. The mean dietary fiber intake was 24.3 ± 10.4 (gm). The percentage of total calories from carbohydrate was more than the American Diabetic Association guidelines (ADA 2007), (48 % of energy), while the percentage of total calories from protein and fat was within the ADA recommendations (10-20% and < 35 % of energy, respectively).

The result of the multiple regression analysis (Table 5) showed that occupation (p-value= 0.037), duration of diabetes (p-value= 0.001), family diabetes history (p-value= 0.014) and education level (p-value= 0.001) were the factors most influencing

Table1: Socioeconomic characteristics, medical and family history of the sample (N=180)

Variables	N (%)
Marital status	
Married	149 (82.8)
Not married	31 (17.2)
Age (years)	
≤ 40	54 (30.2)
41 – 50	89 (49.2)
> 50	37 (21.6)
Occupation	
Housewife	111 (62.9)
Working	69 (38.1)
Education level	
Illiterate	36 (20.0)
Primary and intermediate	58 (32.2)
Secondary	48 (26.7)
Graduate and above	38 (21.1)
Family income (J.D.)	
< 300	20 (11.1)
300 - < 500	78 (43.3)
500 – 800	52 (28.9)
> 800	30 (16.7)
Duration of diabetes (years)	
< 2	39 (21.7)
2 – 5	78 (43.3)
> 5	63 (35.0)
Family history of diabetes	
None	47 (26.1)
Relative 1st degree	129 (71.7)
Relative 2nd degree	4 (02.2)
Related health diseases	
Heart disease	8 (9.9)
Eye disease	21 (25.9)
Stroke	2 (2.5)
Neuropathy	7 (8.6)
Kidney disease	9 (11.1)
Thrombosis	6 (3.3)
Obesity	27 (33.3)
Body mass Index (Kg/m2)	
Normal	32 (17.8)
overweight	90 (50.0)
Obese	58 (32.2)
weight (kg)	79.2 ± 12.1
height (cm)	157 ± 7.1

knowledge. Those factors explained 35.4% of the total variation in knowledge.

DISCUSSION

In the past years many countries, have been experiencing a nutritional transition in food choices from the typical Mediterranean diet to the fast food pattern²¹. The rapid change in physical activity and food habits has led to the coexistence of many nutritional problems that underlies many chronic diseases. As populations become more urbanized, and as lifestyles shift towards reduced physical activity and increased food consumption, the prevalence of obesity is expected to rise.

We found in this study (Table 2) that more than half of patients gave wrong answers for meals recommended for diabetes/day (56.1%) and that should be used with hypoglycemia (44.5%), food rich in protein (47.6%), number of snacks (50.6%) and their role (43.1%), drinks permitted with unlimited amounts (61.4%) and suitable time to drink juice (78.1%). This is comparable to the results of a study conducted in Saudi Arabia by Z. Saadia, et al.²² who reported 90% of patients answered 50% of the knowledge questions correctly. Still a large proportion of population that is almost 40.3% were not able to score above 10. In another study, it was reported that only 40% of diabetic patients had good compliance with their diet²³. Improving the Knowledge of the diabetics in our society will not be an easy task. Great efforts would be needed by health teams to enhance education of the diabetic patient in order to promote compliance.

It is reported that patients had low level of knowledge in Iran²⁴, United Arab Emirates²⁵, Kuwait²⁶ and in Nigeria²⁷. On the other hand, a good level of knowledge was found among patients in Pakistan²⁸, Malaysia²⁹ and in Qatar³⁰. It was found that a good level of knowledge regarding DM, affect patients' adherence to pharmacological therapy^{31,32}, self-care activities³³ and good glycemic control^{11,34,35}.

Diet which contains high fat may cause a problem for people with diabetes³⁶. The more fat there is in the diet, the more difficult it is going to be for insulin to get the glucose into the cells³⁶. Our results showed that the mean dietary fiber intake

Table 2: Dietary Knowledge of diabetic patients (N=180)

Dietary Knowledge Questions	Correct	Answer Incomplete	Wrong or don't know
Which food raises the blood sugar level 1) vegetables 2) dates 3) rice 4) fats 5) don't know	88 (49.1)	59 (32.8)	33 (18.3)
Which food is rich in fat? 1) cucumber 2) Cheese 3) nuts 4) bread 5) don't know	105 (58.4)	50 (27.7)	25 (13.9)
Which food is protein? 1) Fish 2) sweet 3) oil 4) apple 5) don't know	79 (43.7)	15 (8.3)	86 (47.6)
Which food is high in cholesterol? 1) liver 2) kidney 3) eggs 4) all 1,2,3 5) don't know	86 (47.8)	11 (6.1)	83 (46.1)
Which food helps in controlling the blood sugar? 1) Fiber 2) fats 3) protein 4) all 1,2,3 5) don't know	93 (51.7)	76 (42.1)	11 (6.1)
Which food is high in fiber? 1) whole wheat 2) vegetables 3) fruits 4) faba bean 5) all 1,2,3 6) don't know	52 (28.7)	103 (57.1)	25 (13.9)
How many meals are recommended for diabetics/day 1) one 2) 2 3) 3 4) 4 5) don't know	79 (43.9)	0	101 (56.1)
How many times a diabetic is recommended to take snacks between recommended meals? 1) 1 2) 2 – 3 3) > 3 4) don't know	79 (43.7)	0	83 (50.6)
Type of snacks recommended 1) Milk 2) sweets 3) vegetables 4) fruits 5) Sandwich 6) don't know	124 (68.7)	39 (21.4)	17 (9.4)
Role of snack in diabetic diet 1) Prevent hypoglycemia 2) Prevent hyperglycemia 3) Don't know	98 (54.5)	3 (2.3)	78 (43.1)
Drinks permitted with unlimited amounts 1) low fat milk 2) sugar or tea with no sugar 3) juice with no sugar 4) don't know	32 (17.8)	37 (20.6)	111 (61.4)
Suitable time to drink juice 1) Before meals 2) as snack 3) With meals 4) don't know	37 (20.4)	2 (1.1)	141 (78.1)
What are the symptoms of hypoglycemia 1) Thirsty 2) frequent urinating 3) lost of appetite 4) hungry and blood beating 5) don't know	71 (39.2)	47 (26.1)	62 (34.5)
Which food should be used with hypoglycemia 1) Diet Pepsi 2) Juice 3) two-sugar cubes 4) cheese 5) don't know	70 (39.1)	30 (16.7)	80 (44.5)

Table 3: Dietary perceived benefits of diabetic patients (N=180)

	Agree	Disagree	Don't know
Adherence of dieting regime improves sugar	168 (93.5)	8 (4.7)	4 (2.2)
It is importance to regulate time and quantities of meals	117 (65)	41 (22.8)	22 (12.1)
Use brown bread will not raise sugar blood sugar (BS)	100 (55.3)	43 (23.9)	37 (20.8)
Dates will not affect (BS)	27 (15)	148 (82.2)	5 (2.8)
Honey will not affect (BS)	103 (57.4)	19 (10.6)	58 (32.1)
Drinks tea after meals will prevent BS absorption	15 (8.1)	79 (43.9)	86 (47.6)
Diabetic should avoid all starches	92 (51.3)	18 (10)	70 (38.9)
Diabetic should avoid all fruits as it raise BS	18 (10)	124 (68.7)	38 (21.3)
Foods that have bitter and sour taste may benefit diabetes	53 (29.6)	22 (12.2)	105 (58.6)
Regular exercise may benefit diabetes	175 (97.2)	0	5 (2.8)
It is importance to maintain or reduce your weight	173 (96.2)	7 (3.8)	0
Herbals & traditional prescriptions may benefit diabetic management	88 (48.9)	52 (28.9)	40 (22.2)

Table 4: Daily food intake of diabetic patients (N=180)

	Energy (kcal)	Carbohydrates (gm)	Fat (gm)	Protein (gm)	Fiber (gm)
Mean \pm SD	1781.2 \pm 671	264 \pm 83.9	61.4 \pm 37.1	81.5 \pm 31.2	24.3 \pm 10.4

Table 5: Results of multiple regression of factors influencing total knowledge score (N=180)

Factors	Total knowledge	p-value
Education level	1.66	0.001
Family diabetes history	1.21	0.014
Duration of diabetes	2.43	0.001
Occupation	0.878	0.037

was 24.3 \pm 10.4 (gm) (table 4) . Dietary fiber helps to slow the release of sugar into the bloodstream, thus helping to keep the blood sugar levels normal

Diet is an important part in the treatment of a diabetic patient. Following a healthy lifestyle , managing the weight and eating a balanced diet, will improve the patient's health enormously.

CONCLUSION

The dietary knowledge of diabetic patients among female Jordanians is inadequate and need improvement. Education and counseling about all the aspects of diabetes are needed. Studies with wider scope and much large sample size are recommended to confirm findings and explore relevant features .

REFERENCES

1. Al-Rasheedi, A. A. S. The Role of Educational Level in Glycemic Control among Patients with Type II Diabetes Mellitus. *International Journal of Health Sciences, Qassim University*, 8(2), 177-187 2014.
2. The determinants of health. Health Impact Assessment. The World Health Organization website, 2012.
3. Khattaba, M.; Khader, Y.S.; Al-Khawaldehd, A.; Ajlouni, K. Factors associated with poor glycemic control among Jordanian patients with Type 2 diabetes. *Journal of Diabetes and Its Complications*, 24, 84–89 2010.
4. Alqurashi, K.A.; Aljabri, K.S.; Bokhari, S.A. Prevalence of diabetes mellitus in a Saudi community. *Ann Saudi Med.*, 31(1), 19–23 2011.
5. Al-Akour, N. A.; Khader, Y. S.; Alaoui, A. M. Glycemic Control and Its Determinants among Patients with type 2 Diabetes Mellitus Attending a Teaching Hospital. *J Diabetes Metab*, 2, 4-8 2011.
6. Al-Kaabi, J.; Al-Maskari, F.; Saadi, H.; Afandi, B.; Parkar, H.; Nagelkerke, N. Assessment of Dietary Practice Among Diabetic Patients in the United Arab Emirates. *The Review of Diabetic Studies*, 5(2) 2008.
7. Al-Bdour, M.D.; Al-Till, M.I.; Samra, K.M. Risk factors for diabetic retinopathy among Jordanian diabetics. *Middle East Afr J Ophthalmol*, 15(2), 77-80 2008.
8. Jbour, A.S.; Jarrah, N.S.; Radaideh, A.M.; Shegem, N.S.; Bader, I.M.; Bataieha, A.M. Ajlouni, K.M. Prevalence and predictors of diabetic foot syndrome in type 2 diabetes mellitus in Jordan. *Saudi Med J*, 24(7) 761-764 2003.
9. Abdallah, S.; Ahmad, A.; Bataieha, A.; Ajlouni, K. Diabetes mellitus: the leading cause of hemodialysis in Jordan. *East Mediterr Health J*, 13(4) 803-809 2007.
10. Al-Amer, R.M.; Sobeh, M.M.; Zayed, A.A.; Al-Domi, H.A. Depression among adults with diabetes in Jordan: risk factors and relationship to blood sugar control. *J Diabetes Complicat*, 25(4) 247-252 2011.
11. Sawsan HAMMAD, Muhammad DARAWAD, Eman HOURANI, Waddah DEMEH. Predictors of Glycated Hemoglobin among Jordanian Diabetic Patients. *Iran J Public Health*, Vol. 44, No.11, Nov 2015, pp.1482-1491
12. Pastors, J.G.; Warshaw, H.; Daly, A.; Franz, M.; Kulkarni, K. The evidence for the effectiveness of medical nutrition therapy in diabetes management. *Diabetes Care*, 25(3) 608-613 2002.
13. Close, E.J.; Wiles, P.G.; Lockton, J.A.; Walmsley, D.; Oldham, J.; Wales, J.K. The degree of day-to-day variation in food intake in diabetic patients. *Diabet Med*. 10(6) 514 520 1993.
14. Monnier, L.; Grimaldi, A.; Charbonnel, B.; Iannascoli, F.; Lery, T.; Garofano, A. Childs M. Management of French patients with type 2 diabetes mellitus in medical general practice: report of the Mediab observatory. *Diabetes Metab*. 30(1) 35-42 2004.
15. Rivellese, A.A.; Boemi, M.; Cavalot, F.; Costagliola, L.; De Feo, P.; Miccoli, R.; Patti, L.; Trovati, M.; Vaccaro, O.; Zavaroni, I. Dietary habits in type II diabetes mellitus: how is adherence to dietary recommendations? *Eur J Clin Nutr* 2008. 62(5) 660-664 2008.
16. American Diabetes Association. Standards of Medical Care in Diabetes. *Diabetes care*, 36, supplements 1, 2013.
17. Aatur, R. K.; Zaki, N. A.; Mohammad, A. A.; Montaser, A.B.; Ibrahim, A.; Shabbir, A.K. Factors contributing to non-compliance among diabetics attending primary health centers in the Al Hasa district of Saudi Arabia. *J Family Community Med*. 19(1) 26–32 2012.
18. Food Processor SQL. Food Processor nutrition and fitness software. Food Processor SQL Inc., Salem, OR, USA 2008.
19. Pellett, P.L.; Shadarevian. S. Food Composition Tables for Use in the Middle East. American University of Beirut, Beirut, Lebanon, pp: 116 1970.
20. Musaiger, A.O. Food Composition Tables for Kingdom of Bahrain, Arab Center for Nutrition, first edition, Manama., Bahrain. 2011.

21. Musaiger, A.O.; Lloyd, O.L.; Al-Neyadi, S.M.; Bener, A.B. Lifestyle factors associated with obesity among male university students in the United Arab Emirates. *Nutrition & Food Science.*, 33(4) 145-14 2003.
22. Saadia, Z.; Rushdi, S.; Alsheha, M.; Saeed, H.; Rajab, M. A Study Of Knowledge Attitude And Practices Of Saudi Women Towards Diabetes Mellitus. A (KAP) Study In Al-Qassim Region. *The Internet Journal of Health.* 11(2) 2010.
23. Khattab, M.S.; Abofotouh, M.A.; Khan, M.Y.; Humaidi, M.A.; al- Kaldi, Y.M. Compliance and control of diabetes in a family practice setting, Saudi Arabia. *East Mediterr Health J* . 5(4) 755-765 1999.
24. Mansour-Ghanaei, R.; Joukar, F.; Soati, F.; Khanegha, A . Association between knowledge, locus of control and health belief with selfmanagement, HbA1c level and number of attendances in type 1 diabetes mellitus patients. *Int J Clin Exp Med* , 6(6) 470-477 2013.
25. Al-Maskari, F.; El-Sadig, M.; Al-Kaabi, J.; Afandi, B.; Nagelkerke, N.; Yeatts, K. Knowledge, attitude and practices of diabetic patients in the United Arab Emirates. *PLoS One*, 8(1) e52857 2013.
26. Al-Adsani, A.; Moussa, M.; Al-Jasem, L.; Abdella, N.; Al-Hamad, N. The level and determinanat of diabetes knowledge in Kuwaiti adults with type 2 diabetes. *Diabetes Metab J*, 35(2) 121-128 2009.
27. Yusuff, K.; Obe, O.; Joseph, B. Adherence to anti-diabetic drug therapy and self-management practices among type-2 diabetics in Nigeria. *Pharm World Sci*, 30(6) 876-883 2008.
28. Hawthorne, K.; Tomlinson, S. Pakistani muslims with Type 2 diabetes mellitus: effect of sex, literacy skills, known diabetic complications and place of care on diabetic knowledge, reported self-monitoring management and glycemic control. *Diabetic Med*, 16 (7): 591-7 1999.
29. Ng, S.; Chan, K.; Lian, Z.; Chuah, Y.; Waseem, A.; Kadirvelu, A. Reality vs illusion: knowledge, attitude and practice among diabetic patients. *Int J Collab Res Internal Med*, 4(5) 723-732 2012.
30. Mesmar, M.; Eljack, A.; Al-Kuwari, M. Knowledge and practice of type 2 diabetic patients attending primary health care in Qatar. *Middle East J Fam Med*, 9(4) 3-10 2011.
31. Al-Qazaz, H.; Sulaiman, S.; Hassali, M.; Shafie, A.; Sundram, S.; Al-Nuri, R.; Saleem, F. Diabetes knowledge, medication adherence and glycemic control among patients with type 2 diabetes. *Int J Clin Pharm*, 33(6) 1028-1035 2011.
32. Rubin, R. Adherence to pharmacologic therapy in patients with type 2 diabetes mellitus. *Am J Med*, (5A) 27S-34S 2005.
33. Jin, H.; Yuehua, L.; Yan, Z.; Hui, Y. Correlation between self-management and knowledge of and attitude to diabetes in type 2 diabetic patients in Changsha. *J Cent South Univ T*, 38(2) 176-181 2013.
34. Bain, S.; Egede, L. Associations between health literacy, diabetes knowledge, self-care behaviors and glycemic control in a low income population with Type 2 diabetes. *Diabetes Technol Ther*, 13(3) 335-341 2011.
35. Monnier, L.H.; Colette, C.; Aquirre, L.; Orsetti, A.; Combeaux, D. Restored synergistic enterohormonal response after addition of dietary fibre to patients which impaired glucose tolerance and reactive hypoglycemia. *Diabetes Metab.*, 8 217-22 1982.
36. Asif, M. The role of fruits, vegetables, and spices in diabetes. *International Journal of Nutrition Pharmacology Neurological Diseases.* 1(1) 27-35 2011.