



## ENDORSED POLICY STATEMENT

# Dietary Recommendations for Children and Adolescents: A Guide for Practitioners

American Heart Association, Samuel S. Gidding, MD, Chair, Barbara A. Dennison, MD, Cochair, Leann L. Birch, PhD, Stephen R. Daniels, MD, PhD, Matthew W. Gilman, MD, Alice H. Lichtenstein, DSc, Karyl Thomas Rattay, MD, Julia Steinberger, MD, Nicolas Stettler, MD, Linda Van Horn, PhD, RD

The American Heart Association makes every effort to avoid any actual or potential conflicts of interest that may arise as a result of an outside relationship or a personal, professional, or business interest of a member of the writing panel. Specifically, all members of the writing group are required to complete and submit a Disclosure Questionnaire showing all such relationships that might be perceived as real or potential conflicts of interest. A summary of all such reported relationships can be found at the end of this document.

## ABSTRACT

Since the American Heart Association last presented nutrition guidelines for children, significant changes have occurred in the prevalence of cardiovascular risk factors and nutrition behaviors in children. Overweight has increased, whereas saturated fat and cholesterol intake have decreased, at least as percentage of total caloric intake. Better understanding of children's cardiovascular risk status and current diet is available from national survey data. New research on the efficacy of diet intervention in children has been published. Also, increasing attention has been paid to the importance of nutrition early in life, including the fetal milieu. This scientific statement summarizes current available information on cardiovascular nutrition in children and makes recommendations for both primordial and primary prevention of cardiovascular disease beginning at a young age.

IT IS ESTIMATED that 75% to 90% of the cardiovascular disease epidemic is related to dyslipidemia, hypertension, diabetes mellitus, tobacco use, physical inactivity, and obesity; the principal causes of these risk factors are adverse behaviors, including poor nutrition.<sup>1-3</sup> The atherosclerotic process begins in youth, culminating in the risk factor-related development of vascular plaque in the third and fourth decades of life.<sup>4-6</sup> Good nutrition, a physically active lifestyle, and absence of tobacco use contribute to lower risk prevalence and either delay or prevent the onset of cardiovascular disease.<sup>2,3</sup> These observations have established the concept of prevention of the development of cardiovascular risk factors in the first place, now called primordial prevention.<sup>7</sup> Education, with the support of the health care community, combined with health policy and environmental change to support optimal nutrition and physical activity, are central to this health strategy.

This document provides dietary and physical activity recommendations for healthy children; discusses the current content of children's diets; reviews the adverse health consequences of increased intakes of calories (relative to energy expenditure), saturated and trans fat, and cholesterol; and provides age-specific guidelines for implementation of the recommended diet, including the period from before birth to 2 years of age. Medical practitioners are the intended audience, and guidelines to implement recommendations in clinical practice settings are provided. Public health strategies for improving the quality of children's diets are also discussed.

[www.pediatrics.org/cgi/doi/10.1542/peds.2005-2565](http://www.pediatrics.org/cgi/doi/10.1542/peds.2005-2565)

doi:10.1542/peds.2005-2374

This statement was endorsed by the American Academy of Pediatrics on August 24, 2005. All endorsements of statements by the American Academy of Pediatrics automatically expire 5 years after publication unless reaffirmed, revised, or retired at or before that time.

This statement was approved by the American Heart Association Science Advisory and Coordinating Committee on July 22, 2005.

Expert peer review of AHA Scientific Statements is conducted at the AHA National Center. For more on AHA statements and guidelines development, visit [www.americanheart.org/presenter.jhtml?identifier=3023366](http://www.americanheart.org/presenter.jhtml?identifier=3023366). ©2005 American Heart Association. Republished with permission. (*Circulation*. 2005;112:2061-2075). A correction appeared in *Circulation*. 2005;112:2375. This version reflects the corrected text.

### Key Words

AHA scientific statements, adolescents, children, diet, nutrition

### Abbreviations

AHA—American Heart Association  
FDA—Food and Drug Administration  
LDL—low-density lipoprotein

Address correspondence to American Heart Association Scientific Publishing Information. E-mail: [pubauth@heart.org](mailto:pubauth@heart.org)

PEDIATRICS (ISSN Numbers: Print, 0031-4005; Online, 1098-4275).

This scientific statement on optimal cardiovascular nutrition for infants, children, and adolescents revises the 1982 document on the same topic and also builds on the recent consensus statement on optimal nutrition for the prevention of many chronic diseases of adulthood.<sup>8,9</sup> This revision responds to the obesity epidemic that has emerged since the publication of the last statement that addressed children's nutrition from the American Heart Association (AHA) and has new focuses on both total caloric intake and eating behaviors.<sup>10,11</sup> This revision strongly conveys the message that foods and beverages that fulfill nutritional requirements are appropriate for growing and developing infants, children, and adolescents. Calorie-dense foods and beverages with minimal nutritional content must return to their role as occasional discretionary items in an otherwise balanced diet.

A critical component of contemporary guidelines is the strength of the scientific evidence base for recommendations. Whereas the scientific base for understanding the potential harm and benefit of current dietary practices and the relationship to risk factors is strong, the scientific base for recommended interventions is weaker for several reasons: limited number, statistical power, and scope of intervention studies; limited efficacy of attempted interventions; and lack of generalizability of studies of feeding behaviors at younger ages. Historically, most have had small sample size and have not had ethnic diversity among participants. Nonetheless, given the current obesity epidemic, sufficient natural history and prevalence data exist to justify intervention, although continued evaluation is necessary to identify optimal strategies.<sup>12</sup>

## DIETARY RECOMMENDATIONS

The general dietary recommendations of the AHA for those aged 2 years and older stress a diet that primarily relies on fruits and vegetables, whole grains, low-fat and nonfat dairy products, beans, fish, and lean meat.<sup>1,13</sup> These general recommendations echo other recent public health dietary guidelines in emphasizing low intakes of saturated and trans fat, cholesterol, and added sugar and salt; energy intake and physical activity appropriate for the maintenance of a normal weight for height; and adequate intake of micronutrients.<sup>14-16</sup> Tables 1 and 2 provide strategies for implementing healthy cardiovascular nutrition. The recently published Dietary Guidelines for Americans (for those 2 years of age and older) and American Academy of Pediatrics Nutrition Handbook provide important supporting reference information with regard to overall diet composition, appropriate caloric intakes at different ages, macronutrients, micronutrients, portion size, and food choices.<sup>14,17,18</sup> Table 3 provides daily estimated calorie and serving recommendations for grains, fruits, vegetables, and milk/dairy products by age and gender. Consistent with the *Dietary Guidelines for Americans, 2005*,<sup>14,18</sup> nutrient and energy

**TABLE 1 AHA Pediatric Dietary Strategies for Individuals Aged >2 Years: Recommendations to All Patients and Families**

---

Balance dietary calories with physical activity to maintain normal growth
60 min of moderate to vigorous play or physical activity daily
Eat vegetables and fruits daily, limit juice intake
Use vegetable oils and soft margarines low in saturated fat and trans fatty acids
instead of butter or most other animal fats in the diet
Eat whole-grain breads and cereals rather than refined-grain products
Reduce the intake of sugar-sweetened beverages and foods
Use nonfat (skim) or low-fat milk and dairy products daily
Eat more fish, especially oily fish, broiled or baked
Reduce salt intake, including salt from processed foods

---

**TABLE 2 Tips for Parents to Implement AHA Pediatric Dietary Guidelines**

---

Reduce added sugars, including sugar-sweetened drinks and juices
Use canola, soybean, corn oil, safflower oil, or other unsaturated oils in place of solid fats during food preparation
Use recommended portion sizes on food labels when preparing and serving food
Use fresh, frozen, and canned vegetables and fruits and serve at every meal; be careful with added sauces and sugar
Introduce and regularly serve fish as an entrée
Remove the skin from poultry before eating
Use only lean cuts of meat and reduced-fat meat products
Limit high-calorie sauces such as Alfredo, cream sauces, cheese sauces, and hollandaise
Eat whole-grain breads and cereals rather than refined products; read labels and ensure that "whole grain" is the first ingredient on the food label of these products
Eat more legumes (beans) and tofu in place of meat for some entrées
Breads, breakfast cereals, and prepared foods, including soups, may be high in salt and/or sugar; read food labels for content and choose high-fiber, low-salt/low-sugar alternatives

---

contributions from each food group are calculated according to the nutrient-dense forms of foods in each group (eg, lean meats and fat-free milk), with the exception of the guidelines for 1-year-old children, which included 2% fat milk. For youth 3 years of age and older, calorie estimates are based on a sedentary lifestyle. More physically active children and adolescents will require additional calories.<sup>14,17-19</sup> This table is provided as a starting point for dietary counseling; recommendations will need to be individualized in clinical practice. Table 4 provides daily recommended intakes of sodium, potassium, and fiber.<sup>18</sup> More complete guidelines for infants, particularly with regard to the transition from breast/formula-feeding to table foods, will be discussed below.

Emphases different from the past include the allowance of a more liberal intake of unsaturated fat and a focus on ensuring adequate intakes of omega-3 fatty acids. There is an emphasis on foods that are rich in nutrients and that provide increased amounts of dietary fiber. The AHA continues to recommend diets low in saturated and trans fats. Healthy foods include fruits, vegetables, whole grains, legumes, low-fat dairy products, fish, poultry, and lean meats. Fruits, vegetables, and fish are often inadequately consumed by children

**TABLE 3 Daily Estimated Calories and Recommended Servings for Grains, Fruits, Vegetables, and Milk/Dairy by Age and Gender**

	1 y	2–3 y	4–8 y	9–13 y	14–18 y
Kilocalories <sup>a</sup>	900	1000			
Female			1200	1600	1800
Male			1400	1800	2200
Fat, % of total kcal	30–40	30–35	25–35	25–35	25–35
Milk/dairy, cups <sup>b</sup>	2 <sup>c</sup>	2	2	3	3
Lean meat/beans, oz	1.5	2		5	
Female			3		5
Male			4		6
Fruits, cups <sup>d</sup>	1	1	1.5	1.5	
Female					1.5
Male					2
Vegetables, cups <sup>d</sup>	3/4	1			
Female			1	2	2.5
Male			1.5	2.5	3
Grains, oz <sup>e</sup>	2	3			
Female			4	5	6
Male			5	6	7

Calorie estimates are based on a sedentary lifestyle. Increased physical activity will require additional calories: by 0 to 200 kcal/day if moderately physically active and by 200 to 400 kcal/day if very physically active.

<sup>a</sup> For youth 2 years and older; adopted from Tables 2 and 3 and Appendix A-2 in US Department of Health and Human Services, US Department of Agriculture. *Dietary Guidelines for Americans*. 6th ed. Washington, DC: US Government Printing Office; 2005; www.healthier.us.gov/dietaryguidelines. Nutrient and energy contributions from each group are calculated according to the nutrient-dense forms of food in each group (eg, lean meats and fat-free milk).

<sup>b</sup> Milk listed is fat-free (except for children under the age of 2 years). If 1%, 2%, or whole-fat milk is substituted, this will utilize, for each cup, 19, 39, or 63 kcal of discretionary calories and add 2.6, 5.1, or 9.0 g of total fat, of which 1.3, 2.6, or 4.6 g are saturated fat.

<sup>c</sup> For 1-year-old children, calculations are based on 2% fat milk. If 2 cups of whole milk are substituted, 48 kcal of discretionary calories will be utilized. The American Academy of Pediatrics recommends that low-fat/reduced-fat milk not be started before 2 years of age.

<sup>d</sup> Serving sizes are 1/4 cup for 1 year of age, 1/3 cup for 2 to 3 years of age, and 1/2 cup for ≥4 years of age. A variety of vegetables should be selected from each subgroup over the week.

<sup>e</sup> Half of all grains should be whole grains.

**TABLE 4 Daily Recommended Intakes of Fiber, Sodium, and Potassium by Age and Gender**

Gender/Age	Fiber, g <sup>a</sup>	Sodium, mg	Potassium, mg
1–3 y	19	<1500	3000
4–8 y			
Female	25	<1900	3800
Male	25	<1900	3800
9–13 y			
Female	26	<2200	4500
Male	31	<2200	4500
14–18 y			
Female	29	<2300	4700
Male	38	<2300	4700

Adapted from 2005 Dietary Guideline Advisory Committee. Nutrition and your health: dietary guidelines for Americans. Available at: www.health.gov/dietaryguidelines/dga2005/report/HTML/E\_translation.htm.

<sup>a</sup> Total fiber preferred minimum 14 g/1000 kcal. Read labels to determine amounts on all packaged foods

and adolescents. Because the major sources of saturated fat and cholesterol in children's diets are full-fat milk and cheese and fatty meats, use of low-fat dairy products and lean cuts of meat in appropriate portion sizes will be critical in meeting dietary needs and nutrient requirements.<sup>20</sup>

Fish is an important food with growing evidence of potential benefit. However, consumers may have difficulty in distinguishing among several health messages about fish consumption. Although strong data associate cardiovascular disease prevention with increased fish consumption, there are also concerns about potential polycarbonate phenols (PCBs) and mercury contamination.<sup>21,22</sup> The Food and Drug Administration (FDA) and AHA stress that seafood is an important part of a healthy diet and advocate consumption of a wide variety of fish and shellfish. Current FDA recommendations with regard to limiting fish intake pertain to women who may become pregnant or are already pregnant, nursing mothers, and young children. The FDA recommends that people in those categories avoid shark, swordfish, king mackerel, and tilefish because they contain high levels of mercury. Five of the most commonly eaten varieties of fish are low in mercury (shrimp, canned light tuna, salmon, pollack, and catfish). The AHA continues to recommend 2 servings of fish weekly.<sup>23</sup> Recent evidence suggests that commercially fried fish products, likely because they are relatively low in omega-3 fatty acids and high in trans fatty acids (if hydrogenated fat is used for preparation), do not provide the same benefits as other sources of fish.<sup>24</sup>

#### DISCRETIONARY CALORIES

The obesity epidemic has prioritized consideration of the complex issue of matching appropriate energy intake to energy expenditure.<sup>10,11</sup> One approach is the concept of discretionary calories<sup>14</sup> illustrated in Fig 1. Total caloric




FIGURE 1

Concept of discretionary calories by gender. As daily physical activity increases, more energy is needed for normal growth. For sedentary children, only small amounts of discretionary calories can be consumed before caloric intake becomes excessive. Discretionary calories for children aged 4 to 8 years are based on 2 servings of dairy per day. Mod act indicates moderately active. Based on estimated calorie requirements and discretionary calories published in Dietary Guidelines for Americans (2005).<sup>14</sup>

intake is the sum of essential calories, the total energy intake necessary to meet recommended nutrient intakes, and discretionary calories, the additional calories necessary to meet energy demand and for normal growth.<sup>18</sup> The figure shows essential calories and discretionary calories; these increase with age and increasing levels of physical activity. There is a large difference in the discretionary calorie allowance among sedentary, moderately active, and active children, with more physically active children needing more energy from food to maintain normal growth. For young sedentary children, the amount of total energy intake that can come from foods used purely as a source of energy, ~100 to 150 calories, is less than that provided by a usual portion size of most low-nutrient-dense snacks and beverages. With increasing activity, this discretionary calorie amount may increase to 200 to 500 calories, depending on the age and gender of the child and the level of physical activity. The message portrayed by Fig 1 is clear: To be sedentary, have a nutritionally adequate diet, and to avoid excessive caloric intake in contemporary society is difficult.<sup>25</sup> The challenge to health care providers and public health professionals is to translate the complex science-based energy balance message from Fig 1 into effective practice and public health policy.<sup>25a</sup> Consuming diets that include primarily nutrient-dense forms of the foods listed in Table 3, participating in regular moderate to vigorous physical activity most days of the week for at least 1 hour per day, and limiting video screen time to <2 hours per day will help accomplish this goal.

### Scientific Support for Current Dietary Recommendations

The importance of dietary saturated and trans fat and cholesterol to the development of elevated cholesterol and subsequent cardiovascular disease as well as other cardiovascular risk factors has been extensively studied and reviewed. Pathologic evidence demonstrates that as the number of cardiovascular risk factors increases, so does the evidence of atherosclerosis in the aorta and coronary arteries beginning in early childhood.<sup>4,5,26</sup> Evidence of increased carotid artery intima-media thickness and coronary artery calcium measured by electron beam computed tomography among 29- to 39-year-old young adults who have been monitored from childhood further documents that the significant precursors of adult atherosclerosis are obesity, elevated blood pressure, and dyslipidemia.<sup>6,27,28</sup> Epidemiologic data from longitudinal studies provide further evidence that overweight, hypercholesterolemia, and hypertension track over time from childhood into adult life and that lifestyle choices, eg, diet, excess caloric intake, physical inactivity, and cigarette smoking, influence these risk factors.<sup>29–34</sup> Intervention studies aimed at measuring the efficacy and safety of diets reduced in total and saturated fat and cholesterol have also now contributed evidence at both the clinical and school-based levels.<sup>35–37</sup>

A meta-analysis of adult studies of low-saturated fat, low-cholesterol diets suggested that introduction of the diet lowers low-density lipoprotein (LDL) cholesterol an average of 12%, with a 1.93 mg/dL decline in LDL cholesterol for every 1% decline in saturated fat.<sup>38</sup> Further restricting saturated fat from 10% of total energy to 7% (the Therapeutic Lifestyle Change diet) increased the LDL cholesterol reduction to 16%.<sup>38,39</sup> Pediatric confirmation of adult studies showing safety and efficacy of a low-cholesterol and low-saturated fat diet has emerged. The Dietary Intervention Study in Children (DISC) was a randomized trial of a low-saturated fat, low-cholesterol diet conducted over 3 years in US children initially prepubertal and aged 8 to 11 years.<sup>35</sup> The Special Turku Risk Intervention Program (STRIP) was a randomized dietary intervention trial begun at weaning (age ~7 months) with parental dietary education continued through the age of 7 years.<sup>40–42</sup> Both studies achieved diets in intervention groups consistent with current recommendations for therapeutic lifestyle changes to lower elevated cholesterol levels, with total fat <30% of total calories and cholesterol intake <200 mg/day.<sup>39</sup> Saturated fat intake, although not <7% of total calories, was significantly less than in children assigned to usual care. Across a wide array of safety measures, including measures of growth, neurologic development, metabolic function, and nutrient adequacy, no adverse effects of the recommended intervention diets were observed.<sup>40,43–45</sup> LDL cholesterol levels were significantly lower among children receiving dietary intervention in the DISC study and in boys receiving dietary intervention in the STRIP study compared with controls.<sup>35,42</sup> Most importantly, in both studies children receiving dietary intervention were significantly more likely to make healthy food choices.<sup>25</sup> Three-year follow-up of children with severe hyperlipidemia who were following recommended therapeutic lifestyle changes showed no adverse effects on growth and development.<sup>46</sup>

The relationship between obesity and multiple cardiovascular risk factors, including elevated blood pressure, dyslipidemia, low physical fitness, and insulin resistance/diabetes mellitus, is well established.<sup>10,47–49</sup> Both excess caloric intake and physical inactivity are strongly associated with obesity.<sup>50</sup> Studies of weight loss in overweight individuals consistently show improvement in obesity-related comorbidities, particularly when interventions include regular exercise in the treatment program.<sup>51–53</sup> Population-based cross-sectional studies of secular trends in cardiovascular risk have shown strong associations between increasing prevalence of obesity and increasing blood pressure levels but inconsistent trends in dyslipidemia.<sup>47,54,55</sup> Longitudinal studies of secular trends in children, however, have shown strong relationships between increases in adiposity and adverse trends in blood pressure and lipids.<sup>56</sup> Maintaining body mass index (BMI) is beneficial, even without weight

loss, because this prevents worsening of risk status.<sup>57</sup> Maintenance of body weight during normal growth will improve BMI and cardiovascular risk status. Although primary prevention trials of reduction of daily caloric intake in at-risk children are under way, the evidence for harm from excess caloric intake is sufficient to support public health efforts for obesity prevention.<sup>11,58,59</sup>

### What Children Currently Eat

It is important to understand the gap between current dietary practices and recommended diets for infants, children, and adolescents. Sufficient population-based data exist to identify the magnitude of the problem confronting those interested in improving cardiovascular health in youth. Areas to consider include appropriateness of total caloric intake, eating patterns, balance of foods/beverages chosen from each food group, and intake of specific nutrients. Published data evaluate each of these areas with age and gender as important associated considerations.

For infants, it is encouraging that ~76% of mothers have initiated breastfeeding.<sup>60</sup> However, maintenance of breastfeeding for the first 4 to 6 months of life has been less successful. Only 4% of infants participating in the Special Supplemental Nutrition Program for Women, Infants, and Children (WIC) and 17% of the nonparticipants remain exclusively breastfed at 6 months of age. This suggests a strong socioeconomic status gradient in breastfeeding behavior. By 4 to 6 months, 66% of infants have received grain products, 40% vegetables, 42% fruits, 14% meat, and 0.6% some type of sweetened beverages.<sup>61</sup> By 9 to 11 months, 98% of infants have received grain products, 73% vegetables, 76% fruits, 79% meat, and 11% some type of sweetened beverages.<sup>61</sup> Sweetened beverages have been consumed by 28% of the 12- to 14-month-old children, 37% of the 15- to 18-month-old children, and 44% of the 19- to 24-month-old children.<sup>61</sup> During the transition from a milk-based diet to adult foods, the types of vegetables consumed change adversely. Deep-yellow vegetables are consumed by 39% of children at 7 to 8 months and by 13% at 19 to 24 months, whereas French fries become the most commonly consumed vegetable by this age.<sup>61</sup> Similarly, fruit consumption declines to the point where one third of 19- to 24-month-old children consume no fruit, whereas 60% consume baked desserts, 20% candy, and 44% sweetened beverages on a given day.<sup>61</sup>

Significant adverse changes have occurred in older children's food consumption.<sup>62</sup> These include a reduction in regular breakfast consumption, an increase in consumption of foods prepared away from the home, an increase in the percentage of total calories from snacks, an increase in consumption of fried and nutrient-poor foods, a significant increase in portion size at each meal, and an increase in consumption of sweetened beverages, whereas dairy product consumption has decreased, and

a shift away from high-fiber fruits and vegetables as well as a general decline in fruit and vegetable consumption other than potatoes.<sup>62-67</sup> Fried potatoes make up a substantial portion of the vegetable intake.<sup>67</sup> Sugar consumption has increased, particularly in preschool children.<sup>68</sup> With regard to micronutrients, the shift in dietary patterns has resulted in median intakes below recommended values of many important nutrients during adolescence.<sup>69</sup> Sodium intake is far in excess of recommended levels, whereas calcium and potassium intakes are below recommended levels.<sup>69-71</sup>

### Implementation of Dietary Recommendations Including Considerations for Specific Age Groups

This section reviews age-specific pediatric research on cardiovascular and general nutrition. Although in some areas there is a reasonable body of work about which to make useful judgments, in many areas studies have significant methodologic limitations: small sample size, confounding by a variety of factors (including cultural factors), and difficulty of using classic randomized trial designs to answer pertinent research questions. Nevertheless, the current dietary pattern of contemporary children mandates change. The recommendations provided herein are based on expert consensus of emerging evidence. Their purpose is to improve the nutritional quality, amount, and pattern of food consumption by children and their families. Although the narrative emphasizes nutrition to improve cardiovascular risk, it is recognized that optimal nutrition for overall health and normal growth is the preeminent goal.

Recommendations for children's nutrition consider the family and cultural milieu.<sup>72,73</sup> It has been decades since the majority of meals were consumed within the home.<sup>60-67</sup> Sources of nourishment include schools, child-care and after-school youth programs, restaurants, vending machines, convenience stores, work sites, and foods prepared by industry designed for minimal preparation time in the household.<sup>74</sup> Common situations affecting food preparation include households in which both parents work, single-parent households, and work schedules that demand that parents be away from home at mealtime. Likewise, children have complex schedules that demand frequent meals away from home. Schools provide less education on food preparation (eg, home economics) than in the past.

Culture-specific dietary practices can influence the diet both for better and for worse. A specific problem is the folk belief that a fat infant or chubby toddler is healthy. Popular fad diets often mix helpful and harmful components in their educational messages. Layered on top of this is a largely unregulated media dedicated to selling large quantities of a wide array of foods and food products of poor nutritional value. Despite unparalleled availability of nutrition resources, sifting through the food-message bombardment is often the most difficult

task facing a parent interested in providing proper nutrition for his or her family. Teaching those involved in supervision of children's diets to consume a healthful diet themselves and thus provide consistent role model behavior improves diet quality.<sup>73,75</sup> Table 5 provides a summary of areas in which adult influences are most important with regard to childhood nutrition.<sup>76–81</sup> Parents choose the time for meals and snacks and the types of foods and beverages to be served. Children can then choose how much to consume. Parents, guardians, and caregivers must provide appropriate role modeling through their own behavior, that is, influence children to “do as I do” rather than “do as I say.” A similar responsibility falls on those who attempt to provide reliable information to parents and educators in an effort to counterbalance adverse folk/cultural practices, media influences, and other sources of disinformation. Also critical to implementation of nutritional change is the social perception of risk.<sup>82</sup> Unless people believe that certain dietary practices are harmful or food providers believe that their actions endanger their clients, motivation to change will be limited. Increasing social pressure for eating properly can counteract the ubiquitous presence of food and food marketing of energy-dense, nutrient-poor choices.

### *Birth to 2 Years*

There has been considerable interest in the influence of both the intrauterine environment and infant nutrition on future cardiovascular risk.<sup>83,84</sup> It has been hypothesized that “programming” of future cardiovascular responses is established either in the womb or in response to feeding exposures early in life. Animal models support the programming hypothesis, but there are as yet few human experimental data.<sup>85–87</sup> Lower birth weight, because of presumed intrauterine malnutrition and association with rapid postnatal rapid weight gain, is associated with central adiposity, the metabolic syndrome, diabetes mellitus, and cardiovascular disease outcomes in adulthood.<sup>88</sup> Infants large for gestational age, probably through consequences of maternal insulin resistance

and glucose intolerance, are also at higher risk of future obesity.<sup>89,90</sup>

It is important for parents or parents-to-be to obtain a healthy weight, because children whose mothers are obese early in pregnancy are more likely to be overweight as young children.<sup>91</sup> A similar effect is seen in children whose parents are or become obese during their childhood.<sup>31</sup> To ensure optimal growth of the fetus, pregnant women must optimize their nutrition and weight gain during pregnancy, according to the Institute of Medicine guidelines.<sup>92</sup> Excessive maternal weight gain has been associated with a two- to threefold increased risk that the mother will be overweight after a pregnancy.<sup>93</sup> This may increase subsequent offspring risk during adolescence for obesity, impaired glucose tolerance, impaired insulin secretion, and type 2 diabetes. Studies of maternal nutrition, for example, assessments of protein and calcium intake, suggest that maternal diet during pregnancy may influence offspring's blood pressure.<sup>94,95</sup> However, evidence is insufficient to make specific recommendations about nutrition during pregnancy based on future cardiovascular disease.

Human milk is uniquely superior for infant feeding and is the reference against which other infant feeding strategies must be measured.<sup>96</sup> Breast milk is rich in both saturated fat and cholesterol but low in sodium. There has been substantial work on the relationship of breastfeeding to both future cardiovascular events and cardiovascular risk factors. Although pooling estimates from these studies is difficult because of differences in exposure and outcome assessment, recent meta-analyses have suggested no meaningful impact of breastfeeding on subsequent cardiovascular or all-cause mortality in adulthood.<sup>97</sup> Other systematic reviews, however, suggest benefits of breastfeeding, particularly in the prevention of future obesity.<sup>98,99</sup> Several studies suggest that breastfeeding leads to lower blood pressure later in childhood.<sup>100,101</sup> Although breastfeeding is associated with higher blood cholesterol levels at 1 year of age, it may also result in lower blood cholesterol levels in adults.<sup>102</sup> Rapid weight gain during the first 4 to 6 months of life is associated with future risk of overweight<sup>103,106</sup>; studies suggest that partially breastfed and formula-fed infants consume 20% more total calories per day than do exclusively breastfed infants.<sup>104,105</sup> Physicians should identify infants who are gaining weight rapidly and/or whose weight-to-length percentile exceeds the 95th percentile to help correct overfeeding if present.

At least 2 behavioral benefits of breastfeeding may lead to reduced cardiovascular risk, but the impact of these has not been studied in large trials.<sup>107–110</sup> The first potential benefit may be better self-regulation of intake. Compared with parents who bottle-feed, mothers who breastfeed appear to allow the infant to take an active role in controlling intake, possibly promoting maternal feeding practices that can foster better self-regulation of

**TABLE 5** Parent, Guardian, and Caregiver Responsibilities for Children's Nutrition

---

Choose breastfeeding for first nutrition; try to maintain for 12 mo
Control when food is available and when it can be eaten (nutrient quality, portion size, snacking, regular meals)
Provide social context for eating behavior (family meals, role of food in social intercourse)
Teach about food and nutrition at the grocery store, when cooking meals
Counteract inaccurate information from the media and other influences
Teach other care providers (eg, daycare, babysitters) about what you want your children to eat
Serve as role models and lead by example; “do as I do” rather than “do as I say”
Promote and participate in regular daily physical activity

---

energy intake as the child grows up.<sup>108</sup> Children with improved self-regulation may better withstand the current food-surplus environment.<sup>111</sup> The second potential benefit relates to taste preference.<sup>112–116</sup> Both amniotic fluid and breast milk provide flavor exposure to the fetus and infant. These exposures influence taste preference and food choices after weaning. Thus, exposure to healthier foods through maternal food consumption during pregnancy and lactation may improve acceptance of healthy foods after weaning. Because infant responses to taste are different from mature taste, these early exposures may be critical in determining food preference later in life.

A critical social problem for mothers interested in breastfeeding in the United States is the lack of a tolerant social structure.<sup>117</sup> Breastfeeding rates decline rapidly between 2 and 3 months, which is when many mothers return to work or school.<sup>118,119</sup> Full-time employment is consistently associated with shorter periods of breastfeeding.<sup>120–123</sup> In Scandinavian countries, where women routinely receive paid maternity benefits (eg, 42 weeks with full pay or 52 weeks at 80% of salary in Norway), women far surpass the US Healthy People 2010 goals for breastfeeding. In Norway, 97% of women are breastfeeding when they leave the hospital, 80% are breastfeeding at 3 months, and 36% are breastfeeding at 12 months.<sup>124</sup> Most African and Asian countries are highly supportive of breastfeeding. Policies enacted within the workplace and public places can also help to overcome barriers to breastfeeding.<sup>117,125</sup>

The period from weaning to consumption of a mature diet, from 4 to 6 months to ~2 years of age, represents a radical shift in pattern of food consumption (Fig 2),<sup>17,126,126a,129,130</sup> but there has been very little research on the best methods to achieve optimal nutritional intakes during this transition. Infants mature from receiving all nutrition from a milk-based diet to a diet chosen from the range of adult foods, in part self-selected and in part provided by caregivers. Transition to other sources of nutrients should begin at 4 to 6 months of age to ensure sufficient micronutrients in the diet, but the best methods for accomplishing this task are essentially unknown.<sup>15,126</sup> Current feeding practices and guidelines are influenced by small-scale studies of infant feeding behavior, idiosyncratic parental behavior, and popular opinion.<sup>17,60,127,128</sup>

Food-consumption data suggest that infants are currently exposed to a wide variety of “kid” foods that tend to be high in fat and sugar, including excess juice, juice-based sweetened beverages, French fries, and nutrient-poor snacks.<sup>61</sup> Usual food intakes of infants and young children may exceed estimated energy requirements. For infants aged 0 to 6 months, reported intakes exceed requirements by 10% to 20%; for children aged 1 to 4 years, intakes exceed requirements by 20% to 35%. Although some of these reports may reflect overreport-




FIGURE 2

Dramatic change in food sources during the first 2 years of life. Diet is initially based on breast milk at birth and transitions to conventional foods by 2 years, although dairy products remain a major source of energy and nutrition. (Reproduced with permission from Lederman SA, Akabas SR, Moore BJ, et al. Summary of the Presentations at the Conference on Preventing Childhood Obesity, December 8, 2003. *Pediatrics*. 2004;114:1146–1173; adapted from Briefel RR, Reidy K, Karwe V, Devaney B. Feeding infants and toddlers study: improvements needed in meeting infant feeding recommendations. *J Am Diet Assoc*. 2004;104(suppl 1):s31–s37 and Skinner JD, Ziegler P, Ponza M. Transitions in infants' and toddlers' beverage patterns. *J Am Diet Assoc*. 2004;104(suppl 1):s45–s50.)

ing of food intake, these data might also explain the rise in the prevalence of overweight at very young ages.<sup>129,130</sup>

For those participating in public nutrition assistance programs (US Department of Agriculture 2002), the foods supplied for infants and children are limited in variety, reflecting more closely the nutritional concerns of the 1970s, when the program was designed (inadequate calories, protein, vitamin A, vitamin C, and iron), than nutritional concerns today (excess calories, fat, and sugar and inadequate fruits, vegetables, and whole grains).<sup>131</sup> Moreover, beverages provided to most children are not optimal. Children aged 1 to 5 years enrolled in the Special Supplemental Nutrition Program for Women, Infants, and Children receive twice the amount of fruit juice (9.5 fl oz/day) currently recommended, and most participants also receive or choose whole milk.<sup>14,17,131,132</sup> For formula-fed infants, there may also be a role for clearer prescriptive feeding advice for parents to understand their infant's satiety cues and appropriate energy intake than is currently the norm. Table 6 provides a number of strategies to improve general and cardiovascular nutrition during this transitional stage. When normal growth is present, overfeeding may result from arbitrarily increasing amounts fed to achieve specific portion sizes per meal rather than allowing infants and toddlers to self-regulate. New healthy foods may need to be introduced repeatedly, as many as 10 times, to establish taste preferences.<sup>133</sup>

**TABLE 6 Improving Nutritional Quality After Weaning**


---

Maintain breastfeeding as the exclusive source of nutrition for the first 4–6 mo of life
Delay the introduction of 100% juice until at least 6 mo of age and limit to no more than 4–6 oz/day; juice should only be fed from a cup
Respond to satiety clues and do not overfeed; infants and young children can usually self-regulate total caloric intake; do not force children to finish meals if not hungry, because they often vary caloric intake from meal to meal
Introduce healthy foods and continue offering if initially refused; do not introduce foods without overall nutritional value simply to provide calories

---

**Age 2 to 6 Years**

At this age, recommendations for diet content are similar to those for older individuals. Challenges here relate to providing quality nutrient intake and avoiding excess caloric intake. Dairy products are a major source of saturated fat and cholesterol in this age group, and therefore a transition to low-fat milk and other dairy products is important.<sup>34,134</sup> Sweetened beverages and other sugar-containing snacks are a major source of caloric intake.<sup>135,136</sup> Table 7 provides a list of strategies for managing nutrition in young children.<sup>77,78,137–141</sup> Parents should remember that they are responsible for choosing foods that are eaten and when and where they are eaten. The child is responsible for whether he or she wants to eat and how much. Two natural parental impulses, pressuring children to eat and restricting access to specific foods, are not recommended because they often lead to overeating, dislikes, and paradoxical interest in forbidden items.<sup>142,143</sup>

Health care providers must provide useful advice to parents, but they are constrained by time pressures in the typical health maintenance office visit. In addition to the information in Table 7, advice on caloric/energy values of food, particularly nutrient-poor foods, can be provided in a relatively short period of time. At office visits, BMI percentile can be plotted, the appropriateness of weight gain in the last year can be assessed from standard growth curves, and recommendations for opti-

**TABLE 7 Improving Nutrition in Young Children**


---

Parents choose meal times, not children
Provide a wide variety of nutrient-dense foods such as fruits and vegetables instead of high-energy-density/nutrient-poor foods such as salty snacks, ice cream, fried foods, cookies, and sweetened beverages
Pay attention to portion size; serve portions appropriate for the child's size and age
Use nonfat or low-fat dairy products as sources of calcium and protein
Limit snacking during sedentary behavior or in response to boredom and particularly restrict use of sweet/sweetened beverages as snacks (eg, juice, soda, sports drinks)
Limit sedentary behaviors, with no more than 1–2 h/day of video screen/television and no television sets in children's bedrooms
Allow self-regulation of total caloric intake in the presence of normal BMI or weight for height
Have regular family meals to promote social interaction and role model food-related behavior

---

mal weight gain in the next year can be given. Blood pressure screening and cholesterol measurement, if indicated, are begun in this age range.<sup>7</sup>

**Ages 6 Years and Above**

As children grow up, sources of food and influences on eating behavior increase. Social constraints on families may necessitate the presence of multiple caregivers, eating out, and frequent fast food consumption. Many children, because of parental work schedules, are home alone and prepare their own snacks and meals. By early adolescence, peer pressure begins to usurp parental authority, and fad diets may be initiated. Many meals and snacks are routinely obtained outside the home, often without supervision. Sites include schools, friends' homes, child-care centers, and social events. Older children have discretionary funds to use for self-selected foods. Current eating patterns do not at all resemble the "norm" of providing at least breakfast, dinner, and a single snack at home with lunch carried to school or purchased from a health-conscious cafeteria. For example, current diet studies suggest that many children do not eat breakfast and get at least one third of calories from snacks. Sweetened beverage intakes contribute significantly to total caloric intake.<sup>144</sup> Sweetened beverages and naturally sweet beverages, such as fruit juice, should be limited to 4 to 6 oz per day for children 1 to 6 years old, and to 8 to 12 oz per day for children 7 to 18 years old.<sup>144,145</sup> These snacks often contribute to excess consumption of discretionary calories and/or supplant the intake of foods containing essential nutrients.<sup>25,146</sup>

Adolescence is a nutritionally vulnerable developmental stage because growth rate accelerates. Amplified caloric and global nutrition needs due to pubertal growth stimulate appetite. The combination of centrally driven appetite stimulation and an increasingly sedentary lifestyle due to a decline in recreational sports participation augments obesity.<sup>50</sup> Parallel to the psychosocial transition from dependence on parental authority to independent thought processes, food choices and purchases are increasingly made by the adolescent. Peer pressure for conformity, in part driven by media promotion of fast food directly to teens, makes overeating natural. Currently, adolescents have an increased intake of sweetened beverages, French fries, pizza, and fast food entrées, including hamburgers, and a consequent lack of recommended fruits, vegetables, dairy foods, whole grains, lean meats, and fish. This change in eating pattern results in consumption of excess fat, saturated fat, trans fats, and added sugars along with insufficient consumption of micronutrients such as calcium, iron, zinc, and potassium, as well as vitamins A, D, and C and folic acid.<sup>146,147</sup>

Counseling of older children and adolescents must be individualized to accommodate the range of contemporary lifestyles; less success is achieved at older ages.



Current dietary practices and readiness to change must be understood before family-based intervention is attempted.<sup>148</sup> Parental role modeling is important in establishing children's food choices.<sup>53,78,132</sup> Depending on their own food choices, parents can be either positive or negative role models.<sup>81</sup>

### Public Health Issues

Modern life extends the umbrella of social responsibility for provision of appropriate nutrition and nutrition knowledge beyond the home to government, the health professions, schools, the food industry, and the media. It is beyond the scope of this document to evaluate the large public health effort related to overweight and nutrition now being undertaken. Some important areas are highlighted below. Because there is little scientific information to guide current policy directed at changing eating behaviors, it is strongly recommended that evaluation, safety, and efficacy tools be incorporated into policy implementation.

Schools have become a battleground for fighting the obesity epidemic.<sup>149,150</sup> Cafeterias are under attack for serving unhealthy food, yet the food provided is constrained by budgetary and regulatory issues largely external to public health concerns. US Department of Agriculture guidelines require school food programs to provide minimum quantities of specific nutrients over a 3- to 7-day span but do not address maximum food amounts. Vending machines and competing nutrient-poor foods provide excess calories but also provide revenue to support school programs. Table 8 summarizes some strategies currently being implemented in many locales.

Nutrition education in schools is considered useful in improving knowledge about nutrition, but few studies suggest that it is effective in altering eating behaviors in the absence of environmental change.<sup>150</sup> The largest study, the Child and Adolescent Trial for Cardiovascular

Health (CATCH), was a multicenter intervention that included nutrition education, a cafeteria intervention, changes in physical education programs, and parental education. The fat content of school lunch but not school breakfast was modified, and blood cholesterol levels and children's weight status were unchanged.<sup>36,151,152</sup> Other school studies have shown improvements in fruit (but not vegetable) intake.<sup>153-155</sup> High-quality foods sold in vending machines will be selected if they are competitively priced.<sup>62</sup> An intervention that included nutrition education, a cafeteria intervention, changes in physical activity, and a parent component for younger children attending Head Start programs was successful in decreasing children's blood cholesterol levels but did not affect the prevalence of childhood overweight.<sup>156</sup>

Physical education programs are often subject to budget constraints. The percentage of pupils attending physical education classes has decreased. For example, in a study of a representative sample of US high schools, participation rates fell from 79% in ninth grade to 36% in 12th grade.<sup>157</sup> In addition, participation in school-sponsored after-school teams is frequently limited to elite athletes, limiting the opportunity for high school-aged students to engage in regular physical activity.

Media has a pervasive influence on children's food choices. Children, including the very young, are heavily marketed by the food industry. Time spent watching television is directly related to children's food requests. The most frequently advertised foods are high-sugar breakfast cereals, fast food restaurant products, sweetened beverages, frozen dinners, cookies, and candy. Fruits and vegetables are almost never advertised. Watching television during meals is associated with increased frequency of poor food choices and decreased frequency of good choices.<sup>77,158-160</sup> Several European countries now have restrictions on advertising to children as well as school-based marketing.

State and local governments are now becoming active in the effort to control obesity on a wide variety of fronts. For example, several states have adopted legislation mandating that school staff report to parents the BMI status of their children. Changes in food labeling, taxes on certain types of foods, restrictions on foods provided to children in government-sponsored programs, and requiring restaurants to provide nutrition information are examples of regulations under consideration.<sup>11,161</sup> Strategy types are summarized in Table 9. Given the widening discrepancy between recommended dietary guidelines and current dietary intake, a reevaluation of federal agricultural policies may be warranted. Strategies for food subsidies and taxation should reflect health goals. Foods made available and served through public nutrition programs must be consistent with current recommendations.

**TABLE 8 Strategies for Schools**

Identify a "champion" within the school to coordinate healthy nutrition programs
Establish a multidisciplinary team including student representation to assess all aspects of the school environment using the School Health Index (Centers for Disease Control and Prevention) or similar assessment
Identify local, regional, and national nutrition programs; select those proven effective ( <a href="http://www.ActionForHealthyKids.org">www.ActionForHealthyKids.org</a> )
Develop policies that promote student health and identify nutrition issues within the school ( <a href="http://www.nasbe.org/HealthySchools/healthy_eating.html">www.nasbe.org/HealthySchools/healthy_eating.html</a> )
Work to make predominantly healthful foods available at school and school functions by influencing food and beverage contracts, adapt marketing techniques to influence students to make healthy choices, and restrict in-school availability of and marketing of poor food choices
Maximize opportunities for all physical activity and fitness programs (competitive and intramural sports); utilize coaches/teachers as role models
Lobby for regulatory changes that improve a school's ability to serve nutritious food
Ban food advertising on school campuses

**TABLE 9** Types of Legislation Under Consideration to Improve Children's Nutrition

Measurement of BMI by school staff for health surveillance and/or to report information to parents
Restriction of certain types of food and beverages available on school grounds
Taxation of specific foods or sedentary forms of entertainment
Establishment of local school wellness policies using a multidisciplinary team of school staff and community volunteers (mandated for schools participating in federal reimbursable school lunch, breakfast, or milk programs)
Food-labeling regulations, including appropriate descriptions of portion sizes (eg, a medium-sized sugar-containing drink should be 6–8 oz)
Regulation of food advertising directed at children

### Therapeutic Lifestyle Changes for Treatment of Hypertension and Hypercholesterolemia

There are currently established consensus guidelines for the role of diet in the management of children with established cardiovascular risk factors. Cut points for diagnosing dyslipidemia and hypertension are provided in Table 10.<sup>7,39,162,163</sup> The Fourth Pediatric Report of the National High Blood Pressure Education Program recommends a diet consistent with the current recommendations for children with hypertension.<sup>162</sup> For overweight children, weight loss is the initial therapeutic strategy. The Dietary Approaches to Stop Hypertension (DASH) study has recently shown that implementation of a diet rich in fruits, vegetables, nonfat dairy products, and whole grains can effectively lower blood pressure in adults with hypertension.<sup>164</sup> Although there are no comparable clinical trial data in children, there is no reason to suspect that the DASH diet would not be safe to implement in older children and adolescents as long as protein and calorie needs are met.<sup>165,166</sup>

There has not been an update of the Report of the

**TABLE 10** Consensus Guidelines for Diagnosis of Hypertension and Dyslipidemia in Children

Hypertension	Guideline
Prehypertension	Systolic or diastolic blood pressure >90th percentile for age and gender or 120/80 mm Hg, whichever is less
Stage 1 hypertension	Systolic or diastolic blood pressure >95th percentile for age and gender on 3 consecutive visits or 140/90 mm Hg, whichever is less
Stage 2 hypertension	Systolic or diastolic blood pressure >99th percentile +5 mm Hg for age and gender or 160/110 mm Hg, whichever is less
Total cholesterol, mg/dL	
Borderline	≥170
Abnormal	≥200
LDL cholesterol, mg/dL	
Borderline	≥100
Abnormal	≥130
HDL cholesterol, mg/dL	
Abnormal	<40
Triglycerides, mg/dL	
Abnormal	≥200

HDL indicates high-density lipoprotein.

Expert Panel on Blood Cholesterol Levels in Children and Adolescents published since its publication in 1992, but the National Cholesterol Education Program (NCEP) generally recommends restriction of saturated fat intake to <7% of total calories and restriction of cholesterol intake to <200 mg/day for treatment of elevated LDL cholesterol levels.<sup>39,163</sup> There are now data from randomized trials demonstrating that such diets are safe in children as young as 7 months of age.<sup>40,42,44,45</sup> Efficacy is variable, however, and unless the diet is extremely high in saturated fat before changes are made, it is unlikely that diet alone will be sufficient to achieve target levels for LDL cholesterol in those with genetic dyslipidemias and LDL cholesterol ≥190 mg/dL. Increased intake of soluble fiber is recommended as an adjunct to the reduced intakes of saturated fatty acids and cholesterol.<sup>14,167</sup> Recently plant sterols and stanols have been used, often in margarines, to lower LDL cholesterol through inhibition of cholesterol absorption. Adult studies have shown reductions of 4% to 11% without adverse events.<sup>168</sup> One randomized, controlled trial in children showed that 20 g/day of plant sterol-containing margarine lowered LDL cholesterol 8%.<sup>169</sup> These products may be used, although caution is recommended with regard to the potential for decreased absorption of fat-soluble vitamins and  $\beta$ -carotene. Formal recommendation of their use for children awaits clinical trial data.<sup>42,170–177</sup>

### SUMMARY

This scientific statement updates nutrition recommendations for the promotion of cardiovascular health among children. Recommendations have been made with regard to diet composition, total caloric intake, and physical activity. Implementation requires that children and all other members of their households actively make the recommended changes. Adverse recent trends in children's diets have been noted. Cardiovascular nutrition issues surrounding the first 2 years of life have been addressed. Strategies to improve implementation of the recommended diet have been presented. A brief overview of public health issues related to nutrition is included.

### ACKNOWLEDGMENTS

The authors acknowledge the contributions of many additional experts who reviewed portions of the material presented or supplied additional expertise. These include Julie Mennella, PhD, Gary A. Emmett, MD, and Penny Kris-Etherton, PhD. Carol Muscar provided invaluable support in the preparation of the manuscript and its editing. The AHA thanks the Nemours Health and Prevention Services group for providing meeting space and organizational support to the investigators and AHA staff, facilitating the rapid completion of this report.

## Writing Group Disclosures

Writing Group Member	Employment	Research Grant	Other Research Support	Speakers Bureau/Honoraria	Ownership Interest	Consultant/Advisory Board	Other
Samuel Gidding	Nemours Foundation	None	None	None	None	None	None
Barbara Dennison	New York State Department of Health	National Institute of Diabetes and Digestive and Kidney Diseases—grantee USDA—Improving Human Nutrition—grantee	None	None	None	Bassett Healthcare—Research Scientist; Mead Johnson Nutritionals—Consultant Toddler/Children Panel; American Dairy Association—Consultant; University of North Carolina—Consultant	None
Stephen Daniels	Children’s Hospital Cincinnati	None	None	None	None	None	None
Linda Van Horn	Northwestern University—Feinberg School of Medicine	None	None	None	None	<i>Journal of the American Dietetic Association</i> —Editor in Chief	None
Julia Steinberger	University of Minnesota	None	None	None	None	American Phytotherapy Research Lab—Consultant	None
Alice Lichtenstein	Tufts University	None	None	None	None	None	None
Leann Birch <sup>a</sup>	Pennsylvania State University	National Institute of Child Health & Human Development—Grantee; Dairy Management Inc—Grantee; USDA—CSREES—Grantee (coinvestigator)	None	None	None	Institute of Medicine Committee on Prevention of Childhood Obesity in Children and Youth—Chair	None
Nicolas Stettler	University of Pennsylvania School of Medicine	None	None	None	None	European Society for Pediatric Research—Member; International Epidemiological Association—Member; World Heart Federation—Member; Swiss Pediatric Society—Member; Swiss Medical Society—Member; American Society for Nutritional Sciences—Member; American Society for Clinical Nutrition—Member; North American Association for the Study of Obesity—Member; Children Are Our Messengers: Changing the Health Message, International Society on Hypertension in Blacks—Advisory Committee Member; Community Health Centers, Child Health Project, US Department of Health and Human Services—Consultant; Early Origins of Adult Health Working Group, National Children’s Study—Core Member; Society for Pediatric Research—Member; Comprehensive School Nutrition Policy Task Force, US Department of Agriculture/Food—Consultant; National High Blood Pressure Education Program on Blood Pressure in Children and Adolescents; Institute of Medicine Committee on Nutrient Relationships in Seafood	None
Matthew Gillman	Harvard University	None	None	None	None	None	None
Karyl Rattay	Nemours Health and Prevention Services	None	None	None	None	None	None

CSREES indicates Cooperative State Research, Education, and Extension Service. This table represents the relationships of writing group members that may be perceived as actual or reasonably perceived conflicts of interest as reported on the Disclosure Questionnaire that all members of the writing group are required to complete and submit. A relationship is considered “significant” if (1) the person receives  $\geq$ \$10 000 during any 12-month period or  $\geq$ 5% of the person’s gross income or (2) the person owns  $\geq$ 5% of the voting stock or share of the entity or owns  $\geq$ \$10 000 of the fair market value of the entity. A relationship is considered “modest” if it is less than “significant” under the preceding definition.

<sup>a</sup> Modest.

## Reviewer Disclosures

Reviewer	Employment	Research Grant	Other Research Support	Speakers Bureau/Honoraria	Ownership Interest	Consultant/Advisory Board	Other
Frank Greer	University of Wisconsin—Madison	None	None	None	None	None	None
Nancy F. Krebs	The Children’s Hospital	None	None	None	None	None	None
Kristie Lancaster	New York University	None	None	None	None	None	None
William Neal	West Virginia University	None	None	None	None	None	None
Theresa A. Nicklas	Baylor College of Medicine	USDA; National Institutes of Health	Dairy Management Inc; National Cattlemen’s Beef Association; Mars, Inc	National Dairy Council Speakers Bureau; National Cattlemen’s Beef Association Speakers Bureau	None	Brands Global Advisory Council on Health, Nutrition and Fitness; US Potato Board’s Scientific Advisory Panel; Cadbury Scientific Advisory Committee; Grain Foods Foundation Scientific Advisory Board	International Food Information Council Expert Resource to Media; 2005 Dietary Guidelines Advisory Committee

This table represents the relationships of reviewers that may be perceived as actual or reasonably perceived conflicts of interest as reported on the Disclosure Questionnaire that all reviewers are required to complete and submit.

## REFERENCES

- Kavey RE, Daniels SR, Lauer RM, et al. American Heart Association guidelines for primary prevention of atherosclerotic cardiovascular disease beginning in childhood. *Circulation*. 2003;107:1562–1566
- Stamler J, Stamler R, Neaton JD, et al. Low risk-factor profile and long-term cardiovascular and noncardiovascular mortality and life expectancy: findings for 5 large cohorts of young adult and middle-aged men and women. *JAMA*. 1999;282:2012–2018
- Daviglus ML, Stamler J, Pirzada A, et al. Favorable cardiovascular risk profile in young women and long-term risk of cardiovascular and all-cause mortality. *JAMA*. 2004;292:1588–1592
- Relationship of atherosclerosis in young men to serum lipoprotein cholesterol concentrations and smoking: a preliminary report from the Pathobiological Determinants of Atherosclerosis in Youth (PDAY) Research Group. *JAMA*. 1990;264:3018–3024
- Berenson GS, Srinivasan SR, Bao W, Newman WP 3rd, Tracy RE, Wattigney WA. Association between multiple cardiovascular risk factors and atherosclerosis in children and young adults. The Bogalusa Heart Study. *N Engl J Med*. 1998;338:1650–1656
- Mahoney LT, Burns TL, Stanford W, et al. Coronary risk factors measured in childhood and young adult life are associated with coronary artery calcification in young adults: the Muscatine Study. *J Am Coll Cardiol*. 1996;27:277–284
- Williams CL, Hayman LL, Daniels SR, et al. Cardiovascular health in childhood: a statement for health professionals from the Committee on Atherosclerosis, Hypertension, and Obesity in the Young (AHOY) of the Council on Cardiovascular Disease in the Young, American Heart Association. *Circulation*. 2002;106:143–160
- Krauss RM, Eckel RH, Howard B, et al. AHA dietary guidelines: revision 2000—a statement for healthcare professionals from the Nutrition Committee of the American Heart Association. *Circulation*. 2000;102:2284–2299
- Grundy SM, Bilheimer D, Blackburn H, et al. Rationale of the diet-heart statement of the American Heart Association. Report of Nutrition Committee. *Circulation*. 1982;65:839A–854A
- Gidding SS, Leibel RL, Daniels S, Rosenbaum M, Van Horn L, Marx GR. Understanding obesity in youth: a statement for healthcare professionals from the Committee on Atherosclerosis and Hypertension in the Young of the Council on Cardiovascular Disease in the Young and the Nutrition Committee, American Heart Association. Writing Group. *Circulation*. 1996;94:3383–3387
- Institute of Medicine, ed. *Preventing Childhood Obesity: Health in the Balance*. Washington, DC: National Academy Press; 2005
- Olshansky SJ, Passaro DJ, Hershow RC, et al. A potential decline in life expectancy in the United States in the 21st century. *N Engl J Med*. 2005;352:1138–1145
- Fisher EA, Van Horn L, McGill HC Jr. Nutrition and children: a statement for healthcare professionals from the Nutrition Committee, American Heart Association. *Circulation*. 1997;95:2332–2333
- US Department of Health and Human Services, US Department of Agriculture. *Dietary Guidelines for Americans*. 6th ed. Washington, DC: US Government Printing Office; 2005
- US Department of Agriculture. MyPyramid. 2005. Available at: [www.mypyramid.gov](http://www.mypyramid.gov). Accessed June 22, 2005
- Nicklas T, Johnson R. Position of the American Dietetic Association: dietary guidance for healthy children ages 2 to 11 years [published correction appears in *J Am Diet Assoc*. 2004;104:1075]. *J Am Diet Assoc*. 2004;104:660–677
- Kleinman R, ed. *Pediatric Nutrition Handbook*. 5th ed. Elk Grove, IL: American Academy of Pediatrics; 2004
- 2005 Dietary Guideline Advisory Committee. Nutrition and your health: dietary guidelines for Americans. Available at: [www.health.gov/dietaryguidelines/dga2005/report/HTML/E\\_translation.htm](http://www.health.gov/dietaryguidelines/dga2005/report/HTML/E_translation.htm). Accessed January 17, 2005
- Institute of Medicine. *Dietary Reference Intakes for Energy, Carbohydrates, Fiber, Fat, Protein, and Amino Acids (Macronutrients)*. Washington, DC: National Academies Press; 2002. Available at: [www.nap.edu/books/0309085373/html](http://www.nap.edu/books/0309085373/html). Accessed January 31, 2005
- Peterson S, Sigman-Grant M. Impact of adopting lower-fat food choices on nutrient intake of American children. *Pediatrics*. 1997;100(3). Available at: [www.pediatrics.org/cgi/content/full/100/3/e4](http://www.pediatrics.org/cgi/content/full/100/3/e4)
- He K, Song Y, Daviglus ML, et al. Accumulated evidence on fish consumption and coronary heart disease mortality: a meta-analysis of cohort studies. *Circulation*. 2004;109:2705–2711
- US Food and Drug Administration, US Environmental Protection Agency. What you need to know about mercury in fish and shellfish. Available at: [www.fda.gov/oc/opacom/hottopics/mercury/background.html](http://www.fda.gov/oc/opacom/hottopics/mercury/background.html). Accessed December 19, 2005
- Kris-Etherton PM, Harris WS, Appel LJ. American Heart Association Nutrition Committee. Fish consumption, fish oil, omega-3 fatty acids, and cardiovascular disease. *Circulation*. 2002;106:2747–2757
- Mozaffarian D, Lemaitre RN, Kuller LH, Burke GL, Tracy RP, Siscovick DS. Cardiovascular Health Study. Cardiac benefits of fish consumption may depend on the type of fish meal consumed: the Cardiovascular Health Study. *Circulation*. 2003;107:1372–1377
- Van Horn L, Obarzanek E, Friedman LA, Gernhofer N, Barton B. Children's adaptations to a fat-reduced diet: the Dietary Intervention Study in Children (DISC). *Pediatrics*. 2005;115:1723–1733
- 25a. Strong WB, Malina RM, Blimkie CJ, et al. Evidence based physical activity for school-age youth. *J Pediatr*. 2005;146:732–737
- McGill HC Jr, McMahan CA, Zieske AW, Malcom GT, Tracy RE, Strong JP. Effects of nonlipid risk factors on atherosclerosis in youth with a favorable lipoprotein profile. *Circulation*. 2001;103:1546–1550
- Raitakari OT, Juonala M, Kahonen M, et al. Cardiovascular risk factors in childhood and carotid artery intima-media thickness in adulthood: the Cardiovascular Risk in Young Finns Study. *JAMA*. 2003;290:2277–2283
- Li S, Chen W, Srinivasan SR, et al. Childhood cardiovascular risk factors and carotid vascular changes in adulthood: the Bogalusa Heart Study. *JAMA*. 2003;290:2271–2276
- Lauer RM, Lee J, Clarke WR. Factors affecting the relationship between childhood and adult cholesterol levels: the Muscatine Study. *Pediatrics*. 1988;82:309–318
- Lauer RM, Clarke WR. Childhood risk factors for high adult blood pressure: the Muscatine Study. *Pediatrics*. 1989;84:633–641
- Whitaker RC, Wright JA, Pepe MS, Seidel KD, Dietz WH. Predicting obesity in young adulthood from childhood and parental obesity. *N Engl J Med*. 1997;337:869–873
- Janz KF, Dawson JD, Mahoney LT. Tracking physical fitness and physical activity from childhood to adolescence: the Muscatine study. *Med Sci Sports Exerc*. 2000;32:1250–1257
- Feunekes GI, de Graaf C, Meyboom S, van Staveren WA. Food choice and fat intake of adolescents and adults: associations of intakes within social networks. *Prev Med*. 1998;27:645–656

34. Rojas NL, Killen JD, Haydel KF, Robinson TN. Nicotine dependence among adolescent smokers. *Arch Pediatr Adolesc Med.* 1998;152:151–156
35. Obarzanek E, Kimm SY, Barton BA, et al. Long-term safety and efficacy of a cholesterol-lowering diet in children with elevated low-density lipoprotein cholesterol: seven-year results of the Dietary Intervention Study in Children (DISC). *Pediatrics.* 2001;107:256–264
36. Luepker RV, Perry CL, McKinlay SM, et al. Outcomes of a field trial to improve children's dietary patterns and physical activity: the Child and Adolescent Trial for Cardiovascular Health. CATCH collaborative group. *JAMA.* 1996;275:768–776
37. Kaitosaari T, Ronnema T, Raitakari O, et al. Effect of 7-year infancy-onset dietary intervention on serum lipoproteins and lipoprotein subclasses in healthy children in the prospective, randomized Special Turku Coronary Risk Factor Intervention Project for Children (STRIP) study. *Circulation.* 2003;108:672–677
38. Yu-Poth S, Zhao G, Etherton T, Naglak M, Jonnalagadda S, Kris-Etherton PM. Effects of the National Cholesterol Education Program's Step I and Step II dietary intervention programs on cardiovascular disease risk factors: a meta-analysis. *Am J Clin Nutr.* 1999;69:632–646
39. Third Report of the National Cholesterol Education Program (NCEP) Expert Panel on Detection, Evaluation, and Treatment of High Blood Cholesterol in Adults (Adult Treatment Panel III) final report. *Circulation.* 2002;106:3143–3421
40. Lagstrom H, Jokinen E, Seppanen R, et al. Nutrient intakes by young children in a prospective randomized trial of a low-saturated fat, low-cholesterol diet: the STRIP Baby Project. Special Turku Coronary Risk Factor Intervention Project for Babies. *Arch Pediatr Adolesc Med.* 1997;151:181–188
41. Viikari JS, Niinikoski H, Juonala M, et al. Risk factors for coronary heart disease in children and young adults. *Acta Paediatr Suppl.* 2004;93:34–42
42. Talvia S, Lagstrom H, Rasanen M, et al. A randomized intervention since infancy to reduce intake of saturated fat: calorie (energy) and nutrient intakes up to the age of 10 years in the Special Turku Coronary Risk Factor Intervention Project. *Arch Pediatr Adolesc Med.* 2004;158:41–47
43. Obarzanek E, Hunsberger SA, Van Horn L, et al. Safety of a fat-reduced diet: the Dietary Intervention Study in Children (DISC). *Pediatrics.* 1997;100:51–59
44. Lauer RM, Obarzanek E, Hunsberger SA, et al. Efficacy and safety of lowering dietary intake of total fat, saturated fat, and cholesterol in children with elevated LDL cholesterol: the Dietary Intervention Study in Children. *Am J Clin Nutr.* 2000;72:1332S–1342S
45. Rask-Nissila L, Jokinen E, Terho P, et al. Effects of diet on the neurologic development of children at 5 years of age: the STRIP project. *J Pediatr.* 2002;140:328–333
46. Jacobson MS, Tomopoulos S, Williams CL, Arden MR, Deckelbaum RJ, Starc TJ. Normal growth in high-risk hyperlipidemic children and adolescents with dietary intervention. *Prev Med.* 1998;27:775–780
47. Cook S, Weitzman M, Auinger P, Nguyen M, Dietz WH. Prevalence of a metabolic syndrome phenotype in adolescents: findings from the third National Health and Nutrition Examination Survey, 1988–1994. *Arch Pediatr Adolesc Med.* 2003;157:821–827
48. Freedman DS, Dietz WH, Srinivasan SR, Berenson GS. The relation of overweight to cardiovascular risk factors among children and adolescents: the Bogalusa Heart Study. *Pediatrics.* 1999;103:1175–1182
49. Morrison JA, Sprecher DL, Barton BA, Waclawiw MA, Daniels SR. Overweight, fat patterning, and cardiovascular disease risk factors in black and white girls: the National Heart, Lung, and Blood Institute Growth and Health Study. *J Pediatr.* 1999;135:458–464
50. Kimm SY, Glynn NW, Kriska AM, et al. Decline in physical activity in black girls and white girls during adolescence. *N Engl J Med.* 2002;347:709–715
51. Becque MD, Katch VL, Rocchini AP, Marks CR, Moorehead C. Coronary risk incidence of obese adolescents: reduction by exercise plus diet intervention. *Pediatrics.* 1988;81:605–612
52. Rocchini AP, Katch V, Schork A, Kelch RP. Insulin and blood pressure during weight loss in obese adolescents. *Hypertension.* 1987;10:267–273
53. Young EM, Fors SW, Hayes DM. Associations between perceived parent behaviors and middle school student fruit and vegetable consumption. *J Nutr Educ Behav.* 2004;36:2–8
54. Muntner P, He J, Cutler JA, Wildman RP, Whelton PK. Trends in blood pressure among children and adolescents. *JAMA.* 2004;291:2107–2113
55. Ford ES, Mokdad AH, Ajani UA. Trends in risk factors for cardiovascular disease among children and adolescents in the United States. *Pediatrics.* 2004;114:1534–1544
56. Gidding SS, Bao W, Srinivasan SR, Berenson GS. Effects of secular trends in obesity on coronary risk factors in children: the Bogalusa Heart Study. *J Pediatr.* 1995;127:868–874
57. Sinaiko AR, Donahue RP, Jacobs DR Jr, Prineas RJ. Relation of weight and rate of increase in weight during childhood and adolescence to body size, blood pressure, fasting insulin, and lipids in young adults: the Minneapolis Children's Blood Pressure Study. *Circulation.* 1999;99:1471–1476
58. Kumanyika SK, Obarzanek E, Robinson TN, Beech BM. Phase 1 of the Girls Health Enrichment Multi-site Studies (GEMS): conclusion. *Ethn Dis.* 2003;13(suppl 1):S88–S91
59. Caballero B, Clay T, Davis SM, et al. Pathways: a school-based, randomized controlled trial for the prevention of obesity in American Indian schoolchildren. *Am J Clin Nutr.* 2003;78:1030–1038
60. Briefel RR, Reidy K, Karwe V, Devaney B. Feeding infants and toddlers study: improvements needed in meeting infant feeding recommendations. *J Am Diet Assoc.* 2004;104(1 suppl 1):s31–s37
61. Fox MK, Pac S, Devaney B, Jankowski L. Feeding infants and toddlers study: what foods are infants and toddlers eating? *J Am Diet Assoc.* 2004;104(1 suppl 1):s22–s30
62. French SA, Story M, Jeffery RW. Environmental influences on eating and physical activity. *Annu Rev Public Health.* 2001;22:309–335
63. Siega-Riz AM, Popkin BM, Carson T. Trends in breakfast consumption for children in the United States from 1965–1991. *Am J Clin Nutr.* 1998;67:748S–756S
64. Nielsen SJ, Siega-Riz AM, Popkin BM. Trends in energy intake in U.S. between 1977 and 1996: similar shifts seen across age groups. *Obes Res.* 2002;10:370–378
65. Nielsen SJ, Popkin BM. Patterns and trends in food portion sizes, 1977–1998. *JAMA.* 2003;289:450–453
66. Nielsen SJ, Popkin BM. Changes in beverage intake between 1977 and 2001 [published correction appears in *Am J Prev Med.* 2005;28:413]. *Am J Prev Med.* 2004;27:205–210
67. Cavadini C, Siega-Riz AM, Popkin BM. US adolescent food intake trends from 1965 to 1996 [published correction appears in *Arch Dis Child.* 2002;87:85]. *Arch Dis Child.* 2000;83:18–24
68. Kranz S, Smiciklas-Wright H, Siega-Riz AM, Mitchell D. Adverse effect of high added sugar consumption on dietary intake in American preschoolers. *J Pediatr.* 2005;146:105–111
69. Wright JD, Wang CY, Kennedy-Stephenson J, Ervin RB. Dietary intake of ten key nutrients for public health, United States: 1999–2000. *Adv Data.* 2003;(334):1–4
70. Institute of Medicine Panel on Dietary Reference Intakes for

- Electrolytes and Water. *Dietary Reference Intakes for Water, Potassium, Sodium, Chloride, and Sulfate*. Washington, DC: National Academies Press; 2004
71. Ervin RB, Wang CY, Wright JD, Kennedy-Stephenson J. Dietary intake of selected minerals for the United States population: 1999–2000. *Adv Data*. 2004;(341):1–5
  72. Rozin P. Family resemblance in food and other domains: the family paradox and the role of parental congruence. *Appetite*. 1991;16:93–102
  73. Rozin P, Millman L. Family environment, not heredity, accounts for family resemblances in food preferences and attitudes: a twin study. *Appetite*. 1987;8:125–134
  74. Cutler D, Glasser E, Shapiro J. *Why Have Americans Become More Obese? The Economics of Obesity Workshop (April 22, 2003)*. Washington, DC: Economic Research Service, US Department of Agriculture; 2004. E-FAN-04-004
  75. Brown R, Ogdan J. Children's eating attitudes and behaviour: a study of the modelling and control theories of parental influence. *Health Educ Res*. 2004;19:261–271
  76. Videon TM, Manning CK. Influences on adolescent eating patterns: the importance of family meals. *J Adolesc Health*. 2003;32:365–373
  77. Gillman MW, Rifas-Shiman SL, Frazier AL, et al. Family dinner and diet quality among older children and adolescents. *Arch Fam Med*. 2000;9:235–240
  78. Neumark-Sztainer D, Hannan PJ, Story M, Croll J, Perry C. Family meal patterns: associations with sociodemographic characteristics and improved dietary intake among adolescents. *J Am Diet Assoc*. 2003;103:317–322
  79. Kratt P, Reynolds K, Shewchuk R. The role of availability as a moderator of family fruit and vegetable consumption. *Health Educ Behav*. 2000;27:471–482
  80. Fisher JO, Mitchell DC, Smiciklas-Wright H, Birch LL. Parental influences on young girls' fruit and vegetable, micronutrient, and fat intakes. *J Am Diet Assoc*. 2002;102:58–64
  81. Fisher J, Mitchell D, Smiciklas-Wright H, Birch L. Maternal milk consumption predicts the tradeoff between milk and soft drinks in young girls' diets. *J Nutr*. 2001;131:246–250
  82. World Health Organization. *World Health Report 2002: Reducing Risks, Promoting Healthy Life*. Geneva, Switzerland: World Health Organization; 2002. Available at: [www.who.int/whr/2002/en/whr02.en.pdf](http://www.who.int/whr/2002/en/whr02.en.pdf). Accessed April 12, 2005
  83. Rich-Edwards JW, Stampfer MJ, Manson JE, et al. Birth weight and risk of cardiovascular disease in a cohort of women followed up since 1976. *BMJ*. 1997;315:396–400
  84. Curhan GC, Chertow GM, Willett WC, et al. Birth weight and adult hypertension and obesity in women. *Circulation*. 1996;94:1310–1315
  85. Langley-Evans SC. Fetal programming of cardiovascular function through exposure to maternal undernutrition. *Proc Nutr Soc*. 2001;60:505–513
  86. Seckl JR. Prenatal glucocorticoids and long-term programming. *Eur J Endocrinol*. 2004;151(suppl 3):U49–U62
  87. Edwards LJ, Coulter CL, Symonds ME, McMillen IC. Prenatal undernutrition, glucocorticoids and the programming of adult hypertension. *Clin Exp Pharmacol Physiol*. 2001;28:938–941
  88. Eriksson J, Forsen T, Tuomilehto J, Osmond C, Barker D. Size at birth, childhood growth and obesity in adult life. *Int J Obes Relat Metab Disord*. 2001;25:735–740
  89. Hediger ML, Overpeck MD, McGlynn A, Kuczmarski RJ, Maurer KR, Davis WW. Growth and fatness at three to six years of age of children born small- or large-for-gestational age. *Pediatrics*. 1999;104(3). Available at: [www.pediatrics.org/cgi/content/full/104/3/e33](http://www.pediatrics.org/cgi/content/full/104/3/e33)
  90. Frisancho AR. Prenatal compared with parental origins of adolescent fatness. *Am J Clin Nutr*. 2000;72:1186–1190
  91. Whitaker RC. Predicting preschooler obesity at birth: the role of maternal obesity in early pregnancy. *Pediatrics*. 2004;114(1). Available at: [www.pediatrics.org/cgi/content/full/114/1/e29](http://www.pediatrics.org/cgi/content/full/114/1/e29)
  92. Institute of Medicine. *Nutrition During Pregnancy, Weight Gain and Nutrient Supplements: Report of the Subcommittee on Nutritional Status and Weight Gain During Pregnancy*. Washington, DC: National Academy Press; 1990
  93. Gunderson EP, Abrams B, Selvin S. The relative importance of gestational gain and maternal characteristics associated with the risk of becoming overweight after pregnancy. *Int J Obes Relat Metab Disord*. 2000;24:1660–1668
  94. Huh SY, Rifas-Shiman SL, Kleinman KP, Rich-Edwards JW, Lipshultz SE, Gillman MW. Maternal protein intake is not associated with infant blood pressure. *Int J Epidemiol*. 2005;34:378–384
  95. Gillman MW, Rifas-Shiman SL, Kleinman KP, Rich-Edwards JW, Lipshultz SE. Maternal calcium intake and offspring blood pressure. *Circulation*. 2004;110:1990–1995
  96. Gartner LM, Morton J, Lawrence RA, et al. Breastfeeding and the use of human milk. *Pediatrics*. 2005;115:496–506
  97. Martin RM, Davey Smith G, Mangtani P, Tilling K, Frankel S, Gunnell D. Breastfeeding and cardiovascular mortality: the Boyd Orr cohort and a systematic review with meta-analysis. *Eur Heart J*. 2004;25:778–786
  98. Arenz S, Ruckerl R, Koletzko B, von Kries R. Breast-feeding and childhood obesity: a systematic review. *Int J Obes Relat Metab Disord*. 2004;28:1247–1256
  99. Owen CG, Martin RM, Whincup PH, Smith GD, Cook DG. Effect of infant feeding on the risk of obesity across the life course: a quantitative review of published evidence. *Pediatrics*. 2005;115:1367–1377
  100. Martin RM, Gunnell D, Smith GD. Breastfeeding in infancy and blood pressure in later life: systematic review and meta-analysis. *Am J Epidemiol*. 2005;161:15–26
  101. Owen CG, Whincup PH, Gilg JA, Cook DG. Effect of breast feeding in infancy on blood pressure in later life: systematic review and meta-analysis. *BMJ*. 2003;327:1189–1195
  102. Owen CG, Whincup PH, Odoki K, Gilg JA, Cook DG. Infant feeding and blood cholesterol: a study in adolescents and a systematic review. *Pediatrics*. 2002;110:597–608
  103. Stettler N, Zemel BS, Kumanyika S, Stallings VA. Infant weight gain and childhood overweight status in a multi-center, cohort study. *Pediatrics*. 2002;109:194–199
  104. Dewey KG. Nutrition, growth, and complementary feeding of the breastfed infant. *Pediatr Clin North Am*. 2001;48:87–104
  105. Haisma H, Coward WA, Albernaz E, et al. Breast milk and energy intake in exclusively, predominantly, and partially breast-fed infants. *Eur J Clin Nutr*. 2003;57:1633–1642
  106. Dennison BA, Edmunds LE, Stratton H, Pruzek R. Rapid infant weight gain associated with childhood overweight at age 4 years. *Obes Res*. In press
  107. Wright P. Hunger, satiety and feeding behavior in early infancy. In: Boakes RA, Popplewell DA, Burton MJ, eds. *Eating Habits: Food, Physiology and Learned Behavior*. Chichester, United Kingdom: Wiley; 1987
  108. Fisher JO, Birch LL, Smiciklas-Wright H, Picciano MF. Breast-feeding through the first year predicts maternal control in feeding and subsequent toddler energy intakes. *J Am Diet Assoc*. 2000;100:641–646
  109. Taveras EM, Scanlon KS, Birch L, Rifas-Shiman SL, Rich-Edwards JW, Gillman MW. Association of breastfeeding with maternal control of infant feeding at age 1 year. *Pediatrics*. 2004;114(5). Available at: [www.pediatrics.org/cgi/content/full/114/5/e577](http://www.pediatrics.org/cgi/content/full/114/5/e577)
  110. Wright P. Patterns of milk intake in breast- and bottle-fed infants. In: Stratton P, ed. *The Psychology of the Human Newborn*. New York, NY: Wiley; 1982

111. Klesges RC, Coates TJ, Brown G, et al. Parental influences on children's eating behavior and relative weight. *J Appl Behav Anal.* 1983;16:371-378
112. Mennella JA, Griffin CE, Beauchamp GK. Flavor programming during infancy. *Pediatrics.* 2004;113:840-845
113. Mennella JA, Jagnow CP, Beauchamp GK. Prenatal and postnatal flavor learning by human infants. *Pediatrics.* 2001;107(6). Available at: [www.pediatrics.org/cgi/content/full/107/6/e88](http://www.pediatrics.org/cgi/content/full/107/6/e88)
114. Mennella JA, Pepino MY, Reed DR. Genetic and environmental determinants of bitter perception and sweet preferences. *Pediatrics.* 2005;115(2). Available at: [www.pediatrics.org/cgi/content/full/115/2/e216](http://www.pediatrics.org/cgi/content/full/115/2/e216)
115. Gerrish CJ, Mennella JA. Flavor variety enhances food acceptance in formula-fed infants. *Am J Clin Nutr.* 2001;73:1080-1085
116. Liem DG, Mennella JA. Heightened sour preferences during childhood. *Chem Senses.* 2003;28:173-180
117. Li R, Hsia J, Fridinger F, Hussain A, Benton-Davis S, Grummer-Strawn L. Public beliefs about breastfeeding policies in various settings. *J Am Diet Assoc.* 2004;104:1162-1168
118. Visness CM, Kennedy KI. Maternal employment and breastfeeding: findings from the 1988 National Maternal and Infant Health Survey. *Am J Public Health.* 1997;87:945-950
119. Li R, Zhao Z, Mokdad A, Barker L, Grummer-Strawn L. Prevalence of breastfeeding in the United States: the 2001 National Immunization Survey. *Pediatrics.* 2003;111:1198-1201
120. Kurinij N, Shiono PH, Ezrine SF, Rhoads GG. Does maternal employment affect breast-feeding? *Am J Public Health.* 1989;79:1247-1250
121. Gielen AC, Faden RR, O'Campo P, Brown CH, Paige DM. Maternal employment during the early postpartum period: effects on initiation and continuation of breast-feeding. *Pediatrics.* 1991;87:298-305
122. Auerbach KG, Guss E. Maternal employment and breast-feeding: a study of 567 women's experiences. *Am J Dis Child.* 1984;138:958-960
123. Hills-Bonczyk SG, Avery MD, Savik K, Potter S, Duckett LJ. Women's experiences with combining breast-feeding and employment. *J Nurse-Midwifery.* 1993;38:257-266
124. Lande B, Andersen LF, Veierod MB, et al. Breast-feeding at 12 months of age and dietary habits among breast-fed and non-breast-fed infants. *Public Health Nutr.* 2004;7:495-503
125. Ortiz J, McGilligan K, Kelly P. Duration of breast milk expression among working mothers enrolled in an employer-sponsored lactation program. *Pediatr Nurs.* 2004;30:111-119
126. Kramer MS, Kakuma R. The optimal duration of exclusive breastfeeding: a systematic review. *Adv Exp Med Biol.* 2004;554:63-77
- 126a. Lederman SA, Akabas SR, Moore BJ, et al. Summary of the Presentations at the Conference on Preventing Childhood Obesity, December 8, 2003. *Pediatrics.* 2004;114:1146-1173
127. Carruth BR, Skinner JD, Houck KS, Moran JD 3rd. Addition of supplementary foods and infant growth (2 to 24 months). *J Am Coll Nutr.* 2000;19:405-412
128. Mehta KC, Specker BL, Bartholmey S, Giddens J, Ho ML. Trial on timing of introduction to solids and food type on infant growth. *Pediatrics.* 1998;102:569-573
129. Devaney B, Ziegler P, Pac S, Karwe V, Barr SI. Nutrient intakes of infants and toddlers. *J Am Diet Assoc.* 2004;104(suppl 1):s14-s21
130. Skinner JD, Ziegler P, Ponza M. Transitions in infants' and toddlers' beverage patterns. *J Am Diet Assoc.* 2004;104(suppl 1):s45-s50
131. US Department of Agriculture, Food and Nutrition Services. 2002 CFR title 7 volume 4 chapter II (1-1-02 edition). 246-Special Supplemental Nutrition Program for Women, Infants and Children (WIC). Subpart D-Participant benefits. 246.10 Supplemental foods. Available at: [www.fns.usda.gov/wic/lawsandregulations/WICRegulations-7CFR246.pdf](http://www.fns.usda.gov/wic/lawsandregulations/WICRegulations-7CFR246.pdf). Accessed January 31, 2005
132. Dennison BA, Erb TA, Jenkins PL. Predictors of dietary milk fat intake by preschool children. *Prev Med.* 2001;33:536-542
133. Sullivan SA, Birch LL. Infant dietary experience and acceptance of solid foods. *Pediatrics.* 1994;93:271-277
134. Thompson FE, Dennison BA. Dietary sources of fats and cholesterol in US children aged 2 through 5 years. *Am J Public Health.* 1994;84:799-806
135. Dennison BA, Rockwell HL, Baker SL. Excess fruit juice consumption by preschool-aged children is associated with short stature and obesity. *Pediatrics.* 1997;99:15-22
136. Welsh JA, Cogswell ME, Rogers S, Rockett H, Mei Z, Grummer-Strawn LM. Overweight among low-income preschool children associated with the consumption of sweet drinks: Missouri, 1999-2002. *Pediatrics.* 2005;115(2). Available at: [www.pediatrics.org/cgi/content/full/115/2/e223](http://www.pediatrics.org/cgi/content/full/115/2/e223)
137. Drewnowski A. The role of energy density. *Lipids.* 2003;38:109-115
138. Birch LL, Deysher M. Caloric compensation and sensory specific satiety: evidence for self regulation of food intake by young children. *Appetite.* 1986;7:323-331
139. Araya H, Vera G, Alvina M. Effect of the energy density and volume of high carbohydrate meals on short term satiety in preschool children. *Eur J Clin Nutr.* 1999;53:273-276
140. Bandini LG, Vu D, Must A, Cyr H, Goldberg A, Dietz WH. Comparison of high-calorie, low-nutrient-dense food consumption among obese and non-obese adolescents. *Obes Res.* 1999;7:438-443
141. Phillips SM, Bandini LG, Naumova EN, et al. Energy-dense snack food intake in adolescence: longitudinal relationship to weight and fatness. *Obes Res.* 2004;12:461-472
142. Birch LL, Fisher JO. Development of eating behaviors among children and adolescents. *Pediatrics.* 1998;101(3 pt 2):539-549
143. Fisher JO, Birch LL. Restricting access to palatable foods affects children's behavioral response, food selection, and intake. *Am J Clin Nutr.* 1999;69:1264-1272
144. Ludwig DS, Peterson KE, Gortmaker SL. Relation between consumption of sugar-sweetened drinks and childhood obesity: a prospective, observational analysis. *Lancet.* 2001;357:505-508
145. American Academy of Pediatrics, Committee on Nutrition. The use and misuse of fruit juice in pediatrics. *Pediatrics.* 2001;107:1210-1213
146. Bowman SA, Gortmaker SL, Ebeling CB, Pereira MA, Ludwig DS. Effects of fast-food consumption on energy intake and diet quality among children in a national household survey. *Pediatrics.* 2004;113:112-118
147. Putnam J, Allshouse J, Kantor L. *U.S. Per Capita Food Supply Trends: More Calories, Refined Carbohydrates and Fats.* Washington, DC: Economic Research Service, US Department of Agriculture; 2002
148. Epstein LH, Valoski A, Wing RR, McCurley J. Ten-year follow-up of behavioral, family-based treatment for obese children. *JAMA.* 1990;264:2519-2523
149. Hayman LL, Williams CL, Daniels SR, et al. Cardiovascular health promotion in the schools: a statement for health and education professionals and child health advocates from the Committee on Atherosclerosis, Hypertension, and Obesity in Youth (AHOY) of the Council on Cardiovascular Disease in the Young, American Heart Association. *Circulation.* 2004;110:2266-2275

150. Position of ADA, SNE, and ASFSA: school-based nutrition programs and services. *J Am Diet Assoc.* 1995;95:367–369
151. Osganian SK, Ebzery MK, Montgomery DH, et al. Changes in the nutrient content of school lunches: results from the CATCH Eat Smart Food Service Intervention. *Prev Med.* 1996;25:400–412
152. Dwyer JT, Hewes LV, Mitchell PD, et al. Improving school breakfasts: effects of the CATCH Eat Smart Program on the nutrient content of school breakfasts. *Prev Med.* 1996;25:413–422
153. Perry CL, Bishop DB, Taylor GL, et al. A randomized school trial of environmental strategies to encourage fruit and vegetable consumption among children. *Health Educ Behav.* 2004;31:65–76
154. Cullen KW, Ash DM, Warneke C, de Moor C. Intake of soft drinks, fruit-flavored beverages, and fruits and vegetables by children in grades 4 through 6. *Am J Public Health.* 2002;92:1475–1478
155. Birnbaum AS, Lytle LA, Story M, Perry CL, Murray DM. Are differences in exposure to a multicomponent school-based intervention associated with varying dietary outcomes in adolescents? *Health Educ Behav.* 2002;29:427–443
156. Williams CL, Strobino BA, Bollella M, Brotanek J. Cardiovascular risk reduction in preschool children: the “Healthy Start” project. *J Am Coll Nutr.* 2004;23:117–123
157. Levin S, Lowry R, Brown DR, Dietz WH. Physical activity and body mass index among US adolescents: youth risk behavior survey, 1999. *Arch Pediatr Adolesc Med.* 2003;157:816–820
158. Boynton-Jarrett R, Thomas TN, Peterson KE, Wiecha J, Sobol AM, Gortmaker SL. Impact of television viewing patterns on fruit and vegetable consumption among adolescents. *Pediatrics.* 2003;112:1321–1326
159. Dennison BA, Erb TA, Jenkins PL. Television viewing and television in bedroom associated with overweight risk among low-income preschool children. *Pediatrics.* 2002;109:1028–1035
160. Coon KA, Goldberg J, Rogers BL, Tucker KL. Relationships between use of television during meals and children’s food consumption patterns. *Pediatrics.* 2001;107(1). Available at: [www.pediatrics.org/cgi/content/full/107/1/e7](http://www.pediatrics.org/cgi/content/full/107/1/e7)
161. Institute of Medicine. *WIC Food Packages: Time for Change—Executive Summary.* Washington, DC: National Academy of Sciences; 2004. Available at: [www.nap.edu/catalog/11280.html](http://www.nap.edu/catalog/11280.html). Accessed April 30, 2005
162. National High Blood Pressure Education Program Working Group on High Blood Pressure in Children and Adolescents. The fourth report on the diagnosis, evaluation, and treatment of high blood pressure in children and adolescents. *Pediatrics.* 2004;114(2 suppl 4th report):555–576
163. National Cholesterol Education Program (NCEP): highlights of the report of the Expert Panel on Blood Cholesterol Levels in Children and Adolescents. *Pediatrics.* 1992;89:495–501
164. Appel LJ, Moore TJ, Obarzanek E, et al. A clinical trial of the effects of dietary patterns on blood pressure. DASH Collaborative Research Group. *N Engl J Med.* 1997;336:1117–1124
165. Simons-Morton DG, Obarzanek E. Diet and blood pressure in children and adolescents. *Pediatr Nephrol.* 1997;11:244–249
166. Simons-Morton DG, Hunsberger SA, Van Horn L, et al. Nutrient intake and blood pressure in the Dietary Intervention Study in Children. *Hypertension.* 1997;29:930–936
167. Van Horn L. Fiber, lipids, and coronary heart disease: a statement for healthcare professionals from the Nutrition Committee, American Heart Association. *Circulation.* 1997;95:2701–2704
168. Lichtenstein AH, Deckelbaum RJ. AHA Science Advisory. Stanol/sterol ester-containing foods and blood cholesterol levels. A statement for healthcare professionals from the Nutrition Committee of the Council on Nutrition, Physical Activity, and Metabolism of the American Heart Association. *Circulation.* 2001;103:1177–1179
169. Tammi A, Ronnema T, Miettinen TA, et al. Effects of gender, apolipoprotein E phenotype and cholesterol-lowering by plant stanol esters in children: the STRIP study. Special Turku Coronary Risk Factor Intervention Project. *Acta Paediatr.* 2002;91:1155–1162
170. Davidson MH, Maki KC, Umporowicz DM, et al. Safety and tolerability of esterified phytosterols administered in reduced-fat spread and salad dressing to healthy adult men and women. *J Am Coll Nutr.* 2001;20:307–319
171. Homma Y, Ikeda I, Ishikawa T, Tateno M, Sugano M, Nakamura H. Decrease in plasma low-density lipoprotein cholesterol, apolipoprotein B, cholesteryl ester transfer protein, and oxidized low-density lipoprotein by plant stanol ester-containing spread: a randomized, placebo-controlled trial. *Nutrition.* 2003;19:369–374
172. Ntanos FY, Homma Y, Ushiro S. A spread enriched with plant sterol-esters lowers blood cholesterol and lipoproteins without affecting vitamins A and E in normal and hypercholesterolemic Japanese men and women. *J Nutr.* 2002;132:3650–3655
173. Volpe R, Niittynen L, Korpela R, et al. Effects of yoghurt enriched with plant sterols on serum lipids in patients with moderate hypercholesterolaemia. *Br J Nutr.* 2001;86:233–239
174. Maki KC, Davidson MH, Umporowicz DM, et al. Lipid responses to plant-sterol-enriched reduced-fat spreads incorporated into a National Cholesterol Education Program Step I diet. *Am J Clin Nutr.* 2001;74:33–43
175. Miettinen TA, Puska P, Gylling H, Vanhanen H, Vartiainen E. Reduction of serum cholesterol with sitostanol-ester margarine in a mildly hypercholesterolemic population. *N Engl J Med.* 1995;333:1308–1312
176. Hendriks HF, Brink EJ, Meijer GW, Princen HM, Ntanos FY. Safety of long-term consumption of plant sterol esters-enriched spread. *Eur J Clin Nutr.* 2003;57:681–692
177. Nestel P, Cehun M, Pomeroy S, Abbey M, Weldon G. Cholesterol-lowering effects of plant sterol esters and non-esterified stanols in margarine, butter and low-fat foods. *Eur J Clin Nutr.* 2001;55:1084–1090