

Dietary restriction suppresses proteotoxicity and enhances longevity by an *hsf-1*-dependent mechanism in *Caenorhabditis elegans*

Katherine A. Steinkraus,¹ Erica D. Smith,^{1,2} Christina Davis,¹ Daniel Carr,¹ William R. Pendergrass,¹ George L. Sutphin,¹ Brian K. Kennedy² and Matt Kaeberlein¹

Departments of ¹Pathology and ²Biochemistry, University of Washington, Seattle, WA 98195, USA

Summary

Dietary restriction increases lifespan and slows the onset of age-associated disease in organisms from yeast to mammals. In humans, several age-related diseases are associated with aberrant protein folding or aggregation, including neurodegenerative disorders such as Alzheimer's, Parkinson's, and Huntington's diseases. We report here that dietary restriction dramatically suppresses age-associated paralysis in three nematode models of proteotoxicity. Similar to its longevity-enhancing properties, dietary restriction protects against proteotoxicity by a mechanism distinct from reduced insulin/IGF-1-like signaling. Instead, the heat shock transcription factor, *hsf-1*, is required for enhanced thermotolerance, suppression of proteotoxicity, and lifespan extension by dietary restriction. These findings demonstrate that dietary restriction confers a general protective effect against proteotoxicity and promotes longevity by a mechanism involving *hsf-1*.

Key words: *Caenorhabditis elegans*; dietary restriction; *hsf-1*; longevity; proteotoxicity.

Introduction

Dietary restriction (DR) is known to increase lifespan and retard a variety of age-associated pathologies in rodents and invertebrate organisms (Weindruch & Walford, 1988; Masoro, 2005). Among the diseases beneficially impacted by DR in mice are cancer, diabetes, and cardiovascular disease. In addition, DR has been shown to improve outcome in mouse models of neurodegeneration (Martin *et al.*, 2006). The mechanism by which DR confers these benefits remains unknown, but is of high interest.

Dietary restriction can be studied in the nematode *Caenorhabditis elegans* using either environmental or genetic approaches.

In the laboratory, *C. elegans* are typically maintained on a nutrient agar nematode growth medium (Brenner, 1974) and provided *Escherichia coli* OP50 as their dietary food source. Under these standard growth conditions, lifespan extension from bacterial food restriction is maximized when animals are maintained on nutrient agar plates without bacterial food after early adulthood [bacterial food deprivation (BD); also referred to previously as DR through food deprivation or dietary deprivation] (Kaeberlein *et al.*, 2006; Lee *et al.*, 2006). While BD is similar in concept to an alternative method for DR in *C. elegans*, axenic growth (Vanfleteren & Braeckman, 1999), BD is instituted postdevelopmentally and utilizes standard *C. elegans* growth conditions.

An alternative method of DR in *C. elegans* is the use of mutations that reduce food intake, such as loss of function alleles of *eat-2*. *eat-2* mutants have defects in pharyngeal pumping (Avery, 1993) and an adult lifespan 20–40% longer than wild-type N2 animals (Lakowski & Hekimi, 1998). Lifespan extension from BD is nonadditive with the long-lived *eat-2(ad465)* allele (Kaeberlein *et al.*, 2006; Lee *et al.*, 2006), indicating that BD and mutation of *eat-2* are likely to promote longevity via similar or overlapping mechanisms.

Although the mechanism by which DR increases longevity in *C. elegans* is unknown, DR is thought to act through a pathway that is genetically distinct from insulin/IGF-1-like signaling (IIS). Mutations that decrease IIS, such as loss of function alleles of insulin-like receptor *daf-2* (Kenyon *et al.*, 1993; Kimura *et al.*, 1997) or the PI3-kinase *age-1* (Friedman & Johnson, 1988a,b; Morris *et al.*, 1996), increase longevity via a mechanism that is dependent on the FOXO-family transcription factor *daf-16* (Lin *et al.*, 1997; Ogg *et al.*, 1997). In contrast, lifespan extension from DR (whether accomplished by BD, axenic growth, or mutation of *eat-2*) does not require *daf-16* and is additive with mutation of *daf-2* (Lakowski & Hekimi, 1998; Houthoofd *et al.*, 2002; Kaeberlein *et al.*, 2006; Lee *et al.*, 2006). More recently, two transcription factors, *skn-1* and *pha-4*, have been shown to be necessary for lifespan extension from DR (Bishop & Guarente, 2007; Panowski *et al.*, 2007); however, it remains unclear how these factors respond to DR and what the relevant downstream targets for lifespan extension might be.


We have used *C. elegans* to explore the effect of DR on disease processes associated with protein misfolding or aggregation (proteotoxicity). Several transgenic models of proteotoxicity have been developed and characterized in worms. These models show pleiotropic phenotypes (Link, 2001; Brignull *et al.*, 2007), including defective posterior mechanosensation (Parker *et al.*, 2001, 2005), impaired ubiquitin-proteasomal function (Khan

Correspondence

Matt Kaeberlein, Department of Pathology, University of Washington, Seattle, WA 98195, USA. Tel.: (206) 543-4849; fax: (206) 543-3644; e-mail: kaeber@u.washington.edu

Accepted for publication 20 February 2008

Fig. 1 Bacterial food deprivation suppresses proteotoxicity in a nematode model of polyglutamine disease. (A) Age-associated paralysis caused by expression Q_{35} YFP (Q_{35}) is significantly reduced in bacterial food deprived (BD) animals relative to control-fed (CF) animals ($p = 6.2 \times 10^{-12}$). (B) The age-associated paralysis of *unc-22(e66)* mutants is unaltered in BD animals relative to CF animals. (C) BD significantly reduces polyglutamine-associated paralysis when initiated at day 3 ($p = 8.3 \times 10^{-13}$), day 5 ($p = 7.6 \times 10^{-14}$), or day 7 ($p = 5.1 \times 10^{-5}$) of adulthood. (D) BD does not significantly reduce the formation of polyglutamine aggregates. The total amount of aggregates, as measured by the integral of the fluorescent area per animal or the number of discrete aggregates visible as fluorescent spots, of BD animals relative to CF animals were counted. Error bars are standard error of the mean.



et al., 2006), decreased nose touch response (Faber *et al.*, 1999), or age-associated paralysis (Link, 1995; Satyal *et al.*, 2000; Morley *et al.*, 2002; Link *et al.*, 2006). Here we report that both genetic and environmental models of DR are potent suppressors of proteotoxicity and we identify a novel role for the heat shock transcription factor, *hsf-1*, as a key mediator of DR in *C. elegans*.

Results

DR confers protection against polyglutamine proteotoxicity

We first examined the effects of DR on proteotoxicity using a nematode model of polyglutamine disease. Huntington's disease and several other neurodegenerative disorders are caused by polyglutamine tract expansions, although the mechanism of disease progression remains unknown (Bonini & La Spada, 2005). For our studies, we utilized a nematode model of Huntington's disease in which a tract of 35 consecutive glutamine residues is fused to YFP (Q_{35} YFP) and expressed in the body wall muscles (Morley *et al.*, 2002). This tract length has been shown to be at the threshold for age-related toxicity in *C. elegans* (Morley *et al.*, 2002). As previously reported (Morley *et al.*, 2002), we observed an age-dependent paralysis phenotype in Q_{35} YFP animals fed a control diet of abundant UV-killed *E. coli* OP50 (Fig. 1A). The lifespan of these animals was relatively normal (Supplementary Fig. S1A), suggesting that the proteotoxicity caused by the Q_{35} YFP protein, while sufficient to cause paralysis, does not necessarily limit longevity. When placed on a BD regimen at the second day of adulthood, the survival of Q_{35} YFP animals was enhanced in a manner similar

to wild-type controls (Supplementary Fig. S1A). Strikingly, Q_{35} YFP animals on BD were resistant to proteotoxicity ($p = 6.2 \times 10^{-12}$) and remained largely paralysis free (Fig. 2A; Supplementary video files).

To determine whether BD confers general protection against muscular defects independently of the Q_{35} YFP transgene, we examined the effects of BD in two mutant backgrounds that display an age-associated defect in movement similar to that of Q_{35} YFP: *unc-52(e444)* and *unc-22(e66)* (Moerman & Baillie, 1979; Waterston *et al.*, 1980; Rogalski *et al.*, 1995). In both cases, BD failed to improve movement or suppress paralysis, relative to control-fed isogenic animals (Fig. 1B; Supplementary video files). Thus, we conclude that the suppression of paralysis by BD is unlikely to be the result of a general improvement in muscle function or increase in movement.

We have previously reported that BD increases lifespan to a similar extent whether initiated early in adulthood (day 2) or postreproductively (day 10) (Kaerberlein *et al.*, 2006). Accordingly, we wished to determine whether suppression of polyglutamine toxicity is observed by initiating BD at ages later than 2 days of adulthood. Consistent with the effect of BD on longevity, BD initiated at days 3, 5, or 7 of adulthood significantly reduced age-associated paralysis in Q_{35} YFP animals (Fig. 1C; $p = 8.3 \times 10^{-13}$ day 3, 7.6×10^{-14} day 5, 5.1×10^{-5} day 7). However, BD was not sufficient to reverse paralysis. Out of more than 50 individual paralyzed animals examined, none regained movement when transferred from a control-fed diet to BD. These observations can be explained by a model in which BD confers protection against polyglutamine toxicity until a threshold level of toxicity results in paralysis, after which BD is unable to reverse the accumulated cellular damage.




Fig. 2 Bacterial food deprivation (BD) suppresses proteotoxicity in a nematode model of Alzheimer's disease. (A) Relative to control-fed (CF) animals, BD significantly reduces age-associated paralysis in animals expressing A β_{42} ($p = 1.4 \times 10^{-7}$). (B) BD significantly reduces age-associated paralysis in animals expressing the aggregation prone GFP-degron peptide (GFP-deg), relative to control-fed (CF) animals ($p = 1.0 \times 10^{-14}$). (C) By fluorescence microscopy, the steady-state levels of the GFP-degron peptide are not reduced in BD animals as compared to CF animals.

The formation of insoluble aggregates is a hallmark of human polyglutamine diseases such as Huntington's disease (Borrell-Pages *et al.*, 2006). However, whether aggregates are causal in disease progression remains controversial. To address this question, we quantified the abundance of aggregates in aged animals fed either a control diet or maintained on BD from the second day of adulthood. Although paralysis was dramatically reduced in 14-day-old BD animals relative to control-fed animals, there was no significant difference in either the number or size of Q₃₅YFP-aggregates between the two groups (Fig. 1D; see also Supplementary video files and Fig. S2). Consistent with this, the relative amount of Q₃₅YFP protein, when normalized to total protein or to actin (0.96 ± 0.1), was not significantly different in BD animals relative to control-fed animals at day 9 of adulthood (Supplementary Fig. S2C). Thus, BD appears to protect against proteotoxicity without significantly altering the age-associated accumulation of polyglutamine aggregates.

DR is a general suppressor of proteotoxicity

In order to determine whether the suppression of proteotoxicity conferred by BD would be specific to polyglutamine toxicity, we next examined a transgenic model of Alzheimer's disease. A 42-amino acid amyloid-beta peptide (A β_{42}) is expressed in body wall muscle cells of these animals under control of the *unc-54* promoter (Link, 1995). The A β_{42} peptide is a cleavage product of amyloid precursor protein, implicated in Alzheimer's disease (Koo *et al.*, 1999). Similar to the Q₃₅YFP animals, BD was sufficient to largely suppress age-associated paralysis caused by A β_{42} -induced

proteotoxicity (Fig. 2A; $p = 1.4 \times 10^{-7}$). BD also increased both median and maximum survival of A β_{42} animals (Supplementary Fig. S1B; $p = 8.7 \times 10^{-4}$). In multiple independent experiments, we observed that BD caused a notable increase in age-independent early mortality. This may indicate that amyloid-beta toxicity (but not polyglutamine toxicity) causes frailty in response to food restriction among a percentage of the population. Interestingly, the subset of animals that died early also tended to show early paralysis, while the survivors remained largely paralysis free. Further studies will be necessary to address the mechanism behind this observation.

Given that BD was effective at suppressing proteotoxicity in two different nematode models of human disease, we wished to further characterize the generality of this phenomenon. Recently, an aggregation prone form of GFP (GFP-degron) was described that causes progressive paralysis when expressed transgenically in *C. elegans* from a *myo-3* promoter (Link *et al.*, 2006). As observed in the Q₃₅YFP and A β_{42} models, BD significantly suppressed paralysis in animals expressing the GFP-degron peptide (Fig. 2B; $p = 1.0 \times 10^{-14}$). We also assayed whether the levels of aggregated GFP-degron was decreased in worms subjected to BD. As with the Q₃₅YFP animals, age-matched GFP-degron animals did not display a notable loss of the aggregated protein by microscopy (Fig. 2C).

Genetic models of DR suppress proteotoxicity

Although BD is unquestionably a form of food (dietary) restriction, it remains possible that BD could have different or additional




Fig. 3 Genetic models of dietary restriction suppress proteotoxicity. (A) RNAi knockdown of *eat-2* significantly reduces paralysis in Q_{35} YFP animals, relative to animals maintained on empty vector bacteria ($p = 1.0 \times 10^{-3}$). (B) RNAi knockdown of *eat-2* significantly reduces paralysis in $A\beta_{42}$ animals, relative to animals maintained on empty vector bacteria ($p = 1.0 \times 10^{-5}$). (C) RNAi knockdown of *rab-10* significantly reduces paralysis in Q_{35} YFP animals, relative to animals maintained on empty vector bacteria ($p = 1.3 \times 10^{-11}$). (D) RNAi knockdown of *sams-1* or *drr-2* significantly reduces paralysis in Q_{35} YFP animals, relative to animals maintained on empty vector bacteria ($p = 1.3 \times 10^{-14}$ for *sams-1* and $p < 10^{-15}$ for *drr-2*).

properties that are not shared by other methods of DR described in the nematode. To address this possibility, we examined whether decreased *eat-2* function could suppress proteotoxicity. RNAi inhibition of *eat-2* resulted in a significant decrease in paralysis in both the Q_{35} YFP (Fig. 3A; $p = 1.0 \times 10^{-3}$) and $A\beta_{42}$ (Fig. 3B; $p = 1.0 \times 10^{-5}$) models. However, suppression of proteotoxicity by inhibition of *eat-2* did not appear to be as effective as BD. This is similar to the relative effects of BD and mutation of *eat-2* on lifespan (Kaeberlein et al., 2006; Lee et al., 2006), and may suggest that BD represents a level of DR more optimal for longevity and suppression of proteotoxicity. Alternatively, BD may have additional properties that influence aging and proteotoxicity other than those shared by both BD and mutation of *eat-2*.

We next examined genetic models of DR identified from a genome-wide RNAi screen for increased lifespan (Hansen et al., 2005). RNAi knockdown of *rab-10*, *sams-1*, or *drr-2* increases lifespan through a *daf-16*-independent mechanism, but fails to further increase the lifespan of *eat-2(ad1116)* animals (Hansen et al., 2005). Each of these genes are transcriptionally down-regulated in *eat-2* mutants, suggesting that they function downstream of food consumption to modulate lifespan (Hansen et al., 2005). Paralysis was dramatically reduced in Q_{35} YFP animals in response to RNAi inhibition of *rab-10* (Fig. 3C; $p = 1.3 \times 10^{-11}$), *sams-1* (Fig. 3D; $p = 1.3 \times 10^{-14}$) or *drr-2* (Fig. 3D; $p < 1 \times 10^{-15}$). Suppression of proteotoxicity in these three genetic models of DR was comparable in magnitude to BD. Therefore, we conclude that both genetic and environmental models of DR confer a substantial protective effect against proteotoxicity in *C. elegans*.

Bacterial food deprivation suppresses proteotoxicity through a mechanism distinct from IIS

Insulin/IGF-1-like signaling has been previously found to modulate resistance to proteotoxicity in both $A\beta_{42}$ and Q_{35} YFP animals (Hsu et al., 2003; Cohen et al., 2006). We therefore wished to determine the relationship, if any, between DR and IIS in these models. As previously reported for $A\beta_{42}$ animals (Cohen et al., 2006), we observed that RNAi knockdown of the FOXO-family transcription factor, *daf-16*, accelerated paralysis in Q_{35} YFP animals (Fig. 4A; $p = 8.0 \times 10^{-3}$), while RNAi knockdown of the insulin-like receptor, *daf-2*, delayed paralysis (Fig. 4B; $p = 7.6 \times 10^{-7}$). For these experiments, animals were maintained on RNAi bacteria from egg until the second day of adulthood, then transferred to either empty vector RNAi bacteria or BD. Interestingly, RNAi knockdown of *daf-16* did not block the effectiveness of BD (Fig. 4A; $p = 4.5 \times 10^{-9}$). When combined with RNAi knockdown of *daf-2*, BD conferred an even greater suppression of paralysis than was observed in BD animals fed empty vector RNAi (Fig. 4B; $p = 1.0 \times 10^{-14}$). Identical epistasis interactions were observed between IIS and BD in $A\beta_{42}$ animals (Supplementary Fig. S3). These data recapitulate the genetic relationship between DR and IIS with respect to longevity: DR can increase lifespan by a mechanism that is independent of *daf-16* and additive with mutation of *daf-2* (Lakowski & Hekimi, 1998; Houthoofd et al., 2003; Kaeberlein et al., 2006; Lee et al., 2006). Thus, BD and reduced IIS appear to modulate both longevity and proteotoxicity through distinct genetic pathways.




Fig. 4 Bacterial food deprivation (BD) suppresses proteotoxicity by a mechanism distinct from insulin/IGF-1-like signaling. (A) RNAi inhibition of *daf-16* significantly accelerates paralysis in Q_{35} YFP animals relative to growth on empty vector bacteria (8.0×10^{-3}), but does not prevent suppression of paralysis by BD ($p = 4.5 \times 10^{-9}$). (B) RNAi inhibition of *daf-2* significantly reduces paralysis in Q_{35} YFP animals relative to growth on empty vector bacteria ($p = 7.6 \times 10^{-7}$). BD further reduces paralysis of *daf-2* RNAi-treated animals ($p = 1.0 \times 10^{-14}$).

Dietary restriction protects against proteotoxicity through an *hsf-1* dependent mechanism

The heat shock transcription factor, *hsf-1*, regulates expression of many different heat-inducible target genes such as small heat shock proteins, and has been implicated in modulating both longevity (Garigan et al., 2002; Hsu et al., 2003; Morley & Morimoto, 2004) and proteotoxicity (Hsu et al., 2003; Cohen et al., 2006). In order to determine whether *hsf-1* is required for phenotypes associated with DR, we subjected *hsf-1(sy441)* animals to BD and monitored thermotolerance and longevity (Kaeberlein et al., 2006; Lee et al., 2006). The *hsf-1(sy441)* allele contains a point mutation in the *hsf-1* transactivation domain that prevents induction of *hsf-1* target genes in response to heat stress (Hajdu-Cronin et al., 2004). Unlike the case in N2 animals, BD failed to result in enhanced thermotolerance in *hsf-1(sy441)* animals (Fig. 5A). The effect of BD on the longevity of *hsf-1(sy441)* animals was complex: BD failed to significantly increase median lifespan ($p = 0.34$); however, maximum lifespan appeared to be increased (Fig. 5B). The inability of BD to increase the median lifespan of *hsf-1(sy441)* animals is notably different than the increase in both median and maximum lifespan of *daf-16(mu86)* animals in response to BD (Fig. 5B; $p = 3.9 \times 10^{-19}$).

There are two possibilities to explain the observation that *hsf-1* mutation prevents median, but not maximum, lifespan extension from BD. First, it may indicate that lifespan extension from BD involves two different mechanisms, one of which acts through *hsf-1* and one of which is *hsf-1* independent. Alternatively, the residual *hsf-1* function present in *hsf-1(sy441)* animals may be sufficient to partially mediate BD-associated lifespan extension. Consistent with this latter possibility, RNAi knockdown of *hsf-1* completely prevented both median and maximum lifespan extension by BD (Fig. 5C). Thus, mutation of the *hsf-1* transactivation domain prevents induction of thermotolerance and median lifespan extension in response to BD, and RNAi knockdown of *hsf-1* completely abrogates median and maximum lifespan extension by BD.

Finally, we asked whether RNAi knockdown of *hsf-1* would affect suppression of proteotoxicity by BD. Q_{35} YFP animals were maintained on *hsf-1* RNAi from egg until the second day of adulthood, then transferred to either empty vector bacteria or

BD. The onset of paralysis was accelerated when *hsf-1* function was reduced by RNAi, relative to animals maintained on empty vector RNAi from egg (Fig. 6A; $p = 1.1 \times 10^{-11}$). Strikingly, BD failed to significantly suppress paralysis in Q_{35} YFP animals where *hsf-1* was inhibited ($p = 0.5$). RNAi knockdown of *hsf-1* had a similar effect on paralysis in $A\beta_{42}$ animals (Fig. 6B; $p = 0.1$). Thus, we conclude that *hsf-1* activity is required both for lifespan extension and for the general suppression of proteotoxicity associated with BD.

Discussion

Relationship between DR, IIS, and *hsf-1*

Dietary restriction and reduced IIS result in several shared phenotypes, including enhanced thermotolerance, enhanced resistance to proteotoxicity, and increased longevity. In the case of reduced IIS, each of these phenotypes is known to be dependent on both *hsf-1* and *daf-16*, which function to coregulate the expression of a subset of IIS-responsive genes (Hsu et al., 2003; Murphy et al., 2003). In contrast, the effects of DR require *hsf-1*, but *daf-16* is expendable. A model consistent with these observations is one in which BD promotes the activity of a transcription factor that functions in a manner similar to *daf-16* to regulate target genes co-operatively with *hsf-1* (Fig. 7A). In this model, BD would not act solely by inducing HSF-1 activity or expression, but HSF-1 would function together with another transcription factor. Thus, HSF-1 activity would be necessary, but not sufficient, to recapitulate many of the phenotypes associated with BD, consistent with the observation that *daf-16* is required for lifespan extension from overexpression of *hsf-1* in animals fed a normal bacterial diet (Hsu et al., 2003; Morley & Morimoto, 2004).

Recently, the *daf-16* homolog *pha-4* was found to be required for lifespan extension in *eat-2* mutants (Panowski et al., 2007). Expression of *pha-4* is normally low during adulthood, but is increased in *eat-2* mutants (Panowski et al., 2007). Unlike *hsf-1* (Hsu et al., 2003), however, overexpression of *pha-4* is not sufficient to increase lifespan in control-fed animals (Panowski et al., 2007). Thus, it could be that in well-fed animals *hsf-1* activity is limiting for longevity, but in response to nutrient




Fig. 5 Heat shock factor 1 is required for enhanced thermotolerance and increased lifespan in response to bacterial food deprivation (BD). (A) Animals carrying a mutant allele of *hsf-1*, *hsf-1(sy441)*, fail to show increased survival at 38 °C when subjected to BD. Error bars are standard error of the mean. (B) BD fails to increase median lifespan of *hsf-1(sy441)* animals ($p = 0.34$) but significantly increases the lifespan of *daf-16(mu86)* animals. (C) RNAi inhibition of *hsf-1* prevents lifespan extension from BD.




Fig. 6 Heat shock factor 1 is required for suppression of proteotoxicity in response to bacterial food deprivation (BD). (A) BD significantly reduces paralysis of Q_{35} YFP animals grown on empty vector bacteria ($p = 1.1 \times 10^{-11}$), but does not reduce paralysis of Q_{35} YFP animals grown on *hsf-1* RNAi ($p = 0.51$). (B) BD significantly reduces paralysis of $A\beta_{42}$ animals grown on empty vector bacteria ($p = 8.9 \times 10^{-14}$), but does not reduce paralysis of $A\beta_{42}$ animals grown on *hsf-1* RNAi ($p = 0.1$).

deprivation both *hsf-1* and *pha-4* are activated and promote longevity. Since *skn-1* is also required for lifespan extension from DR (Bishop & Guarente, 2007), it may be the case that *hsf-1* and *skn-1* act to coregulate genes important for DR or that *skn-1* acts upstream of *hsf-1* and *pha-4* in response to DR. Similar to the case for *pha-4*, increased expression of *skn-1* is

not sufficient to increase the lifespan of control-fed animals (Bishop & Guarente, 2007). *Skn-1* is important for regulating gene expression in response to oxidative stress (An & Blackwell, 2003; Inoue et al., 2005), but no role for *pha-4* or *skn-1* in modulating the response to proteotoxic stress has been described.




Fig. 7 Models for the role of *hsf-1* in determining longevity and sensitivity to proteotoxicity. (A) The effect of BD is mediated by *hsf-1*. Under conditions of abundant food, the dietary restriction (DR) pathway is inactive and *hsf-1* activity and insulin/IGF-1-like signaling (IIS) are at normal levels. This leads to short lifespan and sensitivity to proteotoxicity. Under DR conditions, *hsf-1* is activated along with other DR-responsive proteins (perhaps *skn-1* and *pha-4*) to increase lifespan and suppress proteotoxicity. In mutants with reduced IIS, the DR pathway is inactive, but lifespan and resistance to proteotoxicity are increased through activation of *daf-16*. (B) BD and *hsf-1* function in parallel to alter protein homeostasis. *hsf-1* activity up-regulates chaperone activity, which decreases the amount of misfolded proteins *in vivo*. At the same time, BD might both increase the rate of protein turnover and decrease the synthesis of new protein. These three factors would combine to improve protein homeostasis, decreasing the rate of aging.

An alternative, but not mutually exclusive, possibility is that BD and *hsf-1* function in parallel to promote longevity by improving protein homeostasis (Fig. 7B). This hypothesis is attractive, since *hsf-1* is known to regulate expression of chaperones, and BD could contribute to protein homeostasis by enhancing protein degradation or by modulating mRNA translation (or both). In *C. elegans*, starvation causes rapid turnover of muscle proteins and induces autophagy (Fostel et al., 2003; Jia & Levine, 2007), suggesting that turnover of damaged proteins is likely to be enhanced by BD. This model also explains that apparent correlation between longevity and resistance to proteotoxicity. For example, a recent RNAi screen identified several genes that play a role in both longevity determination and the formation of polyglutamine aggregates (Curran & Ruvkun, 2007). We have also observed that RNAi of several

previously reported longevity factors (Dillin et al., 2002; Lee et al., 2003; Hamilton et al., 2005; Hansen et al., 2005, 2007; Pan et al., 2007) also suppresses paralysis in Q₃₅YFP and A β ₄₂ animals (K.A.S., C.D., D.C. and M.K., unpublished data). Further studies will be necessary to clarify the precise relationship between *hsf-1* and DR, and the importance of protein homeostasis as a determining factor for longevity.

Conclusion

The data presented here demonstrate a novel and previously unsuspected role for *hsf-1* as a key mediator of DR. This finding is noteworthy, as *hsf-1* orthologs are highly conserved from yeast to humans, as are many of the known *hsf-1* targets. Although it remains unclear whether *hsf-1* is activated by DR

in mammals, DR enhances Hsf1 binding activity in aging rats (Heydari *et al.*, 1996), indicating that the link between DR and *hsf-1* activity may be conserved. Activation of *hsf-1* or specific *hsf-1* target genes may prove therapeutic toward proteotoxic insults leading to neurodegeneration and other age-associated pathologies in people and may also mimic the longevity enhancing effects of DR.

Experimental procedures

Strains and genetics

Standard procedures for *C. elegans* strain maintenance and manipulation were used. Experiments were conducted at 20 °C unless otherwise noted in the text. Nematode strains were obtained from the *Caenorhabditis* Genetics Center, or the laboratories of Dr Richard Morimoto (Northwestern University, Chicago, IL, USA), Dr Chris Link (University of Colorado, Boulder, CO, USA) or Dr Jim Thomas (University of Washington, Seattle, WA, USA). The strains used in this study are listed in Supplementary Table S1.

Bacterial food deprivation

Bacterial food deprivation experiments were carried out essentially as described previously (Kaeberlein *et al.*, 2006). In brief, adult hermaphrodites were allowed to lay eggs on nematode growth medium (NGM) (Brenner, 1974) containing UV-killed *E. coli* OP50. At L4, worms were transferred to fresh NGM + UV-killed OP50 supplemented with 50 μM 5-fluorodeoxyuridine (FUDR) to prevent further eggs from hatching. Unless otherwise stated, worms were then transferred to experimental media at the second day of adulthood: NGM + UV-killed OP50 + 50 μM FUDR (control fed) or NGM + 50 μM FUDR (Kaeberlein *et al.*, 2006). For the experiment shown in Fig. 1(C), adult worms were transferred to experimental media on the days indicated in the figure legend. OP50 were UV-killed by irradiation with a Stratagene UV Stratalinker 2400 (La Jolla, CA, USA) on maximal energy setting. UV killing was periodically verified by streaking UV-irradiated bacteria onto Luna-Bertani (LB) plates and confirming absence of growth.

RNA interference combined with BD

RNA interference (RNAi) experiments were conducted using feeding protocols according to standard procedures. Unless otherwise stated, RNAi feeding strains were obtained from the Ahringer RNAi library (Kamath *et al.*, 2003); library clones were sequenced for verification. RNAi plates consisted of NGM supplemented with 1 mM β -D-isothiothiopyranoside and 25 $\mu\text{g mL}^{-1}$ carbenicillin. Unless otherwise indicated, worms were raised on RNAi bacteria from egg. At L4, they were then transferred to plates containing freshly seeded RNAi bacteria plus 50 μM FUDR. To combine RNAi with BD, worms were maintained on the indicated RNAi bacteria until the second day

of adulthood, at which point they were transferred to experimental media: RNAi plates + 50 μM FUDR seeded with empty vector control bacteria (control fed) or RNAi plates + 50 μM FUDR (Kaeberlein *et al.*, 2006). The *daf-2* RNAi construct used for these studies was provided to us by J. McElwee.

Paralysis and lifespan quantification

The paralysis of worms expressing any of the indicated transgenes (Q35-YFP, A β 1–42, or GFP-degron) was determined by visual analysis. Worms were scored as paralyzed if they were unable to make forward progress on the NGM surface in response to both plate-tapping and tail-prodding. If paralyzed, worms were scored as alive if nose-tapping resulted in nose movement. If no nose movement could be seen, worms were scored as dead.

Thermotolerance

Animals were maintained at 20 °C on either control-fed or BD media as described until the sixth day of adulthood (4 days without food for the BD group). The morning of the sixth day of adulthood, plates (each containing approximately 20 animals) were transferred to 38 °C. Viability was determined periodically by removing one plate at a time and assaying for movement in response to nose-prodding.

Fluorescence microscopy and quantification of Q₃₅YFP aggregates

Movie files were generated using Canon Powershot S31S digital camera (Lake Success, NY, USA) connected to the eyepiece of a Zeiss SteREO Lumar.V12 microscope (Thornwood, NY, USA) equipped with a broad-range GFP filter. Still GFP images were obtained with a Bio-Rad MRC-1024 LSCM (Hercules, CA, USA). This package included a computer-assisted Zeiss inverted microscope and 15 mW KryptonArgon laser (ILT laser, Aachen, Germany) emitting three excitation lines at 488 nm, 568 nm, and 647 nm. The 488-nm line was used to excite the green fluorescent protein. The emitted green fluorescence was filtered through a 522-nm DF 32 band pass filter. Unless otherwise stated, a $\times 10$ objective lens was used for fluorescence measurements. Identical settings were used for all worms analyzed (laser power at 1%; gain at 1500; Iris at 4.0 μm ; scan time at 488 lines/sec). The worms were paralyzed with 25 mM sodium azide and placed on a glass slide under a cover slip to flatten the worm somewhat and optimize viewing. The microscope was focused at the position of maximum fluorescence for each worm. The images were analyzed using Image J 1.341.5.0_07 software [available from National Institutes of Health (NIH) website: <http://rsb.info.nih.gov/ij/java>] and a Macintosh computer with OS X operating system. The 'count objects' command of Image J was used to count the number of fluorescent spots in the images. A constant minimum intensity threshold was set at 75/255 pixels, and a minimum spot size of 3 pixels/spot were used for all worms analyzed.

Quantification of Q₃₅-YFP expression by immunoblot analysis

Worms were prepared for immunoblot analysis using the control-fed and BD protocols described above. At day 9 of adulthood, 1000 worms per group were harvested in 1 mL M9 buffer and concentrated by low speed centrifugation. Worms were washed in M9 twice, then resuspended in phosphate-buffered saline at a concentration of 1 worm per microliter buffer. They were homogenized in a glass homogenizer and spun at 750×g for 3 min; the supernatant was collected as 'total lysate'. Samples were subjected to sodium dodecylsulfate-polyacrylamide gel electrophoresis analysis followed by immunoblot, according to standard protocols. Q₃₅-YFP was detected with a GFP-specific antibody (Roche Applied Science, Indianapolis, IN, USA); protein loading was controlled for using a β-actin-specific antibody (Abcam, Cambridge, MA, USA). The relative abundance of Q₃₅-YFP protein and actin was determined by quantifying the band intensities after probing with the GFP-specific antibody or the β-actin-specific antibody, respectively. When normalized to actin, the relative abundance of Q₃₅-YFP in control-fed to BD animals was 0.96 ± 0.1, based on three biological replicates (independent experiments and independent blots).

Statistical analysis and replication

A Wilcoxon rank sum test (MATLAB 'ranksum' function; Mann-Whitney *U*-test) was used to generate *p*-values to determine statistical significance for lifespan and paralysis assays. This test is a nonparametric comparison of medians between two groups. Each experiment was repeated at least three times with similar results.

Acknowledgments

We would like to thank the R. Morimoto and C. Link for providing the transgenic *C. elegans* strains used in this study. We would also like to thank J. McElwee and J. Thomas for helpful discussion and shared reagents. Some strains used in this work were provided by the *Caenorhabditis* Genetics Center, which is funded by the NIH National Center for Research Resources. This work was conducted while M.K. was an AFAR Research Grant recipient. This work was also supported by a grant from the Ellison Medical Foundation to M. Kaerberlein and B. K. Kennedy. K. A. Steinkraus is supported by a postdoctoral fellowship from the Hereditary Disease Foundation. E. D. Smith was supported by NIH Training Grant P30 AG013280.

References

An JH, Blackwell TK (2003) SKN-1 links *C. elegans* mesendodermal specification to a conserved oxidative stress response. *Genes Dev.* **17**, 1882–1893.
 Avery L (1993) The genetics of feeding in *Caenorhabditis elegans*. *Genetics* **133**, 897–917.

Bishop NA, Guarente L (2007) Two neurons mediate diet-restriction-induced longevity in *C. elegans*. *Nature* **447**, 545–549.
 Bonini NM, La Spada AR (2005) Silencing polyglutamine degeneration with RNAi. *Neuron* **48**, 715–718.
 Borrell-Pages M, Zala D, Humbert S, Saudou F (2006) Huntington's disease: from huntingtin function and dysfunction to therapeutic strategies. *Cell Mol. Life Sci.* **63**, 2642–2660.
 Brenner S (1974) The genetics of *Caenorhabditis elegans*. *Genetics*. **77**, 71–94.
 Brignull HR, Morley JF, Morimoto RI (2007) The stress of misfolded proteins: *C. elegans* models for neurodegenerative disease and aging. *Adv. Exp. Med. Biol.* **594**, 167–189.
 Cohen E, Bieschke J, Perciavalle RM, Kelly JW, Dillin A (2006) Opposing activities protect against age onset proteotoxicity. *Science* **313**: 1604–1610.
 Curran SP, Ruvkun G (2007) Lifespan regulation by evolutionarily conserved genes essential for viability. *PLoS Genet.* **3**, e56.
 Dillin A, Hsu AL, Arantes-Oliveira N, Lehrer-Graiwer J, Hsin H, Fraser AG, Kamath RS, Ahringer J, Kenyon C (2002) Rates of behavior and aging specified by mitochondrial function during development. *Science* **298**, 2398–2401.
 Faber PW, Alter JR, MacDonald ME, Hart AC (1999) Polyglutamine-mediated dysfunction and apoptotic death of a *Caenorhabditis elegans* sensory neuron. *Proc. Natl Acad. Sci. USA* **96**, 179–184.
 Fostel JL, Benner Coste L, Jacobson LA (2003) Degradation of transgene-coded and endogenous proteins in the muscles of *Caenorhabditis elegans*. *Biochem. Biophys. Res. Commun.* **312**, 173–177.
 Friedman DB, Johnson TE (1988a) A mutation in the age-1 gene in *Caenorhabditis elegans* lengthens life and reduces hermaphrodite fertility. *Genetics* **118**, 75–86.
 Friedman DB, Johnson TE (1988b) Three mutants that extend both mean and maximum life span of the nematode, *Caenorhabditis elegans*, define the age-1 gene. *J. Gerontol.* **43**, B102–B109.
 Garigan D, Hsu AL, Fraser AG, Kamath RS, Ahringer J, Kenyon C (2002) Genetic analysis of tissue aging in *Caenorhabditis elegans*: a role for heat-shock factor and bacterial proliferation. *Genetics* **161**, 1101–1112.
 Hajdu-Cronin YM, Chen WJ, Sternberg PW (2004) The L-type cyclin CYL-1 and the heat-shock-factor HSF-1 are required for heat-shock-induced protein expression in *Caenorhabditis elegans*. *Genetics* **168**, 1937–1949.
 Hamilton B, Dong Y, Shindo M, Liu W, Odell I, Ruvkun G, Lee SS (2005) A systematic RNAi screen for longevity genes in *C. elegans*. *Genes Dev.* **19**, 1544–1555.
 Hansen M, Hsu AL, Dillin A, Kenyon C (2005) New genes tied to endocrine, metabolic, and dietary regulation of lifespan from a *Caenorhabditis elegans* genomic RNAi screen. *PLoS Genet.* **1**, 119–128.
 Hansen M, Taubert S, Crawford D, Libina N, Lee SJ, Kenyon C (2007) Lifespan extension by conditions that inhibit translation in *Caenorhabditis elegans*. *Aging Cell* **6**, 95–110.
 Heydari AR, You S, Takahashi R, Gutschmann A, Sarge KD, Richardson A (1996) Effect of caloric restriction on the expression of heat shock protein 70 and the activation of heat shock transcription factor 1. *Dev. Genet.* **18**, 114–124.
 Houthoofd K, Braeckman BP, Lenaerts I, Brys K, De Vreese A, Van Eygen S, Vanfleteren JR (2002) Axenic growth up-regulates mass-specific metabolic rate, stress resistance, and extends life span in *Caenorhabditis elegans*. *Exp. Gerontol.* **37**, 1371–1378.
 Houthoofd K, Braeckman BP, Johnson TE, Vanfleteren JR (2003) Life extension via dietary restriction is independent of the Ins/IGF-1 signalling pathway in *Caenorhabditis elegans*. *Exp. Gerontol.* **38**, 947–954.
 Hsu AL, Murphy CT, Kenyon C (2003) Regulation of aging and

- age-related disease by DAF-16 and heat-shock factor. *Science* **300**, 1142–1145.
- Inoue H, Hisamoto N, An JH, Oliveira RP, Nishida E, Blackwell TK, Matsumoto K (2005) The *C. elegans* p38 MAPK pathway regulates nuclear localization of the transcription factor SKN-1 in oxidative stress response. *Genes Dev.* **19**, 2278–2283.
- Jia K, Levine B (2007) Autophagy is required for dietary restriction-mediated life span extension in *C. elegans*. *Autophagy* **3**, 597–599.
- Kaeberlein TL, Smith ED, Tsuchiya M, Welton KL, Thomas JH, Fields S, Kennedy BK, Kaeberlein M (2006) Lifespan extension in *Caenorhabditis elegans* by complete removal of food. *Aging Cell* **5**, 487–494.
- Kamath RS, Fraser AG, Dong Y, Poulin G, Durbin R, Gotta M, Kanapin A, Le Bot. N, Moreno S, Sohrmann M, Welchman DP, Zipperlin P, Ahringer J (2003) Systematic functional analysis of the *Caenorhabditis elegans* genome using RNAi. *Nature* **421**, 231–237.
- Kenyon C, Chang J, Gensch E, Rudner A, Tabtiang R (1993) A *C. elegans* mutant that lives twice as long as wild type. *Nature* **366**, 461–464.
- Khan LA, Bauer PO, Miyazaki H, Lindenberg KS, Landwehrmeyer BG, Nukina N (2006) Expanded polyglutamines impair synaptic transmission and ubiquitin-proteasome system in *Caenorhabditis elegans*. *J. Neurochem.* **98**, 576–587.
- Kimura KD, Tissenbaum HA, Liu Y, Ruvkun G (1997) *daf-2*, an insulin receptor-like gene that regulates longevity and diapause in *Caenorhabditis elegans*. *Science* **277**, 942–946.
- Koo EH, Lansbury PT Jr, Kelly JW (1999) Amyloid diseases: abnormal protein aggregation in neurodegeneration. *Proc. Natl Acad. Sci. USA* **96**, 9989–9990.
- Lakowski B, Hekimi S (1998) The genetics of caloric restriction in *Caenorhabditis elegans*. *Proc. Natl Acad. Sci. USA* **95**, 13091–13096.
- Lee SS, Lee RY, Fraser AG, Kamath RS, Ahringer J, Ruvkun G (2003) A systematic RNAi screen identifies a critical role for mitochondria in *C. elegans* longevity. *Nat Genet.* **33**, 40–48.
- Lee GD, Wilson MA, Zhu M, Wolkow CA, de Cabo R, Ingram DK, Zou S (2006) Dietary deprivation extends lifespan in *Caenorhabditis elegans*. *Aging Cell* **5**, 515–524.
- Lin K, Dorman JB, Rodan A, Kenyon C (1997) *daf-16*: an HNF-3/forkhead family member that can function to double the life-span of *Caenorhabditis elegans*. *Science* **278**, 1319–1322.
- Link CD (1995) Expression of human β -amyloid peptide in transgenic *Caenorhabditis elegans*. *Proc. Natl Acad. Sci. USA* **92**, 9368–9372.
- Link CD (2001) Transgenic invertebrate models of age-associated neurodegenerative diseases. *Mech. Ageing Dev.* **122**, 1639–1649.
- Link CD, Fonte V, Hiester B, Yerg J, Ferguson J, Csontos S, Silverman MA, Stein GH (2006) Conversion of green fluorescent protein into a toxic, aggregation-prone protein by C-terminal addition of a short peptide. *J. Biol. Chem.* **281**, 1808–1816.
- Martin B, Mattson MP, Maudsley S (2006) Caloric restriction and intermittent fasting: two potential diets for successful brain aging. *Ageing Res. Rev.* **5**, 332–353.
- Masoro EJ (2005) Overview of caloric restriction and ageing. *Mech. Ageing Dev.* **126**, 913–922.
- Moerman DG, Baillie DL (1979) Genetic organization in *Caenorhabditis elegans*: fine-structure analysis of the *unc-22* gene. *Genetics* **91**, 95–103.
- Morley JF, Morimoto RI (2004) Regulation of longevity in *Caenorhabditis elegans* by heat shock factor and molecular chaperones. *Mol. Biol. Cell* **15**, 657–664.
- Morley JF, Brignull HR, Weyers JJ, Morimoto RI (2002) The threshold for polyglutamine-expansion protein aggregation and cellular toxicity is dynamic and influenced by aging in *Caenorhabditis elegans*. *Proc. Natl Acad. Sci. USA* **99**, 10417–10422.
- Morris JZ, Tissenbaum HA, Ruvkun G (1996) A phosphatidylinositol-3-OH kinase family member regulating longevity and diapause in *Caenorhabditis elegans*. *Nature* **382**, 536–539.
- Murphy CT, McCarroll SA, Bargmann CI, Fraser A, Kamath RS, Ahringer J, Li H, Kenyon C (2003) Genes that act downstream of DAF-16 to influence the lifespan of *Caenorhabditis elegans*. *Nature* **424**, 277–283.
- Ogg S, Paradis S, Gottlieb S, Patterson GI, Lee L, Tissenbaum HA, Ruvkun G (1997) The Fork head transcription factor DAF-16 transduces insulin-like metabolic and longevity signals in *C. elegans*. *Nature* **389**, 994–999.
- Pan KZ, Palter JE, Rogers AN, Olsen A, Chen D, Lithgow GJ, Kapahi P (2007) Inhibition of mRNA translation extends lifespan in *Caenorhabditis elegans*. *Aging Cell* **6**, 111–119.
- Panowski SH, Wolff S, Aguilaniu H, Durieux J, Dillin A (2007) PHA-4/Foxa mediates diet-restriction-induced longevity of *C. elegans*. *Nature* **447**, 550–555.
- Parker JA, Connolly JB, Wellington C, Hayden M, Dausset J, Neri C (2001) Expanded polyglutamines in *Caenorhabditis elegans* cause axonal abnormalities and severe dysfunction of PLM mechanosensory neurons without cell death. *Proc. Natl Acad. Sci. USA* **98**, 13318–13323.
- Parker JA, Arango M, Abderrahmane S, Lambert E, Tourette C, Catoire H, Neri C (2005) Resveratrol rescues mutant polyglutamine cytotoxicity in nematode and mammalian neurons. *Nat. Genet.* **37**, 349–350.
- Rogalski TM, Gilchrist EJ, Mullen GP, Moerman DG (1995) Mutations in the *unc-52* gene responsible for body wall muscle defects in adult *Caenorhabditis elegans* are located in alternatively spliced exons. *Genetics* **139**, 159–169.
- Satyal SH, Schmidt E, Kitagawa K, Sondheimer N, Lindquist S, Kramer JM, Morimoto RI (2000) Polyglutamine aggregates alter protein folding homeostasis in *Caenorhabditis elegans*. *Proc. Natl Acad. Sci. USA* **97**, 5750–5755.
- Vanfleteren JR, Braeckman BP (1999) Mechanisms of life span determination in *Caenorhabditis elegans*. *Neurobiol. Aging* **20**, 487–502.
- Waterston RH, Thomson JN, Brenner S (1980) Mutants with altered muscle structure of *Caenorhabditis elegans*. *Dev. Biol.* **77**, 271–302.
- Weindruch RH, Walford RL (1988) *The Retardation of Aging and Disease by Dietary Restriction*. Springfield, IL: Thomas.
- Youngman LD, Park JY, Ames BN (1992) Protein oxidation associated with aging is reduced by dietary restriction of protein or calories. *Proc. Natl Acad. Sci. USA* **89**, 9112–9116.

Supplemental materials

The following supplementary material is available for this article:

Fig. S1 Bacterial food deprivation (BD) increases the survival of proteotoxicity models. The lifespans of the indicated strains were significantly extended on BD relative to control-fed conditions. (A) Q_{35} YFP animals ($p = 1.1 \times 10^{-7}$); (B) $A\beta_{42}$ animals ($p = 8.7 \times 10^{-4}$).

Fig. S2 Suppression of proteotoxicity by bacterial food deprivation (BD) is not correlated with reduced aggregate formation or reduced Q_{35} YFP protein. Aggregates were visualized in 14-day-old Q_{35} YFP animals by fluorescence microscopy and quantified as described in the Experimental procedures. (A) Control-fed animals. (B) BD animals. (C) Immunoblot analysis of total protein lysates from control-fed or BD animals. Equal total protein was loaded per lane and probed with anti-GFP (see the Experimental procedures). The * marks a band of unknown identity specific to the Q_{35} YFP animals. The # marks a doublet running at the predicted size of the Q_{35} YFP transgene.

Fig. S3 Bacterial food deprivation (BD) suppresses amyloid-beta toxicity by a mechanism that is independent of IIS and dependent on *hsf-1*. (A) RNAi inhibition of *daf-16* significantly accelerates paralysis in A β ₄₂ animals relative to growth on empty vector (EV) bacteria in control-fed (CF) animals ($p = 2.6 \times 10^{-7}$), but does not prevent suppression of paralysis by BD ($p = 1.7 \times 10^{-10}$). (B) RNAi inhibition of *daf-2* significantly reduces paralysis in A β ₄₂ animals relative to growth on EV bacteria ($p = 5.3 \times 10^{-4}$). BD further reduces paralysis of *daf-2* RNAi-treated animals ($p = 2.4 \times 10^{-8}$).

Table S1 Lifespan data for Supplementary Fig. S1A

Table S2 Lifespan data for Supplementary Fig. S1B

Table S3 Lifespan data for Fig. 5B

Table S4 Lifespan data for Fig. 5C

Video File 1. Control fed Q₃₅ YFP animals are paralyzed at day 8 of adulthood.

Video File 2. Bacterial food deprivation (BD) suppresses the paralysis of Q₃₅ YFP animals at day 8 of adulthood.

Video File 3. Fluorescence microscopy of control fed Q₃₅ YFP animal at day 8 of adulthood.

Video File 4. Fluorescence microscopy of Q₃₅ YFP animal on Bacterial food deprivation (BD) at day 8 of adulthood.

Video File 5. Thrashing of control fed Q₃₅ YFP animal at day 8 of adulthood.

Video File 6. Thrashing of Q₃₅ YFP animal on Bacterial food deprivation (BD) at day 8 of adulthood.

Video File 7. Control fed unc-52(e444) mutants at day 5 of adulthood.

Video File 8. Unc-52(e444) mutants on bacterial food deprivation (BD) at day 5 of adulthood.

This material is available as part of the online article from: <http://www.blackwell-synergy.com/doi/abs/10.1111/j.1474-9726.2008.00385.x>

(This link will take you to the article abstract).

Please note: Blackwell Publishing are not responsible for the content or functionality of any supplementary materials supplied by the authors. Any queries (other than missing material) should be directed to the corresponding author for the article.