

RESEARCH ARTICLE

# Differences in Hematological Traits between High- and Low-Altitude Lizards (Genus *Phrynocephalus*)

Songsong Lu, Ying Xin, Xiaolong Tang, Feng Yue, Huihui Wang, Yucheng Bai, Yonggang Niu, Qiang Chen\*

Institute of Biochemistry and Molecular Biology, School of Life Science, Lanzhou University, Lanzhou, China

\* [cheng@lzu.edu.cn](mailto:cheng@lzu.edu.cn)



## Abstract

*Phrynocephalus erythrurus* (Lacertilia: Agamidae) is considered to be the highest living reptile in the world (about 4500-5000 m above sea level), whereas *Phrynocephalus przewalskii* inhabits low altitudes (about 1000-1500 m above sea level). Here, we report the differences in hematological traits between these two different *Phrynocephalus* species. Compared with *P. przewalskii*, the results indicated that *P. erythrurus* own higher oxygen carrying capacity by increasing red blood cell count (RBC), hemoglobin concentration ([Hb]) and hematocrit (Hct) and these elevations could promote oxygen carrying capacity without disadvantage of high viscosity. The lower partial pressure of oxygen in arterial blood (PaO<sub>2</sub>) of *P. erythrurus* did not cause the secondary alkalosis, which may be attributed to an efficient pulmonary system for oxygen (O<sub>2</sub>) loading. The elevated blood-O<sub>2</sub> affinity in *P. erythrurus* may be achieved by increasing intrinsic O<sub>2</sub> affinity of isoHbs and balancing the independent effects of potential heterotropic ligands. We detected one  $\alpha$ -globin gene and three  $\beta$ -globin genes with 1 and 33 amino acid substitutions between these two species, respectively. Molecular dynamics simulation results showed that amino acids substitutions in  $\beta$ -globin chains could lead to the elimination of hydrogen bonds in T-state Hb models of *P. erythrurus*. Based on the present data, we suggest that *P. erythrurus* have evolved an efficient oxygen transport system under the unremitting hypobaric hypoxia.

## OPEN ACCESS

**Citation:** Lu S, Xin Y, Tang X, Yue F, Wang H, Bai Y, et al. (2015) Differences in Hematological Traits between High- and Low-Altitude Lizards (Genus *Phrynocephalus*). PLoS ONE 10(5): e0125751. doi:10.1371/journal.pone.0125751

**Academic Editor:** Yang Zhang, University of Michigan, UNITED STATES

**Received:** November 11, 2014

**Accepted:** March 26, 2015

**Published:** May 8, 2015

**Copyright:** © 2015 Lu et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

**Data Availability Statement:** All relevant data are within the paper.

**Funding:** Research funding was provided by the National Natural Science Foundation of China (No. 31272313 and No. 31472005 to Q. Chen). The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

**Competing Interests:** The authors have declared that no competing interests exist.

## Introduction

Animals living in high altitude habitats have to manage certain additional physiological challenges in conditions of reduced oxygen availability and low ambient temperature. Matching O<sub>2</sub> supply (inspired air) with O<sub>2</sub> demand (tissue mitochondria) is necessary and important for both high-altitude natives and animals to acclimate to high altitude [1–3].

To live under high-altitude hypoxia, animals usually adopt some strategies or adjustments in the oxygen transport system. These adjustments should include at least three aspects [1]. Firstly, the highly efficient pulmonary ventilation and pulmonary O<sub>2</sub> diffusion can help

maintain high O<sub>2</sub> partial pressures of arterial blood (PaO<sub>2</sub>). Pulmonary ventilation is mainly affected by the partial pressures of O<sub>2</sub> and CO<sub>2</sub> and the pH of arterial blood. These factors normally stimulate breathing via central and peripheral chemoreceptor [4–6]. Long time acclimatization to high altitude can relieve this hypoxic ventilatory response [5,7]. In addition, pulmonary O<sub>2</sub> diffusion is mainly affected by the thickness and surface area of the pulmonary blood-gas interface [8–10]. Secondly, in order to ensure an adequate supply of O<sub>2</sub> to the cells of aerobically metabolizing tissues, circulatory O<sub>2</sub> delivery and tissue O<sub>2</sub> diffusion can be enhanced by increasing the total cardiac output and the blood-O<sub>2</sub> capacitance coefficient, such as by elevating hemoglobin concentration ([Hb]) and hematocrit (Hct) [1,11]. A moderately increased Hct is conducive to enhancing O<sub>2</sub> carrying capacity of arterial blood, while an excessively increased Hct will result in the increase of blood viscosity to reduce the O<sub>2</sub> carrying capacity by a higher peripheral vascular resistance and hence add more burdens for heart and blood circulation system [12–15]. Finally, fine-tuned adjustments in blood-O<sub>2</sub> affinity are very important for O<sub>2</sub> transport system during high altitude hypoxia. The regulation process may involve changes in intrinsic Hb-O<sub>2</sub> affinity, the sensitivity of Hb to allosteric effectors and compensatory changes in concentration of allosteric effectors (particularly organic and inorganic anions) within the erythrocyte [1,16–19]. In addition, adaptive genetic variations of  $\alpha$ - and  $\beta$ -like globin genes have also identified in many studies [20–24].

The Hb of jawed vertebrates is a heterotetramer which contains two  $\alpha$ -globin chains and two  $\beta$ -globin chains with a heme group in each chain. During the process of vertebrate evolution, the  $\alpha$ - and  $\beta$ -globin gene families have been subjected to repeated rounds of gene duplication and divergence [25,26]. Furthermore, studies have shown that the developmental regulation of Hb synthesis in some reptiles differ from other tetrapod vertebrates [27–29].

The mechanisms underlying the physiological acclimatization and genetic adaptation to high-altitude hypoxia have been studied extensively in birds and mammals. Although these mechanisms have also been found in some reptiles, how reptiles adapt to high-altitude hypoxia still remains largely unknown. Among over 40 species of Asian lizard genus *Phrynocephalus*, several phylogenetic independent lineages (*P. putjatia*, *P. vlangalii vlangalii*, *P. vlangalii pylzowi*, *P. vlangalii nanschanica*, *P. theobaldi theobaldi*, *P. theobaldi orientalis*, *P. erythrurus erythrurus* and *P. erythrurus prava*) could be found on the Qinghai-Tibetan Plateau (QTP) [30–33]. Recently, the toad-headed lizard genus *Phrynocephalus* has drawn the attention of physiological and genetic researchers for its adaptation of broad geographical areas (about 2200–5000 m above the sea level) [33]. Red tail toad-headed lizard (*P. erythrurus*) is considered to be the highest living reptiles in the world (mostly 4500–5000 m above sea level) [34]. Previous study showed that inhibited metabolic, lower anaerobic metabolism, elevated mitochondrial efficiency and a possible higher utilization of fat may effectively compensate for the negative influence of cold and low PO<sub>2</sub> in *P. erythrurus* [35].

In this study, two closely related reptile species based on the biological evolution and phylogeny, *P. erythrurus* and *P. przewalskii* (mostly 1000–1500 m above sea level) were chosen to analyze the physiological and genetic characteristics of the highest living lizard in the following aspects: (1) evaluating oxygen transportation capacity through analyzing the degree of changes in hematological parameters; (2) preliminarily understanding the sequence divergence and expression of  $\alpha$ - and  $\beta$ -like globin genes in these two species; (3) analyzing the structural stability of potentially T-state isoHbs by equilibrium MD simulations. This study may provide important information and new insights into the adaptive mechanism of high-altitude ectothermic vertebrates.

## Materials and Methods

### Animals and sampling

All experiments were carried out according to protocols approved by the Ethics Committee of Animal Experiments at Lanzhou University and in accordance with guidelines from the China Council on Animal Care. *Phrynocephalus erythrurus* with an average weight of  $6.69 \pm 0.13$  g were captured by hand in the wild at Tuotuo River ( $34^{\circ}13'N$ ,  $92^{\circ}13'E$ , 4543 m above sea level), Qinghai province, China, and *P. przewalskii* (the low-altitude sample) with an average weight of  $6.92 \pm 0.14$  g were collected from a semi-desert areas in Minqin ( $38^{\circ}38'N$ ,  $103^{\circ}05'E$ , 1482 m above sea level), Gansu province, China. The Hoh-xil National Nature Reserve and Minqin Desert Control Station are only used for scientific research and the two authorities permitted us to capture the animals used in this study. Our studies did not involve endangered or protected species. All surgery was performed under sodium pentobarbital anaesthesia. Every effort was made to minimize the numbers used and any suffering experienced by the animals in the experiments.

Total 45 adult male lizards of each species were used in this study. Blood samples were obtained from the aortic arch directly in freshly anaesthetized lizard using a heparinized glass capillary tube. After blood collection, both liver and skeletal muscle were harvested by surgery and blotted with filter paper to remove excess liquid. The amount of blood taken from each animal was typically around 150  $\mu$ L. Blood samples for hematological parameters ( $n = 12$ , 50  $\mu$ L), blood gas ( $n = 12$ , 120  $\mu$ L) and organic phosphate (ATP,  $n = 12$ , 150  $\mu$ L) were placed immediately on ice and were measured within 1–2 hour nearby the capture location. Blood samples for the reverse-phase high performance liquid chromatography (RP-HPLC) ( $n = 12$ , the remaining 100  $\mu$ L of hematological measurements) and the liver and skeletal muscle for the rapid amplification of cDNA ends (RACE) PCR ( $n = 36$ , all blood collected lizards) were immediately frozen in liquid nitrogen, and then stored at  $-80^{\circ}C$  prior to use.

Nine lizards of each species from the collecting zone were brought to the laboratory at Lanzhou University ( $36^{\circ}05'N$ ,  $103^{\circ}86'E$ ) within 48 hours of capture. High- and low-altitude lizards were maintained in an air-conditioned room with two self-contained non-pressurized hypoxic chambers (100 cm length, 45 cm width and 45 cm height). In order to minimize the possible effect of changed environments, conditions of chambers were set up to equivalent altitude of 4550m and 1450m (temperature,  $16 \pm 0.5^{\circ}C$ ,  $35 \pm 0.5^{\circ}C$ , respectively, using 60 W bulbs and an air-conditioning system;  $PO_2$ ,  $\sim 92$  and  $\sim 137$  mmHg, respectively, using mixed gas of nitrogen and atmosphere; light: dark, 12h: 12h, using fluorescent lamps; food and water ad libitum) [35]. Blood samples were obtained using above-mentioned method. The determination of blood- $O_2$  affinity was finished within 4 days of the collection. After sampling, all lizards were sacrificed with an overdose of barbiturate.

### Hematological parameters

Hemoglobin concentration ([Hb]) was measured by mixing 10  $\mu$ L of blood with 2.5 mL of Van Kampen-Zijlstra solution and a spectrophotometer (Unico UV-2000) at the wave length of 540 nm. Red blood cell (RBC) count was measured by mixing 10  $\mu$ L of blood into 1.99 mL RBC diluents and the count of erythrocytes was made in hemocytometer under microscope. Hematocrits (HCT) were determined by a modified Guest-Siler (1934) technique [36]; and erythrocyte diameters were measured on dried smears with an ocular micrometer.

### Arterial blood gas analysis

Arterial blood gas and major inorganic ions were measured by a blood gas analyzer (OPTI CCA-TS Analysator, OPTI Medical System Inc., Roswell, GA) [37] with a ComfortSampler

arterial blood gas collection kit and type E-CI BP7559 cassettes. Total 120  $\mu\text{L}$  blood sample was used for arterial blood gas analysis according to the manufacturer's instructions.

### Oxygen dissociation curve and the concentration of ATP in erythrocytes

Oxygen dissociation curves were determined using a Hemox-Analyser (TCS Scientific Corp., USA). Total 40  $\mu\text{L}$  blood sample from each lizard was dissolved in 3.96 mL buffer solution which contained 200  $\mu\text{L}$  NaCl (3 mol/L), 40  $\mu\text{L}$  10% bovine serum albumin (BSA), 400  $\mu\text{L}$  HEPES (0.5 mol/L, pH = 7.3), 40  $\mu\text{L}$  anti-foaming agent and 3.34 mL distilled water. RBCs remained intact throughout the measurement procedure. All samples were analyzed at the temperature of 30°C. The gas mixtures used were 2% CO<sub>2</sub> in air to establish full oxygenation and pure nitrogen for deoxygenation. OECs were directly plotted by software provided with the Hemox-Analyser. P<sub>50</sub> were also obtained from this software. The concentration of ATP in erythrocytes was measured using ATP kits (Nanjing Jiancheng Bioengineering Institute, Jiangsu, China).

### RP-HPLC analysis

Samples were prepared from hemolysate with an Hb concentration of 40 g/L and were diluted further with water (75  $\mu\text{L}$  hemolysate plus 925  $\mu\text{L}$  water). Total 20  $\mu\text{L}$  of diluted samples were used for each assay. Bio-Bond C4 column (5  $\mu\text{m}$ , 250 x 4.6 mm, DIKMA) was used for RP-HPLC analysis. We eluted globin chains with a two-solvent system [solvent A, 200 mL/L acetonitrile and 3 mL/L trifluoroacetic acid (TFA) in water; solvent B, 600 mL/L acetonitrile and 3 mL/L TFA in water] and a 3-step RPLC elution program consisting of a linear gradient of 60%–100% solvent B in 80 min, a linear gradient of 100%–60% solvent B in 10 min, and re-equilibration with 60% solvent B for 10 min. The flow rate was 1 mL/min, eluate was detected at 220 nm [38] and abundance were quantified using Image J [39]. Molecular weight of globins were detected using a MaXis 4G ultra-high resolution time of flight mass spectrometer (Bruker-Daltonics).

### RNA isolation, cDNA synthesis and RACE amplification

Total RNA was extracted and purified from liver and muscle of both two species. RNA concentration and purity was assayed using the NanoDrop 2000 (Thermo Scientific, USA). The integrity of the RNA was confirmed using electrophoresis. Full length cDNA for  $\alpha$ - and  $\beta$ -globin genes was performed using a SMART RACE cDNA Amplification Kit (Clontech Laboratories) and the residue of genomic DNA was executed using Recombinant DNase I according to the manufacturer's instruction. The primers were designed based on the sequences of *Anolis carolinensis* obtained from GenBank as shown in Table 1. The PCR amplification was performed using Touchdown PCR and LA Taq polymerase. PCR products were cloned into pMD18-T Vector (Takara, Dalian, China) and sequenced (Sangon, Shanghai, China). The sequences were deposited into GenBank (Accession number: KP019961-KP019968).

### Preparation of Hb models and simulation setup

Primary structures of the  $\alpha$ - and  $\beta$ -globin polypeptides were deduced from translated DNA sequences and there are potentially three different isoforms in each species. We used MODELLER 9v12 [40] to construct Hb tetramer models in the two species using the *Homo sapiens* deoxyhemoglobin (T-state) 1BZZ as a structural template. Total six Hb models were constructed and the  $\alpha$ - and  $\beta$ -globin subunits composition in the two species as show in Table 2. Missing hydrogen atoms were added by the psfgen plugin of VMD [41]. The starting models

**Table 1. Primer sequence used for RACE amplification.**

Primers name	Primer sequence(5'-3')
Hbα-F	GCTGCGGGTGGACccngknaaytt
Hbα-R	TAACGGTAYTTGGMGTCAGCACRG
Hbβ-F	ATGGTGCACCTGGACCGCCGAAGA
Hbβ-R	TCAGTGGTACCGCGGGACAGG

doi:10.1371/journal.pone.0125751.t001

were immersed in equilibrated TIP3P water boxes. To reflect physiological salt concentrations, NaCl were added to all the six systems (0.17 mol/L NaCl in *P. przewalskii*, 0.16 mol/L NaCl in *P. erythrurus*) using the autoionize plugin of VMD. The total system sizes were almost the same as 5003 atoms with an initial simulation box of  $87 \times 77 \times 79 \text{ \AA}^3$  (hwHb1 of *P. erythrurus*).

All simulations were performed with NAMD2.9 [42] using CHARMM version c35b2 with the all-atom 27 protein force field. The intramolecular bonds involved hydrogen atoms were constrained using the SHAKE algorithm, allowing a 2 fs integration time step. The energy minimizations were performed before the equilibration runs. Waters were melted while others were fixed for 500 ps period, this was followed by 500 ps runs with protein. After they were released, the system was subjected to equilibration runs for 10.5 ns. Simulations were performed with controlling of the constant pressure temperature ( $P = 1 \text{ atm}$ ,  $T = 310 \text{ K}$ ), Periodic boundary conditions were applied, and the electrostatic interactions were calculated by the particle-mesh Ewald method. After simulations, all analysis was used VMD and corresponding Plugs within the final 4 ns.

### Statistical analyses

The data on hematology, blood gas,  $P_{50}$  and ATP concentration were test for normality and homogeneity of the variances before ANOVA. Then data were analyzed using one-way analysis of variance (ANOVA). Values presented as mean  $\pm$  SEM, statistical significance was accepted at  $P < 0.05$ .

## Results

### Hematological parameters and blood gas analysis

The experimental measures of hematological parameters under habitat conditions of both species are presented in Table 3. High-altitude *P. erythrurus* exhibits elevated RBC ( $1.12 \pm 0.04$  and  $0.94 \pm 0.04 \times 10^{12}/L$ , respectively;  $F_{1, 23} = 28.36$ ,  $p < 0.05$ ), [Hb] ( $107.92 \pm 4.32$  and  $92.48 \pm 2.88 \text{ g/L}$ , respectively;  $F_{1, 20} = 19.86$ ,  $p < 0.01$ ) and Hematocrit (HCT,  $32.82 \pm 1.10$  and  $27.70 \pm 0.47\%$ , respectively;  $F_{1, 23} = 25.25$ ,  $p < 0.001$ ) compared with the average values of low-altitude *P. przewalskii*. There was no significant variation in the average values for mean corpuscular hemoglobin concentration (MCHC,  $339.75 \pm 15.68$  and  $328.13 \pm 13.31 \text{ g/L}$ , respectively;  $F_{1, 14} = 0.319$ ,  $p > 0.05$ ), mean corpuscular volume (MCV,  $0.27 \pm 0.01$  and  $0.31 \pm 0.02 \text{ pL}$ , respectively;  $F_{1, 14} = 2.582$ ,

**Table 2. α- and β-globin subunits composition and named of the six Hb models.**

		group 1	group 2	group 3
<i>P. przewalskii</i>	Composition	(αβ1)2	(αβ2)2	(αβ3)2
	Named	hsHb1	hsHb2	hsHb3
<i>P. erythrurus</i>	Composition	(αβ1)2	(αβ2)2	(αβ3)2
	Named	hwHb1	hwHb2	hwHb3

doi:10.1371/journal.pone.0125751.t002

**Table 3. Hematological parameters of *P. erythrurus* and *P. przewalskii*.**

Determination	<i>P. przewalskii</i>	<i>P. erythrurus</i>	p-Value
RBC count × 10 <sup>12</sup> /L	0.94±0.04	1.12±0.04	0.029
Hematocrit (%)	27.70±0.47	32.82±1.10	0.010
Hemoglobin (g/L)	92.48±2.88	107.92±4.32	0.002
MCV (pL)	0.31±0.02	0.27±0.01	0.134
MCH (pg)	100.20±5.92	91.41±5.46	0.293
MCHC (g/L)	328.13±13.31	339.75±15.68	0.581

Data presented as mean ± SEM; RBC, red blood cell; MCV, mean corpuscular volume; MCH, mean cell hemoglobin; MCHC, mean corpuscular hemoglobin concentration.

doi:10.1371/journal.pone.0125751.t003

p>0.05) and mean cell hemoglobin (MCH, 91.41±5.46 and 100.20±5.92 pg, respectively; F<sub>1, 14</sub> = 1.192, p>0.05) between these two species. Blood gas analyzer was applied to the small reptile for the first time, and the results are presented in Table 4. Oxygen partial pressure (PaO<sub>2</sub>, 56.38 ±1.53 and 77.28±2.72 mmHg, respectively; F<sub>1, 22</sub> = 44.90, p<0.001), carbon dioxide partial pressure (PaCO<sub>2</sub>, 27.83±2.20 and 38.13±2.83 mmHg, respectively; F<sub>1, 22</sub> = 8.23, p<0.01), arterial oxygen saturation (SaO<sub>2</sub>, 77.47±1.11 and 83.96±1.40%, respectively; F<sub>1, 22</sub> = 13.16, p<0.01), [HCO<sub>3</sub><sup>-</sup>] (14.30±1.07 and 18.86±1.16 mmol/L, respectively; F<sub>1, 20</sub> = 5.50, p<0.05), [Na<sup>+</sup>] (159.09 ±1.14 and 169.38±1.57 mmol/L, respectively; F<sub>1, 20</sub> = 28.03, p < 0.001) and [Cl<sup>-</sup>] (117.57±1.11 and 123.44±1.02 mmol/L, respectively; F<sub>1, 20</sub> = 15.17, p<0.01) in arterial blood of *P. erythrurus* were significantly lower than that in *P. przewalskii*, while pH (7.33±0.03 and 7.31±0.03, respectively; F<sub>1, 22</sub> = 0.17, p>0.05) and [K<sup>+</sup>] (4.00±0.15 and 4.39±0.29 mmol/L, respectively; F<sub>1, 22</sub> = 1.33, p>0.05) have no significant variation between the species.

### Whole blood Oxygen affinity and the concentration of ATP in erythrocytes


Oxygen equilibrium measurements of the whole blood showed that *P. erythrurus* (green line) exhibits a higher O<sub>2</sub> affinity compared with *P. przewalskii* (blue line) under 30°C and pH 7.3 (Fig 1). There were significant differences in P<sub>50</sub> (51.97±2.64 and 71.27±1.49 mmHg, respectively; F<sub>1, 20</sub> = 49.66, p<0.001) and ATP concentration (200.52 and 91.33 μmol/gHb, respectively; F<sub>1, 20</sub> = 50.68, p<0.05) between these two species, and ATP concentration exhibits two-

**Table 4. Arterial blood gas measurements of *P. erythrurus* and *P. przewalskii*.**

Determination	<i>P. przewalskii</i>	<i>P. erythrurus</i>	p-Value
pH	7.31±0.03	7.33±0.03	0.670
PaCO <sub>2</sub> (mmHg)	38.13±2.83	27.83±2.20	0.009
PaO <sub>2</sub> (mmHg)	77.28±2.72	56.38±1.53	0.000
HCO <sub>3</sub> <sup>-</sup> (mmHg)	18.86±1.16	14.30±1.07	0.028
Na <sup>+</sup> (mmol/L)	169.38±1.57	159.09±1.14	0.000
K <sup>+</sup> (mmol/L)	4.39±0.29	4.00±0.15	0.262
Cl <sup>-</sup> (mmol/L)	123.44±1.02	117.57±1.11	0.001
SaO <sub>2</sub> (%)	83.96±1.40	77.47±1.11	0.001

Data presented as mean ± SEM; PaO<sub>2</sub>, arterial blood oxygen partial pressure; PaCO<sub>2</sub>, arterial blood carbon dioxide partial pressure; SaO<sub>2</sub>, arterial blood oxygen saturation.

doi:10.1371/journal.pone.0125751.t004



**Fig 1. Whole blood oxygen dissociation curves of *P. erythrurus* (green line) and *P. przewalskii* (blue line) at 30°C and pH 7.3.**

doi:10.1371/journal.pone.0125751.g001

fold correlation between them (Table 5). However, we did not detect any difference in oxygen affinity between the sexes in both species.

### RP-HPLC and amino acid sequence analysis

RP-HPLC analysis of hemolysate showed approximately equal amounts of globin peaks in these two species (Fig 2). There are four major peaks A1, A2, A3 and A4 (molecular weight, 15905.2001, 16269.3200, 16063.3959 and 15633.1674 Da, respectively) with a roughly abundance ratio 1.05: 1.04: 1.29: 1.11 in *P. erythrurus*. The four major peaks were also detected including B1, B2, B3 and B4 (molecular weight, 15916.0634, 16288.1647, 15633.0142 and 16061.2838 Da, respectively) with a abundance ratio 0.74: 0.87: 0.99: 1.05 in *P. przewalskii*. Intriguingly, a low-abundance peak (molecular weight, 15909.0636; relative abundance, 0.17) between B1 and B2 was detected in all analysis of *P. przewalskii*.


One  $\alpha$ -globin gene and three  $\beta$ -globin genes were cloned and sequenced in both species. The  $\alpha$ - and  $\beta$ -globin polypeptides were deduced from translated DNA sequences. Alignment amino acid sequences of  $\alpha$ - and  $\beta$ -like globin chains from these two lizard species and five

**Table 5. ATP concentration and  $P_{50}$  of *P. erythrurus* and *P. przewalskii*.**

Determination	<i>P. przewalskii</i>	<i>P. erythrurus</i>	p-Value
RBC [ATP] ( $\mu\text{mol/gHb}$ )	91.33 $\pm$ 9.11	200.52 $\pm$ 28.97	0.040
$P_{50}$ (mmHg)	71.27 $\pm$ 1.49	51.97 $\pm$ 2.64	0.000

Data presented as mean  $\pm$  SEM.

doi:10.1371/journal.pone.0125751.t005



**Fig 2. RP-HPLC chromatograms for erythrocyte hemolysates of *P. erythrurus* (upward profile) and *P. przewalskii* (downward profile).** Four globin chain peaks of *P. erythrurus* (A1–A4) and *P. przewalskii* (B1–B4) were eluted from the C4 RP-HPLC column, and corresponding molecular weight was shown in a red box.

doi:10.1371/journal.pone.0125751.g002


outgroup taxa: human (*Homo sapiens*), chicken (*Gallus gallus*), anole lizard (*Anolis carolinensis*), red-eared slider (*Trachemys scripta*) and painted turtle (*Chrysemys picta*) were shown in Fig 3. Compared with these five taxa, total of 23 and 27 peculiar sites (Alpha and Beta, respectively) were found in these two lizard species. Furthermore, the  $\alpha$ -globin chains of these two species are distinguished only by one amino acid in position of 121 (Ile-Val). For the multiple alignment, a total of 33 site differences were discovered. Then we detected varying degrees of amino acid sequence difference in  $\beta$ 1-,  $\beta$ 2- and  $\beta$ 3-globin chains between these two species (the number of amino acid substitutions are 1, 5 and 24 respectively). The results showed that the  $\beta$ 1-globin chain has the highest sequence identity with only one substitution ( $\beta$ 12Thr-Ser). Identical substitutions of  $\beta$ -globin chains in these two species occur in position  $\beta$ 10,  $\beta$ 14,  $\beta$ 18,  $\beta$ 29,  $\beta$ 32,  $\beta$ 34,  $\beta$ 43 and  $\beta$ 142. There was one particular substitution at site  $\beta$ 22 (Thr-Val) in *P. przewalskii*, meanwhile five particular substitutions at sites  $\beta$ 12 (Thr-Ser),  $\beta$ 13 (Asn-Gly VS Asn-Ser in *P. przewalskii*),  $\beta$ 20 (Val-Leu),  $\beta$ 21 (Pro-Ser VS Pro-Gly in *P. przewalskii*),  $\beta$ 23 (Ile-Val) were found in *P. erythrurus*.

### Structural and dynamical analysis of the six Hb models


We employed equilibrium MD simulations on the six possible  $\alpha_2\beta_2$  tetrameric (hsHb1, hsHb2, hsHb3, hwHb1, hwHb2 and hwHb3) to observe the stability of T-state isoHbs. It was obvious that there are no large structural fluctuations during all the six simulations, as evidence in the time evolution of the backbone root mean square deviation (RMSD) shown in Fig 4. All the six models reach equilibrium within the first 5 ns of the simulations, and average backbone RMSD values were approximately 1.5 Å, so we analyzed the intersubunit contacts in the final 4 ns of the trajectories. Hydrogen bonds and salt bridges at  $\alpha$ 1 $\beta$ 2 and  $\alpha$ 2 $\beta$ 1 interfaces of the six Hb models are shown in Table 6. Total of 24, 28 and 32 hydrogen bonds were found in hsHb1, hsHb2 and hsHb3 of *P. przewalskii* respectively, and hydrogen bonds in *P. erythrurus* (19, 23, 27 in hwHb1, hwHb2, hwHb3, respectively) were significantly less than that in *P. przewalskii*.



Alpha



Beta




**Fig 3. Alignment amino acid sequence of  $\alpha$ - and  $\beta$ -like globin chains from the two lizard species and five outgroup taxa: human (*Homo sapiens*), chicken (*Gallus gallus*), anole lizard (*Anolis carolinensis*), red-eared slider (*Trachemys scripta*) and painted turtle (*Chrysemys picta*).** The  $\alpha$ - and  $\beta$ -globin polypeptides of both species were deduced from translated DNA sequences, peculiar amino acid of the two species were marked with light gray boxes.

doi:10.1371/journal.pone.0125751.g003

We found one salt bridge between  $\alpha 94$ Asp and  $\beta 40$ Arg in all the six Hb models except in hsHb3 which formed an additional salt bridge between  $\alpha 40$ Lys and  $\beta 94$ Asp at  $\alpha 2\beta 1$  interfaces. In group 1, there are 11 hydrogen bonds at  $\alpha 1\beta 2$  interface in both models, simultaneously, the lost 7 hydrogen bonds (Fig 5) and reformed 2 hydrogen bonds at  $\alpha 2\beta 1$  interface were found in hwHb1 of *P. erythrurus* compared with hsHb1 of *P. przewalskii* (Fig 6). Similar results were also found in group 2 and group 3 with the lost 5 hydrogen bonds in isoHb models of *P. erythrurus* compared with *P. przewalskii* (data not shown).

Discussion

Matching  $O_2$  supply with  $O_2$  demand has always been a hot topic in the studies of high altitude adaptation. The highest living lizard *P. erythrurus* has to manage an unremitting hypobaric hypoxia (atmospheric pressure, 587.79 hPa;  $PO_2$ , ~92 mmHg) and cold temperatures while *P.*



**Fig 4. Evolution of root mean square deviation (RMSD) of the atoms in the backbone of the proteins over time from the initial structure for: *P. przewalskii* (hsHb1, hsHb2, hsHb3) and *P. erythrurus* (hwHb1, hwHb2, hwHb3).**

doi:10.1371/journal.pone.0125751.g004


*przewalskii* lives in a relatively mild environment (atmospheric pressure, 863.73 hPa; PO<sub>2</sub>, ~136 mmHg) [35]. The present study offers a snap-shot of hematological characteristics in these two lizard species dwelling at different altitudes for the first time. Our results indicated that *P. erythrurus* has an efficient oxygen transport system by regulating several steps in the O<sub>2</sub> cascade.

When lowland natives ascend to high altitude, many of them can compensate for a reduced O<sub>2</sub> supply by increasing their Hct, [Hb] and RBC. However, an excessive increased Hct will increase blood viscosity and add more budgets for heart, pulmonary and blood circulation system [15,43]. In humans, the available evidence indicates that the optimal Hb concentration at high altitude should be maintained at the typical sea level value and the hypoxia-induced polycythemia is a maladaptive plasticity [1,44]. A moderate increased Hct and [Hb] can be propitious to increase blood O<sub>2</sub>-carrying capacity and to improve tissue oxygenation which must be closer to the optimal values. Comparing our data to available hematologic values in mammals and birds, we found that [Hb] and Hct of both lizards were slightly lower than values reported previously. The optimal Hct for O<sub>2</sub> transport was 40% in dogs [45,46]. [Hb] and hematocrit in imprisoned bar-headed goose (*Anser indicus*) were 17.1±1.24 g/dL and 43.3±3.9%, respectively [47]. In addition, hematological observations have been reported in several reptiles from sea level to 3350 m including RBC (range from 0.955 to 1.37 × 10<sup>12</sup>/L), Hct (25 to 39%) and [Hb] (67 to 114 g/L)

**Table 6. Hydrogen bonds and salt bridges at α1β2 and α2β1 interfaces of the six Hb models.**

		hsHb1	hsHb2	hsHb3	hwHb1	hwHb2	hwHb3
Hydrogen bonds	α1β2	11	18	19	11	14	15
	α2β1	13	10	13	8	9	11
	total	24	28	32	19	23	27
Salt bridges	α1β2	1	1	1	0	0	0
	α2β1	0	0	1	1	1	1
	total	1	1	2	1	1	1

doi:10.1371/journal.pone.0125751.t006




**Fig 5. The loss of 7 hydrogen bonds at  $\alpha 2\beta 1$  interface in hwHb1.** (a1–a5) Hydrogen bonds at  $\alpha 2\beta 1$  interface present in hshb1 and lost in hwHb1, (a) Three-dimensional structure of hshb1 with  $\beta 12$ Met.

doi:10.1371/journal.pone.0125751.g005

[48–51]. The values of both species in this study intervene between the minimum and maximum values of reported reptiles. Meanwhile, Hct of *P. erythrurus* is very close to the calculated optimal values for balancing  $O_2$  carrying capacity and blood viscosity from the lizard *Dipsosaurus dorsalis*, which break the bonds of convention in most small lizards (body mass < 8 g, Hct < 30%) [52]. Unlike the hypoxia-induced maladaptive polycythemia, the elevation of RBC in *P. erythrurus* could promote oxygen carrying capacity without disadvantage of high viscosity. In addition, our previous study indicated a closely related species *P. vlinalgalii* can increase its oxygen carrying capacity in hypoxic acclimatization and adaptation [53]. When acclimatized to environmental hypoxia low-altitude *P. vlinalgalii* exhibited unchanged RBC and elevated Hct and [Hb], MCV and MCHC and similar result was obtained when comparing these parameters in *P. vlinalgalii* living at different altitudes. Our results showed a more propitious Hct in *P. erythrurus* compared to high-altitude *P. vlinalgalii* and a different strategy for *P. erythrurus* to increase oxygen transport efficiency by increasing RBC rather than increasing the volume of red blood cell.

The  $PaO_2$  largely mirror the effectiveness of ventilation and pulmonary diffusion with hypoxia [1,6]. The ambient oxygen partial pressure descend from 136 to 92 mmHg from 1500 m to 4500 m altitude while the  $PaO_2$  of *P. przewalskii* and *P. erythrurus* descend from about 77 to 56 mmHg. Furthermore, a lower  $PaO_2$  in *P. erythrurus* does not caused the secondary alkalosis by accelerated breathing. This result suggest that *P. erythrurus* may have been evolved an efficient pulmonary system for  $O_2$  loading during the prolonged hypoxia. The lower  $PaCO_2$  of *P. erythrurus* may be due to the suppressed aerobic metabolism [35]. The blunted hypoxic ventilatory response in *P. erythrurus* might help to reduce the oxygen cost of breathing and respiratory water loss [6].



**Fig 6. The reformed of 2 hydrogen bonds at  $\alpha 2\beta 1$  interface in hwHb1.** (b1–b2) Hydrogen bonds at  $\alpha 2\beta 1$  interface present in hwHb1 and lost in hsHb1. (b) Three-dimensional structure of hwHb1 with  $\beta 12$ Ser.

doi:10.1371/journal.pone.0125751.g006

Fine-tuned adjustments in blood- $O_2$  affinity play an important role in matching  $O_2$  supply and  $O_2$  demand under high altitude hypoxia. Our result indicated that *P. erythrurus* has an elevated blood- $O_2$  affinity compared with low-altitude *P. przewalskii*. This may be achieved by changes in intrinsic Hb- $O_2$  affinity, the sensitivity of Hb to allosteric cofactors and the concentration of allosteric cofactors. Firstly, Our results demonstrated multiple substitutions of amino acid in Hb. Certain residues from human Hb have been demonstrated for proton binding ( $\alpha 1$ Val,  $\alpha 122$ His,  $\beta 2$ His,  $\beta 82$ Lys,  $\beta 143$ His, and  $\beta 146$ His), chloride ions binding ( $\alpha 1$ Val and  $\alpha 131$ Ser and one  $\beta 1$ Val and  $\beta 82$ Lys) and  $CO_2$  binding (N-terminal  $NH_3^+$  residues) [54–56]. However, checking these three potential binding sites, we did not found any substitutions between two lizard species. Besides, ATP binding site has been described in Hbs of fish including  $\beta 1$ Val,  $\beta 2$ Glu,  $\beta 82$ Lys and  $\beta 143$ Arg [57]. We found that  $\beta$ -globin of both lizards contains His at  $\beta 2$ , but this change may not alter the responsiveness to ATP based on the evidence reported in red-eared slider [27]. Consequently, the elevated blood- $O_2$  affinity in *P. erythrurus* could not be caused by the change in sensitivity of Hb to allosteric cofactors. Secondly, the concentration of ATP in erythrocytes in *P. erythrurus* is over twice than that in *P. przewalskii*. Conversely,  $[Cl^-]$ ,  $[HCO_3^-]$  and Pa $CO_2$  in blood of *P. erythrurus* were significantly lower. However, no significant variation of pH was found between these two species. These results suggest that the elevated blood- $O_2$  affinity in *P. erythrurus* may be attributable to balancing the independent effects of these potential heterotropic ligands under the prevailing conditions. Finally, amino acid substitutions that located at  $\alpha 1\beta 2$  and  $\alpha 2\beta 1$  interfaces of the isoHbs may be critical for controlling Hb- $O_2$  affinity by impact the transformation process from the T-state to the R-state during oxygenation of hemoglobin [18,58,59]. Our results suggest that

isoHbs of *P. erythrurus* may have higher intrinsic Hb-O<sub>2</sub> affinity compared with *P. przewalskii* which may due to the eliminated hydrogen bonds at  $\alpha 1\beta 2$  and  $\alpha 2\beta 1$  interfaces. Structural analysis shows that 2 of these 33 substitutions occurred at  $\alpha 1\beta 2$  or  $\alpha 2\beta 1$  interfaces including  $\beta 34\text{Val-Thr}$  (nonpolar-polar) and  $\beta 101\text{Val-Glu}$  (nonpolar-polar). These substitutions are conducive to form hydrogen bonds with  $\alpha 141\text{Arg}$  and  $\alpha 41\text{Thr}$ . The specific substitutions in position of  $\beta 13(\text{A9})\text{Gly-Ser}$  was also reported in Andean hummingbirds which increased O<sub>2</sub>-affinity in the presence of  $\beta 83\text{Gly}$  and reduced O<sub>2</sub>-affinity in the presence of  $\beta 83\text{Ser}$  (epistasis for Hb-O<sub>2</sub> affinity) [20]. The position  $\beta 83$  of both *Phrynocephalus* lizards was occupied by Gln and the polarity of Gln is obviously closer to Ser. Therefore, this substitution may typically reduce O<sub>2</sub>-affinity of isoHb in *P. przewalskii*. In addition, the similar substitutions in position of  $\beta 142$  have been verified leading to increase in oxygen affinity [60]. All of these examples suggest a higher intrinsic O<sub>2</sub> affinity of isoHbs in *P. erythrurus*. Hence, elevated blood-O<sub>2</sub> affinity in *P. erythrurus* may mainly due to the higher intrinsic Hb-O<sub>2</sub> affinity and concentration-dependent adjustment of allosteric cofactors.

As observed in many birds and nonavian reptiles, the phenomenon of co-express different isoHbs was also found in these two lizard species [20,27–29]. Although the function of each isoHb has not been confirmed, we can speculate that functionally distinct isoHbs exist in these two species. From number of hydrogen bonds in the six models, we can predict the oxygen affinity of isoHbs as follows:  $\text{hwHb1} > \text{hwHb2} > \text{hsHb1} > \text{hwHb3} > \text{hsHb2} > \text{hsHb3}$  (*P. erythrurus* > *P. przewalskii*). A potential mechanism for matching O<sub>2</sub> supply with O<sub>2</sub> demand in *P. erythrurus* could be provided by changes in intra-erythrocytic isoHbs stoichiometry [61–64]. All experiments of MD simulations in this study are based on identified one  $\alpha$ -globin gene and three  $\beta$ -globin genes by RACE-PCR. There may be other homologous globin genes failed to be detected due to scarcity of available sequence in lizard species and increased sequence divergence in two distinct paralogs [25]. In sum, a variety of factors may lead to change of Hb-O<sub>2</sub> affinity, future detailed studies on the relationship between structure and function of isoHbs in these two lizards may reveal novel molecular mechanisms of high altitude adaptation.

## Conclusion

As the highest living lizards in the world, *P. erythrurus* may have evolved an efficient oxygen transport system under an unremitting hypobaric hypoxia. It increases oxygen carrying capacity by increasing RBC and this could promote oxygen carrying capacity without disadvantage of high viscosity. The elevated blood-O<sub>2</sub> affinity in *P. erythrurus* may be achieved by increasing in intrinsic O<sub>2</sub> affinity of isoHbs and balancing the independent effects of potential heterotropic ligands.

## Acknowledgments

The authors thank Xubin Li for field assistance. For technical advice and use of equipment under their care the authors thank Kunping Yan, Feiyun Gao, Zhihui Zhang and Shouliang Dong.

## Author Contributions

Conceived and designed the experiments: SL QC. Performed the experiments: SL YX FY HW YB. Analyzed the data: SL YX. Contributed reagents/materials/analysis tools: SL XT HW YB YN QC. Wrote the paper: SL QC.

## References

1. Storz JF, Scott GR, Cheviron ZA. Phenotypic plasticity and genetic adaptation to high-altitude hypoxia in vertebrates. *The Journal of experimental biology*. 2010; 213(24):4125–36. doi: [10.1242/jeb.048181](https://doi.org/10.1242/jeb.048181) PMID: [21112992](https://pubmed.ncbi.nlm.nih.gov/21112992/)
2. Monge C, Leon-Velarde F. Physiological adaptation to high altitude: oxygen transport in mammals and birds. *Physiological Reviews*. 1991; 71(4):1135–72. PMID: [1924550](https://pubmed.ncbi.nlm.nih.gov/1924550/)
3. Ramirez J-M, Folkow LP, Blix AS. Hypoxia tolerance in mammals and birds: from the wilderness to the clinic. *Annu Rev Physiol*. 2007; 69:113–43. PMID: [17037981](https://pubmed.ncbi.nlm.nih.gov/17037981/)
4. Scott G, Milsom W. Control of Breathing in Birds: Implications for High-Altitude Flight. *Cardio-Respiratory Control in Vertebrates*: Springer; 2009. p. 429–48.
5. Powell F, Milsom W, Mitchell G. Time domains of the hypoxic ventilatory response. *Respiration physiology*. 1998; 112(2):123–34. PMID: [9716296](https://pubmed.ncbi.nlm.nih.gov/9716296/)
6. Powell FL. The influence of chronic hypoxia upon chemoreception. *Respiratory physiology & neurobiology*. 2007; 157(1):154–61.
7. Brutsaert TD. Population genetic aspects and phenotypic plasticity of ventilatory responses in high altitude natives. *Respiratory physiology & neurobiology*. 2007; 158(2):151–60.
8. Hsia CC, Carbayo JJP, Yan X, Bellotto DJ. Enhanced alveolar growth and remodeling in guinea pigs raised at high altitude. *Respiratory physiology & neurobiology*. 2005; 147(1):105–15.
9. Hsia CC, Johnson RL, McDonough P, Dane DM, Hurst MD, Fehmel JL, et al. Residence at 3,800-m altitude for 5 mo in growing dogs enhances lung diffusing capacity for oxygen that persists at least 2.5 years. *Journal of Applied Physiology*. 2007; 102(4):1448–55. PMID: [17218427](https://pubmed.ncbi.nlm.nih.gov/17218427/)
10. Ravikumar P, Bellotto DJ, Johnson RL, Hsia CC. Permanent alveolar remodeling in canine lung induced by high-altitude residence during maturation. *Journal of Applied Physiology*. 2009; 107(6):1911–7. doi: [10.1152/jappphysiol.00552.2009](https://doi.org/10.1152/jappphysiol.00552.2009) PMID: [19833809](https://pubmed.ncbi.nlm.nih.gov/19833809/)
11. Bouverot P. *Adaptation to altitude-hypoxia in vertebrates*: Springer; 1985.
12. Ekblom B, Hermansen L. Cardiac output in athletes. *Journal of Applied Physiology*. 1968; 25(5):619–25. PMID: [4879852](https://pubmed.ncbi.nlm.nih.gov/4879852/)
13. Kanstrup I, Ekblom B. Blood volume and hemoglobin concentration as determinants of maximal aerobic power. *Medicine and science in sports and exercise*. 1984; 16(3):256–62. PMID: [6748923](https://pubmed.ncbi.nlm.nih.gov/6748923/)
14. Ekblom B, Berglund B. Effect of erythropoietin administration on mammal aerobic power. *Scandinavian Journal of Medicine & Science in Sports*. 1991; 1(2):88–93. doi: [10.1136/bjsports-2015-094781](https://doi.org/10.1136/bjsports-2015-094781) PMID: [25878072](https://pubmed.ncbi.nlm.nih.gov/25878072/)
15. Guyton AC, Richardson TQ. Effect of hematocrit on venous return. *Circulation research*. 1961; 9(1):157–64.
16. Weber RE, Fago A. Functional adaptation and its molecular basis in vertebrate hemoglobins, neuroglobins and cytoglobins. *Respiratory physiology & neurobiology*. 2004; 144(2):141–59.
17. Storz JF. Hemoglobin function and physiological adaptation to hypoxia in high-altitude mammals. *Journal of Mammalogy*. 2007; 88(1):24–31.
18. Storz JF, Moriyama H. Mechanisms of hemoglobin adaptation to high altitude hypoxia. *High altitude medicine & biology*. 2008; 9(2):148–57.
19. Weber RE. High-altitude adaptations in vertebrate hemoglobins. *Respiratory physiology & neurobiology*. 2007; 158(2):132–42.
20. Projecto-Garcia J, Natarajan C, Moriyama H, Weber RE, Fago A, Cheviron ZA, et al. Repeated elevational transitions in hemoglobin function during the evolution of Andean hummingbirds. *Proceedings of the National Academy of Sciences*. 2013; 110(51):20669–74. doi: [10.1073/pnas.1315456110](https://doi.org/10.1073/pnas.1315456110) PMID: [24297909](https://pubmed.ncbi.nlm.nih.gov/24297909/)
21. Tufts DM, Natarajan C, Revsbech IG, Projecto-Garcia J, Hoffmann FG, Weber RE, et al. Epistasis constrains mutational pathways of hemoglobin adaptation in high-altitude pikas. *Molecular biology and evolution*. 2014:msu311.
22. Revsbech IG, Tufts DM, Projecto-Garcia J, Moriyama H, Weber RE, Storz JF, et al. Hemoglobin function and allosteric regulation in semi-fossorial rodents (family Sciuridae) with different altitudinal ranges. *The Journal of experimental biology*. 2013; 216(22):4264–71. doi: [10.1242/jeb.091397](https://doi.org/10.1242/jeb.091397) PMID: [24172889](https://pubmed.ncbi.nlm.nih.gov/24172889/)
23. Nagel R, Steinberg M. Hemoglobins of the embryo and fetus and minor hemoglobins of adults. *Disorders of Hemoglobin: Genetics, Pathophysiology, and Clinical Management* Cambridge Univ Press, Cambridge. 2001:197–230. PMID: [12779271](https://pubmed.ncbi.nlm.nih.gov/12779271/)

24. Weber RE, Ostojic H, Fago A, Dewilde S, Van Hauwaert M-L, Moens L, et al. Novel mechanism for high-altitude adaptation in hemoglobin of the Andean frog *Telmatobius peruvianus*. *American Journal of Physiology-Regulatory, Integrative and Comparative Physiology*. 2002; 283(5):R1052–R60. PMID: [12376398](#)
25. Hoffmann FG, Storz JF, Gorr TA, Opazo JC. Lineage-specific patterns of functional diversification in the  $\alpha$ - and  $\beta$ -globin gene families of tetrapod vertebrates. *Molecular biology and evolution*. 2010; 27(5):1126–38. doi: [10.1093/molbev/msp325](#) PMID: [20047955](#)
26. Storz JF, Opazo JC, Hoffmann FG. Gene duplication, genome duplication, and the functional diversification of vertebrate globins. *Molecular phylogenetics and evolution*. 2013; 66(2):469–78. doi: [10.1016/j.ympev.2012.07.013](#) PMID: [22846683](#)
27. Damsgaard C, Storz JF, Hoffmann FG, Fago A. Hemoglobin isoform differentiation and allosteric regulation of oxygen binding in the turtle, *Trachemys scripta*. *American Journal of Physiology-Regulatory, Integrative and Comparative Physiology*. 2013; 305(8):R961–R7. doi: [10.1152/ajpregu.00284.2013](#) PMID: [23986362](#)
28. Weber RE, Fago A, Malte H, Storz JF, Gorr TA. Lack of conventional oxygen-linked proton and anion binding sites does not impair allosteric regulation of oxygen binding in dwarf caiman hemoglobin. *American Journal of Physiology-Regulatory, Integrative and Comparative Physiology*. 2013; 305(3):R300–R12. doi: [10.1152/ajpregu.00014.2013](#) PMID: [23720132](#)
29. Storz JF, Hoffmann FG, Opazo JC, Sanger TJ, Moriyama H. Developmental regulation of hemoglobin synthesis in the green anole lizard *Anolis carolinensis*. *The Journal of experimental biology*. 2011; 214(4):575–81. doi: [10.1242/jeb.050443](#) PMID: [21270305](#)
30. Zhao E-m, Zhao K-t, Zhou K-y. *Fauna Sinica, Reptilia Vol. 2: Squamata, Lacertilia*. Beijing, Science Press: i–xi. 1999; 394:1–8.
31. Jin YT, Brown RP, Liu NF. Cladogenesis and phylogeography of the lizard *Phrynocephalus vlangalii* (Agamidae) on the Tibetan plateau. *Molecular Ecology*. 2008; 17(8):1971–82. doi: [10.1111/j.1365-294X.2008.03721.x](#) PMID: [18363665](#)
32. Jin Y. *Evolutionary studies of Phrynocephalus (Agamidae) on the Qinghai—Xizang (Tibetan) Plateau*. Ph. D. Thesis. Lanzhou University, Lanzhou (in Chinese with English abstract); 2008.
33. Jin Y, Li J, Liu N. Elevation-related variation in life history traits among *Phrynocephalus* lineages on the Tibetan Plateau: do they follow typical squamate ecogeographic patterns? *Journal of Zoology*. 2013; 290(4):293–301.
34. Jin Y-T, Liu N-F. Phylogeography of *Phrynocephalus erythrurus* from the Qiangtang Plateau of the Tibetan Plateau. *Molecular phylogenetics and evolution*. 2010; 54(3):933–40. doi: [10.1016/j.ympev.2009.11.003](#) PMID: [19900565](#)
35. Tang X, Xin Y, Wang H, Li W, Zhang Y, Liang S, et al. Metabolic Characteristics and Response to High Altitude in *Phrynocephalus erythrurus* (Lacertilia: Agamidae), a Lizard Dwell at Altitudes Higher Than Any Other Living Lizards in the World. *PloS ONE*. 2013; 8(8):e71976. doi: [10.1371/journal.pone.0071976](#) PMID: [23951275](#)
36. Guest GM, Siler VE. A centrifuge method for the determination of the volume of cells in blood. *The Journal of Laboratory and Clinical Medicine*. 1934; 19(7):757–68.
37. Freckmann G, Schmid C, Baumstark A, Pleus S, Link M, Haug C. Partial pressure of oxygen in capillary blood samples from the fingertip. *Journal of diabetes science and technology*. 2013; 7(6):1648. PMID: [24351193](#)
38. Zanella-Cleon I, Becchi M, Lacan P, Giordano PC, Wajcman H, Francina A. Detection of a thalassemic  $\alpha$ -chain variant (Hemoglobin Groene Hart) by reversed-phase liquid chromatography. *Clinical chemistry*. 2008; 54(6):1053–9. doi: [10.1373/clinchem.2007.097857](#) PMID: [18420733](#)
39. Abràmoff MD, Magalhães PJ, Ram SJ. Image processing with ImageJ. *Biophotonics international*. 2004; 11(7):36–43.
40. Šali A, Blundell TL. Comparative protein modelling by satisfaction of spatial restraints. *Journal of molecular biology*. 1993; 234(3):779–815. PMID: [8254673](#)
41. Humphrey W, Dalke A, Schulten K. VMD: visual molecular dynamics. *Journal of molecular graphics*. 1996; 14(1):33–8. PMID: [8744570](#)
42. Phillips JC, Braun R, Wang W, Gumbart J, Tajkhorshid E, Villa E, et al. Scalable molecular dynamics with NAMD. *Journal of computational chemistry*. 2005; 26(16):1781–802. PMID: [16222654](#)
43. Connes P, Yalcin O, Baskurt O, Brun J-F, Hardeman M. In health and in a normoxic environment,  $VO_2$  max is/is not limited primarily by cardiac output and locomotor muscle blood flow. *Journal of Applied Physiology*. 2006; 100(6):2099–. PMID: [16714417](#)

44. Villafuerte FC, Cardenas R, Monge-C C. Optimal hemoglobin concentration and high altitude: a theoretical approach for Andean men at rest. *Journal of Applied Physiology*. 2004; 96(5):1581–8. PMID: [14672972](#)
45. Crowell JW, Ford RG, Lewis VM. Oxygen transport in hemorrhagic shock as a function of the hematocrit ratio. *American Journal of Physiology—Legacy Content*. 1959; 196(5):1033–8.
46. Crowell J, Smith E. Determinant of the optimal hematocrit. *Journal of Applied Physiology*. 1967; 22(3):501–4. PMID: [6020234](#)
47. Meir JU, Milsom WK. High thermal sensitivity of blood enhances oxygen delivery in the high-flying bar-headed goose. *The Journal of experimental biology*. 2013; 216(12):2172–5. doi: [10.1242/jeb.085282](#) PMID: [23470665](#)
48. Weathers WW, White FN. Hematological observations on populations of the lizard *Sceloporus occidentalis* from sea level and altitude. *Herpetologica*. 1972:172–5.
49. Troiano JC, Gould EG, Gould I. Hematological reference intervals in argentine lizard *Tupinambis merrianae* (Sauria—Teiidae). *Comparative Clinical Pathology*. 2008; 17(2):93–7.
50. Espinosa-Avilés D, Salomón-Soto VM, Morales-Martínez S. Hematology, blood chemistry, and bacteriology of the free-ranging Mexican beaded lizard (*Heloderma horridum*). *Journal of Zoo and Wildlife Medicine*. 2008; 39(1):21–7. PMID: [18432093](#)
51. Marks SK, Citino SB. Hematology and serum chemistry of the radiated tortoise (*Testudo radiata*). *Journal of Zoo and Wildlife Medicine*. 1990:342–4.
52. Snyder GK. Influence of temperature and hematocrit on blood viscosity. *American Journal of Physiology—Legacy Content*. 1971; 220(6):1667–72.
53. He J, Xiu M, Tang X, Yue F, Wang N, Yang S, et al. The Different Mechanisms of Hypoxic Acclimatization and Adaptation in Lizard *Phrynocephalus vlangalii* Living on Qinghai-Tibet Plateau. *Journal of Experimental Zoology Part A: Ecological Genetics and Physiology*. 2013; 319(3):117–23.
54. Perutz M, Muirhead H, Mazzarella L, Crowther R, Greer J, Kilmartin J. Identification of residues responsible for the alkaline Bohr effect in haemoglobin. *Nature*. 1969; 222:1240–3. PMID: [5789657](#)
55. Lukin JA, Ho C. The structure-function relationship of hemoglobin in solution at atomic resolution. *Chemical reviews*. 2004; 104(3):1219–30. PMID: [15008621](#)
56. Riggs AF. The Bohr effect. *Annual review of physiology*. 1988; 50(1):181–204.
57. Perutz M. Stereochemistry of cooperative effects in fish and amphibian haemoglobins. *Nature*. 1982; 299:421–6. PMID: [7121579](#)
58. Pettigrew DW, Romeo PH, Tsapis A, Thillet J, Smith ML, Turner BW, et al. Probing the energetics of proteins through structural perturbation: sites of regulatory energy in human hemoglobin. *Proceedings of the National Academy of Sciences*. 1982; 79(6):1849–53. PMID: [6952235](#)
59. Dickerson RE, Geis I. Hemoglobin: structure, function, evolution, and pathology: Benjamin-Cummings Publishing Company; 1983.
60. Hirano M, Ohba Y, Imai K, Ino T, Morishita Y, Matsui T, et al. Hb Toyoake: beta 142 (H20) Ala replaced by Pro. A new unstable hemoglobin with high oxygen affinity. *Blood*. 1981; 57(4):697–704. PMID: [7470620](#)
61. Grispo MT, Natarajan C, Projecto-Garcia J, Moriyama H, Weber RE, Storz JF. Gene duplication and the evolution of hemoglobin isoform differentiation in birds. *Journal of Biological Chemistry*. 2012; 287(45):37647–58. doi: [10.1074/jbc.M112.375600](#) PMID: [22962007](#)
62. Hiebl I, Weber RE, Schneeganss D, Kösters J, Braunitzer G. High-Altitude Respiration of Birds. Structural Adaptations in the Major and Minor Hemoglobin Components of adult Rüppell's Griffon (*Gyps rueppellii*, Aegyptiinae): a New Molecular Pattern for Hypoxic Tolerance. *Biological chemistry Hoppe-Seyler*. 1988; 369(1):217–32.
63. Weber RE, Hiebl I, Braunitzer G. High altitude and hemoglobin function in the vultures *Gyps rueppellii* and *Aegypius monachus*. *Biological chemistry Hoppe-Seyler*. 1988; 369(1):233–40.
64. Lutz PL. On the oxygen affinity of bird blood. *American Zoologist*. 1980; 20(1):187–98.