

Different chronological patterns of appearance of blood derived milk components during mastitis indicate different mechanisms of transfer from blood into milk

Olga Wellnitz¹, Christina Zbinden², Johannes Lüttgenau¹, Heinrich Bollwein¹ and Rupert M Bruckmaier^{2*}

¹Vetsuisse Faculty University of Zurich, Clinic of Reproductive Medicine, Switzerland

²Vetsuisse Faculty University of Bern, Veterinary Physiology, Switzerland

Received 15 November 2014; accepted for publication 22 May 2015; first published online 3 July 2015

This study aimed to describe chronological patterns of changes of various candidate blood components in milk during the acute phase of a mammary immune response in detail. Eight dairy cows were challenged with *Escherichia coli* lipopolysaccharide in one udder quarter. Milk from challenged and control quarters and blood samples were taken before, and 1 and 2 h after challenge and then every 15 min until 5 h after challenge. The SCC, serum albumin, immunoglobulin (Ig)G₁, IgG₂, lactate dehydrogenase (LDH), and L-lactate in milk and blood, and α -lactalbumin in blood were analysed. All selected parameters in milk increased in challenged quarters but did not increase in control quarters. Milk IgG₁, IgG₂, serum albumin, and LDH were already significantly increased at 2 h after challenge whereas a significant increase of SCC was detectable at 2.75 h and L-lactate was increased at 2.25 h after challenge. In blood L-lactate was increased at 3.75 h after challenge, however, other factors in blood did not change significantly within the 5 h of experiment. In conclusion, the increase of blood components in milk during inflammation follows two different patterns: There is a rapid increase for IgG₁, IgG₂, or LDH, before the increase of SCC, and their concentrations reach a plateau within 3 h. On the other hand, SCC and L-lactate show a slower but consistent increase not reaching a plateau within 5 h after LPS challenge. L-lactate increases to higher concentrations in milk than in blood. This clearly shows that the increase of blood components follows different patterns and is therefore a controlled and compound-specific process and not exclusively an unspecific type of leakage.

Keywords: dairy cow, mastitis, blood-milk barrier.

In the mammary gland, invaded pathogens induce an immune response by stimulating the innate immune system. Along with this process considerable changes in milk composition occur due to changes in secretion but also through changes of the permeability of the blood-milk barrier leading to an increase of blood components in milk (Burton & Erskine, 2003; Schmitz et al. 2004; Wellnitz & Bruckmaier, 2012). These changes influence the nutritional value and taste, and the further processing of the milk. In addition, the change of milk composition plays an important role in the detection of mastitis through measurement of changes in concentration of mastitis indicators in milk. For this purpose a detailed knowledge of chronological changes and the mechanisms of the transfer of

various factors into milk during early stages of mastitis is essential.

Changes of specific components in milk that have immune function can influence the mammary immune defence. The infiltration of polymorphonuclear neutrophils (PMN), the main effectors of the mammary immune defence, into the mammary gland during mastitis is measured as an increase of the milk somatic cell count (SCC) (Burton & Erskine, 2003; Wellnitz & Bruckmaier, 2012). A low SCC is, therefore, used as a marker of good hygienic milk quality (Harmon, 1994).

Lipopolysaccharide (LPS) is an endotoxin that is released from gram-negative bacteria. Intramammary administration of LPS from *Escherichia coli*, a common mastitis pathogen, is a well-established method to experimentally induce mastitis under defined conditions for studying the immune response of the mammary gland (Bruckmaier et al. 1993; Schmitz et al. 2004; Baumert et al. 2009a; Wellnitz et al. 2011).

*For correspondence; e-mail: rupert.bruckmaier@vetsuisse.unibe.ch

Serum albumin is a blood protein whose appearance in milk can be used as an indicator of blood-milk barrier permeability (Stelwagen et al. 1994) and as an indicator of mastitis (Verhoeff & Smith, 1981). Immunoglobulin (Ig) G is the major immunoglobulin in milk of ruminants (Butler, 1983). The predominant antibody type in milk from healthy udders, although at a very low concentration, is IgG₁ due to its active and selective transport across the blood-milk barrier (Baker et al. 2009). During mastitis IgG₁ and IgG₂ increase in milk and IgG₂ is found in higher concentrations than IgG₁ in mastitis milk (Lehmann et al. 2013). For the mammary immune defence IgG₂ seems to play a more important role than IgG₁ because bovine neutrophils upregulate Fc receptors for IgG₂ when they migrate into infected tissue but not for IgG₁ or IgA (Paape et al. 2000; Burton & Erskine, 2003). In addition, IgM is not a major antibody subclass in normal or mastitic milk (Rainard & Caffin, 1983).

Milk L-lactate and lactate dehydrogenase (LDH) increase in milk during intramammary infection and they have been considered as indicators of mastitis (Davis et al. 2004; Chagunda et al. 2006). However, there is no obvious contribution of these factors to the immune defence.

For all these components a correlation to SCC increase is accepted (Harmon, 1994; Davis et al. 2004; Chagunda et al. 2006). However, a detailed chronological pattern of increase has, to our knowledge, not yet been described in the first 5 h of mastitis. To test the hypothesis that there are different component-specific patterns for the transfer of blood components into milk, this study aimed to investigate the sequence and the pattern of increase of SCC, the blood components IgG₁, IgG₂, serum albumin, L-lactate, and lactate dehydrogenase in milk. To record also a potential transfer in the opposite direction, i.e. from milk into blood, the milk protein α -lactalbumin was measured in blood during the first hours of an endotoxin-induced mastitis.

Material and methods

The experimental procedures followed the Swiss Federal Law on Animal Protection and were approved by the Committee of Animal Experiments of the Canton Fribourg, Switzerland.

Animals and management

Holstein ($n = 6$) and Swiss Fleckvieh ($n = 2$) cows in their 1st to 5th lactation (2.5 ± 0.6) and in month two to nine (5.5 ± 0.8) of lactation were used for this experiment. Cows were housed in tie stall barns during experiments and had constant low SCC below 130 000 cells/ml for at least 5 d before the experiment in all four quarters (morning foremilk samples). None of the udders showed clinical signs of mastitis. An intravenous catheter was inserted into one jugular vein on the day before the experiment to take blood samples.

Experimental design

After morning milking one quarter of each cow was challenged with 200 μ g *Escherichia coli* LPS (serotype O26:B6; Sigma-Aldrich, Buchs, Switzerland), diluted in 10 ml of saline solution after a careful disinfection of teat openings with 70% ethanol. Another quarter within cow was used as control and treated with 10 ml of saline solution. The treated quarters were randomly selected (front or rear, left or right) and equally distributed. Foremilk samples (~10 ml) were taken hourly by hand from LPS challenged and control quarter at 0 h (prior to injection) until 5 h after challenge. In addition, between 2 and 5 h after challenge, samples were taken every 15 min. Both milk samples were taken within 50 s before milk ejection started. Immediately after collection SCC was measured with a DeLaval cell counter (DCC; DeLaval, Tumba, Sweden) and then samples were frozen at -20°C for later analyses.

Blood samples were taken coinciding with milk sampling through the intravenous catheter. Blood was collected into tubes containing EDTA as anticoagulant. Blood samples were stored briefly on ice and then centrifuged for 20 min at 3000 g. Plasma was collected and frozen at -20°C until further analyses.

Concentrations of serum albumin in milk and plasma were measured using a commercial ELISA kit (Bethyl Laboratories, Montgomery, TX, USA) according to the manufacturer's instructions, except for using a higher dilution for milk samples (1 : 12 000) than described in the manual (1 : 5000). Because of high divergent values in treated and control samples, in plate dilutions were performed using a multichannel pipette (one 1 : 2 dilution), to meet the standard curve of the assay.

The immunoglobulin G₁ and G₂ concentrations in whole milk and plasma were measured using a bovine IgG₁ and IgG₂ ELISA quantitation kit (Bethyl Laboratories Inc., Montgomery, TX). The procedure was performed as described in the manufacturer's protocol, except for slight modifications: After coating the plates with anti IgG₁/IgG₂ antibody (50 μ l of capture antibody per 10 ml Coating buffer), the plates were washed with an automatic washing system (microplate washer Columbus Pro, Tecan Group Ltd., Männedorf, Switzerland). To reduce unspecific binding to anti IgG antibodies, the blocking reagent was replaced with 150 μ l/well of 5% fish skin gelatin Solution [5% of cold water fish skin gelatin (G7765; Sigma Aldrich, Steinheim, Germany) in distilled water]. After washing, 100 μ l of milk diluted in washing buffer (1 : 200) were added to each well. Because of the high sensitivity of the assay, in plate dilutions were performed (one serial dilution (1 : 2) using a multichannel pipette) to hit the standard range. Concentrations of IgG₁/IgG₂ were calculated by extrapolating from a standard curve generated by a microplate reader (Thermomax; Molecular Devices, Sunnyvale, CA).

Concentrations of α -lactalbumin in plasma were measured using a bovine α -lactalbumin ELISA quantitation set (Bethyl Laboratories Inc., Montgomery, TX) according to

the manufacturer's instructions. Because of the high sensitivity of the assay, plasma samples had to be diluted 1 : 20. In addition, the samples were serial-diluted once by 1 : 3 on the plate using a multichannel pipette to meet within standard curve of the assay.

For all ELISAs one control sample was added to each plate and was used for the calculation of coefficients of variation. For IgG ELISAs an intra assay variation of 10% and an inter assay variation of 20% was reached, respectively. For all other ELISAs and intra assay variation of 8% and an inter assay variation of 12% was reached, respectively.

Lactate dehydrogenase activity and L-lactate concentrations in milk serum and blood plasma were measured by using the test kits LDH IFCC (Axon Lab AG, Baden, Switzerland) and Lactate PAP (bioMérieux, Marcy l'Etoile, France), respectively, with an automated analyser (COBAS MIRA, Roche Diagnostics, Switzerland) according to the manufacturer's instructions.

Statistical analysis

Data are presented as means \pm SEM. Data of SCC are presented and statistically evaluated at a logarithmic scale (\log_{10}) to achieve normal distribution. Differences between LPS and control within each time point and differences between time points within one group were tested for significance ($P < 0.05$) by ANOVA using PROC MIXED of SAS (SAS Institute Inc., Cary, NC; 2002–2008, Release 9.2). The model included time, treatment (LPS or control) and their interaction as fixed effects and the quarter as repeated subject.

Results

Concentrations of somatic cells, serum albumin, IgG₁, IgG₂, and L-lactate and LDH activity in milk before and until 5 h after LPS challenge in challenged and control quarters are shown in Fig. 1. The SCC in milk was increased in LPS challenged quarters after 2.75 h compared to the initial value before LPS challenge ($P < 0.05$) and increased further until the end of the experiment. No significant changes of SCC were detected in control quarters. Serum albumin, IgG₁, IgG₂, and LDH was low in milk of all quarters at the beginning of the experiment and increased within 2 h in LPS challenged quarters ($P < 0.05$). Concentrations then remained elevated until the end of experiment. In control quarters the concentrations of these factors did not change.

L-lactate was increased ($P < 0.05$) in milk of LPS challenged quarters at 2.25 h and increased further thereafter until the end of experiment. In control quarters L-lactate concentrations did not significantly change within the 5 h of experiment.

Except for L-lactate, no change of any parameter in blood was detectable in the 5 h of experiment. Blood concentration of serum albumin was 4.11 ± 0.11 ng/ml throughout the experiment. Concentrations of IgG₁ and IgG₂ in blood were 5.3 ± 0.2 and 19.7 ± 0.6 mg/l, respectively. The activity of

LDH in blood was 923 ± 15 U/l throughout the entire experiment (mean of all values). Concentrations of α -lactalbumin in blood did not significantly change during the whole experiment, whereas between 2 h after the experiment and the end of the experiment an increase ($P = 0.013$) was detected (Fig. 2). The concentration of L-lactate in blood was increased at 3.75 h after challenge ($P < 0.05$) and further increased until the end of experiment (Fig. 2).

Discussion

The increase of somatic cell count and changes of milk composition during mastitis in dairy cows have been investigated over many decades. However, to our knowledge, the detailed description of the chronological patterns of the changes of blood components in milk within the first hours of mastitis has not been shown before. The hypothesis that different blood components are transferred into milk in different patterns was clearly confirmed by our study. Several blood components follow similar patterns of transfer into milk. This is of high importance for understanding the mechanisms of transfer of these components from blood into milk during mastitis. Furthermore the knowledge of the chronological appearance during the development of mastitis is important if factors are used for the early detection of mastitis and to figure out which is the most suitable parameter.

In LPS challenged quarters the increase of SCC within some hours was expected (Schmitz et al. 2004; Lehmann et al. 2013). This increase is known to be depending on the dosage of LPS (Baumert et al. 2009a) and the SCC before challenge (Baumert et al. 2009b). In this study the SCC increase was significant within 2.75 h.

During a mammary immune response the permeability of tight junctions increases and, therefore, a breakdown of the milk-blood barrier occurs (Kehrli & Shuster, 1994; Burton & Erskine, 2003). This leads to a facilitated passage of several blood components into the milk. The increased appearance of serum albumin in milk is accepted as an indicator for leakage of blood constituents into the milk through the blood-milk barrier (Poutrel et al. 1983; Shuster et al. 1993; Stelwagen et al. 1994) although there is conclusive evidence that the mammary gland also produces albumin and the expression is increased during mastitis (Shamay et al. 2005). The results of the present study agree with results from Shuster et al. (1993) who showed an increase of serum albumin in whey within the first 2 h after LPS infusion before SCC increased. It is possible that there is a retarded influx of cells due to slow diapedesis because the size of the cells necessitates a change in shape to fit through the gaps between endothelial and epithelial cells (Burton & Erskine, 2003). Chemoattractants have to be present and a number of events have to be initiated for an efficient influx of neutrophils into the mammary gland (Burton & Erskine, 2003). For other molecules like IgG, albumin, or sucrose a free transfer through openings within the cell layers is assumed (Nguyen & Neville, 1998).




Fig. 1. Somatic cell count (a), serum albumin (b), IgG₁ (c), IgG₂ (d), LDH (e), and L-lactate (f) in milk within 5 h after LPS challenge in LPS treated and control quarters. Data are presented as means \pm SEM. *: time point of first significant elevation compared to time point 0 until the end of experiment.

Interestingly the albumin concentrations in milk did not further increase after 3 h after challenge in the present study. Obviously, a maximal transfer from blood into milk was already attained at that time and shows a parallel transfer to osmotic regulated influx of water for milk production.

Immunoglobulin G₁ and G₂ concentrations increased in milk similar to albumin concentrations before a significant

increase of SCC was detected. This phenomenon was described before (Guidry et al. 1983). The majority of IgG₂ antibodies in milk are transferred from the blood (Leitner et al. 2000). The early leakage of high levels of blood IgG₂ was discussed to be crucial if pathogen-reactive blood IgG₂ antibodies are present to determine the success of neutrophil defence system against acute clinical mastitis (Burton &




Fig. 2. Alpha-lactalbumin (a) and L-lactate (b) in blood plasma within 5 h after LPS challenge. Data are presented as means \pm SEM. *: time point of first significant elevation compared to time point 0 until the end of experiment.

Erskine, 2003). The results agree with previous findings of parallel increase of albumin and total IgG concentrations in milk (Harmon et al. 1976) or albumin and IgG₁ and IgG₂ concentrations in milk (Rainard & Caffin, 1983) indicating a similar facilitated influx of these factors from blood.

LDH was shown to increase after intramammary LPS challenge mainly due to the leakage from the blood (Lehmann et al. 2013). Although LDH is released from the cytoplasm of cells into the extracellular fluid during cell damage and cell death (Glick, 1969) the rapid increase of LDH concomitantly to serum albumin and immunoglobulins in the present study confirms blood as the main source of milk LDH in the first hours of mammary inflammation. However, an increase of LDH in milk during mammary inflammation could play a role in later stages of the inflammatory process which was not investigated in the present study. The rapid increase of LDH in milk after LPS challenge before SCC increased shows again a rapid opening of the blood-milk barrier that enables a free influx of several blood components compared to slow diapedesis of cells.

For all the above mentioned factors that were increased in milk, no changes in blood could be observed. This is most likely due to the much bigger volume of blood compared to the produced milk during the 5 h of the experiment.

L-lactate concentrations are known to increase in milk during mastitis (Davis et al. 2004) and during LPS challenge (Lehmann et al. 2013). Due to the rapid increase in milk and the small size of the lactate molecule (90 Da) as compared to the other investigated constituents it is assumed that most of the L-lactate in milk in the beginning of mastitis is transferred from blood (Lehmann et al. 2013). However, it was shown before that L-lactate increases in blood in response to intravenous LPS administration (Giri et al. 1990). In addition it increases in milk after intramammary LPS challenge to even higher concentrations compared to those in blood (Lehmann et al. 2013). This clearly shows that L-lactate is not only transferred from blood through an opened blood-milk barrier. Silanikove et al. (2011) found that L-lactate is produced by mammary epithelial cells as a result of a transient shift to anaerobic metabolism during mastitis which is responsible for an increase of L-lactate in milk during the further course of mastitis.

Alpha-lactalbumin in plasma did not increase significantly throughout the 5 h experiment compared to concentrations immediately before LPS challenge. The appearance of α -lactalbumin in blood was expected as this protein, which is produced exclusively in the mammary gland is increased in blood due to the impairment of the blood-milk barrier (McFadden et al. 1988). It was measured to show the chronological sequence of the opening of the blood-milk barrier. However, already high concentrations of α -lactalbumin could be measured at the beginning of the experiment directly after milking with very large individual variations. Cows were in different stages of lactation. As the cows were milked just twice a day, it can be assumed that the udder in cows in earlier lactation has been relatively full immediately before milking and, therefore before the experiment. This opens the mammary tight junctions (Stelwagen et al. 1997). Therefore, it is likely that the α -lactalbumin leaked into the blood until the milking, and although the barrier was closed already at the start of the experiment (minutes after milking) the α -lactalbumin concentrations were still high, because a-lac needs 8 h to return to baseline levels (Stelwagen et al. 1997). On the contrary blood components in milk were removed with the milk during milking and were not transferred into milk in control quarters and in the first hour of the experiment in challenged quarters due to the sealed blood-milk barrier. Thus, serum albumin in milk for example, which is agreed to be a marker of a leaky blood-milk barrier was low in the beginning of the experiment but increased during the experiment which clearly shows the impairment of the barrier. Due to the high levels of alpha lactalbumin in the beginning of the experiment differences between 2 h after challenge and 5 h (end of experiment) were statistically investigated and detected. This shows that alpha-lactalbumin was transferred through the opening of the blood-milk barrier from

milk into the blood in response to LPS challenge, but this effect was superposed in the first 2 h of the experiment by high concentrations in blood before the start of experiment.

Conclusion

In conclusion, through a high frequency of milk sampling during the early stage of induced mammary inflammation the rapid opening of the blood-milk barrier can be demonstrated. However, this early effect is not reflected by the influx of cells from blood into milk. Clearly before the increase of SCC in foremilk samples an increase of several other blood components like serum albumin, IgG₁, IgG₂, or LDH is observed. These components reach a plateau in concentration within 3 h after LPS challenge. On the contrary, the increase of SCC and L-lactate in milk after LPS challenge shows a consistent increase over the experimental period not reaching a plateau within 5 h. While a free influx for several molecules is widely accepted cells are known to move into the milk as a consequence of chemotactic diapedesis. The increase of L-lactate in mastitic milk however, includes other mechanisms than just a facilitated influx from the blood as shown for the other measured factors mechanisms including an additional production in blood and/or milk. These results are of most importance if the measurement of blood components in milk is used for early mastitis detection and show that milk composition can dramatically change before an increase of SCC can be measured.

References

- Baker K, Quiao SW, Kuo T, Kobayashi K, Yoshida M, Lencer WI & Blumberg RS 2009 Immune and non-immune functions of the (not so) neonatal Fc receptor, FcRn. *Seminars in Immunopathology* **31** 223–236
- Baumert A, Bruckmaier RM & Wellnitz O, 2009a Dose dependant SCC changes after intramammary lipopolysaccharide challenge. *Milk Science International* **64** 119–121
- Baumert A, Bruckmaier RM & Wellnitz O 2009b Cell population, viability, and some key immunomodulatory molecules in different milk somatic cell samples in dairy cows. *Journal of Dairy Research* **76** 356–364
- Bruckmaier RM, Schallibaum M & Blum JW 1993 *Escherichia coli* endotoxin-induced mastitis in dairy cows: changes and importance of insulin-like growth factor I and oxytocin. *Milk Science International* **48** 374–378
- Burton JL & Erskine RJ 2003 Immunity and mastitis. Some new ideas for an old disease. *Veterinary Clinics Food Animal Practice* **19** 1–45
- Butler JE 1983 Bovine immunoglobulins: an augmented review. *Veterinary Immunology and Immunopathology* **4** 43–152
- Chagunda MG, Larsen T, Bjerring M & Ingvarsten KL 2006 L-lactate dehydrogenase and N-acetyl-beta-D-glucosaminidase activities in bovine milk as indicators of non-specific mastitis. *Journal of Dairy Research* **73** 431–440
- Davis SR, Farr VC, Prosser CG, Nicholas GD, Turner SA, Lee J & Hart AL 2004 Milk L-lactate concentration is increased during mastitis. *Journal of Dairy Research* **71** 175–181
- Giri SN, Emau P, Cullor JS, Stabenfeldt GH, Bruss ML, Bondurant RH & Osburn BI 1990 Effects of endotoxin infusion on circulating levels of eicosanoids, progesterone, cortisol, glucose and lactic acid, and abortion in pregnant cows. *Veterinary Microbiology* **21** 211–231
- Glick JH 1969 Serum lactate dehydrogenase isoenzyme and total lactate dehydrogenase values in health and disease, and clinical evaluation of these tests by means of discriminant analysis. *American Journal of Clinical Pathology* **52** 320–328
- Guidry AJ, Ost M, Mather IH, Shainline WE & Weinland BT 1983 Sequential response of milk leukocytes, albumin, immunoglobulins, monovalent ions, citrate, and lactose in cows given infusions of *Escherichia coli* endotoxin into the mammary gland. *American Journal of Veterinary Research* **44** 2262–2267
- Harmon RJ 1994 Physiology of mastitis and factors affecting somatic cell count. *Journal of Dairy Science* **77** 2103–2112
- Harmon RJ, Schanbacher FL, Ferguson LC & Smith KL 1976 Changes in lactoferrin, immunoglobulin G, bovine serum albumin, and alpha-lactalbumin during acute experimental and natural coliform mastitis in cows. *Infection and Immunity* **13** 533–542
- Kehrli ME & Shuster DE 1994 Factors affecting milk somatic cells and their role in health of the bovine mammary gland. *Journal of Dairy Science* **77** 619–627
- Lehmann M, Wellnitz O & Bruckmaier RM 2013 Concomitant lipopolysaccharide-induced transfer of blood-derived components including immunoglobulins into milk. *Journal of Dairy Science* **96** 889–896
- Leitner G, Yadlin B, Glickman A, Chaffer M & Saran A 2000 Systemic and local immune response of cows to intramammary infection with *Staphylococcus aureus*. *Research in Veterinary Science* **69** 181–184
- McFadden TB, Akers RM & Capuco AV 1988 Relationship of milk proteins in blood with somatic cell count in milk of dairy cows. *Journal of Dairy Science* **71** 826–834
- Nguyen DD & Neville MC 1998 Tight junction regulation in the mammary gland. *Journal of Mammary Gland Biology and Neoplasia* **3** 233–246
- Paape MJ, Shafer-Weaver K, Capuco A, Van Oostveldt K & Burvenich C 2000 Immune surveillance of mammary tissues by phagocytic cells. *Advances in Experimental Medicine and Biology* **480** 259–277
- Poutrel B, Caffin JP & Rainard P 1983 Physiological and pathological factors influencing bovine serum albumin content of milk. *Journal of Dairy Science* **66** 535–541
- Rainard P & Caffin JP 1983 Sequential changes in serum albumin, immunoglobulin (IgG₁, IgG₂, IgM) and lactoferrin concentrations in milk following infusion of *Escherichia coli* into the udder of immunized and unimmunized cows. *Annales de Veterinaire Research* **14** 271–279
- Schmitz S, Pfaffl MW, Meyer HHD & Bruckmaier RM 2004 Short-term changes of mRNA expression of various inflammatory factors and milk proteins in mammary tissue during LPS-induced mastitis. *Domestic Animal Endocrinology* **26** 111–126
- Shuster DE, Kehrli ME & Stevens MG 1993 Cytokine production during endotoxin-induced mastitis in lactating dairy cows. *American Journal of Veterinary Research* **54** 80–85
- Silanikove N, Rauch-Cohen A, Shapiro F, Blum S, Arieli A & Leitner G 2011 Lipopolysaccharide challenge of the mammary gland in bovine induced a transient glandular shift to anaerobic metabolism. *Journal of Dairy Science* **94** 4468–4475
- Shamay A, Homans R, Fuerman Y, Levin I, Barash H, Silanikove N & Mabjeesh SJ 2005 Expression of albumin in nonhepatic tissues and its synthesis by the bovine mammary gland. *Journal of Dairy Science* **88** 569–576
- Stelwagen K, Politis I, White JH, Zavizion B, Prosser CG, Davis SR & Farr VC 1994 Effect of milking frequency and somatotropin on the activity of plasminogen activator, plasminogen, and plasma in bovine milk. *Journal of Dairy Science* **77** 3577–3583
- Stelwagen K, Farr VC, McFadden HA, Prosser CG & Davis SR 1997 Time course of milk accumulation-induced opening of mammary tight junctions, and blood clearance of milk components. *American Journal of Physiology* **273** R379–R386
- Verhoeff J & Smith JA 1981 Bovine serum albumin and cell counts in the diagnosis of subclinical udder infections. *Veterinary Quarterly* **3** 38–45
- Wellnitz O & Bruckmaier RM 2012 The innate immune response of the bovine mammary gland to bacterial infection. *Veterinary Journal* **192** 148–152
- Wellnitz O, Arnold ET & Bruckmaier RM 2011 Lipopolysaccharide and lipoteichoic acid induce different immune responses in the bovine mammary gland. *Journal of Dairy Science* **94** 5405–5412