

Online Research @ Cardiff

This is an Open Access document downloaded from ORCA, Cardiff University's institutional repository: <https://orca.cardiff.ac.uk/143656/>

This is the author's version of a work that was submitted to / accepted for publication.

Citation for final published version:

Xu, Christine, Rafique, Ashique, Potocky, Terra, Paccaly, Anne, Nolain, Patrick, Lu, Qiang, Iglesias-Rodriguez, Melitza, St John, Gregory, Nivens, Michael C., Kanamaluru, Vanaja, Fairhurst, Jeanette, Ishii, Tomonori, Maldonado, Rafael, Choy, Ernest and Emery, Paul 2021. Differential binding of sarilumab and tocilizumab to IL-6R α and effects of receptor occupancy on clinical parameters. *Journal of Clinical Pharmacology* 61 (5), pp. 714-724.
10.1002/jcph.1795 file

Publishers page: <http://dx.doi.org/10.1002/jcph.1795>
<<http://dx.doi.org/10.1002/jcph.1795>>

Please note:

Changes made as a result of publishing processes such as copy-editing, formatting and page numbers may not be reflected in this version. For the definitive version of this publication, please refer to the published source. You are advised to consult the publisher's version if you wish to cite this paper.

This version is being made available in accordance with publisher policies.

See

<http://orca.cf.ac.uk/policies.html> for usage policies. Copyright and moral rights for publications made available in ORCA are retained by the copyright holders.



Differential binding of sarilumab and tocilizumab to IL-6R α and effects of receptor occupancy on clinical parameters

Christine Xu¹, Ashique Rafique², Terra Potocky², Anne Paccaly², Patrick Nolain³, Qiang Lu¹, Melitza Iglesias-Rodriguez⁴, Gregory St John², Michael C. Nivens², Vanaja Kanamaluru¹, Jeanette Fairhurst², Tomonori Ishii⁵, Rafael Maldonado⁶, Ernest Choy⁷, Paul Emery⁸

¹Sanofi, Bridgewater, NJ, USA, ²Regeneron Pharmaceuticals, Inc., Tarrytown, NY, USA, ³Sanofi Aventis, Montpellier, France, ⁴Sanofi Genzyme, Cambridge, MA, USA, ⁵Tohoku University Graduate School of Medicine, Sendai, Japan, ⁶Pompeu Fabra University, Barcelona, Spain, ⁷CREATE Centre, Section of Rheumatology, Cardiff University School of Medicine, Cardiff, UK, ⁸Leeds Musculoskeletal Biomedical Research Unit, Leeds Teaching Hospitals Trust, Leeds Institute of Rheumatic and Musculoskeletal Medicine, University of Leeds, Leeds, UK

Correspondence to: Christine Xu, Sanofi, Bridgewater, NJ, USA

Email: christine.xu@sanofi.com

Target journal: Rheumatology

Word count: 3425/3500 max (249/250 max for the abstract)

Tables/Figures: 6/6 max

References: 24/50 max

Abstract

Objective. To evaluate IL-6R α signalling inhibition with sarilumab and tocilizumab, the association between IL-6R α receptor occupancy (RO) and C-reactive protein (CRP), and potential clinical relevance of any differences.

Methods. We measured IL-6R α binding and signalling inhibition with sarilumab and tocilizumab *in vitro*, simulated sIL-6R α RO over time for approved sarilumab SC and tocilizumab IV and SC doses, and assessed associations between calculated RO and CRP reduction, DAS28-CRP, and ACR20/50/70 from clinical data.

Results. Sarilumab binds IL-6R α *in vitro* with 15–22-fold higher affinity than tocilizumab, and inhibits IL-6-mediated classical and *trans* signalling via membrane-bound and soluble IL-6R α . Sarilumab 200 and 150 mg SC q2w achieved >90% RO after first and second doses, respectively, maintained through the treatment period. At steady-state trough, RO was greater with sarilumab 200 mg (98%) and 150 mg SC q2w (94%) and tocilizumab 162 mg SC qw (99%) and 8 mg/kg IV q4w (>99%) vs tocilizumab 162 mg SC q2w (84%) and 4 mg/kg IV q4w (60%). Higher RO was associated with greater CRP reduction and DAS28-CRP reduction, and more sarilumab patients achieving ACR20/50/70. Greatest reductions in CRP levels were observed with sarilumab (both doses) and tocilizumab 8 mg/kg IV q4w (reductions proportionally smaller with 4 mg/kg IV q4w).

Conclusion. Higher IL-6R binding affinity translated into higher RO with sarilumab vs tocilizumab 4 mg/kg q4w or 162 mg q2w; tocilizumab required the higher dose or increased frequency to maintain the same degree of RO and CRP reduction. Higher RO was associated with clinical parameter improvements.

Trial registration. MOBILITY Part A (NCT01061736); ASCERTAIN (NCT01768572); SUMMACTA (NCT01194414); BREVACTA (NCT0123256)

Key words: Rheumatoid arthritis, Pharmacology, Cell receptor-ligand interaction and activation, Cytokines and inflammatory mediators, Clinical trials and methods

Rheumatology key messages

- Higher IL-6R α binding with sarilumab translates to higher receptor occupancy vs lower approved tocilizumab doses
- Higher IL-6R receptor occupancy is associated with greater effects on pharmacodynamic/clinical parameters
- The data support sarilumab q2w dosing; both doses resulted in rapid and sustained CRP reduction

Introduction

Rheumatoid arthritis (RA) is a chronic and debilitating autoimmune disease, and interleukin-6 (IL-6) is a pleiotropic cytokine that acts as a critical signalling node in the complex pro-inflammatory cytokine network that underpins RA (1, 2). IL-6 elevations have been noted in serum and synovial fluid in patients with RA and correlate with RA disease activity and joint destruction (3). IL-6 may contribute to comorbidities associated with RA, including mood disorders, cardiovascular disease, diabetes, and osteoporosis (4-6).

IL-6 effects are mediated through interaction with the IL-6 receptor- α (IL-6R α). IL-6 activates classical (*cis*) signalling through membrane-bound IL-6R α (mIL-6R α), expressed on the surface of hepatocytes and haematopoietic cells, and *trans* signalling through soluble IL-6R α (sIL-6R α), found in serum and synovial fluid. Signalling with sIL-6R α occurs after the IL-6/sIL-6R α complex binds to the ubiquitously expressed glycoprotein (gp)130 receptor, thus greatly expanding the spectrum of IL-6-responsive cells (5, 7, 8). Pharmacodynamic (PD) effects of IL-6R blockade include decreased production of inflammatory acute-phase reactant. For instance, C-reactive protein (CRP), which can be considered a surrogate marker of efficacy.

Sarilumab and tocilizumab are monoclonal antibodies (mAbs) that block IL-6 binding to sIL-6R α and mIL-6R α , thereby inhibiting IL-6 signalling through this pathway (9-12). Sarilumab (human mAb) and tocilizumab (humanized mAb) are approved for the treatment of adults with moderately to severely active RA and inadequate responses to disease-modifying antirheumatic drugs. Sarilumab is administered subcutaneously (SC) at 200 mg once every 2 weeks (q2w), with reduction to 150 mg q2w if required to manage laboratory abnormalities (11, 12). In the US, the recommended tocilizumab dose for intravenous (IV) administration is 4 mg/kg every 4 weeks (q4w) or for SC administration it is 162 mg q2w. Up-titration to 8 mg/kg IV q4w or 162 mg SC weekly (qw) (if clinical response is inadequate) is recommended for IV administration and SC administration, respectively. In the European Union, the higher tocilizumab doses of 8 mg/kg IV q4w and 162 mg SC qw are recommended, down-titrating if required to manage laboratory abnormalities (9, 10).

The objectives of this analysis were to evaluate differences in IL-6R α binding profiles *in vitro* and the resultant functional activities of sarilumab and tocilizumab, and then explore, using a pharmacokinetic (PK)/PD modelling approach, how binding translates *in vivo* to receptor occupancy (RO) following recommended dosing of sarilumab and tocilizumab. The association between RO and subsequent changes in clinical efficacy parameters (CRP reduction, 28-joint disease activity score based on CRP [DAS28-CRP], and American College of Rheumatology [ACR]20/50/70 responses), observed in a dose-ranging study in patients with RA, were also evaluated.

Methods

In vitro IL-6 binding and signalling

Kinetic binding analysis

Binding kinetics of sarilumab and tocilizumab to IL-6R α were measured using Surface Plasmon Resonance (SPR; Biacore™ T200). Further details are included in the Supplementary Appendix. Binding kinetics calculated were association rate constant (k_a), dissociation rate constant (k_d), and half-life ($t_{1/2}$). The overall equilibrium dissociation constant (K_D) was calculated from the ratio of k_d to k_a . Further details are included in the Supplementary Appendix.

Blockade of dimeric hIL-6R α binding to IL-6

hIL-6R α binding to IL-6 was assessed using an enzyme-linked immunosorbent assay (ELISA) competition assay. Further details are included in the Supplementary Appendix. Values for the inhibitory concentration at 50% activity (IC_{50}) and effective concentration at 50% activity (EC_{50}) were calculated for hIL-6, sarilumab, and tocilizumab. Further details are included in the Supplementary Appendix.

Inhibition of classical IL-6R α signalling

The activity of sarilumab and tocilizumab in blocking classical IL-6R α signalling was compared *in vitro* in cell proliferation assays using:

- DS-1: a human B lymphocyte cell line that proliferates in response to exogenous hIL-6, and endogenously expresses IL-6R α and gp130
- HepG2: a hepatocytic cell line endogenously expressing IL-6R α and gp130

Further details are included in the Supplementary Appendix.

Inhibition of trans-IL-6R α signalling

The ability of sarilumab to block *trans* signalling stimulated by the IL-6/sIL-6R α complex was assessed in a functional cell-based luciferase assay using a gp130-overexpressing human embryonic kidney/STAT3/luciferase reporter cell line. Further details are included in the Supplementary Appendix.

PK/PD modelling of RO and effects on CRP reduction, DAS28-CRP, and ACR20/50/70

PK model

The PK framework for the sIL-6R α PK/PD models was provided by population PK (PopPK) models for sarilumab SC (13), and tocilizumab IV (14) and SC (15, 16). These models described the PK of sarilumab and tocilizumab by a two-compartmental model with parallel linear and non-linear Michaelis–Menten elimination and with first-order absorption for sarilumab and tocilizumab SC (13-15).

sIL-6R α PK/PD model development

Tocilizumab binding to sIL-6R was described by the PK/PD model previously developed using data from studies evaluating tocilizumab IV (at 4 or 8 mg/kg q4w) for 24 weeks in patients with RA (14). Given the similarity of tocilizumab and sarilumab binding to sIL-6R α and mIL-6R α , the same structure model was used to develop the PK/PD model and describe sarilumab binding to sIL-6R α following SC dosing, using data from MOBILITY Part A (NCT01061736). MOBILITY Part A was a Phase 2, double-blind, placebo-controlled, dose-ranging study in 306 patients with RA, evaluating five sarilumab SC regimens (100 mg q2w, 150 mg q2w, 100 mg qw, 200 mg q2w, 150 mg qw) over 12 weeks (17).

The quasi-steady-state target-mediated drug disposition (TMDD) models describing PK/PD relationships to total sIL-6R α for sarilumab and tocilizumab, including the PD model of inhibiting sIL-6R α elimination, are summarized in Supplementary Fig. 1. The PK/PD model for binding to sIL-6R α was used to predict the time-course of free sIL-6R concentrations for sarilumab and tocilizumab. Only sIL-6R α (not mIL-6R α) was considered in these analyses. The sIL-6R α PK/PD model analysis was performed using NONMEM 7.2.0 (ICON plc, Dublin, Ireland). The quality of the PK/PD model was extensively evaluated using standard goodness-of-fit (GOF) criteria (observations vs individual and population predictions, and weighted residuals), as well as by the condition number. The final PK/PD model was evaluated using a visual predictive check (VPC) to test the robustness of the model and the accuracy of parameter estimates. Model verification was performed by comparing observed data with literature-reported data.

Simulation of sIL-6R α occupancy by sarilumab and tocilizumab

Literature-reported PK and PK/PD models of tocilizumab, and developed PK and PK/PD models of sarilumab were used to profile the time-course of tocilizumab or sarilumab concentrations in serum, and estimate binding to sIL-6R and free sIL-6R concentrations for the approved dosage regimens of sarilumab SC (200 mg and 150 mg q2w), tocilizumab IV (8 mg/kg and 4 mg/kg q4w), and

tocilizumab SC (162 mg qw and q2w). sIL-6R α occupancy dynamic profiles (% RO over time) were calculated based on unbound sIL-6R concentrations:

$$RO=1-(\text{Free sIL-6R}_{\text{post-treatment}}/\text{Free sIL-6R}_{\text{baseline}})$$

RO over 24 weeks was calculated following sarilumab SC and tocilizumab IV regimens from ASCERTAIN (NCT01768572), a Phase 3 safety study in which 202 patients with RA were randomized to sarilumab SC 200 mg or 150 mg q2w, or tocilizumab 4 mg/kg IV q4w for 24 weeks (18). Patients were able to up-titrate their tocilizumab IV dosage to 8 mg/kg in cases of inadequate clinical response (61% required up-titration). Additional simulation of RO for tocilizumab SC regimens was provided based on the Phase 3 randomized, double-blind SUMMACTA and BREVACTA 24-week studies (15) evaluating tocilizumab SC 162 mg qw vs IV 8 mg/kg q4w ($n=1262$) and tocilizumab SC 162 mg q2w vs placebo ($n=656$), respectively (19) (20).

Association between sIL-6R α receptor occupancy and CRP, DAS28-CRP, and ACR response

The association of median RO calculated from observed concentrations of free sIL-6R measured in MOBILITY Part A (17) was plotted against median levels of CRP reduction, median DAS28-CRP reduction, and ACR20/50/70 responses by treatment groups in patients randomized to receive sarilumab 100 mg q2w ($n=51$), sarilumab 150 mg q2w ($n=51$), sarilumab 100 mg qw ($n=50$), sarilumab 200 mg q2w ($n=52$), sarilumab 150 mg qw ($n=50$), or placebo ($n=52$) for 12 weeks.

sIL-6R α RO profiles were compared visually with changes in observed mean CRP levels from ASCERTAIN, described above. To further verify the association observed in ASCERTAIN, sIL-6R α occupancy profiles were compared visually with changes in reported mean CRP levels from SUMMACTA and BREVACTA.

Results

***In vitro* IL-6 binding and signalling**

Sarilumab bound with high affinity to recombinant monomeric and dimeric human IL-6R α in SPR assays, with K_D values of 61.9 pM and 12.8 pM, respectively (Table 1, Supplementary Fig. 2A).

Sarilumab showed 15–22-fold higher affinity than tocilizumab in binding to monomeric and dimeric hIL-6R α forms (Table 1, Supplementary Fig. 2B).

In the ELISA competition assay, sarilumab directly blocked the binding of hIL-6R α -Fc to plate-coated hIL-6 with an IC_{50} of 108 pM (achieving complete blockade to baseline levels), whereas the IgG1 isotype control showed no blocking activity under the same conditions (Supplementary Fig. 3B). A constant concentration of 100 pM IL-6R α -hFc was used in the assay, which bound to hIL-6 with an EC_{50} of 255 pM. At the time, only sarilumab was available for evaluation in this assay.

In the *in vitro* proliferation assay, both sarilumab and tocilizumab inhibited IL-6-mediated proliferation of DS-1 cells (classical signalling, Fig. 1A), with IC_{50} values approximately 3.6-fold more potent for sarilumab than tocilizumab (226 pM *vs* 812 pM, in the presence of 1.0 pM IL-6). hIL-6 had an EC_{50} value of 0.5 pM in this assay. Sarilumab and tocilizumab inhibited IL-6-mediated luciferase activity in the HepG2 cell luciferase reporter assay, indicating inhibition of classical IL-6R α signalling via the STAT-3 pathway (Fig. 1B). Sarilumab was approximately 3.4-fold more potent than tocilizumab with an IC_{50} of 146 pM *vs* 496 pM (in the presence of 50 pM IL-6). hIL-6 had an EC_{50} value of 59 pM in this assay.

In the HEK293 cell line, IL-6 was shown to activate gp130 receptor *trans* signalling in the presence of 1 nM or 10 nM soluble IL-6R α -mmH, with EC_{50} values of 1.8 nM and 0.7 nM, respectively; signalling could not be activated by IL-6 alone (Fig. 1C and 1D). Sarilumab blocked *trans* signalling with an IC_{50} of 0.9 nM in the presence of 1 nM sIL-6R α -mmH and 10 nM IL-6, and an IC_{50} of 8.9 nM in the presence of 10 nM sIL-6R α -mmH and 10 nM IL-6 (Fig. 1C and 1D). Again, at the time, only sarilumab was available for evaluation in this assay.

PK/PD modelling of RO and effects on clinical efficacy parameters

Parameter estimates of the sIL-6R α PK/PD models are presented in Supplementary Table 1. GOF evaluation indicated that the final sIL-6R α PK/PD model was consistent with the observed data (Supplementary Fig. 4) and the VPC showed that the time-course profiles with the observed concentrations (2.5th, 50th, and 97.5th percentiles) fitted the predicted parameters well (Supplementary Fig. 5).

Simulation of sIL-6R α occupancy by sarilumab and tocilizumab

Simulated sIL-6R α occupancy profiles over 24 weeks for sarilumab SC vs tocilizumab IV and sarilumab SC vs tocilizumab SC are shown in Fig. 2A and 2B. Sarilumab 200 mg SC q2w achieved >90% RO after the first dose, maintained over the dosing interval and through the 24-week simulated treatment period (Fig. 2A). Simulated RO for sarilumab SC 150 mg q2w decreased to 74% towards the end of the first dosing interval, but from the third dose onward was maintained at >90% over the full dosing interval.

The tocilizumab IV 8 mg/kg q4w and the SC 162 mg qw dose regimens achieved >90% simulated RO from the first dose, maintained over the dosing interval and through the 24-week simulated treatment period. In contrast, with the tocilizumab IV 4 mg/kg q4w and the SC 162 mg q2w dose regimens, trough RO values below the 90% threshold were predicted by the end of each dosing interval over the 24-week period.

At Week 24, steady-state trough IL-6R α occupancy levels were 98% and 94% for sarilumab SC 200 mg and 150 mg q2w regimens, respectively, 99% and 60% for tocilizumab IV 8 mg/kg q4w and 4 mg/kg q4w regimens, respectively (Fig. 2A), and >99% (99.6%) and 84% for tocilizumab SC 162 mg qw and q2w regimens, respectively (Fig. 2B).

Association between sIL-6R α occupancy and clinical efficacy parameters

In MOBILITY Part A, the association of sIL-6R α RO was investigated for the following clinical efficacy parameters: percentage CRP reduction, DAS28-CRP score, and ACR20/50/70. Week 12 RO was calculated for placebo and sarilumab 100 and 150 mg qw and sarilumab 100, 150, and 200 mg q2w, and plotted against Week 12 efficacy data. Higher RO was associated with greater CRP reduction and consequently with larger reductions in DAS28-CRP scores (Fig. 3A and 3B). There was also an apparent association between higher RO and ACR20/50/70 responses (Fig. 3C).

In ASCERTAIN, where patients were randomized to sarilumab or tocilizumab, sarilumab induced rapid (from Week 4: first assessment after first dose) and sustained reduction of CRP throughout the dosing period. At Week 24, the greatest reduction in CRP levels was observed in patients receiving sarilumab SC (at either dose), or in patients who escalated their tocilizumab dose to 8 mg/kg IV q4w (Fig. 4). The magnitude of CRP reduction was lower in patients who remained on tocilizumab 4 mg/kg IV q4w. Comparing observed CRP levels for SC qw and q2w tocilizumab from SUMMACTA and BREVACTA (Fig. 5), the inverse relationship between sIL-6R α occupancy and CRP appeared to hold true for these SC tocilizumab regimens.

Discussion

In vitro experiments reported here demonstrate that sarilumab has a higher relative binding affinity to sIL-6R α compared to tocilizumab and inhibits IL-6-induced cellular responses (i.e. cell proliferation and STAT signalling) with higher potency and at lower concentrations than tocilizumab. Consistently, PK/PD modelling using data from Phase 2 and 3 studies in patients with RA, indicated higher and more sustained IL-6R α RO with sarilumab SC q2w dose regimens than with tocilizumab 4 mg/kg IV q4w or 162 mg SC q2w. Higher receptor occupancy was associated with better clinical parameters.

IL-6 activates cells via a signalling mechanism that requires two receptor components, the IL-6R α and gp130. IL-6 forms a heterodimer with IL-6R α that subsequently binds with high affinity to gp130, forming a heterotrimeric complex (5, 7, 8). IL-6R α exists in both membrane-bound and

soluble forms, with sIL-6R α generated through cleavage of mIL-6R α or alternative splicing. Classical (*cis*) signalling through mIL-6R α is limited to the few cell types that express mIL-6R α , that is hepatocytes, monocytes/macrophages, neutrophils, and some T cell subsets. *Trans* signalling through sIL-6R α may occur in virtually any nucleated cell type (including those that lack IL-6R α expression) because of the ubiquitous expression of membrane gp130 [8].

Previously, Mihara and colleagues showed tocilizumab bound to sIL-6R α , inhibited IL-6 binding in a dose-dependent manner, and dissociated IL-6 and sIL-6R α from their preformed complex. Tocilizumab suppressed the IL-6/sIL-6R complex-induced proliferation of human gp130-transfected cells (BAF-h130) and bound human IL-6R expressing COS-7 cells. It also suppressed growth of the IL-6-dependent myeloma cell line, KPMM2 (21). In the present *in vitro* studies, sarilumab was shown to bind directly to IL-6R α , but with a binding affinity for both monomeric and dimeric IL-6R α approximately 15- to 22-fold higher than that of tocilizumab. The studies confirm the ability of sarilumab to block both classical- and *trans*-mediated signalling. Sarilumab blocked IL-6-induced growth of the human B cell line DS-1 and inhibited IL-6-induced STAT signalling in the human hepatocyte cell line HepG2 at concentrations approximately 3-fold lower than tocilizumab. Sarilumab was found to completely inhibit activation in a *trans* signalling assay in which cells expressing gp130 were stimulated by the soluble form of the sIL-6R α complex.

Additional factors can influence binding affinity *in vivo*, including baseline receptor concentration, receptor turnover, receptor distribution, antibody concentration, and antibody distribution (22). Therefore, a PK/PD model was developed that incorporated these parameters to profile sIL-6R α occupancy for approved dose regimens of sarilumab SC (200 and 150 mg q2w) and tocilizumab IV (4 and 8 mg/kg q4w) and SC (162 mg qw and q2w). This model used previously published and validated PopPK parameters to describe sarilumab and tocilizumab drug concentrations in serum for the dosing regimens tested. Sarilumab and tocilizumab are both eliminated by parallel linear and non-linear pathways, with the linear, non-saturable proteolytic pathway predominating at higher concentrations (9, 12).

Sarilumab and tocilizumab bind to both the sIL-6R and mIL-6R forms. It was shown that the TMDD system with two targets can be approximated by equations that describe both sarilumab or tocilizumab and sIL-6R concentrations, and include two target-mediated elimination terms (with different maximum elimination rate [V_{max}] and Michaelis–Menten constant [K_m] parameters). However, mIL-6R was not measured in the clinical studies and insufficient data did not allow for separation of the two different V_{max} and K_m parameters. The *in vitro* K_D values for tocilizumab–sIL-6R (0.11 $\mu\text{g/mL}$ [0.75 nmol/L]) and tocilizumab–mIL-6R binding (0.38–0.43 $\mu\text{g/mL}$ [2.5–2.9 nmol/L]) suggest a similar range of binding affinity for the two forms of the target (14).

RO assays applied in both non-clinical and clinical studies provided an insight into PK/PD relationships for binding to receptors. RO on circulating cells has been used as a PD biomarker for nivolumab and etrolizumab (23). RO simulations generated from the PK/PD models in the current study indicate that sarilumab 200 mg SC dosing regimens and the tocilizumab IV 8 mg/kg q4w and SC 162 mg qw regimens are able to achieve and maintain target sIL-6R α occupancy (>90%) for the entire dosage interval over a 24-week treatment period (for sarilumab 150 mg SC dosing Week 4 onwards). For the lower tocilizumab dose of 4 mg/kg IV q4w and the less frequent 162 mg SC q2w regimen, which comprise the US-recommended tocilizumab starting doses, RO fell below 90% towards the end of each dosing period. These RO findings are consistent with the higher sIL-6R α binding affinity and slower dissociation kinetics of sarilumab compared with tocilizumab observed in earlier *in vitro* assays, and with serum tocilizumab trough concentration being 134-fold lower with the 4 vs 8 mg/kg IV regimen and 10.5-fold lower with the q2w vs qw regimen (9, 15).

Besides being of scientific interest, the importance of the current findings becomes apparent with the associations between RO, PD parameters, and clinical parameters. Concentrations of sIL-6R in serum measured in MOBILITY Part A, the dose-ranging portion of that study indicated an association between the degree of RO and the degree of CRP reduction, DAS28-CRP improvement, and/or ACR20/50/70 response at Week 12. ASCERTAIN was the first double-blind multiple-dose safety study in patients with RA to include sarilumab SC and tocilizumab IV dosing regimens within the same study (24), allowing PK/PD comparisons to be made. The IL-6R α occupancy results

generated with the present models are inversely associated with observed changes in CRP levels over the 24-week treatment period of ASCERTAIN, confirming the expected association between RO and clinically-relevant PD markers. Rapid dose-related reduction of CRP sustained over 24 weeks was noted for sarilumab SC 200 and 150 mg q2w regimens, whereas little or no CRP reduction was evident at Week 4 in the tocilizumab IV 4 mg/kg q4w group. As anticipated based on RO simulations, patients on the 4 mg/kg q4w dose who up-titrated their tocilizumab IV dose to 8 mg/kg q4w, because of insufficient clinical response, experienced reductions in CRP levels similar to those observed in the sarilumab groups.

It is not possible to directly compare SC regimens of sarilumab and tocilizumab in terms of the association between sIL-6R α occupancy and CRP, as ASCERTAIN did not include a SC tocilizumab dose group as none was available at that time. When the RO results generated for the qw and q2w tocilizumab regimens were evaluated, the inverse relationship between sIL-6R α occupancy and CRP appeared to hold true for the SC tocilizumab regimens, in addition to the observed CRP data for these regimens in SUMMACTA and BREVACTA (15). PK/PD analysis of these two studies showed a more gradual decline of PD responses (both CRP and erythrocyte sedimentation rate) over time for the SC q2w regimen, compared with the tocilizumab IV q4w and SC qw regimens. Other limitations include the possible effects on CRP by a number of other factors, most notably infection. Also, the findings presented here are restricted to the sIL-6R α isoform, because mIL6-R α was not measured in either the tocilizumab or sarilumab trials.

These findings on the effect of RO on clinical efficacy parameters support selection of dosing regimens of sarilumab (150 mg and 200 mg q2w SC) as providing the required occupancy of IL-6R α to elicit optimal reduction of clinical markers of IL-6 activity (CRP). These findings might have important clinical implications because they suggest that for tocilizumab, the lower 4 mg/kg IV q4w and 162 mg SC q2w dosing regimens may not provide adequate occupancy of IL-6R α to elicit the desired clinical effect. This is consistent with *in vitro* experiments that show sarilumab has a higher relative binding affinity for IL-6R α and is more potent at inhibiting IL-6-mediated signalling at lower concentrations in serum than tocilizumab.

Acknowledgements

The authors would like to thank Ching-Ha (Vicki) Lai for providing support in the bioanalytical analyses of PK data. Medical writing assistance was provided under the direction of the authors by Peter Tran, PhD, at Adelphi Communications Ltd (Macclesfield, UK) in accordance with Good Publication Practice (GPP3) guidelines.

Funding: The studies described herein were sponsored by Sanofi and Regeneron Pharmaceuticals, Inc. Funding for medical writing assistance with this manuscript was provided by Sanofi and Regeneron Pharmaceuticals, Inc.

Disclosure statement: C. Xu, P. Nolain, Q. Lu, M. Iglesias-Rodriguez, and V. Kanamaluru are employees of Sanofi Genzyme, and may hold stock and/or stock options in the company; A. Paccaly, A. Rafique, T. Potocky, G. St John, M. Nivens, and J. Fairhurst are employees of Regeneron Pharmaceuticals, Inc., and may hold stock and/or stock options in the company; R. Maldonado has received research grants or consulting fees from Aelis, Almirall, Boehringer Ingelheim, BrainCo, Esteve, Ferrer, GlaxoSmithKline, Grünenthal, GW Pharmaceuticals, Janus, Lundbeck, Pharmaleads, PhytoPlant, Rhodes, Sanofi, Spherium, Union de Pharmacologie Scientifique Appliquée, Upjohn, and Uriach; T. Ishii has received research grants or consulting fees or participated in speakers' bureaus from/for AbbVie, Asahi, Astellas, Chugai, Daiichi Sankyo, Eisai, GlaxoSmithKline, Janssen, Kasei Pharma, Mitsubishi Tanabe, Ono, Pfizer, Sanofi, Takeda, Teijin, and UCB; E. Choy has received research grants or consulting fees or participated in advisory boards or speakers' bureaus from/for AbbVie, Amgen, AstraZeneca, Bio-Cancer, Biogen, Boehringer Ingelheim, Bristol-Myers Squibb, Celgene, Chelsea Therapeutics, Chugai Pharma, Eli Lilly, Ferring Pharmaceuticals, GlaxoSmithKline, Hospira, ISIS, Janssen, Jazz Pharmaceuticals, MedImmune, Merck Sharp & Dohme, Merrimack Pharmaceuticals, Napp, Novartis, Novimmune, ObsEva, Pfizer, Regeneron Pharmaceuticals Inc., Roche, R-Pharm, Sanofi-Aventis, Sanofi Genzyme, SynAct Pharma, Tonix, and UCB; P. Emery has received grant/research support from AbbVie, Merck, Pfizer and Roche, and is a consultant for AbbVie, Bristol-Myers Squibb, Lilly, Merck, Novartis, Pfizer, Roche, and Samsung Bioepis.

Institutional review board approval and ethics: For each of the clinical studies described in this manuscript, institutional review board and ethics committee approval of the protocols were obtained prior to study initiation, and all patients provided written informed consent.

Availability of data and materials: Qualified researchers may request access to patient-level data and related study documents including the clinical study report, study protocol with any amendments, blank case report form, statistical analysis plan, and dataset specifications. Patient-level data will be anonymized and study documents will be redacted to protect the privacy of trial participants. Further details on Sanofi's data-sharing criteria, eligible studies, and process for requesting access can be found at: <https://www.clinicalstudydatarequest.com>

References

1. Calabrese LH, Rose-John S. IL-6 biology: implications for clinical targeting in rheumatic disease. *Nat Rev Rheumatol*. 2014;10(12):720-7.
2. Dayer JM, Choy E. Therapeutic targets in rheumatoid arthritis: the interleukin-6 receptor. *Rheumatology (Oxford)*. 2010;49(1):15-24.
3. Choy E. Understanding the dynamics: pathways involved in the pathogenesis of rheumatoid arthritis. *Rheumatology (Oxford)*. 2012;51(Suppl 5):v3-11.
4. Raimondo MG, Biggioggero M, Crotti C, Becciolini A, Favalli EG. Profile of sarilumab and its potential in the treatment of rheumatoid arthritis. *Drug Des Devel Ther*. 2017;11:1593-603.
5. Choy EHS, Calabrese LH. Neuroendocrine and neurophysiological effects of interleukin 6 in rheumatoid arthritis. *Rheumatology (Oxford)*. 2018;57(11):1885-95.
6. Nicolau J, Lequerre T, Bacquet H, Vittecoq O. Rheumatoid arthritis, insulin resistance, and diabetes. *Joint Bone Spine*. 2017;84(4):411-6.
7. Narazaki M, Kishimoto T. The Two-Faced Cytokine IL-6 in Host Defense and Diseases. *Int J Mol Sci*. 2018;19(11):3528.
8. Rose-John S. IL-6 trans-signaling via the soluble IL-6 receptor: importance for the pro-inflammatory activities of IL-6. *Int J Biol Sci*. 2012;8(9):1237-47.
9. ACTEMRA (Tocilizumab). US prescribing information 2019 [Available from: https://www.gene.com/download/pdf/actemra_prescribing.pdf].
10. RoACTEMRA (Tocilizumab). Summary of product characteristics 2019 [updated November 14, 2019. Available from: https://www.ema.europa.eu/en/documents/product-information/roactemra-epar-product-information_en.pdf].
11. KEVZARA (Sarilumab). US Prescribing information 2018 [updated April 2018. Available from: https://www.accessdata.fda.gov/drugsatfda_docs/label/2018/761037s001lbl.pdf].
12. KEVZARA (Sarilumab). Summary of product characteristics 2019 [updated May 2019. Available from: https://www.ema.europa.eu/en/documents/product-information/kevzara-epar-product-information_en.pdf].
13. Xu C, Su Y, Paccaly A, Kanamaluru V. Population Pharmacokinetics of Sarilumab in Patients with Rheumatoid Arthritis. *Clin Pharmacokinet*. 2019;58:1455-67
14. Gibiansky L, Frey N. Linking interleukin-6 receptor blockade with tocilizumab and its hematological effects using a modeling approach. *J Pharmacokinet Pharmacodyn*. 2012;39(1):5-16.
15. Abdallah H, Hsu JC, Lu P, Fettner S, Zhang X, Douglass W, et al. Pharmacokinetic and Pharmacodynamic Analysis of Subcutaneous Tocilizumab in Patients With Rheumatoid Arthritis From 2 Randomized, Controlled Trials: SUMMACTA and BREVACTA. *J Clin Pharmacol*. 2017;57(4):459-68.

16. Food and Drug Administration. ACTEMRA (Tocilizumab): Clinical Pharmacology and Biopharmaceutics review(s) [Last accessed November 13, 2019]. Available from: https://www.accessdata.fda.gov/drugsatfda_docs/nda/2013/125472Orig1s000ClinPharmR.pdf.
17. Huizinga TW, Fleischmann RM, Jasson M, Radin AR, van Adelsberg J, Fiore S, et al. Sarilumab, a fully human monoclonal antibody against IL-6 α in patients with rheumatoid arthritis and an inadequate response to methotrexate: efficacy and safety results from the randomised SARIL-RA-MOBILITY Part A trial. *Ann Rheum Dis*. 2014;73(9):1626-34.
18. Emery P, Rondon J, Parrino J, Lin Y, Pena-Rossi C, van Hoogstraten H, et al. Safety and tolerability of subcutaneous sarilumab and intravenous tocilizumab in patients with rheumatoid arthritis. *Rheumatology (Oxford)*. 2018;58:849–58.
19. Burmester GR, Rubbert-Roth A, Cantagrel A, Hall S, Leszczynski P, Feldman D, et al. A randomised, double-blind, parallel-group study of the safety and efficacy of subcutaneous tocilizumab versus intravenous tocilizumab in combination with traditional disease-modifying antirheumatic drugs in patients with moderate to severe rheumatoid arthritis (SUMMACTA study). *Ann Rheum Dis*. 2014;73(1):69–74.
20. Kivitz A, Olech E, Borofsky M, Zazueta BM, Navarro-Sarabia F, Radominski SC, et al. Subcutaneous tocilizumab versus placebo in combination with disease-modifying antirheumatic drugs in patients with rheumatoid arthritis. *Arthritis Care Res (Hoboken)*. 2014;66(11):1653-61.
21. Mihara M, Kasutani K, Okazaki M, Nakamura A, Kawai S, Sugimoto M, et al. Tocilizumab inhibits signal transduction mediated by both mIL-6R and sIL-6R, but not by the receptors of other members of IL-6 cytokine family. *International immunopharmacology*. 2005;5(12):1731-40.
22. Gibiansky L, Gibiansky E. Target-mediated drug disposition model: approximations, identifiability of model parameters and applications to the population pharmacokinetic-pharmacodynamic modeling of biologics. *Expert opinion on drug metabolism & toxicology*. 2009;5(7):803-12.
23. Liang M, Schwickart M, Schneider AK, Vainshtein I, Del Nagro C, Standifer N, et al. Receptor occupancy assessment by flow cytometry as a pharmacodynamic biomarker in biopharmaceutical development. *Cytometry Part B, Clinical cytometry*. 2016;90(2):117-27.
24. Emery P, Rondon J, Parrino J, Lin Y, Pena-Rossi C, van Hoogstraten H, et al. Safety and tolerability of subcutaneous sarilumab and intravenous tocilizumab in patients with rheumatoid arthritis. *Rheumatology (Oxford)*. 2018.

FIGURES AND TABLES

FIG 1 Blockade of classical IL-6R α signalling by sarilumab and tocilizumab in (A) proliferation assay in DS-1 cells, and (B) STAT3 signalling in HepG2/STAT3-Luc cells, and sarilumab blockade of *trans* IL-6R α signalling in HEK293/gp130/STAT3-Luc cells exposed to 10 nM hIL-6 and (C) 1 nM hIL-6R α or (D) 10 nM hIL-6R α

FIG 2 Simulated receptor occupancy (RO) profile through Week 24 for (A) sarilumab SC vs tocilizumab IV and (B) sarilumab SC vs tocilizumab SC


FIG 3 Relationship between receptor occupancy (RO) and (A) CRP reduction, (B) DAS28-CRP reduction, and (C) ACR20/50/70 responses with placebo and sarilumab (MOBILITY Part A)

FIG 4 Observed mean CRP levels (\pm SE) in patients with RA treated with sarilumab SC or tocilizumab IV (ASCERTAIN)

FIG 5 Mean changes from baseline in CRP for patients treated with tocilizumab SC and IV (SUMMACTA, BREVACTA), reprinted under CC BY-NC-ND 4.0 from Fig. 3 by *The Journal of Clinical Pharmacology*


TABLE 1 Pharmacokinetic and binding parameters for sarilumab and tocilizumab to monomeric (mmH-tagged) and dimeric (mFc-tagged) human IL-6R α proteins

FIG. 1 Blockade of classical IL-6R α signalling by sarilumab and tocilizumab in (A) proliferation assay in DS-1 cells, and (B) STAT3 signalling in HepG2/STAT3-Luc cells, and sarilumab blockade of *trans* signalling in HEK293/gp130/STAT3-Luc cells exposed to 10 nM hIL-6 and (C) 1 nM hIL-6R α or (D) 10 nM hIL-6R α




EC₅₀, effective concentration at 50% activity; HepG2, human hepatocellular carcinoma cell line HepG2; hIgG1, human immunoglobulin G 1; hIL-6, human interleukin-6; IC₅₀ inhibitory concentration at 50% activity; IL-6R α , interleukin-6 receptor alpha subunit; O.D., optical density; RLU, relative luminescence units; STAT3, signal transducer and activator of transcription 3.

FIG. 2 Simulated receptor occupancy (RO) profile through Week 24 for (A) sarilumab SC vs tocilizumab IV and (B) sarilumab SC vs tocilizumab SC



IV, intravenous; q2w, every two weeks; q4w, every four weeks; SC, subcutaneous.


FIG. 3 Relationship between receptor occupancy (RO) and (A) CRP reduction, (B) DAS28-CRP reduction, and (C) ACR20/50/70 responses with placebo and sarilumab (MOBILITY Part A)



Patients received placebo or sarilumab 100 mg q2w, 150 mg q2w, 200 mg q2w, 100 mg qw, or 150 mg q2w in MOBILITY Part A. Changes in free sIL-6R levels were used to calculate receptor occupancy, and associations were assessed against clinical parameters.

ACR, American College of Rheumatology; ACR20/50/70, 20%/50%/70% improvement in American College of Rheumatology criteria; CRP, C-reactive protein; DAS28-CRP, Disease Activity Score of 28 joints using C-reactive protein; RO, receptor occupancy.


FIG. 4 Observed mean CRP levels (\pm SE) in patients with RA treated with sarilumab SC or tocilizumab IV (ASCERTAIN)



Number of patients	0	4	8	12	24
Sarilumab 150 mg	49	47	45	44	39
Sarilumab 200 mg	51	47	51	43	39
Tocilizumab 4 mg/kg	40	37	37	37	35
Tocilizumab 4 to 8 mg/kg	39	39	39	39	39

^aPatients started at 4 mg/kg followed by an increase to 8 mg/kg, based on clinical response; ASCERTAIN (NCT01768572); CRP, C-reactive protein; IV, intravenous; q2w, every two weeks; q4w, every four weeks; RA, rheumatoid arthritis; SC, subcutaneous; SE, standard error.

FIG. 5 Observed mean CRP levels (\pm SE) in patients with RA treated with tocilizumab SC and IV (SUMMACTA, BREVACTA) (Abdallah H, et al. 2017), is reprinted under CC BY-NC-ND 4.0 from Fig. 3 by *The Journal of Clinical Pharmacology*



Mean CRP levels following treatment with SC or IV tocilizumab from Fig. 3 by *The Journal of Clinical Pharmacology* is licensed under CC BY-NC-ND 4.0; SUMMACTA (NCT01194414); BREVACTA (NCT1232569); CRP, C-reactive protein; IV, intravenous; RA, rheumatoid arthritis; SC, subcutaneous; SE, standard error.

TABLE 1 Pharmacokinetic and binding parameters for sarilumab and tocilizumab to monomeric (mmH-tagged) and dimeric (mFc-tagged) human IL-6R α proteins, determined by SPR-Biacore

Antigen	Antibody	ka (1/Ms)	kd (1/s)	K _D (M)	t _{1/2} (h)
Human IL-6R α monomeric	Sarilumab	8.56e ⁵	5.30e ⁻⁵	6.19e ⁻¹¹	3.6
	Tocilizumab	1.60e ⁵	2.14e ⁻⁴	1.34e ⁻⁹	0.9
Human IL-6R α dimeric	Sarilumab	4.02e ⁵	5.16e ⁻⁶	1.28e ⁻¹¹	37.3
	Tocilizumab	7.48e ⁴	1.47e ⁵	1.96e ⁻¹⁰	13.1

Supplementary appendix

Christine Xu et al. Differential binding of sarilumab and tocilizumab to IL-6R α and effects of receptor occupancy on clinical parameters

Supplementary methodology

Supplementary FIG S1. Sarilumab and tocilizumab sIL-6R α receptor occupancy models

Supplementary FIG S2. Ligand-binding properties of sarilumab and tocilizumab. Representative sensograms of (A) sarilumab binding to monomer human IL-6R α -mmH 10, 5, 2.5, 1.25, and 0.625 nM, and (B) tocilizumab binding to monomer human IL-6R α -mmH 20, 10, 5, 2.5, 1.25, and 0.625 nM, are shown as black lines. The data were globally fit to a 1:1 binding interaction model using T200 evaluation software 2.0. Kinetic fits from the analyses were overlaid on the binding data in red

Supplementary FIG S3. Blockade of dimeric hIL-6R α binding to IL-6 (ELISA competition assay). (A) Dose–response of hIL-6R α -hFc binding to hIL-6 and (B) concentration-dependent blockade of hIL-6R α -hFc binding to hIL-6 by sarilumab

Supplementary FIG S4. Basic goodness-of-fit plots with locally weighted scatterplot smoothing (LOWESS) (red lines) for the final model. $|iWRES|$ individual weighted residuals

Supplementary FIG S5. Final model visual predictive check after multiple doses of sarilumab 100 mg q2w, 150 mg q2w, 200 mg q2w, 100 mg qw or 150 mg qw

Supplementary Table S1. Parameter estimates of the sIL-6R α population PK/PD model

Supplementary methodology

Kinetic binding analysis

The binding kinetics of sarilumab and tocilizumab to IL-6R α were measured using Surface Plasmon Resonance (SPR; Biacore™ T200). Sarilumab and tocilizumab were captured on an anti-human Fc-coupled chip surface and human IL-6R α (hIL-6R α) flowed across the surface (at concentrations of 20 nM to 1.25 nM, depending on the antigen used). Antigen-dependent changes in resonance units (reflecting binding to the captured antibody) were monitored, from which binding kinetics were calculated, including association rate constant (k_a), dissociation rate constant (k_d), and half-life ($t_{1/2}$). The overall equilibrium dissociation constant (K_D) was calculated from the ratio of k_d to k_a .

Blockade of dimeric hIL-6R α binding to IL-6 (ELISA competition assay)

A 3-fold dilution series of sarilumab or an isotype control IgG1 antibody (30 nM to 0.5 pM) were pre-incubated for 1 hour with 100 pM of dimeric hIL-6R α with a C-terminal human IgG1 Fc tag (IL-6R α -hFc), after which the mixtures were transferred to 96-well microtitre plates onto which hIL-6 (2 μ g/mL) had been immobilized. Bound IL-6R α -hFc/IL-6 complexes were detected with a horseradish peroxidase-conjugated anti-hIL-6 Fc antibody (Jackson ImmunoResearch). Plates were visualized with 3,3', 5,5' tetramethylbenzidine (BD Biosciences) and absorbance determined using a Victor Multilabel counter (Perkin Elmer). IC₅₀ and EC₅₀ values, the concentration of drug resulting in half-maximal inhibition or response, respectively, were determined (GraphPad Prism™ v6) using a four-parameter logistic model.

Inhibition of classical IL-6R α signalling (cell proliferation and STAT3 response element activation assays)

The activity of sarilumab and tocilizumab in blocking classical IL-6R α signalling was compared *in vitro* in cell proliferation assays using:

- DS-1: a human B lymphocyte cell line that proliferates in response to exogenous hIL-6 and endogenously expresses IL-6R α and gp130

- HepG2: a hepatocytic cell line endogenously expressing IL-6R α and gp130

Serial dilutions of sarilumab (100 nM to 1.7 pM) and tocilizumab (100 nM to 1.7 pM) were added to DS-1 cells (ATCC, CRL-11102), followed by hIL-6 (1 pM). Plates were incubated (37°C, 5% CO₂ for 4 days) and then visualized using AlamarBlue (Biosource) or WST-8 (Dojindo).


In the second assay, HepG2 (ATCC, HB-8065) was transiently transfected with a STAT3-luciferase reporter plasmid. Dilution series of sarilumab (100 nM to 1.7 pM) and tocilizumab (100 nM to 1.7 pM) were added to the transfected cells seeded in 96-well plates, followed by hIL-6 (50 pM). Plates were incubated (6 hours at 37°C, 5% CO₂) and visualized using Steady-Glo or One-Glo luciferase substrate.

In both assays, plates were read on a Victor X5 multilabel counter, and EC₅₀ and IC₅₀ values calculated as described above.

Inhibition of *trans*-IL-6R α signalling (luciferase assay)


The ability of sarilumab to block *trans* signalling stimulated by a soluble complex of IL-6 and IL-6R α was assessed in a functional cell-based luciferase assay using a gp130-overexpressing human embryonic kidney/STAT3/luciferase reporter cell line. Serial dilutions of sarilumab (200 nM–3.4 pM) were pre-incubated with 1 nM of monomeric sIL-6R α with a C-terminal myc-myc-hexahistidine tag (IL-6R α -mmH), and transferred together with hIL-6 (12.5 nM) to 96-well plates seeded with HEK293/gp130/STAT3-Luc cells (ATCC CRL-1573). For the dose–response curve, hIL-6 concentrations ranging from 15 nM to 2.5 pM were used along with human monomeric sIL-6R α -mmH (1 nM). Following incubation (5 hours at 37°C, 5% CO₂), response was measured using One-Glo luciferase substrate (Promega) and read on the Victor X5 multilabel counter.

Supplementary FIG S1 Sarilumab and tocilizumab sIL-6R α receptor occupancy models



C, concentration; CL_0 , apparent clearance; F, bioavailability; K_{deg} , degradation rate constant; K_{int} , internalization rate constant; K_m , Michaelis–Menten constant; K_{off} , dissociation rate constant; K_{on} , association rate constant; K_{syn} , synthesis rate constant; mAb, monoclonal antibody; sIL-6R, soluble form of the interleukin-6 receptor; V_c , apparent volume of central compartment; V_m , maximum elimination rate.


Supplementary FIG S2 Ligand-binding properties of sarilumab and tocilizumab. Representative sensograms of (A) sarilumab binding to monomer human IL-6R α -mmH 10, 5, 2.5, 1.25, and 0.625 nM, and (B) tocilizumab binding to monomer human IL-6R α -mmH 20, 10, 5, 2.5, 1.25, and 0.625 nM, are shown as black lines. The data were globally fit to a 1:1 binding interaction model using T200 evaluation software 2.0. Kinetic fits from the analyses were overlaid on the binding data in red.



Sensogram of binding kinetics of sarilumab binding to monomer IL-6R α proteins; (107 RU) sarilumab was captured on an anti-human Fc-coupled chip surface. Human IL-6R α proteins were tested in duplicate in a 2-fold dilution series, the association phase of human IL-6R α was monitored at 50 μ l/min for 5 minutes over each of the captured surfaces.

IL-6R, interleukin-6 receptor; RU, resonance units; SPR, surface plasmon resonance.


Supplementary FIG S3 Blockade of dimeric hIL-6R α binding to IL-6 (ELISA competition assay). (A) Dose-response of hIL-6R α -hFc binding to hIL-6 and (B) concentration-dependent blockade of hIL-6R α -hFc binding to hIL-6 by sarilumab



hIgG1, human immunoglobulin G1; hIL-6, human interleukin-6; hIL-6R α -hFc, recombinant extracellular domain of human IL-6R α generated with an N-terminal amino acid linker sequence comprising the Fc region of human immunoglobulin G 1 (IgG1); IL-6R α , interleukin-6 receptor alpha subunit.


Supplementary FIG S4 Basic goodness-of-fit plots with locally weighted scatterplot smoothing (LOWESS) (red lines) for the final model. $|iWRES|$ individual weighted residuals

Basic goodness-of-fit plots



Supplementary FIG S5 Final model visual predictive check after multiple doses of sarilumab

100 mg q2w, 150 mg q2w, 200 mg q2w, 100 mg qw or 150 mg qw



In these figures, solid and dashed red lines connect the median and bounds (i.e., 2.5th and 97.5th percentiles) of observed concentrations at each time point. Red and blue rectangles represent the median and bounds, respectively, of predicted concentrations at each time point.

Supplementary TABLE S1 Parameter estimates of the sIL-6R population PK/PD model

Parameter	Estimate	RSE %	95% CI
K_{syn} (ng/mL/day)	40.1	7.79	33.8–46.3
K_{dge} (1/h)	0.734	9.16	0.600–0.871
I_{max}	0.891	0.650	0.879–0.902
K_{ss} (μ g/mL)	0.264	22.4	0.146–0.381
Inter-individual variability (CV %)			
K_{syn}	47.8	19.3	37.4–56.2
K_{ss}	158	49.3	186–223
Residual variability (CV %)			
Proportional	30.9	10.8	27.4–34.1

CI, confidence interval; CV, coefficient of variation; I_{max} , maximum intensity; K_{deg} , degradation rate constant; K_{ss} , steady-state constant; K_{syn} , synthesis rate constant; mAb, monoclonal antibody; RSE, relative standard error; sIL-6R, soluble form of the interleukin-6 receptor.