

INFORME BREVE

Differential effects of two strains of *Rhizophagus intraradices* on dry biomass and essential oil yield and composition in *Calamintha nepeta*

Roxana P. Colombo^{a,*}, Alicia E. Martínez^a, Agustina Fernández di Pardo^a,
Laura Fernández Bidondo^a, Catalina van Baren^b, Paola di Leo Lira^b, and Alicia M. Godeas^a

^aDepartamento de Biodiversidad y Biología Experimental, Facultad de Ciencias Exactas y Naturales (FCEyN), Universidad de Buenos Aires (UBA), Ciudad Autónoma de Buenos Aires, Argentina

^bCátedra de Farmacognosia-IQUIMEFA (UBA-CONICET), Facultad de Farmacia y Bioquímica, Universidad de Buenos Aires (UBA), Ciudad Autónoma de Buenos Aires, Argentina

Received 25 September 2012 ; accepted 15 May 2013

KEYWORDS

Arbuscular mycorrhizal symbiosis;
Calamintha nepeta;
Essential oil yield;
Essential oil composition

Abstract

The aim of this work was to determine the effects of two geographically different strains of *Rhizophagus intraradices* (M3 and GA5) on the total biomass and essential oil (EO) yield and composition of *Calamintha nepeta*, with or without phosphorus (P) fertilization, under greenhouse conditions. The plant biomass was not significantly affected by any of the treatments, showing higher values in control plants. Strains had a differential response in their root colonization rates: M3 reduced these parameters while GA5 did not modify them. Both strains affected EO yield in absence of P fertilization: M3 promoted EO yield in *C. nepeta* plants and GA5 resulted in negative effects. The percentage composition of EO was not significantly modified by either strain or P fertilization. M3 strain could be a potential fungal bioinoculant for production and commercialization of *C. nepeta* in the aromatic plant market.

© 2013 Asociación Argentina de Microbiología. Published by Elsevier España, S.L. All rights reserved.

PALABRAS CLAVE

Simbiosis micorrízica arbuscular;
Calamintha nepeta;
Rendimiento de aceites esenciales;
Composición de aceites esenciales

Efecto diferencial de dos cepas de *Rhizophagus intraradices* sobre la biomasa y el rendimiento y composición de aceites esenciales de *Calamintha nepeta*

Resumen

El objetivo de este trabajo fue determinar, bajo condiciones de invernadero, el efecto de dos cepas geográficamente diferentes de *Rhizophagus intraradices* (M3 y GA5) sobre la biomasa total y el rendimiento y composición de aceites esenciales (AE) de *Calamintha nepeta*, con fertilización fosforada (P) o sin esta. La biomasa de la planta no fue signifi-

*Corresponding author.

E-mail address: rcolombo@soilgene.net (R.P. Colombo).

cativamente afectada por ningún tratamiento, y se observaron valores más altos en las plantas control. Las cepas mostraron diferencias en sus tasas de colonización y en las respuestas a la fertilización con fósforo: M3 redujo sus valores de colonización, mientras que GA5 no los modificó. En ausencia de fertilización fosforada, las plantas colonizadas por ambas cepas presentaron rendimientos de AE diferentes a aquellos de las plantas control: M3 los aumentó y GA5 los disminuyó. La composición porcentual de AE no fue modificada significativamente por ninguno de los tratamientos. M3 podría ser considerada como un posible bioinoculante fúngico para la producción de *C. nepeta* destinada al mercado de las plantas aromáticas.

© 2013 Asociación Argentina de Microbiología. Publicado por Elsevier España, S.L. Todos los derechos reservados.

Calamintha nepeta subsp. *nepeta* (Lamiaceae), which is native to the Mediterranean basin, is an introduced species of aromatic plants in Argentina. It is a domesticated plant used for herbal infusions¹⁰ under the same commercial name of the native species *Minthostachys mollis*, 'peperina', due to their similar flavor and aroma¹⁵. *M. mollis* is nowadays becoming increasingly scarce because of its overexploitation¹⁴. The essential oils (EO) of aromatic plants such as *C. nepeta*, are volatile lipophilic mixtures of secondary plant compounds, mainly used in food, cosmetic and pharmaceutical industries². EO yield is strongly influenced by biotic and abiotic factors⁶.

It has been shown that arbuscular mycorrhizal (AM) symbiosis induces changes in secondary compounds which act as signal molecules in plant-AM fungal interaction¹². Previous studies have been focused on different AM fungal effects on several aromatic plants: biomass production^{7,13}, EO yield and composition⁶ and nutrient uptake^{7,13}. It has been demonstrated that different species and isolates of AM fungi have diverse effects on mycorrhizal plants and that different plant species or varieties react differently to the same AM isolates⁸. However, to our knowledge there are no reports about the potential effect of AM mycorrhization on EO yield and composition in *C. nepeta* plants.

The purpose of this work was to evaluate the effect of two geographically different strains of *R. intraradices* on plant growth response and EO yield and composition of *C. nepeta*, with or without phosphorous (P) fertilization, under greenhouse conditions. We also aim at improving the quality and quantity of the EO of *C. nepeta* in order to promote its commercial production, protecting the endangered native plant species.

A *C. nepeta* plant used for the assay was collected in Jujuy province, Argentina. A single healthy and vigorous plant of *C. nepeta* was chosen. Cuttings (10 cm) were collected from the mother plant in order to avoid genetic variability in the experiment. Cuttings were established in sterile perlite until the differentiation of adventitious roots. Rooted cuttings were transplanted [1 plant per 500 ml pot with 300 ml autoclaved substrate (100 °C for 1 h, three consecutive days)] into a mixture of soil:perlite (1:2) as growth substrate. The soil used had the following features: clay loam soil; pH 7.1; total C: 12.08 and N: 1.1 (g/Kg); P: 34.2 (mg/Kg); K: 0.9, Ca 7.5, Mg: 1.7 and Na: 0.2 (cmol/Kg).

Each plant was placed in a hole where the inoculum had been previously added. The inoculum, consisted of colonized roots of *Trifolium* sp. (over 70% of AM fungal colonization)

and soil with hyphae and spores, previously quantified in order to use the same amount for each pot (approx. 50 spores per plant). Two different strains of *R. intraradices* were used in this experiment: GA5 [Banco de Glomeromycota in Vitro (BGIV) collection of the Laboratorio de Microbiología del Suelo, Facultad de Ciencias Exactas y Naturales, UBA, (<http://www.bgiv.com.ar/strains/Rhizophagus-intraradices/ga5>)] and M3 [collection of the Laboratorio de Microbiología del Suelo, Facultad de Ciencias Exactas y Naturales, UBA]. The latter strain was isolated from the same geographic region of *C. nepeta*.

Pots were grown in a random design for 60 days in a greenhouse with controlled temperature and humidity conditions at the Facultad de Ciencias Exactas y Naturales, Universidad de Buenos Aires. All the plants were watered daily. Ten milliliters (10 ml) of PO₄H₂Na (0.05 g/l) were added to the corresponding treatment once a week. The P solution concentration is consistent with that routinely used in Hewitt's nutrient solution⁵. Treatments were: 1- C (Control): perlite-sterilized soil substrate; 2- CP: perlite-sterilized soil substrate and P fertilization; 3- M3: perlite-sterilized soil substrate inoculated with M3; 4- M3P: perlite-sterilized soil substrate inoculated with M3 and P fertilization; 5- GA5: perlite-sterilized soil substrate inoculated with GA5; 6- GA5P: perlite-sterilized soil substrate inoculated with GA5 and P fertilization. Three replicates (each replicate was composed by ten pots) were performed for each treatment.

Root samples were taken from each treatment after 30 days of growth to evaluate the success of mycorrhizal colonization. After 60 days of growth, plants were harvested and air-dried in a dark room for their final processing and measurement of biomass yield. Biomass was significantly higher in control plants than in mycorrhized ones ($F = 12.822$, $p < 0.05$), particularly when P fertilization was performed (C: 12.65 g, CP: 14.96 g). Mean values correspond to plants inoculated with M3 strains (M3: 11.28 g, M3P: 10.85 g), while the lowest biomass yields were those of plants colonized by GA5, especially when P fertilization was performed (GA5: 10.47 g, GA5P: 8.33 g). Nevertheless, the observed differences between strains resulted non-significant when they were statistically analyzed (Fig. 1A).

AM fungal colonization was also evaluated at this time. Root samples were cleared in KOH (10%) for 24 h at room temperature and then stained with trypan blue in lactic acid (0.02%) for 24 h at room temperature (modified procedure of




Figure 1 Effect of phosphorus fertilization and inoculation of arbuscular mycorrhizal strains, M3 and GA5, on: (A) biomass yield, and (B) essential oil yield, in *Calamintha nepeta*. C represents values for control plants. Black bars correspond to non-phosphorus fertilized treatments. Grey bars correspond to phosphorus-fertilized treatments. Mean values with error bars. Different letters represent significant differences ($p < 0.05$). EO: essential oil.

Phillips and Hayman's technique¹¹). Mycorrhization was detected in all treatments, except in control plant roots. Intraradical colonization was quantified from 60 randomly selected root segments (1 cm long) per treatment. Frequency (%F) and intensity (%I) of mycorrhizal colonization was calculated according to Declerck *et al.*³ A Nikon light binocular microscope (model: Optiphot-2) was used for this purpose.

Our results showed that P fertilization had different effects on root colonization by both *R. intraradices* strains tested. Mycorrhization frequency (Fig. 2A) reached values above 70% but no significant differences between treatments were found ($F = 0.355$, $p = 0.556$). The GA5 strain showed the highest colonization frequency, particularly when P fertilization was performed (GA5: 80.65%; GA5P: 85.35%). P-fertilized roots inoculated with the M3 strain presented the lowest mycorrhization frequency (M3: 72.50%; M3P: 68.75%). Mycorrhization intensity (Fig. 2B) resulted significantly higher in plants inoculated with GA5 ($F = 29.361$, $p < 0.05$), especially when plants were not P-fertilized (GA5: 67.69%; GA5P: 65.70%). The lowest intensity percentages corresponded to P-fertilized plants inoculated with the M3 strain (M3: 48.78%; M3P: 26.42%), coinciding with mycorrhization frequency results. As expected, the intensity of root colonization by M3 decreased with P fertilization; however this difference was not observed in GA5 treatments.

Ryan *et al.* observed that P levels in soil and host plant were responsible for the degree of root colonization by AM fungi. It is possible that genetic variations between *R. intraradices* strains could result in significant functional variations⁸ and, therefore, in a differential response to P fertilization.

In the present study, we observed that the lowest value of plant biomass coincided with the highest level of AM fungal colonization. This may be a negative effect of the inoculation of GA5 on *C. nepeta* plants. Similar results were observed by Koch *et al.*⁸ when two genetically different *R. intraradices* strains negatively influenced the growth of *Daucus carota* transformed roots. These authors related this observation to the cost of AM fungal colonization for the plant⁸.

After natural drying, the plant stems were subjected to hydrodistillation for 3 h using a Clevenger-type trap. Extracted essences were dried with anhydrous sodium sulphate and stored in vials at -20°C for qualitative and quantitative analysis. EO yield of *C. nepeta* plants was calculated as: $(\text{EO volume} / \text{Plant dry weight}) \times 100$ ¹.

M3 strain significantly promoted EO yield (2.13%, $F = 43.346$, $p < 0.05$) (Fig. 1B). However plants inoculated with this strain and P-fertilized showed similar values to those of control plants (0.87%). The highest value coincided with the lowest frequency and intensity of AM fungal colonization. Otherwise, the lowest yield was obtained when *C. nepeta* was inoculated with GA5 (0.53%); these plants presented the highest values of fungal colonization.

Our results suggested a positive correlation between the level of M3 colonization and EO yield in *C. nepeta*. The opposite correlation was observed when GA5 was inoculated. Since it is known that different strains of *R. intraradices* have either favorable or harmful effects in host plant development⁸, this could be a reasonable explanation for the results in this study.

For qualitative and quantitative analysis of the EO, hyphenated gas chromatography with a flame ionization detector and mass spectrometer (GC-FID-MS) was performed on a Perkin Elmer Autosystem gas chromatograph (model: Clarus 500). GC conditions: Mobile phase: Helium 1.87 ml/min; autosampler connected to a single injector split (split relation: 1:100) connected by a flow splitter to two fused silica capillary columns (5% phenyl-95% dimethylpolysiloxane and polyethyleneglycol 20 000, both of 60 m x 0.25 μm with a film thickness of 0.25 μm); temperature program: 50 $^{\circ}\text{C}$, then 3 $^{\circ}\text{C}/\text{min}$ until 225 $^{\circ}\text{C}$ (15 min); injector and detector temperature FID: 255 $^{\circ}\text{C}$ and 275 $^{\circ}\text{C}$, respectively; transfer line temperature: 180 $^{\circ}\text{C}$; ion source temperature: 150 $^{\circ}\text{C}$; total run time: 70 min; scanned mass range: 40-400 m/z; injected volume: 0.2 μl of a 10% dilution in ethanol.

EO constituents were identified on the basis of their retention index and their MS, which were compared with those found in literature data and standards of the Cátedra de Farmacognosia (Facultad de Farmacia y Bioquímica,




Figure 2 Colonization of *Calamintha nepeta* roots by arbuscular mycorrhizal strains GA5 and M3 expressed as: (A) percentage (%) of intradical colonization frequency, and (B) percentage (%) of intradical colonization intensity. Black bars correspond to non-phosphorus fertilized treatments. Grey bars correspond to phosphorus-fertilized treatments. Mean values with error bars. Different letters represent significant differences ($p < 0.05$).

Universidad de Buenos Aires). The percentage composition was determined by the percentage of areas of each peak, without the correction factor. The lowest response for each component was taken among those obtained from each column used. A homologous series of methyl esters of fatty acids (C4-C18) was co-chromatographed with the oil for determination of the retention index.

The main compound contents in the EO of *C. nepeta* are shown in Table 1. Even when the total contents of each main compound seem to differ among treatments, these differences were not statistically significant. The percentage of menthone was always higher when P fertilization was performed with respect to the same treatments without P supplementation (e.g. C: 31.7%; CP: 33.2%). The opposite was observed with isomenthone (e.g. C: 14.8%; CP: 14.1%) and pulegone (e.g. C: 27%; CP: 23.9%). Finally, neomenthol followed the menthol tendency, except for treatments involving M3 inoculation. Menthone and isomenthone reached their highest values in control plants, while the percentage composition of neomenthol and pulegone was improved by GA5 and M3 inoculation respectively. These

percentage changes in EO composition resulted in a variation in the menthone/pulegone ratio. While non-mycorrhized and mycorrhized plants with P supplementation had a menthone/pulegone ratio greater than 1, the inoculated ones with no P supplementation presented a value lower than 1.

P fertilization had no influence in EO yield or composition in plants without fungal colonization. The supplementation of this nutrient, as performed in the present study, was not a limiting factor for these variables in *C. nepeta*. Our study partially supports Hammer *et al.*⁴ results, where P fertilization does not change the physical and chemical properties of EO in *C. nepeta*, but, the pulegone content is increased at high P rates. Nell *et al.*⁹ concluded that P fertilization improves biomass yield and secondary metabolite concentration in *Salvia officinalis* (garden sage), reaching higher values than AM-inoculated plants. Phosphorus is an important constituent of nucleic acids and phospholipids. For this reason, plants required it in large amounts for biosynthesis of primary and secondary metabolites. Besides P fertilization, AM fungal symbiosis

Table 1 Total content (μl) of each main compound in the essential oil extracted from *Calamintha nepeta* according to the different treatments

Main compounds	Total content in the extract (μl) [percentage composition (%)] ^a					
	C	CP	M3	M3P	GA5	GA5P
Menthone	41.21 [31.7]	39.84 [33.2]	81.76 [29.2]	32.7 [32.7]	17.34 [28.9]	17.46 [29.1]
Isomenthone	19.24 [14.8]	16.92 [14.1]	37.24 [13.3]	13 [13]	7.26 [12.1]	7.08 [11.8]
Neomenthone	12.87 [9.9]	17.4 [14.5]	26.6 [9.5]	9 [9]	7.38 [12.3]	7.92 [13.2]
Pulegone	35.1 [27]	28.68 [23.9]	98.28 [35.1]	30.8 [30.8]	18.24 [30.4]	13.56 [22.6]
EO total volume (μl)	130	120	280	100	60	60

EO, essential oil.

^aNumbers in brackets correspond to the percentage composition (%) of each main compound in the extract. No significant differences were observed among values ($p > 0.05$).

positively affects the P status of the plant⁹. In this context we expected that AM inoculation and P fertilization would have similar effects on EO yield and composition in *C. nepeta*. Although our results differ from those of Hammer *et al.*⁴ and Nell *et al.*⁹, further experiments with higher rates of P fertilization should be performed.

Two-way ANOVA was performed for all data. Treatment effects were analyzed using the Tukey HSD test ($p < 0.05$). Homogeneity of variances and normal distribution assumptions were checked with the Levene test and distribution of within-self residuals, respectively. All statistical analyses were performed with STATISTICA, version 7 StatSoft, Inc. (2004).

To our knowledge, this is the first study on the effect of AM fungal inoculation in *C. nepeta*. Further studies with more species and strains of AM fungi on this host plant are necessary to find out possible mutualistic interactions that promote both EO and dry biomass yields. The effect of AM fungi colonization on the EO of aromatic plants was studied in different host plants and, in every case, the response to colonization was different^{2,6,12,13}.

According to the interactions observed between the two fungal strains tested and *C. nepeta* plants, we conclude that some strains of *R. intraradices*, in this case the M3 strain, could be used as bioinoculants in *C. nepeta* production due to their positive inoculation effect on EO yield and non-modification on fragrance properties, without affecting total biomass. Furthermore, the commercialization of *M. molli* 'peperina' in the aromatic plants market is done by potting¹⁰, thus facilitating AM fungi inoculation.

Conflicts of interest

The authors declare that they have no conflicts of interest.

Acknowledgments

We wish to thank UBA, CONICET and ANCYPT for financial support. We also acknowledge Dra. Vanesa Silvani (Facultad de Ciencias Exactas y Naturales, UBA, Argentina), Dr. Arnaldo Bandoni (Cátedra de Farmacognosia-IQUIMEFA, UBA, Argentina), and Dr. Alexander N. Schmidt-Lebuhn (Institut für Systematische Botanik, Zurich, Switzerland), for their technical assistance.

References

1. Bandoni AL, Lopez MA, Juarez MA, Elechosa MA, Baren C, di Leo Lira P. Seasonal variations in the composition of the essential

- oil of *peperina* (*Minthostachys mollis* (Kunth) Griseb.) from a local population of the Province of Córdoba, Argentina. *Essenze Derivati Agrumari*. 2002;72:11-4.
2. Cabello M, Irrazabal G, Bucsinzky AM, Saparrat M, Schalamuk S. Effect of an arbuscular mycorrhizal fungus, *Glomus mosseae*, and a rock - phosphate - solubilizing fungus, *Penicillium thomii*, on *Mentha piperita* growth in a soilless medium. *J Basic Microb*. 2005;45:182-9.
3. Declerck S, Strullu DG, Plenchette C. *In vitro* mass-production of the arbuscular mycorrhizal fungus *Glomus versiforme* associated with Ri T-DNA transformed carrot roots. *Mycol Res*. 1996;100:1237-42.
4. Hammer K, Laghetti G, Pistrick K. *Calamintha nepeta* (L.) Savi and *Micromeria thymifolia* (Scop.) Fritsch cultivated in Italy. *Genet Resour Crop Ev*. 2004;52:215-20.
5. Hewitt EJ. Sand water culture methods used in the study of plant nutrition. Commonwealth Agriculture Bureau. 1952; Techn Commun. 22.
6. Kapoor R, Giri B Mukerji KG. *Glomus macrocarpum*: a potential bioinoculant to improve essential oil quality and concentration in Dill (*Anethum graveolens* L.) and Carum [*Trachyspermum ammi* (Linn.) Sprague]. *World J Microbiol Biotechnol*. 2002;18:459-63.
7. Karagiannidis N, Thomidis T, Lazari D, Panou-Filotheou E, Karagiannidou C. Effect of three Greek arbuscular mycorrhizal fungi in improving the growth, nutrient concentration, and production of essential oils of oregano and mint plants. *Sci Hort*. 2011;129:329-34.
8. Koch AM, Croll DP, Sanders IR. Genetic variability in a population of arbuscular mycorrhizal fungi causes variation in plant growth. *Ecology Letters*. 2006;9:103-10.
9. Nell M, Vötsch M, Vierheilig H, Steinkellner S, Zitterl-Eglseer K, Chlodwig F, Novak J. Effect of phosphorus uptake on growth and secondary metabolites of garden sage (*Salvia officinalis* L.). *J Sci Food Agric*. 2009;89:1090-6.
10. Ordóñez A, Baldoncini S, Berioli G, Chaves G, Bled L, Massuh Y, Liébana C, Torres L, Ojeda M. Domestication of native aromatic plants. *Mol Med Chem*. 2006;11:58-9.
11. Phillips JM, Hayman DS. Improved procedures for clearing roots and staining parasitic and vesicular-arbuscular mycorrhizal fungi for rapid assessment of infection. *Trans Br Mycol Soc*. 1970;55:158-60.
12. Ponce MA, José M. Scervino, Rosa Erra-Balsells, Juan A. Ocampo, Alicia M. Godeas. New flavonoids isolated from shoots and roots of *Trifolium repens* (white clover) grown with and without the mycorrhizal (AM) fungus *Glomus intraradices*. *Phytochemistry*. 2004;65:1925-30.
13. Rasouli-Sadaghian M, Hassani A, Barin M, Danesh YR, Sefidkon F. Effects of arbuscular mycorrhizal (AM) fungi on growth, essential oil production and nutrients uptake in basil. *J Med Plant Res*. 2010;4:2222-8.
14. Ryan MH, Small DR, Ash JE. Phosphorus controls the level of colonization by arbuscular mycorrhizal fungi in conventional and biodynamic irrigated dairy pastures. *Aust J Exp Agr*. 2000;40:663-70.
15. Scandaliaris M, Fuentes, E, Lovey RJ. Dos especies de Lamiáceas comercializadas en Córdoba (Argentina) bajo el nombre de "Peperina". *Multequina*. 2007;16:73-81.