

DIFFERENTIAL FORMS, WEITZENBÖCK FORMULAE AND FOLIATIONS

HANSKLAUS RUMMLER

Abstract

The Weitzenböck formulae express the Laplacian of a differential form on an oriented Riemannian manifold in local coordinates, using the covariant derivatives of the form and the coefficients of the curvature tensor. In the first part, we shall describe a certain "differential algebra formalism" which seems to be a more natural frame for those formulae than the usual calculations in local coordinates.

In this formalism, there appear some interesting differential operators which may also be used to characterize local geometric properties of foliations. That is the topic of the second part.

I. Differential Forms and Weitzenböck Formulae

1. Introduction.

Relations between the curvature tensor of a compact oriented Riemannian manifold and its topology play an important role in global differential geometry. The oldest and still most beautiful theorem of this kind is of course the global Gauß-Bonnet formula. Another theorem of this type is the following one by S. Bochner (see [3],[1]): If all sectional curvatures are positive, then the first Betti number must vanish. To prove this theorem, one can for instance show that there are no non-trivial harmonic 1-forms, and in order to derive this from the positivity of sectional curvature, one uses the so-called Weitzenböck formulae, expressing the Laplacian of a differential form in local coordinates by an expression which contains coefficients of the curvature tensor. In order to see, how the curvature tensor comes in, we shall develop an appropriate formalism in the next two paragraphs.

The author would like to thank the organizers of the Differential Geometry Semester in autumn 1988 at the Centre de Recerca Matemàtica of the Institut d'Estudis Catalans, where this article was written.

2. Ordinary and Vector-valued Differential Forms.

Let M be a differentiable (i.e. C^∞) manifold, TM its tangent bundle and T^*M the cotangent bundle. A *vector-valued* p -form Φ on M is a differentiable (i.e. C^∞) section in the bundle $TM \otimes \wedge^p T^*M$. Locally, Φ may be written as

$$\Phi = \sum_{i=1}^n \varphi_i \otimes X_i$$

with ordinary p -forms φ_i and vector fields X_i . For $x \in M$ one can interpret $\Phi(x)$ as a p -linear alternating map from $T_x M \times \dots \times T_x M$ to $T_x M$.

The vector space of these forms will be denoted by $\vec{A}^p = \vec{A}^p(M)$, and we shall write $A^p = A^p(M)$ for that of ordinary p -forms. $\vec{A} = \bigoplus_{p=0}^n \vec{A}^p$ is in a natural way an A -module, i.e. there are exterior products

$$\wedge : A^p \times \vec{A}^q \longrightarrow \vec{A}^{p+q} \quad \text{and} \quad \wedge : \vec{A}^q \times A^p \longrightarrow \vec{A}^{p+q},$$

with the obvious commutation rules.

Now let M be a Riemannian manifold. Then the Riemannian metric induces also an "exterior inner product"

$$\langle \quad , \quad \rangle : \vec{A}^p \times \vec{A}^q \longrightarrow A^{p+q},$$

and there is also a Hodge \star -operator

$$\star : \vec{A}^p \longrightarrow \vec{A}^{n-p},$$

locally defined by $\star(\sum \varphi_i \otimes X_i) = \sum (\star \varphi_i) \otimes X_i$.

Let D denote the Levi-Civita connection on M . It can be interpreted as a map $D : \vec{A}^0 \longrightarrow \vec{A}^1$, and this map is extended in a unique way by maps $D : \vec{A}^p \longrightarrow \vec{A}^{p+1}$, such that the product rule

$$D(\varphi \wedge \Psi) = d\varphi \wedge \Psi + (-1)^p \varphi \wedge D\Psi$$

holds for $\varphi \in A^p$ and $\Psi \in \vec{A}^q$.

This can be done for any linear connection on TM , but the Levi-Civita connection satisfies also the product rule for our "exterior inner product":

$$d\langle \Phi, \Psi \rangle = \langle D\Phi, \Psi \rangle + (-1)^p \langle \Phi, D\Psi \rangle$$

for $\Phi \in \vec{A}^p$ and $\Psi \in \vec{A}^q$.

In this formalism, the curvature tensor is simply given by

$$R_{XY}Z = D^2Z(X, Y) \quad \text{for} \quad X, Y, Z \in \vec{A}^0,$$

and also the torsion (which vanishes in one case) has a simple interpretation: $T_D = \text{D}id_{TM}$, where the identity on the tangent bundle TM is interpreted as a vector-valued 1-form.

The Riemannian metric induces also several maps between ordinary and vector-valued differential forms, which we shall now describe: The first one consist of maps

$$j : A^p \longrightarrow \bar{A}^{p-1}$$

which extend the canonical isomorphism between 1-forms and vector fields. j is simply defined by $\langle j\omega, X \rangle = i_X\omega$ for $\omega \in A^p$ and $X \in \bar{A}^0$, where i_X denotes the interior derivative. j is a derivation of degree -1 , i.e. it satisfies the product rule

$$j(\varphi \wedge \psi) = j\varphi \wedge \psi + (-1)^p \varphi \wedge j\psi$$

for $\varphi \in A^p$ and $\psi \in A^q$. We can as well extend the inverse isomorphism $\alpha = j^{-1} : \bar{A}^0 \longrightarrow A^1$ by

$$\alpha\Phi(X_1, \dots, X_{p+1}) = \sum_{i=1}^{p+1} (-1)^{i+1} \langle X_i, \Phi(X_1, \dots, \hat{X}_i, \dots, X_{p+1}) \rangle.$$

The next map is the contraction with respect to the first argument, i.e. the map "trace" $tr : \bar{A}^p \longrightarrow A^{p-1}$, defined by

$$tr \Phi(X_1, \dots, X_{p-1}) = \text{trace} (\Phi(-, X_1, \dots, X_{p-1}) : TM \longrightarrow TM),$$

and the last map is a sort of a "diagonal" $diag : A^p \longrightarrow \bar{A}^{p+1}$, an extension of the map from A^0 to \bar{A}^1 that sends the constant 1 to the identity id_{TM} .

We shall not derive all the relations between those maps, but we mention that the following two diagrams commute up to sign:

$$\begin{array}{ccc} A^p & \longleftarrow & \longrightarrow & A^{n-p} & & A^p & \longleftrightarrow & A^{n-p} \\ & & & & \text{and} & & & \\ \bar{A}^{p+1} & \longleftrightarrow & \bar{A}^{n-p-1} & & & \bar{A}^{p-1} & \longleftarrow & \longrightarrow & \bar{A}^{n-p+1} \end{array}$$

3. Gradient, Divergence and Laplacian.

In our language, the classical operators $grad$ and div may be expressed by

$$grad = j d : A^0 \longrightarrow \bar{A}^0 \quad \text{and} \quad div = tr D : \bar{A}^0 \longrightarrow A^0.$$

As by definition j vanishes on A^0 and tr on \bar{A}^0 , we may as well use the "more symmetric" formulae

$$grad = j d + D j \quad \text{and} \quad div = tr D + d tr,$$

and we use these formulae to define gradient and divergence in any degree p for $0 \leq p \leq n$.

The reason for adding the terms $D j$ and $d tr$ which are "hidden" in degree zero is of course not only an esthetic one, but it is the following theorem:

Theorem 1. *The operators grad and div satisfy the following relations, by which they are uniquely determined:*

- (1) $f \in A^0 \implies \text{grad } f$ is the ordinary gradient
- (2) $f \in A^0 \implies \text{grad } df = D \text{ grad } f$
- (3) $\varphi \in A^p, \psi \in A^q \implies \text{grad}(\varphi \wedge \psi) = (\text{grad } \varphi) \wedge \psi + \varphi \wedge \text{grad } \psi$
- (4) $X \in \vec{A}^0 \implies \text{div } X$ is the ordinary divergence
- (5) $\varphi \in A^p, \psi \in \vec{A}^q \implies \text{div}(\varphi \wedge \Psi) = (\text{grad } \varphi, \Psi) + \varphi \wedge \text{div } \Psi$

We shall not prove this theorem – nor the next one – but only indicate that this can be done using local coordinates which are geodesic at a given point $x \in M$. Observe also that property (2) is not valid in degree $p > 0$, where we have $D \text{ grad } \omega - \text{grad } d\omega = D^2 j\omega$, which does not vanish in general if the curvature tensor is non-zero.

The properties (1) – (3) of the gradient and the well-known rules for covariant derivation imply:

Corollary. *For $\varphi \in A^p$ and $X \in \vec{A}^0$,*

$$\langle X, \text{grad } \varphi \rangle = D_X \varphi, \quad \text{the covariant derivative.}$$

This characterizes grad φ completely.

So the gradient $\text{grad } \omega$ contains the information about all the covariant derivatives of the p -form ω , and is thus metrically equivalent to the tensor $D\omega$, defined by $D\omega(X; X_1, \dots, X_p) = (D_X \omega)(X_1, \dots, X_p)$. So why not simply use this tensor instead of the vector-valued form $\text{grad } \omega$? One good reason is that this tensor is only alternating in the last p arguments, and by changing the variance of the first argument we exhibit its special role. (Antisymmetrizing $D\omega$ would give the ordinary exterior derivative $d\omega$, because the Levi-Civita connection D is torsion-free.)

Theorem 2. *Between the operators grad, div and the classical operators $d, \star, \delta = (-1)^{np+n} \star d \star$, there hold the following relations:*

- (6) $d = \alpha \text{ grad}$
- (7) $\delta = \text{tr grad}$
- (8) $\star \text{ grad} = \text{grad } \star$
- (9) $\star \text{ div} = \text{div } \star$

Now let us try – for the sake of curiosity – whether we have also the relation $\Delta = \text{div grad}$ as in degree zero: With the relations (6) and (7) of the above theorem and the fact that $\text{tr } j = 0$ we get easily

$$\text{div grad} = (\text{tr } D + d \text{tr})(j d + D j) = \delta d + d \delta + \text{tr } D^2 j = \Delta + \text{tr } D^2 j.$$

So we don't have $\Delta = \text{div grad}$, but we make the following definition:

Definition. On an oriented Riemannian manifold M , we set

$$\Delta_0 := \operatorname{div} \operatorname{grad} \quad \text{and} \quad \operatorname{Ric} := \operatorname{tr} D^2 j.$$

Thus we have the

Weitzenböck Formula.

$$\Delta = \Delta_0 - \operatorname{Ric}$$

Of course, this is not really interesting as long as we don't know something about at least two of these three operators! To begin with, let us consider the operator Ric :

One checks easily, that Ric commutes with \star , and that it is A^0 -linear, i.e. it is a tensorial operator: for $\omega \in A^p$ the value $\operatorname{Ric}\omega(x)$ at a point $x \in M$ depends only on $\omega(x)$. The name "Ric" for this operator is justified by the following: For two vector fields X and Y , we have for the 1-form $\omega = \alpha X$:

$$(\operatorname{Ric}\omega)(Y) = (\operatorname{tr} D^2 X)(Y) = \operatorname{trace} R_{\cdot, Y} X = \operatorname{Ric}(Y, X) = \operatorname{Ric}(X, Y),$$

the well-known Ricci tensor on M !

The operator Δ_0 commutes also with \star , so its kernel

$$H_0 := \ker \Delta_0 = \bigoplus_{p=0}^n H_0^p$$

is a \star -invariant subspace of $A = \bigoplus A^p$. As we shall see in the next paragraph, it is even a finite-dimensional subalgebra. For the moment, we mention the product rule

$$\Delta_0(\varphi \wedge \psi) = \Delta_0 \varphi \wedge \psi + \varphi \wedge \Delta_0 \psi + 2\langle \operatorname{grad} \varphi, \operatorname{grad} \psi \rangle.$$

4. Compact manifolds.

The reason for using differential forms rather than arbitrary, non-alternating tensors is that we can integrate differential forms. So let M be in this paragraph a compact connected oriented Riemannian manifold of dimension n .

There are global inner products on A^p and \bar{A}^p , obtained by integration of the pointwise inner products:

$$(\varphi, \psi) := \int_M \varphi \wedge \star \psi \quad \text{for } \varphi, \psi \in A^p$$

and

$$(\Phi, \Psi) := \int_M \langle \Phi, \star \Psi \rangle \quad \text{for } \Phi, \Psi \in \bar{A}^p.$$

The following lemma will be the key formula for the further properties we shall derive for Δ_0 :

Lemma. *With respect to the global inner products, grad and $-\text{div}$ are adjoints, i.e. we have for $\varphi \in A^p$ and $\Psi \in \bar{A}^p$*

$$(\text{grad } \varphi, \Psi) = -(\varphi, \text{div } \Psi).$$

Proof:

$$\begin{aligned} (\text{grad } \varphi, \Psi) + (\varphi, \text{div } \Psi) &= \int_M ((\text{grad } \varphi, \star \Psi) + \varphi \wedge \star \text{div } \Psi) \\ &= \int_M ((\text{grad } \varphi, \star \Psi) + \varphi \wedge \text{div } \star \Psi) \\ &= \int_M \text{div}(\varphi \wedge \star \Psi), \end{aligned}$$

but $\text{div}(\varphi \wedge \star \Psi) = d \text{tr}(\varphi \wedge \star \Psi)$, because the second term of the divergence, $\text{tr } D(\varphi \wedge \star \Psi)$, vanishes since $\varphi \wedge \star \Psi$ is of maximum degree n . Thus, by Stokes' theorem, the result follows.

This simple integral formula has some interesting consequences. The following one is obvious:

Proposition. $(\Delta_0 \omega, \omega) \leq 0$ for all $\omega \in A^p$.

(Observe that with our sign convention for δ the same holds for the Laplacian Δ .)

Theorem 3.

$$\Delta_0 \omega = 0 \iff \text{grad } \omega = 0 \iff \Delta \omega = 0 \text{ and } \text{Ric } \omega = 0.$$

Proof: By definition of Δ_0 , $\text{grad } \omega = 0$ implies trivially $\Delta_0 \omega = 0$.

So let $\Delta_0 \omega = 0$. Then $(\text{grad } \omega, \text{grad } \omega) = -(\Delta_0 \omega, \omega) = 0$, and so $\text{grad } \omega = 0$, which implies by theorem 2 also $d\omega = 0$ and $\delta\omega = 0$, hence $\Delta \omega = 0$, which, together with the hypothesis $\Delta_0 \omega = 0$, gives also $\text{Ric } \omega = 0$. The remaining part is again trivial.

An immediate consequence is the following corollary, if we remember that grad is a derivation of degree zero:

Corollary. $H_0 = \bigoplus_{p=0}^n H_0^p$, the kernel of Δ_0 , is a subalgebra of $A = \bigoplus A^p$.

Moreover $H_0^p \subset H^p := \ker \Delta \cap A^p$, the space of harmonic p -forms.

In particular, the space H_0^p must be finite-dimensional, as H^p is. But there is another proof of this fact, which gives still more information about the structure of the space H_0^p :

Theorem 4. For $\varphi, \psi \in H_0^p$,

$$\langle \varphi, \psi \rangle = \text{const.} = \frac{1}{\text{vol } M} \langle \varphi, \psi \rangle,$$

where $\langle \cdot, \cdot \rangle$ is the pointwise scalar product on p -forms, induced by the Riemannian metric.

Proof: $\langle \varphi, \psi \rangle = \star(\varphi \wedge \star\psi)$, and thus

$$d\langle \varphi, \psi \rangle = \pm \star \delta(\varphi \wedge \star\psi) = \pm \star \text{tr grad}(\varphi \wedge \star\psi),$$

but this expression vanishes by the product rule and \star -invariance of grad .

This theorem says, that for any $x \in M$ the evaluation map

$$\text{eval}_x : H_0^p \longrightarrow \bigwedge^p T_x^* M$$

is an isometric injection, up to the constant factor $1/\text{vol}M$. As a consequence, H_0^p is of dimension $\leq \binom{n}{p}$.

If M is flat, we have $D^2 = 0$, hence $\text{Ric} = 0$, and thus $\Delta = \Delta_0$. So we get in this case $\dim H^p < \infty$ without using the ellipticity of Δ . (Of course, we cannot prove the regularity in this way: if we admit harmonic forms with coefficients in a Sobolev space, we are not sure that the C^∞ ones are already all of them.) The same property as for flat manifolds holds still in the case where Ric is positive semi-definite:

Theorem 5. If the operator $\text{Ric} : A^p \longrightarrow A^p$ is positive semi-definite, then $H_0^p = H^p$.

If $\text{Ric} : A^p \longrightarrow A^p$ is even positive definite, then $H_0^p = H^p = 0$.

Proof: In the expression

$$(\Delta \omega, \omega) = (\Delta_0 \omega, \omega) - (\text{Ric} \omega, \omega)$$

all three terms are ≤ 0 . So, $\Delta \omega = 0$ implies $(\Delta_0 \omega, \omega) = 0$, and that implies $\text{grad} \omega = 0$, hence $\omega \in H_0^p$. But $(\text{Ric} \omega, \omega)$ must also vanish, and thus $\omega = 0$ if Ric is positive definite.

The special case $p = 1$ yields of course the theorem by Bochner that we mentioned in the introduction. Even if the sectional curvatures are only positive in almost all plane directions at almost all points, the ordinary Ricci tensor is positive definite, which means exactly that our $\text{Ric} : A^1 \longrightarrow A^1$ is so.

II. The Geometry of Foliations

1. The characteristic form of a foliation.

We shall be interested in local geometric properties of foliations on Riemannian manifolds. Therefore we look at the distribution in TM tangent to the leaves. So, *foliation* means in the following simply *integrable distribution* (i.e. a C^∞ -subbundle) $F \subset TM$. For the sake of simplicity everything is oriented: the manifold and the foliation.

Consider the situation at a point $x \in M$: The p -dimensional linear subspace $F_x \subset T_x M$ is an element of the Grassmannian manifold of all p -dimensional oriented linear subspaces of $T_x M$. But things become simpler in our case, where $T_x M$ is a Euclidean space: In this case, the Grassmannian becomes in a natural way a (generating) submanifold of the linear space $\bigwedge^p T_x M$, and F_x is represented by $u_1 \wedge \dots \wedge u_p$, where (u_1, \dots, u_p) is a positively oriented orthonormal basis of F_x . As analysts are more used to consider the space $\bigwedge^p T_x^* M$ of p -forms than that of p -vectors, we can still use the Euclidean structure of $T_x M$ to identify $\bigwedge^p T_x M$ with $\bigwedge^p T_x^* M$: To F_x corresponds now the p -form $\chi \in \bigwedge^p T_x^* M$ with

$$\chi(u_1, \dots, u_p) = \begin{cases} 1 & \text{for any pos. oriented orthonormal basis of } F_x, \\ 0 & \text{if one of the } u_i\text{'s is perpendicular to } F_x. \end{cases}$$

Doing this at every point $x \in M$ simultaneously, we get the *characteristic form* χ_F of the foliation or distribution F . In the case of a foliation, the restriction of χ_F to a leaf is simply the volume form for that leaf.

For calculations with the characteristic form of the distribution F , the following simple representation is often useful: Let (X_1, \dots, X_n) be a local positively oriented orthonormal frame of vector fields on the open subset $U \subset M$, such that X_1, \dots, X_p span F . If $(\omega_1, \dots, \omega_n)$ denotes the dual frame of 1-forms, we have

$$\chi_F = \omega_1 \wedge \dots \wedge \omega_p,$$

and as $\omega_i(X_j) = \delta_{ij}$, we get for the differentials of the ω_i simply

$$\begin{aligned} d\omega_i(X_j, X_k) &= -\omega_i([X_j, X_k]) \\ &= \langle X_i, [X_k, X_j] \rangle \\ &= \langle X_i, D_{X_k} X_j \rangle - \langle X_i, D_{X_j} X_k \rangle \\ &= \langle D_{X_j} X_i, X_k \rangle - \langle D_{X_k} X_i, X_j \rangle. \end{aligned}$$

2. Geometric properties of a foliation and differential properties of its characteristic form.

To begin with, and to give also an idea of what kind of calculations one has to do, let us try to find out what it means that χ_F is closed:

With the above notations that means

$$d\chi_F(X_{i_1}, \dots, X_{i_{p+1}}) = 0 \quad \text{for } 1 \leq i_1 \dots i_{p+1} \leq n.$$

It is clear that $d\chi_F(X_{i_1}, \dots, X_{i_{p+1}}) = 0$ if more than two indices are $> p$. So there remain two cases to be interpreted:

$$(1) \quad d\chi_F(X_1, \dots, X_p, X_j) = 0 \quad \text{for } p+1 \leq j \leq n$$

and

$$(2) \quad d\chi_F(X_1, \dots, \hat{X}_i, \dots, X_p, X_j, X_k) = 0 \quad \text{for } 1 \leq i \leq p < j < k \leq n$$

For the first case one finds easily

$$d\chi_F(X_1, \dots, X_p, X_j) = (-1)^p \sum_{i=1}^p \langle X_i, D_{X_j} X_j \rangle = (-1)^{p+1} p \langle H, X_j \rangle,$$

where H is the mean curvature vector field of the foliation. Thus (1) means that all leaves are minimal submanifolds. Observe that (1) says simply that the restriction of χ_F to any $(p+1)$ -dimensional submanifold tangent to the leaves is closed.

For the second case one gets

$$d\chi_F(X_1, \dots, \hat{X}_i, \dots, X_p, X_j, X_k) = (-1)^i \langle X_i, [X_j, X_k] \rangle,$$

and (2) means simply that the orthogonal complement F^\perp is integrable. This applies also if F itself is not integrable. Together with the fact that $\star\chi_F = \chi_{F^\perp}$, we have thus the following theorem and the two corollaries:

Theorem 6. *The characteristic form of the foliation F is closed if and only if all the leaves are minimal submanifolds and the orthogonal complement F^\perp is integrable.*

Corollary 1. *If the characteristic form χ_F of the distribution F is co-closed, i.e. if $\delta\chi_F = 0$, then F is integrable.*

However, this condition is not necessary for the integrability of F : If $\dim F = 1$, we have $\delta\chi_F = \operatorname{div} X$, where X is the normed vector field describing F . So $\delta\chi_F = 0$ means in this case that the corresponding flow is volume preserving.

Corollary 2. *On a compact oriented Riemannian manifold, the characteristic form χ_F of the distribution F is harmonic if and only if F and F^\perp are foliations by minimal submanifolds.*

In order to check the properties of $\operatorname{grad} \chi_F$, one needs essentially only the following lemma, the proof of which is trivial:

Lemma. *With the notations introduced in the last section, locally*

$$\text{grad } \chi_F = \sum_{i=1}^p \omega_1 \wedge \dots \wedge \omega_{i-1} \wedge \text{grad } \omega_i \wedge \omega_{i+1} \wedge \dots \wedge \omega_p,$$

with $\langle \text{grad } \omega_i(X_j), X_k \rangle = \langle D_{X_k} X_i, X_j \rangle$ for $1 \leq i \leq p$ and $1 \leq j, k \leq n$.

Before using this lemma, let us make two definitions:

Definition. *A distribution $F \subset TM$ is said to be geodesic if $D_X Y$ is tangent to F whenever the vector fields X and Y are tangent to F .*

The distribution is said to be parallel if $D_X Y$ is tangent to F whenever the vector field Y is tangent to F and the vectorfield X is perpendicular to F .

Let us make some remarks concerning these definitions:

- (1) By definition, these notions are dual: F is geodesic if and only if F^\perp is parallel.
- (2) If F is geodesic, it is integrable and all the leaves are totally geodesic submanifolds. The converse is also true: A foliation with totally geodesic leaves is geodesic in the sense of the above definition. However, if a distribution F is totally geodesic in the sense that any geodesic tangent to F at some time remains tangent to F forever, the distribution F needs not be integrable, and so it is not necessarily geodesic in the sense of the above definition.
- (3) If F is a parallel foliation, then the Riemannian metric is bundle-like for F , in the sense of B. Reinhart, but the converse is only true if F^\perp is integrable.
- (4) The terminology is inspired by the following example in $\mathbb{R}^2 \setminus \{0\}$: The foliation F by straight lines passing through the origin is geodesic, and the foliation F^\perp by concentric circles around the origin is parallel.

The following two theorems are now easy consequences of the lemma, and we leave their proof as an exercise:

Theorem 7. *The distribution F is*

$$\begin{aligned} \text{geodesic} &\iff \text{grad } \chi_F \perp F, \text{ i.e. all values are perpendicular to } F, \\ \text{parallel} &\iff \text{grad } \chi_F \parallel F, \text{ i.e. all values are tangent to } F. \end{aligned}$$

Theorem 8. *Let F be a foliation. Then the given Riemannian metric on M is bundle-like for F if and only if*

$$i_X \text{grad } \chi_F \perp X \quad \text{for every } X \perp F.$$

3. Foliations of compact Riemannian manifolds.

If the manifold M is compact, then $\text{grad } \chi_F = 0 \iff \Delta_0 \chi_F = 0$, and we may apply the results of the last section of the first part to get some informations about geodesic and parallel foliations. The first result is a trivial consequence of the fact that $H_0^p \subset H^p$:

Theorem 9. *If the Betti number $b_p(M) = 0$, then there is no p -dimensional foliation on M which is at the same time geodesic and parallel, whatever is the Riemannian metric given on M .*

In the case $p = 1$ we can also say something about the set of all geodesic and parallel flows (i.e. 1-dimensional foliations) on M :

Theorem 10. *Let M be a compact, connected, oriented Riemannian manifold, and let $p := \dim H_0^1 > 0$. Then there is a differentiable action*

$\Phi : \mathbf{R}^p \times M \rightarrow M$ *such that*

- (1) \mathbf{R}^p acts on M by isometry ;
- (2) The orbits of this action are the leaves of a geodesic and parallel p -dimensional foliation ;
- (3) The projection of \mathbf{R}^p onto any orbit is a local isometry, where \mathbf{R}^p carries the standard Euclidean structure ;
- (4) Any geodesic and parallel flow on M is the restriction of this action to a 1-dimensional linear subspace of \mathbf{R}^p .

Proof: Recall that the evaluation maps $eval_x : H_0^1 \rightarrow T_x^*M$ are isometric injections, up to a constant factor, and that also $j : T_x^*M \rightarrow T_xM$ is an isometry. Thus, by choosing an appropriate basis in H_0^1 , we get a linear map

$$X : \mathbf{R}^p \rightarrow \bar{A}^0$$

with the following properties:

- (i) $eval_x \circ X : \mathbf{R}^p \rightarrow T_x^*M$ is an isometric injection for any $x \in M$, where \mathbf{R}^p carries the standard Euclidean structure ;
- (ii) The 1-form $\omega_u = j^{-1}(X_u)$ has vanishing gradient for any $u \in \mathbf{R}^p$.

Now, if $\|u\| = 1$, ω_u is the characteristic form for the flow defined by the vector field X_u , and this flow must be geodesic and parallel. Conversely, the characteristic form of any geodesic and parallel flow on M is in H_0^1 and hence of the form ω_u for some $u \in \mathbf{R}^p$ with $\|u\| = 1$. This will prove (4) once we have shown the other properties.

We define now the action of \mathbf{R}^p on M by these flows, i.e. by

$$\Phi(u, x) := \exp_x(X_u(x)).$$

To prove that this is a well-defined action, we must show that the flows defined by the vectorfields X_u commute. So let $u, v \in \mathbf{R}^p$. Then

$$D_{X_u}X_v = DX_v(X_u) = (Dj\omega_v)(X_u) = grad\omega_v(X_u) - j d\omega_v(X_u) = 0,$$

because with $grad\omega_v = 0$ we have also $d\omega_v = 0$. Thus $[X_u, X_v] = D_{X_u}X_v - D_{X_v}X_u = 0$.

A similar calculation shows that the vector fields X_u are Killing fields, i.e. that \mathbf{R}^p acts on M by isometries.

If e_1, \dots, e_p are the standard basis vectors of \mathbf{R}^p , the distribution F on M defined by the map $X : \mathbf{R}^p \rightarrow \bar{A}^0$ has just the characteristic form

$$\chi_F = \omega_{e_1} \wedge \dots \wedge \omega_{e_p},$$

which has vanishing gradient. So F is a geodesic and parallel foliation, and by construction its leaves are the orbits of the action Φ .

Finally, $eval_x \circ X : \mathbf{R}^p \rightarrow F_x$ is an isometry for any $x \in M$, and this map is just the tangent map at the origin of the map $\Phi_x : \mathbf{R}^p \rightarrow M$, so this map is a local isometry from \mathbf{R}^p onto the orbit of the point x .

4. Final remarks.

The last two theorems suggest that on a compact oriented Riemannian manifold the spaces H_0^p are in general very small, much smaller than the spaces H^p of harmonic forms. In fact, the last theorem shows that the flat torus is the only compact oriented surface admitting non-trivial 1-forms ω with $\Delta_0 \omega = 0$. But this is not necessarily so in higher dimensions: On a compact connected Lie group endowed with an invariant Riemannian metric, all invariant forms are already in H_0 , so in this case $H_0^p = H^p$ in any degree p . But $H_0^p \neq 0$ for $p > 1$ does not imply the existence of geodesic and parallel p -dimensional foliations.

References

1. GOLDBERG, S.I., "Curvature and Homology," New York-London, 1962.
2. KOBAYASHI, S. AND K. NOMIZU, "Foundations of Differential Geometry, Vol. 1," New York-London, 1963.
3. YANO, K. AND S. BOCHNER, "Curvature and Betti Numbers," Princeton, 1953.

Institut de Mathématiques
l'Université de Fribourg
Chemin du Musée 23
CH-1700 Fribourg, SUISSE

Rebut el 3 de Novembre de 1988