

# PHYSICAL REVIEW LETTERS

VOLUME 60

29 FEBRUARY 1988

NUMBER 9

## Diffraction-Free Beams Remain Diffraction Free under All Paraxial Optical Transformations

Kurt Bernardo Wolf

*Instituto de Investigaciones en Matemáticas Aplicadas y en Sistemas/Cuernavaca,  
Universidad Nacional Autónoma de México, 01000 México DF, México*

(Received 11 January 1988)

It is pointed out that the nondiffracting  $J_0$  beams of Durnin, Miceli, and Eberly are subgroup-reduced basis functions for the group of paraxial (symplectic, linear) transformations of optical phase space. This allows the action of the group on such and similar beams to be computed through  $2 \times 2$  matrix algebra.

PACS numbers: 02.20.+b, 03.50.-z, 03.65.-w

The diffraction-free beams predicted by Durnin<sup>1</sup> and observed by him together with Miceli and Eberly<sup>2</sup> have attracted attention for their properties of being extremely narrow and (almost paradoxially) nondiffracting. They have been called  $J_0$  beams and have the form<sup>2</sup>:

$$\begin{aligned} \Phi_\alpha(q_x, q_y, z; \kappa) &= e^{i\beta z} J_0(\alpha q), \quad 0 < \alpha < \kappa, \\ \alpha^2 + \beta^2 &= \kappa^2, \quad q_x^2 + q_y^2 = q^2, \end{aligned} \quad (1)$$

oscillating in time with a factor  $\exp(-i\kappa ct)$  (i.e., a phase  $\kappa ct$ ). The half width of the central intensity peak is nearly  $1/\alpha$ , and is independent of  $z$ . They are separated solutions of the wave equation in cylindrical coordinates. As for plane waves (also nondiffracting), (1) are not square integrable. Indeed, the necessarily finite extent of the experimental wave front brings in diffraction in the form of a shadow zone<sup>1</sup>; this was one of the determining properties measured in Ref. 2.

In this Letter I want to point out that the  $J_0$  beams (1) are a class of functions with the property of self-reproducing under *paraxial optical transformations*: small angles, thin lenses, parabolic-profile graded media, and similar mirrors. This is a consequence of the fact that (1) are subgroup-classified representation functions of  $\text{Sp}(2, \mathcal{R}) = \text{SL}(2, \mathcal{R})$ , the group of paraxial transformations by axis-symmetric optical systems.

Bacry and Cadilhac<sup>3</sup> derived the (double cover of)  $\text{Sp}(4, \mathcal{R})$  as symmetry group of the wave equation in the

paraxial limit. The recognition that the optical transfer functions are *canonical transform* kernels<sup>4</sup> is due to Nazarathy and co-workers<sup>5</sup>; a monographic account of canonical transforms in paraxial wave optics may be found in Castaño, López-Moreno, and Wolf.<sup>6</sup> It is to be expected (*post factum*—with apology) that the  $J_0$  beams should be identified with a clear group-theoretic structure.

In 1980 Basu and Wolf<sup>7</sup> studied all representations of the  $\text{SL}(2, \mathcal{R})$  group in all subgroup reductions. I consider the oscillator representation of the full  $\text{Sp}(4, \mathcal{R})$  group of linear (asymmetric) transformations of optical phase space, reduced<sup>6</sup> by the symmetry  $\text{O}(2)$  of rotations and inversions around the optical  $z$  axis, as  $\text{Sp}(4, \mathcal{R}) \supset \text{Sp}(2, \mathcal{R}) \otimes \text{O}(2)$ . The three generators of the “radial”  $\text{Sp}(2, \mathcal{R})$  are given by

$$K_0^\gamma = \frac{1}{4} (-d^2/dq^2 + \gamma/q^2 + q^2), \quad (2a)$$

$$K_0^\zeta = -(\frac{1}{2}i)(q d/dq + \frac{1}{2}), \quad (2b)$$

$$K_0^\eta = K_0^\zeta + K_0^\gamma + \frac{1}{2} (-d^2/dq^2 + \gamma/q^2), \quad (2c)$$

$$K_0^\chi = K_0^\zeta - K_0^\gamma = \frac{1}{2} q^2. \quad (2d)$$

The Casimir invariant is  $-\gamma/4 + \frac{3}{16} = k(1-k)$ , for  $\gamma = (2k-1)^2 - \frac{1}{4}$ , where  $k$  is Bargmann's label<sup>8</sup> for the (positive) discrete representations  $D_k^\pm$ ,  $k = \frac{1}{2}, 1, \frac{3}{2}, \dots$ ; this range is related by integer  $|m| = 2k-1$  to the representations of the conjugate  $\text{O}(2)$  subgroup.

On  $\mathcal{L}^2(R^+)$ , with inner product

$$(f, g) = \int_0^\infty dq f(q)^* g(q),$$

eigenfunctions of  $K_0^\zeta$  are harmonic oscillator  $+ \gamma/q^2$  potential wave functions; those of  $K_0^\eta$  are similar repulsive oscillators; those of  $K_0^\xi$  are the Mellin transform basis functions; and those of  $K_0^\zeta$  are  $\delta_a(q) = \delta(q - a)$ , the generalized basis of  $q$ -localized states. Our interest here is in the  $\mathcal{L}^2(R^+)$  generalized eigenbasis of  $K_0^\zeta$ , since

they are Bessel functions:

$$\Phi_a^k(q) = e^{i\pi k} (aq)^{1/2} J_{2k-1}(aq), \tag{3}$$

with eigenvalue  $a^2/2 \in R^+$ ; they are Dirac orthonormal and complete.

The group  $Sp(2, R)$  generated by the second-order differential operators (2) is a group of unitary integral transforms<sup>5</sup> on  $\mathcal{L}^2(R^+)$  of the Hankel type.<sup>7</sup> The matrix representation and optical elements of the various subgroups are

$$\exp(i\theta K_0^\xi) \leftrightarrow \begin{pmatrix} \cos\theta/2 & -\sin\theta/2 \\ \sin\theta/2 & \cos\theta/2 \end{pmatrix} \leftrightarrow \begin{cases} SO(2) \\ \text{quadratic fibers.} \end{cases} \tag{4a}$$

$$\exp(i\beta K_0^\eta) \leftrightarrow \begin{pmatrix} \exp(-\beta/2) & 0 \\ 0 & \exp(\beta/2) \end{pmatrix} \leftrightarrow \begin{cases} SO(1, 1) \text{ of pure} \\ \text{magnifiers.} \end{cases} \tag{4b}$$

$$\exp(ib K_0^\zeta) \leftrightarrow \begin{pmatrix} 1 & -b \\ 0 & 1 \end{pmatrix} \leftrightarrow \begin{cases} E(2) \text{ parabolic group} \\ \text{of free flight } z = Kb. \end{cases} \tag{4c}$$

$$\exp(ic K_0^\zeta) \leftrightarrow \begin{pmatrix} 1 & 0 \\ C & 1 \end{pmatrix} \leftrightarrow \begin{cases} E(2) \text{ of thin lenses} \\ \text{of power } C. \end{cases} \tag{4d}$$

The paraxial group composition law is that of matrix multiplication (here no metaplectic sign). (See Ref. 7 with the notation correspondence of its footnote 1.) The group action is the integral transform  $C^k(\mathbf{M})$ ,  $\det \mathbf{M} = 1$ ,

$$\left[ C^k \begin{pmatrix} a & b \\ c & d \end{pmatrix} f \right] (q) = \int_0^\infty dq' C^k \begin{pmatrix} a & b \\ c & d \end{pmatrix} (q, q') f(q'), \tag{5a}$$

with a kernel that is the optical transfer function

$$C^k \begin{pmatrix} a & b \\ c & d \end{pmatrix} (q, q') = e^{-i\pi k} b^{-1} (qq')^{1/2} \exp \left[ \frac{i(dq^2 + aq'^2)}{2b} \right] J_{2k-1}(qq'/b). \tag{5b}$$

In the singular case  $b = 0$ , the action is geometrical:

$$\left[ C^k \begin{pmatrix} a & 0 \\ c & a^{-1} \end{pmatrix} f \right] (q) = a^{-1/2} \exp \left[ \frac{icq^2}{2a} \right] f(q/a), \quad a > 0. \tag{5c}$$

Using the matrix representation (4) and products thereof, one can see that a paraxial transformation (5) applied to the  $K_0^\zeta$  (free flight) eigenbasis  $\Phi_a^k$  reproduces the form of the wave function:

$$\left[ C^k \begin{pmatrix} a & b \\ c & d \end{pmatrix} \Phi_a^k \right] (q) = \left[ C^k \begin{pmatrix} a & 0 \\ c & a^{-1} \end{pmatrix} \exp(-iba^{-1} K_0^\zeta) \Phi_a^k \right] (q) = a^{-1/2} \exp \left[ \frac{-iba^2}{2a} \right] \exp \left[ \frac{icq^2}{2a} \right] \Phi_a^k(q/a). \tag{6}$$

In words: A thin lens (4d) will multiply the diffraction-free states  $\Phi_a^k$  by a phase  $\exp(icq^2/2)$  (with no effect on the intensity); pure magnifiers (4b) will yield  $a^{-1/2} \Phi_a^k(q/a) = a^{1/2} \Phi_{a/a}^k(q)$ , a diffraction-free state of a different width; a length of quadratic-profile fiber (4a) will produce a line of integral transforms of the (Fourier) Hankel type. Finally, free propagation (4c) by  $b = z/\kappa$  multiplies  $\Phi_a^k$  by the phase factor  $\exp(-iza^2/2\kappa)$ . This phase is the paraxial approximation to the phase anomaly factor of Durnin's function (1) for  $a \ll \kappa$ ,  $\beta \approx \kappa - a^2/2\kappa - a^4/8\kappa^3 - \dots$ . The paraxial approximation with integral transforms (5) does violence to the relation  $\beta = (\kappa^2 - a^2)^{1/2}$  that keeps a minimum beam width: It frees the conjugate optical Hamiltonian mo-

mentum from the constraint of taking values only inside a circle of radius  $n$ , the refractive index of the medium.<sup>6</sup> This is the price of the application of Schrödinger quantization to model paraxial optics. The cutoff to  $a < \kappa$  comes from the restriction of having a finite smallest wavelength in the spectrum of optical wave functions.

Modulo these caveats, we identify Durnin's diffraction-free  $J_0$  states (1) with the  $K_0^\zeta$ -classified eigenbasis (3) of the group of paraxial transformations, for  $k = \frac{1}{2}$  ( $\gamma = -\frac{1}{4}$ ): thus

$$\Phi_a(q_x, q_y, 0; \kappa) \leftrightarrow \Phi_a^{1/2}((q_x^2 + q_y^2)^{1/2}). \tag{7}$$

The  $(aq)^{1/2}$  factors in (3) and (5b) yield the proper

measure  $\int_0^\infty q' dq'$  . . . to integrate over the circles' radii. We should not overlook that also  $k \neq \frac{1}{2}$  transforms exist and they originate from wave syntheses [Eq. (2) in Ref. 1] with azimuthal dependence  $A(\phi) \sim e^{im\phi}$ .

The title of the Letter refers to the self-reproducing form of Eq. (6) with respect to  $\Phi_a^k$ . Paraxial optical systems normally end with free flight  $\zeta$ , i.e., the image on the screen is the canonical transform  $\mathbb{C}^k(\mathbf{M}_\zeta)$  of the object  $J_k$  state corresponding to the matrix

$$\mathbf{M}_\zeta = \begin{pmatrix} 1 & \zeta \\ 0 & 1 \end{pmatrix} \begin{pmatrix} a & b \\ c & d \end{pmatrix} = \begin{pmatrix} a + \zeta c & b + \zeta d \\ c & d \end{pmatrix}. \quad (8)$$

Since the 1-1 element of  $\mathbf{M}_\zeta$  depends on  $\zeta$ , the half-width  $a$  of the outgoing beam will depend on  $\zeta$ . This converging or diverging "diffraction-free" beam differs only by scale and quadratic phase from the object  $J_k$  state; the transversal illumination pattern, however, will be the same, up to scale.

Self-reproducing wave functions in this sense<sup>9</sup> are all the subgroup-reduced eigenfunctions including radial harmonic and repulsive oscillator functions, and, complexifying, Gaussians (with a modified Bessel function in  $aq$  in place of the usual exponential factor), and Barut-Girardello coherent states.<sup>10</sup> The last two are, respectively,

$$G_\omega(q-a) = \omega^{-1} (aq)^{1/2} e^{-a^2/2\omega} I_{2k-1}(aq/\omega) e^{-q^2/2\omega} = \left[ \mathbb{C}^k \begin{pmatrix} i\omega & 1 \\ -1 & 0 \end{pmatrix} \Phi_a^k \right] (q), \quad (9a)$$

$$Y_a(q) = (2\pi)^{1/2} \left[ \mathbb{C}^k \begin{pmatrix} 1/\sqrt{2} & -i/\sqrt{2} \\ -i/\sqrt{2} & 1/\sqrt{2} \end{pmatrix} \delta_a \right] (q) = (2n)^{1/2} \left[ \mathbb{C}^k \begin{pmatrix} 1/\sqrt{2} & -i/\sqrt{2} \\ -i/\sqrt{2} & 1/\sqrt{2} \end{pmatrix} \mathbb{C}^k \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix} \Phi_a^k \right] (q). \quad (9b)$$

<sup>1</sup>J. Durnin, J. Opt. Soc. Am. A **4**, 651-654 (1987).

<sup>2</sup>J. Durnin, J. J. Miceli, and J. H. Eberly, Phys. Rev. Lett. **58**, 1499-1501 (1987).

<sup>3</sup>H. Bacry and M. Cadilhac, Phys. Rev. A **23**, 2533-2536 (1981).

<sup>4</sup>K. B. Wolf, *Integral Transforms in Science and Engineering* (Plenum, New York, 1979), Pt. 4.

<sup>5</sup>M. Nazarathy and J. Shamir, J. Opt. Soc. Am. **72**, 356-364 (1982); M. Nazarathy, A. Hardy, and J. Shamir, J. Opt. Soc. Am. A **3**, 1360-1369 (1986).

<sup>6</sup>O. Castaños, E. López-Moreno, and K. B. Wolf, in *Lie Methods in Optics*, edited by J. Sánchez Mondragón and K. B. Wolf, Lecture Notes in Physics Vol. 250 (Springer-Verlag, Heidelberg, 1986).

<sup>7</sup>D. Basu and K. B. Wolf, J. Math. Phys. **23**, 189-205 (1982).

<sup>8</sup>V. Bargmann, Ann. Math. **48**, 568-640 (1947).

<sup>9</sup>K. B. Wolf, J. Math. Phys. **18**, 1046-1051 (1977).

<sup>10</sup>A. O. Barut and L. Girardello, Commun. Math. Phys. **21**, 41-55 (1971).