

Digestion and absorption of refractory carbon from the plant *Spartina alterniflora* by the oyster *Crassostrea virginica*

Roger I. E. Newell¹ & Christopher J. Langdon²

¹ Center for Environmental and Estuarine Studies, Horn Point Laboratories, University of Maryland, PO Box 775, Cambridge, Maryland 21613, USA

² Hatfield Marine Science Center, Oregon State University, Newport, Oregon 97365, USA

ABSTRACT: The vascular plant *Spartina alterniflora* Loisel., grown in an atmosphere containing ¹⁴CO₂, was chemically treated to produce a lignocellulosic, crude fiber material containing 0.1 % lipid, no measurable protein, 4.4 % starch, 85.4 % cellulose and 7.7 % lignin. This material was introduced, via syringe through the mouth, into the stomachs of oysters *Crassostrea virginica*; these were then held for 24 h in 13 ppt filter-sterilized seawater at 25 °C. A second group of *C. virginica* was treated identically, except for the addition of 5 mg l⁻¹ of the antibiotics chloramphenicol and rifampicin to both food material and seawater. Direct enumeration (DAPI) of bacteria demonstrated that treatment with antibiotics eliminated bacteria from the oysters' stomach fluid. The ¹⁴C specific activities of cell-free hemolymph and tissue samples taken 24 h after feeding indicated that oysters were able to digest and absorb carbon from the *S. alterniflora* material with a mean efficiency of 1.3 %. There were no significant differences (ANOVA, p > 0.05) in the digestion and absorption of ¹⁴C material between antibiotic treated and untreated oysters. Furthermore, *in vitro* cellulolytic activities of tissue homogenates of antibiotic treated oysters were not significantly different (ANOVA, p > 0.05) from those of untreated oysters. These results indicate that oysters are able to digest only small amounts of refractory cellulosic material and that this process is not enhanced by bacteria present in the stomach. *In vitro* characterization of the cellulolytic enzymes of the crystalline style of *C. virginica* under aseptic conditions indicated the presence of β-1,4-glucanase (Cx cellulase) activity that released oligosaccharides from the *S. alterniflora* material. Neither C₁ cellulase, capable of degrading crystalline cellulose, nor β-glucosidase activities were detected, using cotton fibre and cellobiose as substrates, respectively. These results suggest that the principal function of style cellulase activity is the partial depolymerisation of refractory amorphous cellulose, perhaps aiding the digestion of algal cells and detritus.

INTRODUCTION

Spartina alterniflora Loisel., a dominant vascular plant in salt marshes of the Eastern United States (McIntire & Dunstan 1976), is a major source of organic carbon to adjacent waters (Teal 1962). Living *S. alterniflora* is composed of ca 70 to 82 % carbohydrate by dry weight (Squiers & Good 1974, Smith et al. 1979), and 99 % of this carbohydrate is cellulosic structural material (McIntire & Dunstan 1976). The refractory cellulosic components may slowly decompose *in situ* after senescence of *S. alterniflora* (Smith et al. 1979, Mountford 1980), but some of the litter may be exported to adjacent estuaries during periods of high

tides, unusual storms (Pickral & Odum 1976) or as the result of ice rafting (Heinle & Flemer 1976).

In order for this refractory cellulose to be directly utilized as a source of carbon by estuarine molluscs, such as the oyster *Crassostrea virginica* (Gmelin), they must possess a suite of 3 cellulolytic enzymes: (1) C₁ cellulase which acts only on crystalline, native cellulose; (2) Cx or β-1,4-glucanases which act on amorphous cellulose and soluble cellulose derivatives; (3) β-glucosidases (cellobiase) acting on cellobioses and, to a lesser extent, on other β-oligosaccharides (Mandels et al. 1976).

It is often technically difficult to determine if the complete complement of cellulases is present in living

organisms. C_1 cellulase activity has rarely been clearly demonstrated in a bivalve mollusc because researchers have not used truly crystalline cellulose substrates to test digestive enzymes. Many investigators (e.g. Payne et al. 1972) have used cellulose filter paper as a substrate for C_1 cellulase, but have not allowed for the fact that filter paper, like many other insoluble celluloses, is a complex substrate of both amorphous and crystalline cellulose (Mandels et al. 1976). Halliwell (1966) suggested the use of cotton as a crystalline cellulose substrate for determining the activity of C_1 cellulase. Koopmans (1970) found that style extracts from the bivalve *Cardium edule* only weakly converted cotton to reducing sugars, compared with breakdown of reprecipitated cellulose or carboxymethyl cellulose. The presence of β -1,4-glucanase and β -glucosidase activities has been demonstrated in many bivalve species (Kristensen 1972, Wojtowicz 1972, Mathers 1973, Morton 1983, Lucas & Newell 1984, Mayasich & Smucker 1986).

Cellulolytic enzymes present in shipworms (Teredinidae) are exogenously derived from bacteria associated with the gland of Deshayes (Crosby & Reid 1971, Waterbury et al. 1983), and gut bacteria have also been implicated in cellulose digestion by the sea urchin *Strongylocentrotus droebachiensis* (Fong & Mann 1980). Conversely, Payne et al. (1972) found cellulolytic activity in the bivalve *Scrobicularia plana* but no evidence of cellulolytic bacteria in the clam's gut. Crosby & Reid (1971) used histological techniques to demonstrate the presence of cellulases in the tubules of the diverticula of several bivalve species and asserted that this cellulolytic activity was partly endogenous in origin and not dependent on the presence of gut bacteria.

In this study we have examined the digestion and utilization of a refractory cellulosic material, derived by chemical extraction from *Spartina alterniflora*, by the American oyster *Crassostrea virginica*, a bivalve mollusc commonly found in estuaries draining salt marshes. The role of the crystalline style in extracellular cellulose digestion has been characterized *in vitro* using a range of cellulose substrates. The overall digestion of cellulose and absorption of carbon into the oyster's hemolymph have been measured *in vivo* using ^{14}C tracer techniques and compared in antibiotic treated and untreated oysters so as to determine the importance of enteric bacteria in cellulose digestion.

METHODS

Preparation and characterization of ^{14}C -cellulose from *Spartina alterniflora*. Young *Spartina alterniflora* plants (short form) with associated soil were collected

from the Canary Creek marsh, Lewes, Delaware in late May 1985 and transplanted to 41 plastic pots. Leaves and stems were trimmed to soil level and re-growth allowed to take place over a period of 2 wk. Four pots of plants were transferred to an air-tight 'Plexiglas' chamber of the type designed by Ferguson & Williams (1974) for labelling *S. alterniflora* with $^{14}\text{CO}_2$. This method has been shown to produce *S. alterniflora* with a proximate composition very similar to that of marsh-grown plants (Coston-Clements & Ferguson 1985).

The plants were labelled with ^{14}C by growing them in an atmosphere of 0.03 % v/v CO_2 which was partly made up with $^{14}\text{CO}_2$. Preliminary experiments indicated that the plants completely removed 0.03 % v/v CO_2 from the chamber's atmosphere over a period of 14 h illumination and 10 h darkness. Caseous $^{14}\text{CO}_2$ was produced by adding a solution of $\text{NaH}^{14}\text{CO}_3$ to 250 ml of 1 M citric acid solution contained in a stirred flask in the chamber. Five mCi of $\text{NaH}^{14}\text{CO}_3$ (specific activity $30 \mu\text{Ci mmol}^{-1}$) were added to the chamber over a 5 d period, followed by a further 5 d period in which the plants were grown in a $^{14}\text{CO}_2$ -free atmosphere of 0.03 % CO_2 . The plants were then removed from the chamber and the aerial parts harvested and dried at 60°C for 24 h.

Crude fiber was prepared from dry *Spartina alterniflora*

- (1) Grind 20 g dry plant material in a ball-mill (Crescent 'Wig-L-Bug')
- ▼
- (2) Extract lipid with 150 ml of a solution of 1 vol. chloroform and 2 vol. methanol while mixing thoroughly. (Repeat $\times 5$)
- ▼
- (3) Extract protein with 100 ml of a solution containing 25 ml 10 % Na_2CO_3 , 25 ml 0.8 N NaOH, 25 ml water and 25 ml 10 % lauryl sulphate. ($\times 4$)
- ▼
- (4) Extract with 100 ml of 0.9 % H_2SO_4 at 95°C for 30 min. Wash material with 100 ml distilled H_2O . ($\times 1$)
- ▼
- (5) Extract with 100 ml of 12.5 % (w/v) NaOH at 95°C for 30 min. Wash material with 100 ml distilled H_2O . ($\times 1$)
- ▼
- (6) Dry material at 60°C
- ▼
- (7) Regrind material in a ball-mill. Resuspend in 1 M NaCl; sieve with a $20 \mu\text{m}$ 'Nitex' screen and collect material $< 20 \mu\text{m}$. Re-dry, grind and sieve material $> 20 \mu\text{m}$
- ▼
- (8) Wash sieved material with 100 ml sterile seawater; then wash with 100 ml distilled H_2O . ($\times 2$)
- ▼
- (9) Freeze dry material and store at -25°C

Fig. 1. Preparation of cellulosic material from ^{14}C -labelled *Spartina alterniflora*

flora by acid/alkali extraction (Fig. 1), based on the method of Strickland & Parsons (1972). The method was more extreme than commonly used for the preparation of cellulose (e.g. Whistler 1963) in order to minimize contamination by protein, lipid, starch and hemicelluloses. ^{14}C specific activity of the extracted material was $0.24 \mu\text{Ci mg}^{-1}$ dry weight, as determined by complete oxidation and measurement of $^{14}\text{CO}_2$ trapped in a liquid scintillation medium (Benner et al. 1984). Composition and distribution of ^{14}C in the extracted material was determined using a combination of chemical and enzymatic assays:

(a) A modified Klason hydrolysis (Benner et al. 1984) was used to determine the distribution of ^{14}C between the lignin and polysaccharide fractions. This involved hydrolyzing 20 mg samples of the *Spartina alterniflora* material in 0.5 ml of 72 % sulphuric acid at 30°C for 1 h with continuous shaking. The resulting suspension was then diluted with 14 ml distilled water and autoclaved at 15 psi and 121°C for 1 h. This procedure hydrolyses polysaccharides, whereas lignin condenses and remains insoluble (Benner et al. 1984). The tubes were centrifuged at 1200 g for 10 min and the supernatant filtered through a $0.2 \mu\text{m}$ Nuclepore filter. The pellet was resuspended in 10 ml distilled water at 80°C and the suspension refiltered with a $0.2 \mu\text{m}$ Nuclepore filter. The filter was washed with a further 5 ml of distilled water. ^{14}C activity in the hydrolysate was expressed as a percentage of the combined radioactivities of the pellet and filtrates.

(b) The percentage of ^{14}C in the cellulose fraction from 10 mg of *Spartina alterniflora* material was estimated by enzyme digestion with a mixture of cellulases from *Trichoderma viride* (Sigma type V) and *Aspergillus niger* (Sigma type II) made up in 0.1M acetate buffer (pH 5.0) containing 20 ppm Thimerosal. The cellulase activity of this mixture was composed of equal activities of *T. viride* and *A. niger* cellulases and was sufficient to break down completely 10 mg of pure cellulose to glucose in 2 h at 37°C . Assay tubes were incubated by shaking for 24 h at 25°C . Control tubes containing only acetate buffer and cellulosic material were used to estimate leaching rates. The amount of ^{14}C label solubilized by the cellulases, minus that released by leaching, was expressed as a percentage of the combined radioactivities of the pellet and filtrates.

(c) The proportion of ^{14}C in the form of starch was estimated by incubating 20 mg of the ^{14}C cellulosic material with α -amylase (*Bacillus* sp., Sigma type II-A) and determining the release of soluble ^{14}C . The α -amylase preparation was found to have detectable cellulolytic activity when incubated with highly purified cellulose (Sigmacell, Sigma). Potential liberation of ^{14}C from the cellulose component of the *Spartina alterniflora* material by this enzyme preparation

was minimized by the addition of excess non-radioactive cellulose to the assays (40:1 W/W, Sigmacell: ^{14}C *S. alterniflora* material) to saturate the contaminating cellulase activity. Addition of this quantity of cellulose had no effect on α -amylase activity.

The α -amylase was made up in $0.22 \mu\text{m}$ filtered, autoclaved seawater (sterile seawater) at salinity 13 to 14 ppt, pH 6.9 and containing 20 ppt of the antibiotic Thimerosal. The amount of enzyme used in the assay was sufficient to convert completely 20 mg of starch to maltose in 1 min at 37°C and pH 6.9.

After digestion of the *Spartina alterniflora* material for 24 h at 25°C (with continuous shaking), the suspension was centrifuged at 1200 g for 10 min and the supernatant filtered through a $0.2 \mu\text{m}$ Nuclepore filter. *S. alterniflora* material suspended in sterile seawater was used as a control for leaching. The percentage ^{14}C solubilized by β -amylase was expressed as in (b).

(d) Lipids were extracted with chloroform/methanol (Holland & Gabbott 1971). The specific activity of the lipid extract was expressed as a percentage of the specific activities of the extract and pellet combined.

(e) Protein was extracted by boiling in 0.5M NaOH for 10 min. After centrifugation (1000 g) for 10 min, the protein content of the supernatant was determined using the method of Lowry et al. (1951) as modified by Rice (1982) to correct for interference by phenolic compounds from associated lignin. Bovine serum albumen was used as a standard.

***In vitro* estimates of oyster cellulase activity.** *Preparation of style and digestive gland extracts.* *Crassostrea virginica* (shell height 10 to 12 cm; mean $[\pm \text{SE}]$ dry tissue weight = 1.35 ± 0.031 g) were collected in October from the Choptank River, Maryland and held in the laboratory for 10 to 20 d in a constant flow of unfiltered seawater pumped from the Choptank River estuary (ambient temperature 16 to 17°C , salinity 13 to 14 ppt). Styles were removed from oysters and immediately added to chilled sterile seawater, containing 20 ppt Thimerosal, resulting in a concentration of 0.65 styles ml^{-1} . Styles were homogenized in a glass Potter Elvehjem homogeniser to aid dissolution after which the preparation was centrifuged at 1000 g for 10 min to remove insoluble particulate matter. The pH of sterile seawater was adjusted to 6.9 with dilute HCl in order to approximate the pH of the oyster stomach and style sac (Morton 1983).

The digestive glands of 21 oysters were removed and gently homogenized in 100 ml of sterile seawater (pH 6.9). The homogenate was vacuum filtered at <5 mm Hg through a $30 \mu\text{m}$ Nitex screen. Both digestive gland and style homogenates were stored at 4°C for no more than 2 h before use. The protein contents of both homogenates were determined using the method of Lowry et al. (1951).

Assay of style cellulase activity. Four cellulose substrates were used to test style cellulase activity: (1) pharmaceutical grade cotton balls, pre-washed in hexane to remove any wax or oil coating; (2) non- ^{14}C labelled *Spartina alterniflora*, prepared as detailed in Fig. 1; (3) cellulose azure (Sigma Type I) which releases a dye (absorbance 595 nm) as a result of breakdown by cellulases; (4) cellobiose (Sigma) to test for the presence of β -glucosidase. All substrates were added at a concentration of 0.5 % W/V to the enzyme preparations together with an antibiotic mix of 20 ppm Thimerosal and 2 % V/V toluene to inhibit bacterial activity. The tubes were incubated on a shaker table at 25 °C for 24 h, a period that ensured measurable levels of enzyme product. Each assay was performed in triplicate.

Assays were terminated by immersing them in a boiling water bath to denature the enzymes. Each tube was centrifuged at 1400 g for 10 min and the supernatant filtered through a 0.2 μm Nuclepore filter. Glucose concentration was determined using a hexokinase enzymatic test kit (Sigma 16 UV), standardized with α -D(+) glucose. Total carbohydrate concentration was determined using the phenol-sulphuric acid method of Dubois et al. (1956), also standardized with glucose. The relation between cellulose azure dye release and total carbohydrate release was determined and described by the following regression equation:

$$A = (0.000462 \times C) + 0.00728 \quad (n = 17; r = 0.943)$$

where A = absorbance at 595 nm; C = concentration of α -D(+) glucose in $\mu\text{g ml}^{-1}$.

Initial glucose, total carbohydrate and azure concentrations were determined in control tubes boiled immediately upon addition of the enzyme to the substrate. Additional control tubes, containing only enzyme or substrate in sterile water, were boiled at time zero and after the 24 h incubation to determine the rate of release of product from sources other than cellulase acting on the cellulose substrate.

Assay of digestive gland cellulase activity. Substantial release of glucose and soluble carbohydrate from digestive gland autolysis masked the *in vitro* release of these end-products by cellulase activity. Therefore, cellulose azure was used as a substrate to avoid this difficulty. The experimental protocol was identical to that described above for the style, except that the incubation period was increased to 168 h to ensure a measurable concentration of released azure. Digestive gland samples turned cloudy upon boiling and so could not be used as controls. Instead, assay tubes containing only digestive gland or cellulose azure in sterile water with 2 % V/V toluene and 20 ppm Thimerosal were used as controls. At the end of the incubation period, samples were centrifuged at 1400 g

for 10 min, the supernatant filtered (Whatman GF/C), and the amount of azure released determined.

In Vivo ^{14}C absorption experiment. *Oyster feeding.* Thirty-six *Crassostrea virginica* (shell height 10 to 12 cm) collected and held as above were scrubbed clean of epibionts. Shell-boring polychaetes were removed by 2 consecutive, 5 min immersions in 0.1 % V/V domestic hypochlorite solution ('Chlorox'). Ventral margins of both left and right valves of the oysters were partially removed using a high-speed diamond saw to expose gills and labial palps. The hood formed by anterior fusion of the 2 mantle folds (Galtsoff 1964) was cut to expose the mouth. Fluid from oesophagus and stomach was collected using a Pasteur pipette inserted through the mouth and stored at 5 °C with 1 % W/V (final concentration) formalin added, for estimation of bacteria concentrations (see below). The described manipulation of the oysters did not cause fatal injury because oysters survived and after several weeks showed signs of shell repair.

One hundred μl of a suspension of 38.1 mg ml^{-1} ^{14}C -labelled *Spartina alterniflora* material (9.14 $\mu\text{Ci ml}^{-1}$) made up in sterile seawater were introduced by syringe through the mouth and into the stomach of each of 12 oysters. A second group of 12 oysters was similarly fed the same amount of material made up in sterile seawater containing 5 mg l^{-1} each of the antibiotics chloramphenicol and rifampicin. Langdon & Bolton (1984) found that this antibiotic mixture added to seawater had no adverse effects on either the filtration rate or growth of *Crassostrea virginica*. Control groups of 4 oysters had either 100 μl of sterile seawater or 100 μl of sterile water containing both antibiotics introduced into their stomachs.

Antibiotic or non-antibiotic treated oysters were placed in separate 0.22 μm filtered, 20 l volumes of seawater (13 ppt, 25 °C). Five mg l^{-1} of each of the antibiotics chloramphenicol and rifampicin were added to the seawater in which antibiotic-treated oysters were held. Seawater in both aquaria was replaced after 15 h of incubation with fresh, 0.22 μm filtered seawater.

Hemolymph collection and analysis. In order to estimate the rate and time course of ^{14}C absorption into the circulatory system of *Crassostrea virginica*, 500 μl samples of hemolymph were periodically withdrawn by syringe from the sinus of the adductor muscle. Hemolymph samples were taken from oysters at the start of the experiment to estimate background radioactivity. Six fed oysters from each of the antibiotic and non-antibiotic treatments were sampled at 5 and 24 h after the beginning of the experiment and 6 were sampled after 15 and 24 h. Unfed oysters were sampled at 15 h and 24 h.

Hemolymph was drawn into 500 μl of artificial sea-

water (salinity 12 ppt; free of Ca^{++} and Mg^{++}) containing 2 W/V sodium ethylenediaminetetracetate (Fisher & Newell 1986). The hemolymph sample was centrifuged at 1000 g for 10 min and the supernatant removed. The pellet was washed twice with 1.0 ml of sterile seawater and resuspended in 0.5 ml of sterile seawater. The ^{14}C specific activities of the pellet and supernatant were determined.

Digestion and incorporation of ^{14}C *S. alterniflora* material. Digestion and incorporation of ^{14}C into oyster tissue was determined by homogenizing individual body tissues in 10 ml of sterile seawater, extracting 1 ml of the homogenate with acid and alkali, followed by centrifugation (Fig. 2). The ^{14}C specific activity of 1.0 ml of the supernatant solutions and all pellets were determined. The supernatant radioactivity came from ^{14}C incorporated in oyster tissues, as well as ^{14}C that had been digested but not absorbed at the time the experiment was terminated. In addition, the acid/alkali extraction solubilised 35.1 % of the ^{14}C from the undigested cellulosic food material present in the oyster's digestive system (Table 1). The pellet contained 64.9 % of the remaining ^{14}C undigested food material. The specific activities of oyster tissue extracts were therefore corrected for this solubilization by acid/alkali treatment of food remaining in the gut. Specific activities were corrected for quenching and expressed on a per oyster basis.

Cellulose breakdown by bacteria present in tissue homogenates from individual oysters that were either treated or not treated with antibiotics was assessed using cellulose azure as a substrate. The protocol was the same as that described above for digestive gland




Fig. 2. Determination of ^{14}C in oyster tissues

assays, except that styles were not removed prior to homogenization.

Bacteria concentrations. In the feeding experiment bacteria concentrations were estimated by staining the bacteria with 4,6-diamidino-2-phenylindole (DAPI; Sigma) and directly counting the bacterial cells under UV illumination (Porter & Feig 1980). Concentrations of bacteria were determined in samples of stomach fluids from both antibiotic and non-antibiotic treated oysters at the beginning and end of the feeding experiment. Concentrations of bacteria present in seawater samples from the 20 l aquaria were also determined just prior to the water being changed at 15 and 24 h.

Statistical analyses. The General Linear Models procedure (Statistical Analysis System Institute, Inc.) for an unbalanced ANOVA was used to analyze these data.

RESULTS

Distribution of ^{14}C in extracted *Spartina alterniflora*

Enzymatic digestion indicated that 85.4 % of the ^{14}C was present in material susceptible to cellulolytic breakdown (Table 1). Klason hydrolysis indicated that 92.3 % of ^{14}C was present in the polysaccharide fraction and 7.7 % in the lignin fraction. Protein and lipid contamination of the extracted material was negligible and 4.4 % of the ^{14}C present was available as a substrate for α -amylase.

In vitro determination of cellulase activity

Style and digestive gland cellulases liberated oligosaccharides from the extracted *Spartina alterniflora* material and from cellulose azure, indicating the

Table 1. *Spartina alterniflora*. Characterization of acid/alkali extracted, ^{14}C -labelled plants. All values are means \pm SE. Number of replicates was 3, except where indicated

Particle diameter	< 20 μm
Percent ^{14}C solubilised by	
(a) 24 h incubation in sterile seawater	1.6 \pm 0.28
(b) second acid/alkali extraction (see Fig. 1)	35.1 \pm 1.34 (n = 5)
Percent distribution of ^{14}C	
Lignin (Klason hydrolysis)	7.7 \pm 0.31
Polysaccharide (Klason hydrolysis)	92.3 \pm 0.31
Cellulose (cellulase digestion)	85.4 \pm 0.40
Starch (α -amylase digestion)	4.4 \pm 0.15
Lipid (methanol/chloroform extraction)	0.1 \pm 0.001
Protein	nd*
* nd = not detectable	

Table 2. *Crassostrea virginica*. Mean (\pm SE) μg glucose and μg total carbohydrate (glucose equivalents) released per mg protein in 24 h by style and digestive gland preparations. All assays were performed with 20 mg l^{-1} Thimerosal and 2 % toluene added. Concentration of all substrates was 0.5 % W/V. Number of replicates was 4, except where otherwise indicated

Substrate	μg glucose released mg^{-1} protein 24 h^{-1}	μg carbohydrate released mg^{-1} protein 24 h^{-1}
Style		
<i>Spartina alterniflora</i>	1.64 ± 0.48	88.3 ± 11.43
Cellulose azure	2.67 ± 0.38	517.3 ± 8.37
Cotton	0	0
Cellobiose	0	
Digestive gland		
Cellulose azure		0.16 ± 0.004 (n = 6)

presence of β -1,4-glucanases (Table 2). β -1,4-glucanase activity was also present in diverticula preparations as demonstrated by the breakdown of cellulose azure. In contrast, rates of glucose liberation from all substrates by style β -1,4-glucanases were negligible. No glucose was liberated from cellobiose by the style indicating an absence of β -glucosidase activity. Furthermore, the style preparation was not capable of breaking down cotton fibers under the conditions of the assay, suggesting an absence of C_1 cellulase.

In vivo digestion and absorption of ^{14}C

Five h after being fed ^{14}C labelled cellulosic material, oysters had ^{14}C specific activities significantly greater than background in both hemolymph plasma (ANOVA, $p < 0.002$) and pellet fractions (ANOVA, $p < 0.01$). The plasma ^{14}C activities (Table 3) did not significantly change between 5 and 24 h after feeding (ANOVA, $p > 0.05$). However, hemolymph pellet ^{14}C activities did significantly differ (ANOVA, $p = 0.018$) between the 3 sampling times (Table 3), with significantly higher activity at 5 h compared to 15 and 24 h (t-test; $p < 0.05$). Unfed control oysters held in the same 20 l volume of seawater as fed oysters had hemolymph plasma and pellet ^{14}C specific activities slightly above background. This accumulation of ^{14}C by unfed oysters may have been due to uptake of dissolved organic material (DO^{14}C) leaching into the seawater (Table 1) from particulate ^{14}C cellulosic material fed to the other oysters. Specific activities of DO^{14}C in $0.22 \mu\text{m}$ filtered seawater samples taken from both aquaria at 15 h and at the end of the 24 h incubation period (Table 4) were not significantly different (ANOVA, $p > 0.05$). Overall, the ^{14}C activities in plasma and pellet hemolymph fractions of ^{14}C fed oysters were significantly greater (ANOVA, $p = 0.002$ [plasma]; $p = 0.017$ [pellet]) than those of unfed oysters (Table 3).

The ^{14}C specific activities of tissue extracts of fed oysters was significantly greater (ANOVA, $p < 0.0001$) than those of unfed controls (Table 5). The ^{14}C specific activities of tissue extracts from unfed controls held in the same seawater as fed oysters were above background at the end of the 24 h incubation (Table 5), again indicating slight uptake of DO^{14}C .

^{14}C specific activities of hemolymph or tissue samples from oysters treated or untreated with antibiotics were not significantly different at the end of the feeding experiment (Tables 3 & 5). Antibiotics were effective in maintaining low concentrations of bacteria in the $0.22 \mu\text{m}$ filtered seawater of the experimental aquarium (Table 4). Bacteria were not detected in stomach fluid samples of antibiotic treated oysters, in contrast to a concentration of 1.62 ± 0.11 (SE) $\times 10^7$ bacteria ml^{-1} in stomach fluid samples of non-antibiotic treated oysters at the end of the 24 h feeding experiment.

Treatment of live oysters with the antibiotics rifampicin and chloramphenicol had no significant effect on cellulase activity of tissue homogenates, measured using cellulose azure as the substrate (Table 6; ANOVA, $p = 0.13$). When the antibiotics toluene and Thimerosal were included in enzyme assays, cellulolytic activity was significantly enhanced (ANOVA, $p < 0.0001$).

DISCUSSION

A Klason hydrolysis of the extracted ^{14}C labelled *Spartina alterniflora* indicated that 92.3 % of the ^{14}C was present in the polysaccharide fraction and that 7.7 % of the ^{14}C was incorporated in lignin (Table 1). Treatment with cellulase showed that 92.5 % of the ^{14}C in the polysaccharide fraction (as defined by Klason hydrolysis), or 85.4 % of the total particulate ^{14}C , was bound in material susceptible to cellulase action.

Table 3. *Crassostrea virginica*. Radioactivity (mean dpm \pm SE) ml⁻¹ hemolymph from oysters fed extracted ¹⁴C labelled *Spartina alterniflora* or from unfed oysters. Unfed and fed oysters were kept in the same aquarium during the experiment. Five mg l⁻¹ each of antibiotics rifampicin and chloramphenicol were added to filter-, sterilized seawater in which antibiotic treated oysters were held. Number of replicates in parenthesis. Dpm values corrected for background radioactivity

Time (h) bled after feeding	Radioactivity (DPM \pm SE)			
	Plasma of hemolymph		Pellet of hemolymph	
	Antibiotic- treated	Non-antibiotic- treated	Antibiotic- treated	Non-antibiotic- treated
Fed oysters				
5	732.2 \pm 240.1 (6)	813.2 \pm 150.1 (6)	34.2 \pm 8.1 (6)	88.7 \pm 33.9 (6)
15	1171.4 \pm 318.2 (5)	1247.8 \pm 735.2 (5)	20.8 \pm 5.4 (5)	18.4 \pm 5.8 (6)
24	1076.1 \pm 418.3 (11)	739.1 \pm 141.9 (11)	10.6 \pm 13.5 (11)	23.9 \pm 10.6 (11)
Unfed oysters				
15	12.3 \pm 1.8 (4)	12.5 \pm 1.4 (4)	12.8 \pm 1.5 (4)	14.5 \pm 1.5 (4)
24	6.5 \pm 1.7 (4)	5.4 \pm 1.5 (4)	-2.8 \pm 4.7 (4)	-1.5 \pm 2.2 (4)
ANOVA table for hemolymph plasma and pellet data presented above:				
Source	DF	SS	F	p
Plasma				
Fed/unfed	1	10 589 780	15.77	0.0002
Antibiotic treatment	1	51 571	0.08	0.783
Interaction	1	50 278	0.07	0.785
Error	57	38 266 187		
Pellet				
Fed/unfed	1	9 959	5.98	0.017
Antibiotic treatment	1	1 662	1.0	0.322
Interaction	1	1 067	0.64	0.427
Error	57	109 764		

Biologically degradable cellulose therefore formed a major component of the extracted *S. alterniflora*.

Our *in vitro* assays of style preparations indicated an absence of C₁ cellulase activity and the presence of β -1,4-glucanase activity. The style cellulases are therefore capable of breaking down amorphous cellulose but not crystalline cellulose. Lucas & Newell (1984) have also reported the presence of β -1,4-glucanase in style preparations of *Crassostrea virginica*. We found no β -glucosidase activity in the style, which is consistent with the findings of Mayasich & Smucker (1986) for *C. virginica*, however the latter researchers found β -glucosidase activity in the digestive diverticula.

Although an enzymatic approach to the study of cellulose digestion can demonstrate the presence or absence of cellulases, it is difficult to use the results to assess the quantitative importance of cellulase activity in the supply of carbon to the living oyster. A more realistic determination of the oyster's ability to digest

refractory cellulosic material and of the quantitative importance of this source of carbon to the animal's nutrition can be obtained by means of ¹⁴C radiotracer studies.

The results of our ¹⁴C radiotracer study demonstrate that *Crassostrea virginica* can digest and absorb small amounts of carbon from the cellulosic material prepared from *Spartina alterniflora*. ¹⁴C specific activities of hemolymph and pellet samples were significantly higher (ANOVA, $p = 0.0002$ and $p = 0.017$, respectively) in oysters fed ¹⁴C material than in unfed oysters (Table 3). These results demonstrate that the ¹⁴C labelled material was digested and absorbed into the hemolymph. Specific activities of ¹⁴C in tissue extracts of *C. virginica* were also significantly higher (ANOVA, $p < 0.0001$) in fed than in unfed oysters (Table 5).

Mean digestion efficiency of 1.3 % for oysters fed on the *Spartina alterniflora* material was low and even

this low level may have been due to the action of either cellulase and/or amylase enzymes (Table 1). However, our experimental oysters were pre-fed on natural ses-

ton. We are currently studying the possibility that there may be enhanced cellulose digestion in oysters pre-conditioned by feeding them a diet rich in cellulose.

Table 4. Bacteria numbers ml⁻¹ (mean ± SE) and radioactivity ml⁻¹ (mean dpm ± SE) in the seawater in which *Crassostrea virginica* fed ¹⁴C *Spartina alterniflora* material were held. Dpm values corrected for background activity. Number of replicates in parenthesis

Time (h) after feeding oysters	Aquarium	
	With antibiotics	Without antibiotics
Bacteria numbers		
15	7.4 ± 0.52 × 10 ⁴ (n = 16)	5.43 ± 0.55 × 10 ⁶ (n = 7)
24	8.1 ± 1.2 × 10 ⁴ (n = 8)	5.87 ± 0.63 × 10 ⁶ (n = 8)
Radioactivity		
15	13.0 ± 2.54 (n = 3)	13.8 ± 5.31 (n = 3)
24	13.8 ± 0.58 (n = 3)	11.0 ± 0.81 (n = 3)

Table 5. *Crassostrea virginica*. Total radioactivity of body tissue extracts of individual oysters (dpm oyster⁻¹). Oysters were either fed 2 016 333 dpm ¹⁴C labelled *Spartina alterniflora* or unfed. Unfed and fed oysters were kept in the same aquarium during the experiment. Five mg l⁻¹ each of antibiotics rifampicin and chloramphenicol were added to filter-sterilized seawater in which antibiotic-treated oysters were held. Dpm values corrected for background radioactivity

	Antibiotic-treated oysters		Non-antibiotic-treated oysters	
	DPM	Digestion efficiency	DPM	Digestion efficiency
Fed oysters				
	9 330	0.46	12 005	0.60
	9 468	0.47	12 696	0.63
	14 553	0.72	14 289	0.71
	15 357	0.76	18 866	0.90
	22 468	1.11	22 545	1.12
	25 706	1.27	24 988	1.24
	27 156	1.35	25 545	1.27
	27 468	1.36	34 477	1.71
	30 786	1.53	35 486	1.76
	41 047	2.04	36 334	1.80
	<u>53 025</u>	<u>2.63</u>	<u>50 822</u>	<u>2.50</u>
Mean ± SE	25 124 ± 4 025	1.25 %	26 186 ± 3 641	1.30 %
Unfed oysters				
	600		283	
	128		107	
	268		453	
	<u>372</u>		<u>436</u>	
Mean ± SE	342 ± 99		319 ± 80	
ANOVA table for data presented above:				
Source	DF	SS	F	p
Fed/unfed	1	3 762 373 174	30.18	0.0001
Antibiotic treatment	1	1 585 861	0.01	0.911
Interaction	1	1 724 499	0.01	0.907
Error	29	3 241 517 907		

Table 6. Cellulase activity measured by cellulose azure breakdown; expressed as μg carbohydrate released in 24 h per 100 mg (dry wt) of homogenized tissue (including style) from *Crassostrea virginica* treated or not treated with 5 mg l⁻¹ each of antibiotics rifampicin and chloramphenicol. Enzyme assays performed with or without 20 mg l⁻¹ Thimerosal and 2 % V/V toluene added to incubation tubes. Values are means \pm SE. Number of replicates in parenthesis

	Carbohydrate liberated (μg 100 mg ⁻¹ oyster tissue 24 h ⁻¹)			
	Antibiotic-treated		Non-antibiotic-treated	
Toluene/Thimerosal in assay tubes	1 506.2 \pm 115.2 (11)		999.0 \pm 140.9 (12)	
No toluene/Thimerosal in assay tubes	577.5 \pm 56.6 (11)		741.9 \pm 130.5 (12)	
ANOVA table for data presented above:				
Source	DF	SS	F	P
Oyster antibiotic treatment	1	337 197.7	2.44	0.1259
Toluene/Thimerosal in assay tubes	1	4 034 977.0	29.19	0.0001
Interaction	1	1 294 822.4	9.37	0.0038
Error	42	138 253.8		

Dietary induction of carbohydrases has been clearly demonstrated in other invertebrates, including copepods (Cox & Willason 1981), mixed samples of zooplankton (Mayzaud & Conover 1984) and amphipods (Borowsky 1984).

Treatment of oysters with antibiotics reduced bacterial populations in oyster stomach fluids below the concentrations detectable using direct counting techniques. However, antibiotic-treated oysters did not show a significantly reduced digestion or absorption of ¹⁴C from *Spartina alterniflora* into hemolymph and tissue fractions, compared with untreated oysters (Tables 3 & 5). These results indicate that the digestion of cellulosic plant material was not dependent on bacteria in the stomach.

Further evidence that bacteria did not significantly contribute to cellulose digestion in oysters can be seen in the measured cellulase activities of tissue homogenates from oysters used in the feeding experiment (Table 6). No significant differences in the cellulase activities of antibiotic-treated oysters and of untreated oysters were observed. Furthermore, incubation of tissue homogenates with cellulose azure in the absence of Thimerosal and toluene indicated that bacteria from the oyster did not enhance cellulose breakdown *in vitro*; indeed, cellulose breakdown was significantly greater (ANOVA, $p < 0.0001$) when toluene and Thimerosal were added to the assays to inhibit bacterial activity. This is possibly due to bacteria in the non-antibiotic treated assay tubes degrading cellulolytic enzymes from the oyster.

Lucas & Newell (1984) calculated that style carbohydrases acting directly on plant detrital material may provide up to 40 % of the carbon requirements of *Crassostrea virginica*, but their calculations did not take into account the possible rate-limiting step of β -glucosidase activity in the oyster's diverticula (Maya-

sich & Smucker 1986) nor the availability of refractory carbon to oysters in their natural environment. Therefore, we have calculated the potential contribution that refractory carbon can make to the carbon requirement of *C. virginica*, using data collected for a population of oysters in Broad Creek, a sub-estuary of Chesapeake Bay, Maryland.

At 25 °C under ambient conditions, an oyster of 1 g dry tissue weight has a filtration rate of 5 l h⁻¹ and an oxygen consumption of 1.0 ml O₂ h⁻¹ (Newell unpubl.) which is equivalent to 0.537 mg C respired h⁻¹. Thus the oyster's total carbon requirement is 0.714 mg h⁻¹, assuming that respiratory carbon requirements account for 75 % of the total carbon requirement (Bayne & Newell 1983, Lucas & Newell 1984). The mean concentration of detrital carbon in the seston available to oysters in Broad Creek is ca 0.3 mg l⁻¹ (Berg & Newell 1986). This was estimated from the carbon concentration present during the winter when the carbon-to-chlorophyll ratio was 275. Values greater than 100 for this ratio indicate that only a small proportion of the carbon originates from living phytoplankton (Chervin et al. 1981). Therefore, an oyster filtering 5 l h⁻¹ will ingest 1.5 mg detrital C h⁻¹. Assuming a digestion efficiency of 1.27 %, as determined for refractory cellulosic material, the oyster will absorb 0.019 mg C h⁻¹. This will only provide 2.7 % of the oyster's total carbon requirement.

These calculations indicate that it is unlikely that carbon from refractory plant detritus provides a large proportion of the oyster's carbon requirements. This suggests that the primary function of the oyster's cellulase enzymes may be to aid the digestion of other nutrients that are either contained within the cellulosic cell walls of algae or adsorbed onto cellulosic detrital particles.

The importance of carbon from *Spartina alterniflora*

detritus to estuarine animals, including *Crassostrea virginica*, has been studied using stable carbon isotope ratios by Haines (1977), Montague et al. (1981) and Hughes & Sherr (1983). They concluded that there was little evidence that the plant detritus was being directly assimilated by subtidal suspension-feeders in the Georgia marshes. Recently, Peterson et al. (1985) used the combined stable isotope ratios of carbon, nitrogen and sulphur to show that the ribbed mussel *Geukensia demissa*, living high within the marsh, may derive up to 80 % of its diet from *S. alterniflora*. According to our results, however, the stable isotope ratios of *C. virginica* living in salt marshes would not reflect direct utilization of refractory *S. alterniflora* detritus as a food source because of its poor digestibility by the oyster. Indirect utilization of *S. alterniflora* material via digestion and assimilation of microheterotrophic decomposers either free living or attached to plant detrital particles in the seston (Newell & Field 1983) is likely to be a more significant source of nutrition for oysters. Studies to determine the exact contributions that cellulose makes to the seasonal carbon requirements of natural populations of *C. virginica* and *G. demissa* are currently being undertaken in our laboratories.

Acknowledgements. We are grateful to Mrs Shelley Sulkin and Ms Linda Franklin for technical assistance, and to Dr. Ron Benner for determining the specific activity of the ^{14}C labeled cellulosic material. We thank Dr. Ray Thompson for constructive review of our manuscript. This research was supported by a grant (# OCE-8000264) from the Biological Oceanography division of the National Science Foundation to Drs. R. I. E. Newell, C. J. Langdon and H. Ducklow. The U.S. Government is authorized to produce and distribute reprints for governmental purposes, notwithstanding any copyright notation that may appear hereon.

LITERATURE CITED

- Bayne, B. L., Newell, R. C. (1983). Physiological energetics of marine molluscs. In: Saleuddin, A. S. M., Wilbur, K. M. (ed.) *The Mollusca*, Vol. 4, Physiology, Part 1. Academic Press, New York, p. 407–515
- Benner, R., Maccubbin, A. E., Hodson, R. E. (1984). Preparation, characterization and microbial degradation of specifically radiolabeled [^{14}C] lignocelluloses from marine and freshwater macrophytes. *Appl. environ. Microbiol.* 47: 381–389
- Berg, J. A., Newell, R. I. E. (1986). Temporal and spatial variations in the composition of seston available to the suspension feeder *Crassostrea virginica*. *Estuar. coast. Shelf. Sci.* 23: in press
- Borowsky, R. (1984). Environmental control of amylase phenotype in amphipods of the genus *Gammarus*. *Biol. Bull. mar. biol. Lab., Woods Hole* 167: 647–657
- Chervin, M. B., Malone, T. C., Neale, P. J. (1981). Interactions between suspended organic matter and copepod grazing in the plume of the Hudson River. *Estuar. coast. Shelf. Sci.* 13: 169–184
- Coston-Clements, L., Ferguson, R. L. (1985). Proximate composition and distribution of radioactive label in laboratory cultured *Spartina alterniflora* Loisel. *J. exp. mar. Biol. Ecol.* 93: 259–271
- Cox, J. L., Willason, S. W. (1981). Laminarase induction in *Calanus pacificus*. *Mar. Biol. Lett.* 2: 307–311
- Crosby, N. D., Reid, R. G. (1971). Relationships between food, physiology and cellulose digestion in the Bivalvia. *Can. J. Zool.* 49: 617–622
- Dubois, M., Gilles, K. A., Hamilton, J. K., Rebers, P. A., Smith, F. (1956). Colorimetric method for the determination of sugars and related substances. *Analyt. Chem.* 28: 350–356
- Ferguson, R. L., Williams, R. B. (1974). A growth chamber for production of ^{14}C -labelled salt marsh plant and its application to smooth cordgrass, *Spartina alterniflora* Loisel. *J. exp. mar. Biol. Ecol.* 14: 251–259
- Fisher, W. S., Newell, R. I. E. (1986). Salinity effects on granular hemocytes of the American oyster, *Crassostrea virginica*. *Biol. Bull. mar. biol. Lab., Woods Hole* 170: 122–134
- Fong, W., Mann, K. H. (1980). Role of gut flora in the transfer of amino acids through a marine food chain. *Can. J. Fish. aquat. Sci.* 37: 88–96
- Galtsoff, P. S. (1964). The American oyster *Crassostrea virginica* Gmelin. *Fish. Bull.* 64: 1–480
- Haines, E. B. (1977). The origin of detritus in Georgia salt marshes. *Oikos* 29: 254–260
- Halliwell, G. (1966). Solubilization of native and derived forms of cellulose by cell-free microbial enzymes. *Biochem. J.* 100: 315–320
- Heinle, D. R., Flemer, D. A. (1976). Flows of materials between poorly flooded tidal marshes and an estuary. *Mar. Biol.* 35: 359–373
- Holland, D. L., Gabbot, P. A. (1971). A micro-analytical scheme for the determination of protein, carbohydrate, lipid and RNA levels in marine invertebrate larvae. *J. mar. biol. Ass. U.K.* 51: 659–668
- Hughes, E. H., Sherr, E. B. (1983). Subtidal food webs in a Georgia estuary: ^{13}C analysis. *J. exp. mar. Biol. Ecol.* 67: 227–242
- Koopmans, J. J. C. (1970). Cellulases in molluscs. I. The nature of the cellulases in *Helix pomata* and *Cardium edule*. *Neth. J. Zool.* 20: 445–463
- Kristensen, J. H. (1972). Carbohydrates of some marine invertebrates with notes on their food and on the natural occurrence of the carbohydrates studied. *Mar. Biol.* 14: 130–142
- Langdon, C. J., Bolton, E. T. (1984). A microparticulate diet for a suspension feeding bivalve mollusc, *Crassostrea virginica* (Gmelin). *J. exp. mar. Biol. Ecol.* 82: 239–258
- Lowry, O. H., Josenbrough, N. J., Farr, A. L., Randall, R. J. (1951). Protein measurement with the folin phenol reagent. *J. biol. Chem.* 193: 265–275
- Lucas, M. I., Newell, R. C. (1984). Utilization of saltmarsh detritus by two estuarine bivalves: Carbohydrase activity of crystalline style enzymes of the oyster *Crassostrea virginica* (Gmelin) and the mussel *Geukensia demissa* (Dillwyn). *Mar. Biol. Lett.* 5: 275–290
- Mandels, M., Andreotti, R., Roche, C. (1976). Measurement of saccharifying cellulase. *Biotechnol. Bioeng. Symp.* 6: 21–33
- Mathers, N. F. (1973). A comparative histochemical survey of enzymes associated with the processes of digestion in *Ostrea edulis* and *Crassostrea angulata* (Mollusca: Bivalvia). *J. Zool. Lond.* 169: 169–179
- Mayasich, S. A., Smucker, R. A. (1986). Glycosidases in the

- American oyster, *Crassostrea virginica* Gmelin, digestive tract. *J. exp. mar. Biol. Ecol.* 95: 95-98
- Mayzaud, P., Conover, R. J. (1984). Distribution of digestive enzymes in zooplankton during the spring bloom in a Nova Scotia inlet. *Can. J. Fish. Aquat. Sci.* 41: 245-252
- McIntire, G. L., Dunstan, W. M. (1976). Non-structural carbohydrates in *Spartina alterniflora* Loisel. *Botanica mar.* 19: 93-96
- Montague, C. L., Bunker, S. M., Haines, E. B., Pace, M. L., Wetzel, R. L. (1981). Aquatic macroconsumers. In: Pomeroy, L. R., Wiegert, R. G. (ed.) *The ecology of a salt marsh*. Springer-Verlag, New York, p. 69-86
- Morton, B. (1983). Feeding and digestion in Bivalvia. In: Saleuddin, A. S. M., Wilbur, K. M. (ed.) *The Mollusca*, Vol. 5., Physiology, Part 2. Academic Press, New York, p. 65-147
- Mountford, K. (1980). Aspects of the ecology of a small estuarine embayment. *Mar. Biol.* 61: 53-67
- Newell, R. C., Field, J. G. (1983). The contribution of bacteria and detritus to carbon and nitrogen flow in benthic community. *Mar. Biol. Lett.* 4: 249-257
- Payne, D. W., Thorpe, N. A., Donaldson, E. M. (1972). Cellulolytic activity and a study of the bacterial population in the digestive tract of *Scrobicularia plana* (Da Costa). *Proc. malac. Soc. Lond.* 40: 147-160
- Peterson, B. J., Howarth, R. W., Garritt, R. H. (1985). Multiple stable isotopes used to trace the flow of organic matter in estuarine food webs. *Science* 227: 1361-1363
- Pickral, J. C., Odum, W. E. (1976). Benthic detritus in a saltmarsh tidal creek. In: Wiley, M. (ed.) *Estuarine processes*, Vol. II. Academic Press, New York, p. 280-292
- Porter, K. G., Feig, Y. S. (1980). The use of DAPI for identifying and counting aquatic microflora. *Limnol. Oceanogr.* 25: 943-948
- Rice, D. L. (1982). The detritus nitrogen problem: new observations and perspectives from organic geochemistry. *Mar. Ecol. Prog. Ser.* 9: 153-162
- Smith, K. K., Good, R. E., Good, N. F. (1979). Production dynamics for above and below ground components of a New Jersey *Spartina alterniflora* tidal marsh. *Estuar. coast. Shelf Sci.* 9: 189-201
- Squiers, E. R., Good, R. E. (1974). Seasonal changes in the productivity, caloric content and chemical composition of a population of salt marsh cordgrass (*Spartina alterniflora*). *Chesapeake Sci.* 15: 63-71
- Strickland, J. D. H., Parsons, T. R. (1972). A practical handbook of seawater analysis. *Bull. Fish. Res. Bd Can.* 167: 1-310
- Teal, J. M. (1962). Energy flow in the salt marsh ecosystem of Georgia. *Ecology* 43: 614-624
- Waterbury, J., Calloway, C. B., Turner, R. D. (1983). A cellulolytic nitrogen-fixing bacterium cultured from the gland of *Deshayes* in shipworms (Bivalvia: Teredinidae). *Science* 221: 1401-1403
- Whistler, R. L. (1963). *Methods in carbohydrate chemistry*, Vol III, Cellulose. Academic Press, New York
- Wojtowicz, M. B. (1972). Carbohydrases of the digestive gland and the crystalline style of the Atlantic deep-sea scallop (*Placopecten magellanicus* Gmelin). *Comp. Biochem. Physiol.* 43 A: 131-141