



This is a repository copy of *Dilatonic black holes in higher curvature string gravity. II. Linear stability.*

White Rose Research Online URL for this paper:
<http://eprints.whiterose.ac.uk/102592/>

Version: Submitted Version

Article:

Kanti, P., Mavromatos, N.E., Rizos, J. et al. (2 more authors) (1998) Dilatonic black holes in higher curvature string gravity. II. Linear stability. PHYS REV D, 57 (10). pp. 6255-6264. ISSN 0556-2821

<https://doi.org/10.1103/PhysRevD.57.6255>

Copyright 1998 American Physical Society. Published version, Phys. Rev. D 57, 6255, available at <http://dx.doi.org/10.1103/PhysRevD.57.6255>

Reuse

Unless indicated otherwise, fulltext items are protected by copyright with all rights reserved. The copyright exception in section 29 of the Copyright, Designs and Patents Act 1988 allows the making of a single copy solely for the purpose of non-commercial research or private study within the limits of fair dealing. The publisher or other rights-holder may allow further reproduction and re-use of this version - refer to the White Rose Research Online record for this item. Where records identify the publisher as the copyright holder, users can verify any specific terms of use on the publisher's website.

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.



eprints@whiterose.ac.uk
<https://eprints.whiterose.ac.uk/>

Dilatonic Black Holes in Higher-Curvature String Gravity II: Linear Stability

P. Kanti^a, N.E. Mavromatos^b, J. Rizos^a,
K. Tamvakis^a and E. Winstanley^b

^a Division of Theoretical Physics, Physics Department,
University of Ioannina, Ioannina GR 451 10, Greece,

^b University of Oxford, Department of Physics (Theoretical Physics),
1 Keble Road, Oxford. OX1 3NP U.K.

Abstract

We demonstrate linear stability of the dilatonic Black Holes appearing in a string-inspired higher-derivative gravity theory with a Gauss-Bonnet curvature-squared term. The proof is accomplished by mapping the system to a one-dimensional Schrödinger problem which admits no bound states. This result is important in that it constitutes a linearly stable example of a black hole that bypasses the ‘no-hair conjecture’. However, the dilaton hair is *secondary* in the sense that it is not accompanied by any new quantum number for the black hole solution.

March 1997

1 Introduction

In ref. [1] we have presented analytic arguments in favour of, and demonstrated numerically, the existence of dilatonic black hole solutions with non-trivial scalar hair, in a string-inspired higher-derivative gravity theory with a Gauss Bonnet (GB) curvature-squared term. The numerical solutions clearly demonstrated the existence of a regular event horizon and asymptotic flatness of the four-dimensional spherically-symmetric space-time configurations considered in the analysis. There is a non-trivial dilaton (global) charge, which however is related to the ADM mass of the black hole, and hence the hair is of secondary type [2]: the gravitational field acts as a source for the scalar hair and one does not obtain a new independent set of quantum numbers characterizing the black hole¹. Subsequent to the work of [1], other researchers confirmed these results by discussing the internal structure of the solutions behind the horizon, and demonstrating numerically the existence of curvature singularities [5]. Also an extension of the analysis of ref. [1] to incorporate gauge fields became possible [6, 7].

The key feature for the existence of such hairy black holes is the bypass of the no-scalar-hair theorems [8] due to the fact that, as a result of the GB term, the scalar field stress tensor becomes *negative* near the horizon [1], thereby violating one of the main assumptions in the proof of the no-hair theorem [8]. An additional element was the fact that the higher-derivative GB term provides a sort of ‘repulsion’ that balances the gravitational attraction of the standard Einstein terms, and a black hole is formed. In this respect the GB term plays a similar rôle to the *non-Abelian* gauge field kinetic terms in Einstein-Yang-Mills-Higgs theories [4], which are also notable exceptions of the no-scalar-(Higgs)-hair theorem.

An important question which arises concerns the stability of the dilatonic black holes. In the Yang-Mills case, the structures are similar to the sphaleron solutions of flat-space Yang-Mills theories, and thus unstable [4]. This may be easily understood by the fact that the black hole solutions owe their existence to a delicate balance between the gravitational attraction and the Yang-Mills repulsive forces. On the other hand, the dilatonic black hole solutions are entirely due to the existence of a *single* force, that of gravity. This already

¹A similar situation occurs in the Higgs hair of the Einstein-Yang-Mills-Higgs systems [3, 4], where the Yang-Mills field act as a source for the non-trivial configurations of the Higgs field outside the horizon.

prompts one to think that such structures might be stable.

It is the purpose of this article to argue that the dilatonic black holes are indeed stable under linear time-dependent perturbations of the classical solutions. To this end, we shall map the system of gravitational-dilaton equations for spherically-symmetric solutions into a one-dimensional Schrödinger problem, where the instabilities are equivalent to bound states. We shall prove that our dilaton-graviton system admits *no bound states*. This result is important, since it constitutes an example of a hairy black hole structure that appears to be, at least linearly, stable. Its importance is also related to the fact that such higher-curvature gravity theories are effective theories obtained from superstrings, which may imply that there is plenty of room in the gravitational sector of string theory to allow for physically sensible situations that are not covered by the no-hair theorem as stated [8]. Unfortunately, at present non-linear stability of the dilaton-graviton-GB system cannot be checked analytically, and is left for future investigations.

2 Relevant Formalism

We start by considering the action of the Einstein-Dilaton-Gauss-Bonnet (EDGB) theory:

$$S = \int d^4x \sqrt{-g} \left(\frac{R}{2} + \frac{1}{4} \partial_\mu \phi \partial^\mu \phi + \frac{\alpha' e^\phi}{8g^2} \mathcal{R}_{GB}^2 \right) \quad (1)$$

where

$$\mathcal{R}_{GB}^2 = R_{\mu\nu\rho\sigma} R^{\mu\nu\rho\sigma} - 4R_{\mu\nu} R^{\mu\nu} + R^2 \quad (2)$$

The spherically-symmetric ansatz for the metric takes the form

$$ds^2 = e^{\Gamma(r,t)} dt^2 - e^{\Lambda(r,t)} dr^2 - r^2 (d\theta^2 + \sin^2 \theta d\varphi^2) \quad (3)$$

The equations of motion derived from (1) are:

$$\begin{aligned} \phi'' + \phi' \left(\frac{\Gamma'}{2} - \frac{\Lambda'}{2} + \frac{2}{r} \right) - e^{\Lambda-\Gamma} \left[\ddot{\phi} + \frac{\dot{\phi}}{2} (\dot{\Lambda} - \dot{\Gamma}) \right] = \\ \frac{\alpha' e^\phi}{g^2 r^2} \left\{ \Gamma' \Lambda' e^{-\Lambda} - \dot{\Lambda}^2 e^{-\Gamma} + (1 - e^{-\Lambda}) \left[\Gamma'' + \frac{\Gamma'}{2} (\Gamma' - \Lambda') \right] \right. \\ \left. - (1 - e^{-\Lambda}) e^{\Lambda-\Gamma} \left[\ddot{\Lambda} + \frac{\dot{\Lambda}}{2} (\dot{\Lambda} - \dot{\Gamma}) \right] \right\} \quad (4) \end{aligned}$$

$$\begin{aligned} \Lambda' \left[1 + \frac{\alpha' e^\phi \phi'}{2g^2 r} (1 - 3e^{-\Lambda}) \right] &= \frac{r}{4} (\phi'^2 + e^{\Lambda-\Gamma} \dot{\phi}^2) + \frac{1 - e^\Lambda}{r} \\ &+ \frac{\alpha' e^\phi}{g^2 r} (1 - e^{-\Lambda}) \left[\phi'' + \phi'^2 - \frac{\dot{\phi} \dot{\Lambda}}{2} e^{\Lambda-\Gamma} \right] \end{aligned} \quad (5)$$

$$\begin{aligned} \Gamma' \left[1 + \frac{\alpha' e^\phi \phi'}{2g^2 r} (1 - 3e^{-\Lambda}) \right] &= \frac{r}{4} (\phi'^2 + e^{\Lambda-\Gamma} \dot{\phi}^2) + \frac{e^\Lambda - 1}{r} \\ &+ \frac{\alpha' e^\phi}{g^2 r} (1 - e^{-\Lambda}) e^{\Lambda-\Gamma} \left[\ddot{\phi} + \dot{\phi}^2 - \frac{\dot{\phi} \dot{\Gamma}}{2} \right] \end{aligned} \quad (6)$$

$$\dot{\Lambda} \left[1 + \frac{\alpha' e^\phi \phi'}{2g^2 r} (1 - 3e^{-\Lambda}) \right] = \frac{r \dot{\phi} \phi'}{2} + \frac{\alpha' e^\phi}{g^2 r} (1 - e^{-\Lambda}) \left(\dot{\phi} \phi' + \dot{\phi} - \frac{\dot{\phi} \Gamma'}{2} \right) \quad (7)$$

$$\begin{aligned} \Gamma'' + \frac{\Gamma'}{2} (\Gamma' - \Lambda') + \frac{(\Gamma' - \Lambda')}{r} - e^{\Lambda-\Gamma} \left[\ddot{\Lambda} + \frac{\dot{\Lambda}}{2} (\dot{\Lambda} - \dot{\Gamma}) \right] &= \\ + \frac{1}{2} (e^{\Lambda-\Gamma} \dot{\phi}^2 - \phi'^2) + \frac{8\alpha'}{r} e^{-\Lambda} \left\{ f' \Gamma'' + f'' \Gamma' + \frac{f' \Gamma'}{2} (\Gamma' - 3\Lambda') \right. \\ \left. + e^{\Lambda-\Gamma} \left[\Lambda' \ddot{f} - 2\dot{f}' \dot{\Lambda} + \frac{f' \dot{\Lambda}}{2} (\Gamma' + \Lambda') - f' \ddot{\Lambda} + (f' \dot{\Lambda} - \Lambda' \dot{f}) \frac{(\dot{\Gamma} + \dot{\Lambda})}{2} \right] \right\} \end{aligned} \quad (8)$$

where $f = e^\phi / 8g^2$.

For later use we note that for the derivatives of the dilaton field at the horizon in the static case [1] one has the following behaviour:

$$\phi'_h = \frac{g^2}{\alpha'} r_h e^{-\phi_h} \left(-1 \pm \sqrt{1 - \frac{6(\alpha')^2 e^{2\phi_h}}{g^4 r_h^4}} \right) \quad (9)$$

which implies that black hole solutions exists only if:

$$e^{\phi_h} \leq \frac{g^2 r_h^2}{\alpha' \sqrt{6}} \quad (10)$$

We now notice that [1] only one of the two branches of solutions in (9), the one with the + sign, leads to asymptotic flatness of the fields, and this is the branch we shall consider here. The equation for ϕ'' near r_h is:

$$\phi'' = -\frac{1}{2} \frac{(\frac{\alpha'}{g^2} e^\phi \phi' + 2r)(6\frac{\alpha'}{g^2} e^\phi + \frac{\alpha'}{g^2} e^\phi \phi'^2 r^2 + 2\phi' r^3)}{-6\frac{\alpha'^2}{g^4} e^{2\phi} + \frac{\alpha'}{g^2} e^\phi \phi' r^3 + 2r^4} \Gamma' + O(1) \quad (11)$$

which is finite $O(1)$, as a result of (9). The asymptotic form of the dilaton field and the metric components near the even horizon $r \simeq r_h$ are:

$$\begin{aligned} e^{-\Lambda(r)} &= \lambda_1(r - r_h) + \lambda_2(r - r_h)^2 + \dots \\ e^{\Gamma(r)} &= \gamma_1(r - r_h) + \gamma_2(r - r_h)^2 + \dots \\ \phi(r) &= \phi_h + \phi'_h(r - r_h) + \phi''_h(r - r_h)^2 + \dots \end{aligned} \quad (12)$$

where:

$$\lambda_1 = 2/(\alpha' e^{\phi_h} \phi'_h / g^2 + 2r_h) , \quad (13)$$

and γ_1 is an arbitrary finite *positive* integration constant, which cannot be fixed by the equations of motion, since the latter involve only $\Gamma'(r)$ and not $\Gamma(r)$. This constant is fixed by the asymptotic limit of the solutions at infinity. At infinity, one uses the following asymptotic behaviour:

$$\begin{aligned} e^{\Lambda(r)} &= 1 + \frac{2M}{r} + \frac{16M^2 - D^2}{4r^2} + O\left(\frac{1}{r^3}\right) \\ e^{\Gamma(r)} &= 1 - \frac{2M}{r} + O\left(\frac{1}{r^3}\right) \\ \phi(r) &= \phi_\infty + \frac{D}{r} + \frac{MD}{r^2} + O\left(\frac{1}{r^3}\right) \end{aligned} \quad (14)$$

which guarantees asymptotic flatness of the space time. Above, M denotes the ADM mass of the black hole, and D the dilaton charge. The numerical analysis of [1] has shown that M and D are not independent quantities, thereby leading to the secondary nature of the dilaton hair [1, 2].

The black hole solutions of [1] are characterized uniquely by two parameters (ϕ_h, r_h) . Note, however, that the equations of motion remain invariant

under a shift $\phi \rightarrow \phi + \phi_0$ as long as it is accompanied by a radial rescaling $r \rightarrow r e^{\phi_0/2}$. Due to the above invariance it is sufficient to vary only one of r_h and ϕ_h . In the present analysis we choose to keep r_h fixed ($r_h = 1$), and to vary ϕ_h . A typical family of solutions has dilaton configurations (outside the horizon) of the form depicted in Figure 1. The solutions are characterized by negative ϕ_h , and *monotonic, non-intersecting* behaviour from r_h until infinity. These are essential features of the solutions, that we shall make use of in our linear stability analysis.

3 Linear Stability Analysis

We now consider perturbing the equations (4)-(8) by time-dependent *linear* perturbations of the form:

$$\begin{aligned}\Gamma(r, t) &= \Gamma(r) + \delta\Gamma(r, t) = \Gamma(r) + \delta\Gamma(r)e^{i\sigma t} \\ \Lambda(r, t) &= \Lambda(r) + \delta\Lambda(r, t) = \Lambda(r) + \delta\Lambda(r)e^{i\sigma t} \\ \phi(r, t) &= \phi(r) + \delta\phi(r, t) = \phi(r) + \delta\phi(r)e^{i\sigma t}\end{aligned}\tag{15}$$

where the variations $\delta\Gamma$, $\delta\Lambda$ and $\delta\phi$ are assumed small (bounded), and the quantities without a δ prefactor denote classical time-independent solutions of the equations (4)-(8). The above *harmonic* time dependence is sufficient for a *linear stability* analysis, since by assumption the linear variations are characterized by a well-defined Fourier expansion in time t [4, 9]. The linear stability analysis proceeds by mapping the algebraic system of variations of the equations of motion under consideration to a stationary one-dimensional Schrödinger problem, in an appropriate potential well, in which the ‘squared frequencies’ σ^2 will constitute the energy eigenvalues. In the present problem, the ‘wavefunction’ turns out to be the dilaton linear variation $\delta\phi(r)e^{i\sigma t}$. Instabilities, then, correspond to negative energy eigenstates (‘bound states’), i.e. imaginary frequencies σ . As we shall show in this article, for the system of variations corresponding to (4)-(8),(15) the corresponding Schrödinger problem admits *no bound states*, thereby proving the linear stability of the EDGB hairy black holes.

As we shall discuss below, some technical complications arise, as usual [4], in the above process due to the fact that the ‘naive’ stationary Schrödinger equation with respect to the original coordinate r is not well defined at some

points of the domain of $r \in [r_h, \infty)$. This necessitates a change of coordinates in such a way that the resulting Schrödinger problem is well defined. A convenient choice is provided by the so-called ‘tortoise’ coordinate r^* [4, 9], which is defined in such a way so that the domain $[r_h, \infty)$ is extended over the entire real axis $r \rightarrow r^* \in (-\infty, \infty)$. In our specific problem, we shall define the ‘tortoise’ co-ordinate r^* as [4]:

$$\frac{dr^*}{dr} = e^{-(\Gamma-\Lambda)/2} \quad (16)$$

As we shall show, then, the associated stationary Schrödinger equation, pertaining to the dilaton variation in (15), will be of the form:

$$p_*^2 u + \mathcal{V}(r^*)u(r^*) = -\sigma^2 u(r^*) \quad ; \quad p_* \equiv \frac{d}{dr^*} \quad (17)$$

where $u(r^*)$ is related to the dilaton variation $\delta\phi(r)$ in (15), and the potential \mathcal{V} is well defined over the domain of validity of r^* .

Let us now proceed with our analysis. The perturbed equations (4)-(8), under the variations (15) read:

$$\begin{aligned} & \delta\phi'' + \delta\phi' \left(\frac{\Gamma'}{2} - \frac{\Lambda'}{2} + \frac{2}{r} \right) - \delta\phi \left[\phi'' + \phi' \left(\frac{\Gamma'}{2} - \frac{\Lambda'}{2} + \frac{2}{r} \right) \right] - e^{\Lambda-\Gamma} \delta\ddot{\phi} \\ & + \delta\Gamma' \left\{ \frac{\phi'}{2} - \frac{\alpha'e^\phi}{g^2 r^2} \left[\Lambda'e^{-\Lambda} + (1 - e^{-\Lambda}) \left(\Gamma' - \frac{\Lambda'}{2} \right) \right] \right\} + \delta\Lambda \frac{\alpha'e^\phi}{g^2 r^2} \left\{ \Gamma'\Lambda'e^{-\Lambda} \right. \\ & \left. - e^{-\Lambda} \left[\Gamma'' + \frac{\Gamma'}{2} (\Gamma' - \Lambda') \right] \right\} - \delta\Lambda' \left\{ \frac{\phi'}{2} + \frac{\alpha'e^\phi}{g^2 r^2} \left[\Gamma'e^{-\Lambda} - (1 - e^{-\Lambda}) \frac{\Gamma'}{2} \right] \right\} \\ & + \frac{\alpha'e^\phi}{g^2 r^2} (1 - e^{-\Lambda}) e^{\Lambda-\Gamma} \delta\ddot{\Lambda} - \frac{\alpha'e^\phi}{g^2 r^2} (1 - e^{-\Lambda}) \delta\Gamma'' = 0, \end{aligned} \quad (18)$$

$$\begin{aligned} & \delta\Lambda' \left[1 + \frac{\alpha'e^\phi\phi'}{2g^2 r} (1 - 3e^{-\Lambda}) \right] + \delta\phi' \left[-\frac{r\phi'}{2} + \frac{\alpha'e^\phi}{2g^2 r} \Lambda' (1 - 3e^{-\Lambda}) \right] \\ & + \delta\Lambda \left\{ \frac{e^\Lambda}{r} + \frac{\alpha'e^{\phi-\Lambda}}{g^2 r} \left[\frac{3\phi'\Lambda'}{2} - (\phi'' + \phi'^2) \right] \right\} + \delta\phi \frac{\alpha'e^\phi}{2g^2 r} \phi' \Lambda' (1 - 3e^{-\Lambda}) \end{aligned}$$

$$-\frac{\alpha'e^\phi}{g^2r}(1-e^{-\Lambda})[\delta\phi''+2\phi'\delta\phi'+\delta\phi(\phi''+\phi'^2)]=0, \quad (19)$$

$$\begin{aligned} & \delta\Gamma' \left[1 + \frac{\alpha'e^\phi\phi'}{2g^2r}(1-3e^{-\Lambda}) \right] + \delta\Lambda \left[-\frac{e^\Lambda}{r} + \frac{3\alpha'e^\phi}{2g^2r}\phi'\Gamma'e^{-\Lambda} \right] \\ & + \delta\phi \frac{\alpha'e^\phi}{2g^2r}\phi'\Gamma'(1-3e^{-\Lambda}) + \delta\phi' \left[-\frac{r\phi'}{2} + \frac{\alpha'e^\phi}{2g^2r}\Gamma'(1-3e^{-\Lambda}) \right] \\ & - \frac{\alpha'e^\phi}{g^2r}(1-e^{-\Lambda})e^{\Lambda-\Gamma}\delta\ddot{\phi} = 0 \end{aligned} \quad (20)$$

$$\delta\dot{\Lambda} \left[1 + \frac{\alpha'e^\phi\phi'}{2g^2r}(1-3e^{-\Lambda}) \right] = \frac{r\phi'}{2}\delta\dot{\phi} + \frac{\alpha'e^\phi}{g^2r}(1-e^{-\Lambda}) \left(\delta\dot{\phi}\phi' + \delta\dot{\phi}' - \delta\dot{\phi}\frac{\Gamma'}{2} \right) \quad (21)$$

$$\begin{aligned} & \delta\Gamma'' \left(1 - \frac{\alpha'e^{\phi-\Lambda}}{g^2r}\phi' \right) + \frac{\delta\Gamma'}{2}(\Gamma' - \Lambda') + (\delta\Gamma' - \delta\Lambda') \left(\frac{\Gamma'}{2} + \frac{1}{r} \right) - e^{\Lambda-\Gamma}\delta\ddot{\Lambda} \\ & - \frac{\alpha'e^{\phi-\Lambda}}{g^2r} \left\{ \delta\phi'\Gamma'' + (\delta\phi'' + 2\phi'\delta\phi')\Gamma' + \delta\Gamma'(\phi'' + \phi'^2) + \frac{\delta\phi'\Gamma'}{2}(\Gamma' - 3\Lambda') \right. \\ & \left. + \frac{\phi'\delta\Gamma'}{2}(\Gamma' - 3\Lambda') + \frac{\phi'\Gamma'}{2}(\delta\Gamma' - 3\delta\Lambda') + e^{\Lambda-\Gamma}(\Lambda'\delta\ddot{\phi} - \phi'\delta\ddot{\Lambda}) \right\} + \phi'\delta\phi' \\ & - \frac{\alpha'e^{\phi-\Lambda}}{g^2r} \left[\phi'\Gamma'' + \Gamma'(\phi'' + \phi'^2) + \frac{\phi'\Gamma'}{2}(\Gamma' - 3\Lambda') \right] (\delta\phi - \delta\Lambda) = 0 \end{aligned} \quad (22)$$

We can integrate eq.(21) to obtain

$$\begin{aligned} \delta\Lambda \left[1 + \frac{\alpha'e^\phi\phi'}{2g^2r}(1-3e^{-\Lambda}) \right] & = \frac{\alpha'e^\phi}{g^2r}(1-e^{-\Lambda}) \left(\delta\phi\phi' + \delta\phi' - \delta\phi\frac{\Gamma'}{2} \right) \\ & + \frac{r\phi'}{2}\delta\phi + \mu(r) \end{aligned} \quad (23)$$

where $\mu(r)$ is an arbitrary function of r which can be set equal to zero if we require that $\delta\Lambda = 0$ when $\delta\phi = 0$. An independent check of this assertion can be obtained as follows: first we differentiate (23) with respect to r and then use eq.(19), as well as the time-independent equations of motion. Thus, we obtain the following differential equation for $\mu(r)$

$$\mu'(r) + \mu(r) \left(\frac{\Gamma'}{2} - \frac{\Lambda'}{2} + \frac{1}{r} \right) = 0 \quad (24)$$

This can be integrated to give $\mu(r) \sim e^{(\Lambda-\Gamma)/2}/r$. When $r \rightarrow r_h$, $e^{(\Lambda-\Gamma)/2} \rightarrow \infty$ and $\mu \rightarrow \infty$, which is incompatible with the assumption of a small $\delta\Lambda(r)$ required for the linear stability analysis. Rejecting this solution, we are left with the trivial one, $\mu = 0$. For calculational convenience we also set $\alpha'/g^2 = 1$, from now on, which is achieved by an appropriate rescaling of the dilaton field.

Rearranging the above equations, one obtains, after a tedious but straightforward procedure, an equation for $\delta\phi$ which has the following structure:

$$A \delta\phi'' + 2B \delta\phi' + C \delta\phi + \sigma^2 E \delta\phi = 0 \quad (25)$$

where A , B , C , and E are rather complicated functions of ϕ , ϕ' , ϕ'' , Λ , Λ' , Γ , Γ' and Γ'' . In the limit $r \rightarrow r_h$ these coefficients take the form

$$A = \frac{2 \sqrt{1 - \frac{6e^{2\phi_h}}{r_h^4}}}{1 + \sqrt{1 - \frac{6e^{2\phi_h}}{r_h^4}}} + O(r - r_h) \quad (26)$$

$$B = \frac{\sqrt{1 - \frac{6e^{2\phi_h}}{r_h^4}}}{1 + \sqrt{1 - \frac{6e^{2\phi_h}}{r_h^4}}} \frac{1}{(r - r_h)} + O(1) \quad (27)$$

$$C = \frac{2 e^{2\phi_h}}{r_h^4 (1 + \sqrt{1 - \frac{6e^{2\phi_h}}{r_h^4}})} \frac{1}{(r - r_h)^2} + O\left(\frac{1}{r - r_h}\right) \quad (28)$$

$$E = \frac{r_h \sqrt{1 - \frac{6e^{2\phi_h}}{r_h^4}}}{\gamma_1} \frac{1}{(r - r_h)^2} + O\left(\frac{1}{r - r_h}\right) \quad (29)$$

where we have used the asymptotic behaviour (12) near the event horizon.

On the other hand, when $r \rightarrow \infty$ we obtain

$$A = 1 + \frac{e^{\phi_\infty} DM}{r^4} + O\left(\frac{1}{r^5}\right) \quad (30)$$

$$B = \frac{1}{r} + \frac{M}{r^2} + O\left(\frac{1}{r^5}\right) \quad (31)$$

$$C = \frac{D^2}{2r^4} + O\left(\frac{1}{r^5}\right) \quad (32)$$

$$E = 1 + \frac{4M}{r} + \frac{4M^2}{r^2} + \frac{e^{\phi_\infty} DM}{r^4} + O\left(\frac{1}{r^5}\right) \quad (33)$$

where we have used the asymptotic behaviour (14) near infinity.

As we can see from (26)–(29) the coefficients of the Schrödinger equation (25) are not finite at the boundary $r = r_h$, where the variation $\delta\phi$ is bounded. As mentioned previously, to arrive at a well-defined Schrödinger problem, one can use the ‘tortoise’ coordinate (16). Then, the perturbed equation for the dilaton field takes the form

$$A \frac{d^2 \delta\phi}{dr^{*2}} + \left[2B e^{(\Gamma-\Lambda)/2} - \frac{A}{2} \frac{d(\Gamma-\Lambda)}{dr^*} \right] \frac{d\delta\phi}{dr^*} + e^{\Gamma-\Lambda} (C + \sigma^2 E) \delta\phi = 0 \quad (34)$$

or

$$\mathcal{A} \frac{d^2 \delta\phi}{dr^{*2}} + 2\mathcal{B} \frac{d\delta\phi}{dr^*} + (\mathcal{C} + \sigma^2 \mathcal{E}) \delta\phi = 0 \quad (35)$$

where

$$\mathcal{A} = A \quad , \quad \mathcal{B} = B e^{(\Gamma-\Lambda)/2} - \frac{A}{4} \frac{d(\Gamma-\Lambda)}{dr^*}, \quad (36)$$

$$\mathcal{C} = e^{\Gamma-\Lambda} C \quad , \quad \mathcal{E} = e^{\Gamma-\Lambda} E \quad (37)$$

Note that near the horizon

$$e^{(\Gamma-\Lambda)/2} = \sqrt{\gamma_1 \lambda_1} (r - r_h) + O(r - r_h)^2 \quad (38)$$

$$\frac{d(\Gamma-\Lambda)}{dr^*} = 2\sqrt{\gamma_1 \lambda_1} + O(r - r_h) \quad (39)$$

where λ_1 is given by (13). As a result, all the coefficients in eq. (34) are now well-behaved near the horizon r_h . In order to eliminate the term proportional to $\delta\phi'$, we first divide eq. (34) by \mathcal{A} and then we use the function

$$F = \exp\left(\int_{-\infty}^{r^*} \frac{\mathcal{B}}{\mathcal{A}} dr^{*'}\right) \quad (40)$$

Then, the equation for $\delta\phi$ takes the form

$$p_*^2 u + \left[\frac{\mathcal{C}}{\mathcal{A}} + \sigma^2 \frac{\mathcal{E}}{\mathcal{A}} - \frac{\mathcal{B}^2}{\mathcal{A}^2} - p_* \left(\frac{\mathcal{B}}{\mathcal{A}} \right) \right] u = 0 \quad (41)$$

where

$$p_* \equiv \frac{d}{dr^*} \quad (42)$$

and we have set $u = F \delta\phi$.

It is straightforward to see that

$$\frac{\mathcal{B}}{\mathcal{A}} \rightarrow 0 \quad \text{for } r \rightarrow r_h \quad \text{and } r \rightarrow \infty \quad (43)$$

independently of r_h, ϕ_h . In addition

$$\frac{\mathcal{B}}{\mathcal{A}} \frac{dr^*}{dr} = \text{finite} \quad \text{for } r \rightarrow r_h \quad (44)$$

Moreover, as shown in Fig. 2, the function \mathcal{B}/\mathcal{A} is well-behaved over the entire domain outside the horizon, implying the integrability of the function F . Also, the quantity \mathcal{E}/\mathcal{A} is finite and always *positive* outside the horizon of the numerical black-hole solutions of [1] (see Fig. 3). It is also immediately seen that the eigenfunction u_0 , corresponding to the eigenvalue $\sigma = 0$, can be constructed out of the difference of any two curves in Fig. 1. From the *monotonic* and *non-intersecting* nature of the various members of the family of the numerical solutions of Fig. 1, then, one can conclude that u_0 has *no nodes* in the domain $r^* \in (-\infty, \infty)$, and that $p_*^2 u_0 / u_0 = e^{\Gamma - \Lambda} [\frac{1}{2}(\Gamma' - \Lambda')u_0' + u_0''] / u_0$ is *finite*. This, together with the finite and smooth form of \mathcal{B}/\mathcal{A} (Fig. 2), implies, on account of (41), the finiteness of the coefficient \mathcal{C}/\mathcal{A} outside the horizon, without the need for an explicit numerical computation. Thus, equation (41) assumes the form of an ordinary Schrödinger with *regular* coefficients over the entire domain of r^* .

From (40), (43) one obtains on the boundaries:

$$p_* u_0|_{r^*=\pm\infty} = \frac{\mathcal{B}}{\mathcal{A}} u_0|_{r^*=\pm\infty} + F p_* \delta\phi|_{r^*=\pm\infty} \quad (45)$$

The $\delta\phi$ remains bounded at $r^* = \pm\infty$. From the asymptotic behaviour (14) it becomes clear that, at the $r = \infty$ boundary, $\mathcal{B}/\mathcal{A} \rightarrow B/A \sim 1/r$, independently of ϕ_h . Thus, $F(r^* \rightarrow \infty) = e^{\int_{-\infty}^r \frac{B}{A} dr} \sim r$ for $r \rightarrow \infty$. Hence, the first term in (45) becomes $\delta\phi_\infty + \mathcal{O}(\frac{1}{r})$, whilst the second term vanishes as $1/r$, for $r \rightarrow \infty$ (see (14)). Hence, at the $r = \infty$ boundary the boundary values of $p_* u_0$ are proportional to those of u_0 :

$$p_* u_0|_{r^*=\infty} = \frac{u_0}{r}|_{r^*=\infty} \quad (46)$$

On the other hand, from (40) and (44) it becomes clear that at the boundary $r^* = -\infty$ (horizon), $F(r^* = -\infty) = 1$. Moreover, $p_* u_0|_{r^*=-\infty}$ is given by:

$$p_* u_0|_{r^*=-\infty} = (r - r_h)(\phi^{(2)'} - \phi^{(1)'})|_{r=r_h} = \left((r - r_h) \frac{\partial \phi_h'}{\partial \phi_h} u_0 \right) |_{r=r_h} \quad (47)$$

since, in our construction, each family of solutions is uniquely characterized [1] by the value ϕ_h for fixed r_h (we remind the reader that here we chose to keep $r_h = 1$ (fixed) and vary ϕ_h). From (9) one easily observes that for linear variations, $\delta\phi \equiv \phi^{(2)} - \phi^{(1)}$, the difference $\phi_h^{(2)'} - \phi_h^{(1)'}$ is *finite*. Hence,

$$p_* u_0|_{r^*=-\infty} \rightarrow 0 \quad (48)$$

We now discuss the $\sigma^2 < 0$ unstable modes u_σ . It can be easily seen from (30)-(33) that the asymptotic form of equation (41) at the $r^* = \infty$ boundary reads:

$$p_*^2 u_\sigma = -\sigma^2 u_\sigma \quad (49)$$

Hence, the bound-state solution u_σ behaves as ²:

$$u_\sigma(r^* = \infty) = e^{-|\sigma|r^*} \rightarrow 0 \quad (50)$$

²Here, and in the following, we insist on bounded, or- at most- linearly divergent u_σ , at the boundaries $r^* = \pm\infty$. This is due to the fact that, since $u = F\delta\phi$, and F is independent of σ and at most linearly divergent at $r^* = \infty$, then it is only for such a behaviour that the variation $\delta\phi$ remains bounded, as required by the linear stability analysis. An exponentially divergent $u_\sigma \sim e^r$ at the boundaries, would imply $\delta\phi \sim e^r/F$, and, hence, is not acceptable.

On the other hand, from (26)-(29) it is obvious that, on the horizon, equation (41) assumes the form (for the case $r_h = 1$):

$$p_*^2 u_\sigma + k^2 u_\sigma = 0 \quad (51)$$

$$k^2 \equiv \frac{2\gamma_1 e^{2\phi_h}}{(1 + \sqrt{1 - 6e^{2\phi_h}})\sqrt{1 - 6e^{2\phi_h}}} + \sigma^2 = k_0^2 + \sigma^2 \quad (52)$$

From (52) one can see two possibilities near the horizon of the black hole:

- (i) The ‘total energy’ is such that $0 > \sigma^2 > -\frac{2\gamma_1 e^{2\phi_h}}{(1 + \sqrt{1 - 6e^{2\phi_h}})\sqrt{1 - 6e^{2\phi_h}}}$. Taking into account the asymptotic form at $r^* = \infty$ (49) one also observes that in this case the spectrum of the respective Schrödinger equation is *continuous* and *non degenerate*. The general solution of the perturbation u_σ near the horizon is, therefore, oscillatory (unbound state):

$$u_\sigma^\pm \sim e^\pm ikr^* \quad r^* \sim -\infty \quad (53)$$

Such a continuum of states *cannot exist* in our case by continuity. Indeed, due to the non-degenerate nature of the eigenvalue problem, the limiting case $\sigma \rightarrow 0$ should yield the solution u_0 . However, in the limit $\sigma^2 \rightarrow 0$, and in terms of r , one obtains

$$u_0^\pm \sim \cos\left(\frac{e^{\phi_h}}{(1 - 6e^{2\phi_h})^{1/4}} \ln(r - r_h) + \varphi_0\right) \quad (54)$$

where φ_0 is a constant phase shift, and we have taken u_0 to be the real part of eq. (53). Then,

$$p_* u_0 \sim -k_0 \sin\left(\frac{e^{\phi_h}}{(1 - 6e^{2\phi_h})^{1/4}} \ln(r - r_h) + \varphi_0\right) \quad (55)$$

The above result is not in agreement with eq. (48), thereby contradicting the non-degenerate nature of these solutions, which, in turn, implies the absence of such solutions in the problem (41).

- (ii) This leaves one with the second possibility of a *discrete spectrum* of *bound states*, which would occur for:

$$\sigma^2 < -\frac{2\gamma_1 e^{2\phi_h}}{(1 + \sqrt{1 - 6e^{2\phi_h}})\sqrt{1 - 6e^{2\phi_h}}} < 0 \quad (56)$$

As we shall show below this is also *not realized* due to the special form of u_0 .

To this end, one first observes that such bound states would *vanish* exponentially at the $r^* = -\infty$ boundary:

$$u_\sigma(r^* = -\infty) \sim e^{|k|r^*} \rightarrow 0 \quad (57)$$

Thus, on account of (46,48,50,57) the Wronskian of any two solutions u_1, u_2 of the equation (41) with $\sigma^2 \leq 0$ vanishes at the boundaries:

$$W = (u_1 p_* u_2 - u_2 p_* u_1)|_{r^* = \pm\infty} = 0 \quad (58)$$

To count the unstable gravitational modes of the original problem, one needs to count the nodes of the wave function u of the one-dimensional Schrödinger problem (41). Fortunately, this can be done without detailed knowledge of the solutions. As we shall discuss below, all one needs to observe is the monotonic and non-intersecting nature of the dilaton curves in Fig. 1. To this end, one first observes that a standard ‘node rule’ for the discrete spectrum of the equation (41) applies, which is a direct consequence of Fubini’s theorem of ordinary differential equations [10]. This theorem can be stated as follows: consider two differential equations:

$$u'' + 2p_1 u' + q_1 u = 0 \quad (59)$$

$$u'' + 2p_2 u' + q_2 u = 0 \quad (60)$$

If,

$$p'_2 + p_2^2 - q_2 \leq p'_1 + p_1^2 - q_1 \quad (61)$$

throughout the interval $[a, b]$, then, between *any two consecutive zeroes of a solution of (59)*, in the interval $[a, b]$, *there is at least one zero of a solution of (60)*.

In our case, we can apply this theorem for two-different eigenfunctions, u_1, u_2 , corresponding to eigenvalues σ_1^2 and σ_2^2 of (41). The interval $[a, b]$ is the entire domain of validity of the solutions of (41), $(-\infty, \infty)$, including the boundaries at infinity. In this case,

$$p_i = 0, \quad q_i = \frac{\mathcal{C}}{\mathcal{A}} + \sigma_i^2 \frac{\mathcal{E}}{\mathcal{A}} - \frac{\mathcal{B}^2}{\mathcal{A}^2} - p_* \left(\frac{\mathcal{B}}{\mathcal{A}} \right), \quad i = 1, 2 \quad (62)$$

Then, the (sufficient) condition for a ‘node rule’ (61) reads simply:

$$\frac{\mathcal{E}}{\mathcal{A}}\sigma_2^2 \geq \frac{\mathcal{E}}{\mathcal{A}}\sigma_1^2 \quad (63)$$

The positivity of \mathcal{E}/\mathcal{A} (Fig. 3) implies that the condition (63) becomes simply:

$$\sigma_2^2 \geq \sigma_1^2 \quad (64)$$

This special version of the theorem is known as Sturm’s theorem [11]. As a corollary of Fubini/Sturm’s theorem one obtains the standard ‘node rule’ for the number of zeroes of the eigenfunctions in the discrete spectrum of bound states, according to which if the eigenfunctions are ranked in order of increasing energy, then, the n -th eigenfunction has $n - 1$ nodes (excluding the boundary zeroes) [11] (‘node rule’).

Consider, now, the case where σ_2 corresponds to the zero eigenvalue of (41), $\sigma_0 = 0$. As can be seen from the numerical solution of Fig. 1, the *monotonic* and *non-intersecting* character of the dilaton curves in the entire domain outside the horizon implies that the solution $u_2 = u_0$, which, as we have mentioned earlier, can be constructed out of the difference of two such solutions, has *no nodes* in the domain $r^* \in (-\infty, \infty)$. Since any solution u_n from the *discrete set* of negative eigenvalues $\sigma_n^2 < 0$ (unstable modes) has at least two nodes at the boundaries, according to Fubini/Sturm’s theorem, u_0 should have at least one in the domain $r^* \in (-\infty, \infty)$. This contradicts the fact that u_0 is nodeless. Thus, the only consistent situation is the one *without* such *negative-energy* modes. This, in turn, implies *linear stability* for the dilaton-GB black holes of ref. [1]. The reader might worry about the divergent boundary conditions of u_0 at $r^* = \infty$, which makes it *not an ordinary* eigenfunction of a Schrödinger problem. In the appendix we argue that this is not an obstacle. In fact, we present an explicit proof of the absence of bound states in our case, following the same spirit used in the proof of Fubini/Sturm’s theorem. The crucial element, which allows the standard proof to go through, is the special boundary condition for the Wronskian (58), which is valid for the entire spectrum of eigenfunctions of (41) with $\sigma^2 \leq 0$, including the non-standard one u_0 .

The above considerations can be extended straightforwardly to the case where $r_h \rightarrow 0$. All the coefficients of (41) are still well-defined in this case, which implies that the stability in principle does not change. However, according to the analysis of ref. [1], the case $r_h \rightarrow 0$ corresponds to a singular

curvature scalar

$$R \sim \frac{2}{r_h^2} \quad r_h \rightarrow 0 \quad (65)$$

which implies the absence of ‘particle-like’ solutions in the EDGB system [1, 12]. From the condition for the existence of black hole solutions (10), one observes that the only consistent value of ϕ_h for $r_h \rightarrow 0$ is [1] $\phi_h \rightarrow -\infty$. The limit is taken in such a way that $6\alpha'e_h^\phi/g^2r_h^2 = 1$. This implies that in such a case the GB term in the action (1) becomes irrelevant, and one is left with the standard Einstein term, which admits only Schwarzschild black holes, known to be stable. This stability is confirmed by the above smooth limit of the coefficients in (41) as $r_h \rightarrow 0$.

4 Remarks and Outlook

Above we have demonstrated linear stability of the dilatonic black hole solutions in the EDGB system, found in ref. [1]. This result is important, since it constitutes an example of a stable, albeit secondary, hair that bypasses the no-hair conjecture [4, 8]. Non-linear stability of the EDGB system, however, although expected, still remains an open issue.

Before closing we would like to compare our semi-analytic results on linear stability with some remarks in favour of stability by virtue of a catastrophe theory approach made in ref. [6]. As usual [4, 13], catastrophe theory can only indicate *relative changes* of stability, and hence cannot constitute a ‘proof’ of stability. In ref. [6] a numerical solution was found for the (+)-branch of solutions (9) which indicated the existence of a ‘turning point’ (TP) in the r_h - M (or equivalently ϕ_h - M) graph. The TP occurs at the ‘critical point’ for the existence of black hole solutions, which is the point where the black hole acquires a minimal mass, below which no solution is found. In the numerical solution of ref. [6] a continuation beyond this critical point emerged, which ends at a point (‘singular point’) where a singularity appears in the square of the Riemann tensor, as well as in ϕ_h'' .

The part of the solution from the critical point to the singular point was argued in ref. [6] to be *relatively unstable*, as compared to the regular branch discussed here. Such a change in stability manifests itself as a cusp in an appropriate catastrophe theory diagram [4, 13]. In ref. [6] such a diagram has been chosen to be the diagram of the thermodynamic entropy [14] versus

the mass of the black hole.

In our numerical solutions [1], used here, we found no evidence for such a continuation of the solution after the critical point, of minimum mass, which in our analysis is also the end point, where the two branches of (9) meet (see Figure 3). Our black hole solutions are uniquely specified by the pair (r_h, ϕ_h) , which was essential in our linear stability analysis above. Moreover, for finite r_h , the quantity ϕ_h'' is finite [1]. In this respect we are in agreement with the results of ref. [5], where a branching of solutions was found only *inside* the event horizon. These authors have also given a graph of the entropy versus the mass of the black hole solution of [1] outside the horizon, and found, as expected, a smooth curve, with no cusps. In this article we have proven *analytically* the stability under linear perturbations of this (unique) branch of the black hole solutions, which in the $r_h - M$ graphs of Figs. 4,5 terminates at the minimum-mass critical point. Of course, this result is in agreement with the relative stability of this branch *suggested* by the catastrophe theory analysis of ref. [6], but, as we said, the form of their numerical solution appears to be different, as it continues beyond the critical point. Nevertheless, we consider this as an *indication* for the stability of the solutions beyond the linear approximation. This, however, still remains an open issue.

As a final remark we would like to mention that the effects of gauge fields on the dilatonic black hole GB solutions have been considered in ref. [6, 7]. From the point of view of stability, one expects that, in the case of ‘coloured’ black holes, involving non-Abelian gauge fields, *instabilities* occur in *both* the gauge and gravitational sectors of the solutions. Instabilities in the gauge sector are of sphaleron type [4]. Those in the gravitational sector can be studied in a similar way as for coloured black holes in Einstein-Yang-Mills theories [4]. One can go beyond linear stability analysis in such systems, by invoking catastrophe theory [4, 13], which is capable of giving the relative stability of various branches of solutions for the coloured EDGB black holes [6, 7]. However, analytic methods can still be combined with the catastrophe theory approach [4] in order to *count* the *unstable modes* in both sectors, gauge and gravitational, by invoking appropriate maps of the system of perturbations into one-dimensional stationary Schrödinger problems [4, 9]. We hope to return to a detailed analytic study of these issues in a future publication.

Acknowledgements

Three of us (P.K., J.R., and K.T.) acknowledge traveling support from the EU-TMR Network ‘Beyond the Standard Model’. P.K. and K.T. wish to thank the Greek Ministry of Research and Technology for financial support (PENED 95). P.K. also thanks the Department of (Theoretical) Physics of Oxford University for the hospitality, and partial financial support, during the final stages of this work. The work of N.E.M. is supported by a P.P.A.R.C. (UK) Advanced Research Fellowship. E.W. thanks Oriel College, Oxford, for financial support.

Appendix: Absence of bound states

In this Appendix we prove the absence of bound states in the problem (41) by following a Wronskian treatment for the entire set of eigenmodes with $\sigma^2 \leq 0$. This justifies the validity of Fubini/Sturm’s theorem in our case. We stress that the crucial point in the proof is the special boundary conditions of the Wronskian (58). These allow a standard Wronskian treatment to go through for the u_0 solution of (41), despite its (linear) divergence at the $r^* = \infty$ boundary, which makes it not an ordinary eigenfunction of a Schrödinger problem.

To this end, we consider first the two solutions of (41), u_0 , and u_b - the ground state, at the bottom of the discrete spectrum. Both of these have *no nodes* in the interior domain of r^* , excluding the boundaries (the node structure of u_b follows from the ‘node rule’). We, then, employ properties of the Wronskian of the solutions as follows: first we multiply each equation with the other eigenfunction. Next, we subtract the resulting system of equations, and then, integrate it over the entire domain of $r^* \in (-\infty, \infty)$. In this way one obtains, in a standard fashion [11]:

$$\Delta W|_{r^*=\pm\infty} = (\sigma_b^2 - \sigma_0^2) \int_{-\infty}^{\infty} dr^* \frac{\mathcal{E}}{\mathcal{A}} u_b u_0 \quad (66)$$

where the left-hand-side denotes the change in the Wronskian between the two boundaries. From (58) this *vanishes*. Moreover, as we have mentioned previously, \mathcal{E}/\mathcal{A} is *positive definite* for the entire domain of $r^* \in (-\infty, \infty)$ (Figure 3). Since u_b, u_0 have *no nodes* in the domain $(-\infty, \infty)$, excluding

the boundaries, one obtains from (66) that the *only consistent* case is the degenerate one $\sigma_b^2 = \sigma_0^2$. But $\sigma_0^2 = 0$, whilst $\sigma_b^2 < 0$ by assumption; this implies a contradiction, excluding σ_b from the spectrum.

One repeats the construction, using u_0 and any of the higher eigenfunctions of the discrete spectrum, u_n , corresponding to $\sigma_n^2 < 0$. The change in the Wronskian between $-\infty$ and the first encountered zero of u_n , at $r^* = z_0$, is then given by:

$$\Delta W|_{r^*=-\infty}^{z_0} = -u_0 p_* u_n|_{z_0} = (\sigma_n^2 - \sigma_0^2) \int_{-\infty}^{z_0} dr^* \frac{\mathcal{E}}{\mathcal{A}} u_n u_0 \quad (67)$$

Without loss of generality, one may assume that $u_n > 0$ in the interval $(-\infty, z_0)$. Then $p_* u_n(z_0) < 0$. It is immediate to see that there is a contradiction in (67). The middle part has the sign of u_0 , whilst the right-hand-side has the opposite sign of u_0 . The case of a zero of u_n , at a point z_0 , such that $p_* u(z_0) = 0$ is dealt with similarly. In that case the contradiction lies in the fact that the left-hand side vanishes, whilst the right-hand side is a negative number (for $u_n > 0$ in the interval). These results exclude the possibility of bound-state eigenfunctions with zeroes in $(-\infty, \infty)$.

The above analysis, therefore, implies the *absence of negative energy modes* (bound states) in the problem (41), which, in turn, leads to *linear stability* for the Dilaton-Gauss-Bonnet black holes of ref. [1].

References

- [1] P. Kanti, N.E. Mavromatos, J. Rizos, K. Tamvakis and E. Winstanley, Phys. Rev. D54 (1996), 5049.
- [2] S. Coleman, J. Preskill and F. Wilczek, Nucl. Phys. B378 (1992), 175.
- [3] B.R. Greene, S.D. Mathur and C.M. O' Neill, Phys. Rev. D47 (1993), 2242.
- [4] N.E. Mavromatos and E. Winstanley, Phys. Rev. D53 (1996), 3190.
- [5] S.O. Alexeev and M.V. Pomazanov, Phys. Rev. D56 (1996), 2110.
- [6] T. Torii, H. Yajima and K. Maeda, Phys. Rev. D55 (1996), 739.
- [7] P. Kanti and K. Tamvakis, Phys. Lett. B392 (1996), 30.
- [8] J. Bekenstein, Phys. Rev. D5 (1972), 1239; Phys. Rev. D51 (1995), 6608;
A. Mayo and J. Bekenstein, Phys. Rev. D54 (1996), 5059.
- [9] P. Boschung, O. Brodbeck, F. Moser, N. Straumann and M.S. Volkov, Phys. Rev. D50 (1994), 3842;
M.S. Volkov and D.V. Gal'tsov, Phys. Lett. B341 (1995), 279;
M.S. Volkov, O. Brodbeck, G. Lavrelashvili, and N. Straumann, Phys. Lett. B349 (1995), 438;
E. Winstanley and N.E. Mavromatos, Phys. Lett. B352 (1995), 242;
O. Brodbeck and N. Straumann, Phys. Lett. B324 (1994); gr-qc/9411058 (1994);
N. Straumann and Z.H. Zhou, Phys. Lett. B237 (1990), 353;
M. Heusler, S. Droz, and N. Straumann, Phys. Lett. B285 (1992), 21;
G. Lavrelashvili and D. Maison, Phys. Lett. B343 (1995), 214.
- [10] G. Birkhoff and G-C. Rota, *Ordinary Differential Equations*, (Wiley 1989).
- [11] See for instance, A. Messiah, *Quantum Mechanics*, Vol. I (North-Holland Publishing Co., Amsterdam 1970).

- [12] E.E. Donets and D.V. Gal'tsov, Phys. Lett. B352 (1995), 261.
- [13] T. Torii, K. Maeda and T. Tachizawa, Phys. Rev. D51 (1995), 1510;
ibid. 4054.
- [14] G. 't Hooft, Nucl. Phys. B256 (1985), 727.




Figure 1: Dilaton configurations for a family of black hole solutions, corresponding to fixed $r_h = 1$, for various values of ϕ_h . Notice the monotonic behaviour of the solutions, and the fact that the curves do not intersect.




Figure 2: The graph depicts the coefficients B/A and \mathcal{B}/\mathcal{A} , for a typical member of the family of the black hole solutions of Fig.1 corresponding to $\phi_h = -1$, $r_h = 1$. It is clear that the coefficient \mathcal{B}/\mathcal{A} , incorporating the tortoise coordinate, is finite in the entire domain outside the horizon, thereby implying that the quantity F is well-defined and integrable.




Figure 3: This diagram depicts the coefficients E/A and \mathcal{E}/\mathcal{A} for a typical member of the family of the black hole solutions depicted in Fig. 1 ($\phi_h = -1$, $r_h = 1$); the coefficient E/A diverges at the horizon as $1/(r - r_h)^2$. On the other hand, $\mathcal{E}/\mathcal{A} = e^{\Gamma - \Lambda} E/A$, appearing in (40), is finite at the horizon. The positive-definiteness of both coefficients is clear.




Figure 4: The graph depicts r_h versus the ADM Mass $2M$ of the black hole, for a fixed value of $\phi_h = -1$. The emergence of the asymptotic critical point, $r_h^4 \simeq 6\alpha'^2 e^{2\phi_h}/g^4$, below which there are no solutions, is apparent. At this point the mass becomes minimal.




Figure 5: The magnification around the critical point, depicted in the above figure, shows clearly the abrupt (almost vertical) slope with which this point is approached. No turning point is found.