

Dimensional Renormalization and the Action Principle

P. Breitenlohner and D. Maison*

Max-Planck-Institut für Physik und Astrophysik, D-8000 München 40, Federal Republic of Germany

Abstract. Dimensional renormalization is defined in such a way that the renormalized action principle holds. It is shown that this leads to a minimal, additive renormalization. The derivation of Ward-Takahashi identities and Callan-Symanzik equations from the action principle is exemplified.

I. Introduction

Dimensional regularization [1, 2] has become an almost indispensable tool to the perturbative treatment of non-abelian gauge theories. Nevertheless the first complete proof of the so-called Slavnov-Taylor identities, guaranteeing the unitarity of the S-matrix, has been given recently with help of the “Normal Product Algorithm” (NPA) based on BPHZ renormalization [3]. The reason lies in the validity of the renormalized action principle in that formalism. The action principle specifies the change of the Green's functions under an infinitesimal variation of fields or parameters entering the Lagrangian. In our opinion it provides a much more efficient approach to the proof of Ward-Takahashi or Slavnov-Taylor identities—or other structural properties—than the algebraic manipulations performed on individual diagrams advocated by the pioneers in that field [4]. The approach of Ref. [3] suffers, however, from the fact that the subtraction method in the BPHZ renormalization is not compatible with the identities to be proved. This is overcome by appropriately chosen asymmetric finite counterterms added to the Lagrangian. Apart from the difficulty to prove that such counterterms can really be found in all orders of perturbation theory the method appears much too clumsy for practical calculation.

Therefore, it is only natural to make an effort to establish the renormalized action principle in the framework of dimensional renormalization. This is one of the main objectives of the present work. The second is to show that dimensional renormalization meets the requirements of a minimal, additive renormalization

* Present address: II. Institut für Theoretische Physik der Universität Hamburg, Luruper Chaussee 149, D-2000 Hamburg 50, Federal Republic of Germany

in the sense of Hepp [5]. Although there exists already a proof for that by E. R. Speer [6], who reduces dimensional renormalization to standard BPH renormalization (= minimal Taylor series subtractions) performing a suitable finite renormalization, we have included a direct proof without recourse to BPH renormalization. *In fact our subtractions differ by finite terms from those proposed by Speer*, which proves to be necessary to establish the renormalized action principle and only the latter guarantees the validity of Ward identities etc.¹

Our proof of the renormalized action principle proceeds in two steps:

i) We demonstrate its validity in the regularized theory. The essential property to show is that the application of the kinetic differential operator to the propagator corresponding to some line ℓ in a Feynman graph (i.e. $\square + m^2$ for a scalar line) is equivalent to the contraction of that line to a point.

ii) We prove that the subtractions corresponding to all renormalization parts (= superficially divergent 1PI subgraphs) before and after contraction of the line are identical, although the respective classes of renormalization parts are generally different.

We have decided to devote some space especially to the latter point, since there does not seem to exist an adequate discussion of that problem in the literature². As far as the treatment of covariants in the dimensionally regularized theory is concerned, especially the notorious γ_5 , we follow essentially t'Hooft and Veltman [1], with somewhat more emphasis on a consistent formalism. We try to convince the reader that there is no better choice for the definition of γ_5 .

In Appendix A we have collected all relevant definitions and properties concerning Feynman graphs.

II. Dimensional Regularization

II.1. Feynman Amplitudes

Assigning propagators Δ_ℓ to lines ℓ and vertex parts \hat{X}_i to vertices V_i of some Feynman graph G we may construct Feynman amplitudes \mathcal{T}_G as formal expressions of the type³ (Definitions are given in Appendix A)

$$\mathcal{T}_G(x_1, \dots, x_M) = \prod_{i \in V_G} \hat{X}_i \prod_{\ell \in L_G} \Delta_\ell \left(\sum_{i \in V_G} e_{\ell i} x_i \right), \quad x_i \in \mathbb{R}^n, \quad n \text{ pos. integer} \quad (1)$$

with

$$\begin{aligned} \hbar^{-1} \Delta_\ell(x) &= \lim_{\varepsilon \rightarrow 0} Z_\ell(-i \partial/\partial u) \int_0^\infty d\alpha (-2i\alpha)^{-n/2} \\ &\cdot \exp \{ -(i/4\alpha)(x+u)^2 - i\alpha(m_\ell^2 - i\varepsilon) \}|_{u=0}. \end{aligned} \quad (2)$$

$Z_\ell(-i \partial/\partial u)$ are some constant coefficient diff. operators containing in general covariant algebraic objects like γ -matrices etc. and $\hbar \hat{X}_i = X_i(-i \partial/\partial x_i)$ are again constant coeff. diff. operators like the Z_ℓ .

¹ A related work by Collins [7] has recently appeared

² The proof of Y.M.P. Lam Ref. [8] is not complete in our mind

³ The \hat{X}_i and Δ_ℓ have to be appropriately ordered, which is understood to be done whenever necessary

Taking the $n \times M$ -dimensional Fourier transform of $\mathcal{T}_G(x)$ and performing the Gaussian integrations over x_i we obtain⁴

$$\begin{aligned}
\mathcal{T}_G(p_1, \dots, p_M) &= (2\pi)^{-nM/2} \int e^{ipx} \mathcal{T}_G(x) dx \\
&= \lim_{\varepsilon \rightarrow 0} \hbar^{h_G - 1} (2\pi)^{n/2} \delta(\sum p_i) t^{h_G} (2i)^{-nh_G/2} \int_0^\infty \prod_{\ell \in \mathcal{L}_G} d\alpha_\ell I_\varepsilon(p, u, \underline{\alpha})|_{u=0} ; \\
I_\varepsilon(p, u, \underline{\alpha}) &= d(\alpha)^{-n/2} \prod_{i \in \mathcal{V}_G} X_i(p_i - i \partial/\partial u_\ell) \prod_{\ell \in \mathcal{L}_G} Z_\ell(-i \partial/\partial u_\ell) \exp iW(p, u, \underline{\alpha}) ; \\
W(p, u, \underline{\alpha}) &= (p^+ - u^+(2\alpha)^{-1} e_E, u^+) \begin{pmatrix} A_E^{-1} & 0 \\ 0 & -(4\alpha)^{-1} \end{pmatrix} \begin{pmatrix} p - e_E^+(2\alpha)^{-1} u \\ u \end{pmatrix} \\
&\quad - \sum_{\ell \in \mathcal{L}_G} \alpha_\ell (m_\ell^2 - ie) \\
&= (p^+, u^+) \begin{pmatrix} 0 & -2e_E^+ \\ -2e_E & -4\alpha \end{pmatrix}^{-1} \begin{pmatrix} p \\ u \end{pmatrix} - \sum_{\ell \in \mathcal{L}_G} \alpha_\ell (m_\ell^2 - ie) .
\end{aligned} \tag{3}$$

This expression is the starting point for the dimensional regularization. It is achieved in two steps:

i) we interpret the parameter n appearing in (3) as a complex variable,

ii) we give a definition of the algebraic objects (like γ -matrices, momentum 4-vectors etc.) appearing in (3) which makes no direct use of the concept of dimension (see Sect. II.2).

II.2. “ n -Dimensional” Lorentz Covariants

For the validity of canonical Ward-Takahashi identities etc. for the dimensionally regularized Feynman amplitudes resp. Green’s functions it is essential to find an extension of the Lorentz covariants (γ_μ , p_μ , $g_{\mu\nu}$ etc.) which allows one to manipulate these objects like in 4-dimensions. This will be achieved by treating them as formal objects obeying certain algebraic identities. We shall make no use of the fact that indices can assume values, but treat objects like γ_μ as just one entity (i.e. we shall never use γ_2 etc.). The only place where one runs into troubles with that prescription is where the $\varepsilon_{\kappa\lambda\mu\nu}$ -tensor comes into play, since it is a genuine 4-dimensional object (like ε_{ijk} is a 3-dimensional one). This is expressed by its property that in 4 dimensions the value of an arbitrary Lorentz index is equal to exactly one of the values of the indices of ε . Correspondingly there are identities involving ε depending crucially on 4-dimensionality as

$$\varepsilon_{\mu_1 \dots \mu_4} \varepsilon_{\nu_1 \dots \nu_4} = - \sum_{\pi \in S_4} \text{sign } \pi \prod_{i=1}^4 g_{\mu_i \nu_{\pi(i)}} \tag{4}$$

and as a consequence

$$\sum_{\pi \in S_5} \text{sign } \pi \varepsilon_{\mu_{\pi(1)} \dots \mu_{\pi(4)}} g_{\mu_{\pi(5)} v} = 0 . \tag{5}$$

⁴ Using the relation $[p_i + e_{i\ell}^+ (-i \partial/\partial u_\ell)] I_\varepsilon(p, u, \underline{\alpha}) = 0$ the vertex factors $X_i(p_i)$ can be included in the factors $Z_\ell(-i \partial/\partial u_\ell)$

Two possibilities are to be considered:

i) no use is made of (4), i.e. there is no possibility to reduce a product of several ε -tensors in the formal algebra,

ii) in addition to the " n -dimensional" covariants also "(n -4)-dimensional" ones (or equivalently 4-dimensional ones) are introduced.

Possibility ii) allows a further simplification of covariants containing a product of several ε -tensors; as a consequence $\gamma_5^2 = -1$ can be proved.

Equation (5), however, has the disadvantage that it prevents one from finding easily a *unique* normal form for the covariants (see below).

We use the usual symbols⁵ $g_{\mu\nu}$, p_μ , γ_μ , $\mathbb{1}$, $\varepsilon_{\kappa\lambda\mu\nu}$ which can be added, multiplied etc. as usual, obeying the additional identities⁶ (all symbols except γ_μ are assumed to commute)

$$\begin{aligned} g_{\mu\nu}g_{\nu\lambda} &= g_{\mu\lambda}, \quad g_{\mu\nu} = g_{\nu\mu}, \\ g_{\mu\nu}p_\nu &= p_\mu, \quad g_{\mu\nu}\gamma_\nu = \gamma_\mu, \quad p_\mu q_\mu = p \cdot q, \\ g_{\mu\nu}\varepsilon_{\nu\varrho\sigma\tau} &= \varepsilon_{\mu\varrho\sigma\tau}, \\ \{\gamma_\mu, \gamma_\nu\} &= \gamma_\mu\gamma_\nu + \gamma_\nu\gamma_\mu = 2g_{\mu\nu}\mathbb{1}, \quad \mathbb{1}\gamma_\mu = \gamma_\mu\mathbb{1} = \gamma_\mu, \\ g_{\mu\mu} &= n, \\ \text{Tr } \mathbb{1} &= 4. \end{aligned} \tag{6}$$

In order to be able to formulate Equation (4) we introduce additional symbols (as discussed above) $\hat{g}_{\mu\nu}$, \hat{p}_μ , $\hat{\gamma}_\mu$ obeying the identities

$$\begin{aligned} g_{\mu\nu}\hat{g}_{\nu\lambda} &= \hat{g}_{\mu\nu}\hat{g}_{\nu\lambda} = \hat{g}_{\mu\lambda}, \quad \hat{g}_{\mu\nu} = \hat{g}_{\nu\mu}, \\ \hat{g}_{\mu\nu}p_\nu &= \hat{p}_\mu, \quad \hat{g}_{\mu\nu}\gamma_\nu = \hat{\gamma}_\mu; \end{aligned} \tag{7}$$

$$\varepsilon_{\mu_1\dots\mu_4}\varepsilon_{\nu_1\dots\nu_4} = -\sum_{\pi\in S_4} \text{sign } \pi \prod_{i=1}^4 (g_{\mu_i\nu_{\pi(i)}} - \hat{g}_{\mu_i\nu_{\pi(i)}}) \tag{4'}$$

where S_4 denotes the permutation group of 4 objects. From these assumptions we can prove for example

$$\begin{aligned} g_{\mu\nu}\hat{\gamma}_\nu &= \hat{g}_{\mu\nu}\hat{\gamma}_\nu = \hat{\gamma}_\mu, \\ \{\gamma_\mu, \hat{\gamma}_\nu\} &= \{\hat{\gamma}_\mu, \hat{\gamma}_\nu\} = 2\hat{g}_{\mu\nu}\mathbb{1}, \\ \varepsilon_{\kappa\lambda\mu\nu}\hat{g}_{\nu\varrho} &= 0, \\ \varepsilon_{\mu_1\dots\mu_4} &= \text{sign } \pi \varepsilon_{\mu_{\pi(1)}\dots\mu_{\pi(4)}}, \\ \hat{g}_{\mu\mu} &= n-4, \\ \text{Tr } \gamma_\mu &= 0. \end{aligned} \tag{8}$$

Before we give a definition for γ_5 , we will try to convince the reader, that there is no better choice. In order to do so, we observe, that γ_5 can be defined for $n=4$ by its two properties

i) $\text{Tr}(\gamma_5\gamma_{\mu_1}\dots\gamma_{\mu_4}) = \varepsilon_{\mu_1\dots\mu_4} \text{Tr } \mathbb{1}$

⁵ This list may be enlarged if necessary by other covariant objects

⁶ There are a lot of obvious rules to be observed, like: no index μ can appear more than twice in one product etc., which we do not specify here

which will be maintained and

$$\text{ii)} \quad \{\gamma_\mu, \gamma_5\} = 0$$

which has to be given up, because the two are not compatible for $n \neq 4$. As an alternative we might try to maintain ii), but this leads immediately to

$$n(n-2)(n-4) \text{Tr}(\gamma_5 \gamma_{\mu_1} \dots \gamma_{\mu_4}) = 0$$

and there is no smooth limit $n \rightarrow 4$ which reproduces i). Nevertheless, this approach strongly supports the result that axial anomalies can always be avoided except in matrix elements where i) becomes essential.

Let us define $\gamma_5 = (4!)^{-1} \epsilon_{\mu_1 \dots \mu_4} \gamma_{\mu_1} \dots \gamma_{\mu_4}$. With that definition we can prove

$$\text{Proposition 1. } \{\gamma_\alpha, \gamma_5\} = \{\hat{\gamma}_\alpha, \gamma_5\} = 2\hat{\gamma}_\alpha \gamma_5. \quad (9)$$

Using the abbreviations $\hat{g}_{\mu\nu} = g_{\mu\nu} - \hat{g}_{\mu\nu}$, $\hat{\gamma}_\mu = \gamma_\mu - \hat{\gamma}_\mu$ etc. we show first

$$\text{Lemma 1. } \sum_{\pi \in S_5} \text{sign } \pi \epsilon_{\mu_{\pi(1)} \dots \mu_{\pi(4)}} \hat{g}_{\mu_{\pi(5)} \alpha} = 0. \quad (10)$$

$$\begin{aligned} \text{Proof. } & \sum_{\pi \in S_5} \text{sign } \pi \epsilon_{\mu_{\pi(1)} \dots \mu_{\pi(4)}} \hat{g}_{\mu_{\pi(5)} \alpha} \\ &= -1/6 \sum_{\pi \in S_5} \text{sign } \pi \epsilon_{\mu_{\pi(1)} \dots \mu_{\pi(4)}} \epsilon_{\mu_{\pi(5)} v_1 v_2 v_3} \epsilon_{\alpha v_1 v_2 v_3} \\ &= 4 \epsilon_{\alpha v_1 v_2 v_3} \sum_{\pi \in S_5} \text{sign } \pi \hat{g}_{\mu_{\pi(1)} \mu_{\pi(5)}} \hat{g}_{\mu_{\pi(2)} v_1} \hat{g}_{\mu_{\pi(3)} v_2} \hat{g}_{\mu_{\pi(4)} v_3} = 0 \end{aligned}$$

since $\hat{g}_{\mu_{\pi(1)} \mu_{\pi(5)}}$ is symmetric under the exchange of 1 and 5.

Proof of Proposition 1. Using the above Lemma we get

$$\begin{aligned} 0 &= (1/4!)(\gamma_{\mu_1} \dots \gamma_{\mu_5} + \gamma_{\mu_5} \dots \gamma_{\mu_1}) \sum_{\pi \in S_5} \text{sign } \pi \epsilon_{\mu_{\pi(1)} \dots \mu_{\pi(4)}} \hat{g}_{\mu_{\pi(5)} \alpha} \\ &= (5/4!)(\hat{\gamma}_\alpha \gamma_{\mu_1} \dots \gamma_{\mu_4} + \gamma_{\mu_4} \dots \gamma_{\mu_1} \hat{\gamma}_\alpha) \sum_{\pi \in S_4} \text{sign } \pi \epsilon_{\mu_{\pi(1)} \dots \mu_{\pi(4)}} \\ &\quad + (20/3!) \hat{g}_{\nu\alpha} (\gamma_{\mu_1} \dots \gamma_{\mu_3} + \gamma_{\mu_3} \dots \gamma_{\mu_1}) \sum_{\pi \in S_3} \text{sign } \pi \epsilon_{\mu_{\pi(1)} \mu_{\pi(2)} \mu_{\pi(3)} \nu} \\ &= 5! \{\hat{\gamma}_\alpha, \gamma_5\} \quad \text{and hence} \quad \{\gamma_\alpha, \gamma_5\} = \{\hat{\gamma}_\alpha, \gamma_5\} \end{aligned}$$

where use is made of the fact that the permutation (54321) is even, whereas (321) is odd. The second part of Equation (9) is trivial to prove.

From Proposition 1 one can immediately compute

$$\text{Tr}(\{\gamma_\alpha, \gamma_5\} \gamma_\lambda \gamma_\mu \gamma_\nu \gamma_\rho) = 8(n-4) \epsilon_{\lambda\mu\nu\rho}$$

which is important for the derivation of the Adler anomaly.

Similarly one may prove

$$\text{Proposition 2. } \gamma_5^2 = -\mathbb{1}. \quad (11)$$

Remark. All the quantities appearing are treated as real, treating charges by doubling the number of components.

The following **proposition** can be proved:

The identities above can be used as simplification rules for syntactically correct expressions, leading to a normal form (NF) which is unique insofar as (4) is neglected—otherwise there remain further linear relations with n -independent coefficients between expressions in NF derived from Equation (5).

As for the proof we execute the following transformations (simplifications) leading to the NF:

- (i) we eliminate γ_s by its definition;
- (ii) we eliminate Tr by applying $\{\gamma_\mu, \gamma_\nu\} = 2g_{\mu\nu}$, $\text{Tr}\gamma_\mu = 0$, $\text{Tr}\mathbf{1} = 4$, $\text{Tr}AB = \text{Tr}BA$;
- (iii) we antisymmetrize products of γ 's by $\{\gamma_\mu, \gamma_\nu\} = 2g_{\mu\nu}$;
- [iv] we eliminate ε -tensors by (4');
- (v) we eliminate $g_{\mu\nu}$'s (resp. $\hat{g}_{\mu\nu}$'s) by $g_{\mu\nu}p_\nu = p_\mu$ (resp. $\hat{g}_{\mu\nu}p_\nu = g_{\mu\nu}\hat{p}_\nu = \hat{g}_{\mu\nu}\hat{p}_\nu = \hat{p}_\mu$) etc.;
- (vi) we replace $p_\mu q_\mu$ by $p \cdot q$ (resp. $\hat{p}_\mu q_\mu = p_\mu \hat{q}_\mu = \hat{p}_\mu \hat{q}_\mu$ by $\widehat{p \cdot q}$) etc. and use $g_{\mu\mu} = n$ (resp. $\hat{g}_{\mu\mu} = n - 4$).

It is easy to see, that [without (4')] any further application of any of the identities (6) no longer changes the expression obtained after the application of (i) to (vi).

Having given a meaning to the Lorentz covariants contained in Equation (3) we can now come to the definition of the dimensionally regularized Feynman amplitude corresponding to the formal expression (3):

This is done in two steps.

Step 1: Interpret the covariants contained in the X_i 's resp. Z_ℓ 's (i.e. also u_ℓ and $\partial/\partial u_\ell$) as elements of the algebra described above. The exponential function defining $I_\varepsilon(p, u, \alpha)$ is considered as a formal power series. Next execute all derivatives $\partial/\partial u_\ell$ and put $u_\ell = 0$. Reduce the expressions obtained to their NF. This displays explicitly the n -dependence of the covariants and therefore can be used for the definition of the counterterms (see Sect. III).

Step 2: Once the counterterms are determined, we interpret the covariants in the NF as 4-dimensional (specifically we put all $\hat{O} = 0$, i.e. $\hat{g}_{\mu\nu} = 0$, $\hat{p}_\mu = 0$ etc.) and set all of the $M - 1$ momenta p_i to zero which correspond to internal vertices. After that the Feynman integrand can be considered as a distribution in the remaining $M' - 1$ momenta over $\mathcal{S}(\mathbb{R}^{4(M'-1)})$ and depends parametrically on α and n . Taking $\text{Re } n \ll 0$ all α -integrals converge absolutely. For $\min\{m_\ell\} \neq 0$ the limit $\varepsilon \rightarrow 0$ exists in $\mathcal{S}'(\mathbb{R}^{4(M'-1)})$ and defines a distribution $\mathcal{T}_G(p, n)$ over $\mathcal{S}(\mathbb{R}^{4M'})$ [9]. $\mathcal{T}_{G,\varepsilon}(p, n)$ can be continued analytically to the whole complex n -plane as a meromorphic function, the singularities of which lie on the manifolds

$$\omega_H(n) = nh_H - 2L_H + \sum_{\mathcal{L}_H} r_\ell = k, \quad k = 0, 1, 2, \dots, H \subset G$$

where r_ℓ is the degree of Z_ℓ [10]. For values of n not on any of these manifolds and $\varepsilon \geq 0$ $\mathcal{T}_{G,\varepsilon}(p, n)$ is continuous in ε . Therefore, $\mathcal{T}_G(p, n)$ is again a meromorphic function of n with singularities on the same manifolds.

II.3. The Action Principle

Most of the equations characterizing the structural properties of a Lagrangian *QFT* like equations of motion, Ward-Takahashi identities, Callan-Symanzik equations etc. can be understood as a consequence of the “action principle” [11]. Let us consider a Lagrangian $\mathcal{L} = \mathcal{L}_0 + \mathcal{L}_{\text{int}}$ depending on a set of commuting or anticommuting quantized fields generically called $\phi(x)$, unquantized “external” fields $a(x)$ and space-time independent parameters λ . Then we may ask how the Green’s functions of the theory change under an infinitesimal variation of these quantities. The answer, given by the quantum action principle, is most efficiently expressed in terms of the generating functional $Z(a, \lambda)$ for the Green’s functions⁷ given by the Gell-Mann-Low series

$$Z(a, \lambda) = \langle \exp i\hbar^{-1} \int \mathcal{L}_{\text{int}}(\phi(x), a(x), \lambda) dx \rangle_{+,c}^0. \quad (12)$$

$Z(a, \lambda)$ is a well defined formal power series in \hbar using dimensionally regularized Feynman amplitudes corresponding to the graphical expansion of $\langle \exp i\hbar^{-1} \int \mathcal{L}_{\text{int}} \rangle_{+,c}^0$. $\langle \rangle_{+,c}^0$ designates the connected vacuum expectation value of the time ordered product, evaluated in the free theory given by $L_0 = \frac{1}{2} \phi D\phi$, i.e. with propagator $i\hbar D^{-1}$. About D we make the assumption that it is a real, second order differential operator with constant coefficients, i.e.

$$D_{ij} = a_{ij} + b_{ij}^\mu \partial_\mu + c_{ij}^{\mu\nu} \partial_\mu \partial_\nu \quad \text{with real coefficients ,}$$

$$a_{ij} = \pm a_{ji}, \quad b_{ij}^\mu = \mp b_{ji}^\mu, \quad c_{ij}^{\mu\nu} = \pm c_{ji}^{\mu\nu} \quad \text{the } \pm \text{ sign referring}$$

to Bose resp. Fermi components of $\phi = (\phi_i)$. The constants a, b, c contain in general algebraic objects like $\gamma_\mu, g_{\mu\nu}$ etc. We assume, however, that D^{-1} is the same algebraic expression as in 4 dimensions, i.e. $D(-i\partial/\partial u_\mu) Z_\mu(-i\partial/\partial u_\mu) = -(\square_u + m_\mu^2)$, which can easily be proved for the standard representations like $-(\square + m^2)$, $-i\gamma_\mu \partial_\mu + m$ etc., using the algebraic rules of Section II.2.

The action principle for the respective kinds of variations takes the following form:

i) variations of the quantized field $\phi(x)$

$\delta\phi(x) = P(\phi(x))\delta\varepsilon(x)$ with some polynomial P leave $Z(a, \lambda)$ invariant:

$$0 = \langle \delta\mathcal{L} \exp i\hbar^{-1} \int \mathcal{L}_{\text{int}}(x) dx \rangle_{+,c}^0 \quad (13)$$

where $\delta\mathcal{L}$ is the linear part of $\mathcal{L}(\phi + \delta\phi) - \mathcal{L}(\phi)$ in $\delta\varepsilon$. Neglecting ordering of factors we may write

$$\begin{aligned} \delta\mathcal{L} = & \int \{ (\delta\mathcal{L}/\delta\phi(x)) P(\phi(x)) + (\delta\mathcal{L}/\delta\partial_\mu\phi(x)) \partial_\mu P(\phi(x)) \\ & - \partial_\mu ((\delta\mathcal{L}/\delta\partial_\mu\phi(x)) P(\phi(x))) \} \delta\varepsilon(x) dx \end{aligned} \quad (13a)$$

resp.

$$\delta\mathcal{L} = \int P(\phi(x)) ((\delta\mathcal{L}/\delta\phi(x)) - \partial_\mu (\delta\mathcal{L}/\delta\partial_\mu\phi(x))) \delta\varepsilon(x) dx ; \quad (13b)$$

ii) variations of external fields result in

$$-i\hbar \delta Z/\delta a(x) = \langle (\delta\mathcal{L}/\delta a(x)) \exp i\hbar^{-1} \int \mathcal{L}_{\text{int}}(x) dx \rangle_{+,c}^0 ; \quad (14)$$

⁷ We assume that \mathcal{L}_{int} contains a “source term” $\int J(x)\phi(x)dx$ which allows to express Green’s functions as functional derivatives of Z

iii) variations of parameters give

$$-i\hbar \partial Z/\partial \underline{\lambda} = \langle (\partial \mathcal{L}/\partial \underline{\lambda}) \exp i\hbar^{-1} \int \mathcal{L}_{\text{int}}(x) dx \rangle_{+,c}^0. \quad (15)$$

In the following we try to show that Equations (13)–(15) hold in the sense of formal power series in \hbar in the dimensionally regularized theory corresponding to \mathcal{L} .

The only difficulty of the proof lies in the asymmetrical treatment of \mathcal{L}_0 and \mathcal{L}_{int} in the definition of Z via Feynman graphs: \mathcal{L}_0 determines the lines, \mathcal{L}_{int} the vertices. We have to demonstrate how the variations in both constituents conspire to give the simple result of Equations (13)–(15). Two kinds of properties are responsible for this conspiracy:

- i) algebraic properties of the Feynman integrands $I(p, u, \alpha)$,
- ii) combinatorial properties of the graphical expansion of

$$\langle \exp i\hbar^{-1} \int \mathcal{L}_{\text{int}}(x) dx \rangle_{+,c}^0.$$

We shall restrict ourselves to the discussion of point i), since ii) is straightforward and has nothing to do with dimensional regularization. This immediately settles Equation (14), because we have assumed that there is no external field dependence in \mathcal{L}_0 and therefore we may write Equation (14) in the form

$$-i\hbar \partial Z/\partial \underline{a}(x) = \langle (\delta \mathcal{L}_{\text{int}}/\delta \underline{a}(x)) \exp i\hbar^{-1} \int \mathcal{L}_{\text{int}}(x) dx \rangle_{+,c}^0, \quad (16)$$

the proof of which is essentially combinatorial.

Equation (13) needs somewhat more work. Splitting \mathcal{L} into \mathcal{L}_0 and \mathcal{L}_{int} we may rewrite it as

$$\begin{aligned} & \langle P(\phi(x)) D\phi(x) \exp i\hbar^{-1} \int \mathcal{L}_{\text{int}}(x) dx \rangle_{+,c}^0 \\ &= -\langle P(\phi(x)) (\delta \mathcal{L}_{\text{int}}/\delta \phi(x)) \exp i\hbar^{-1} \int (x) dx \rangle_{+,c}^0. \end{aligned} \quad (17)$$

Reducing Equation (17) to Feynman graphs it boils down to the following property: replacing $Z_\ell(-i\partial/\partial u_\ell)$ in Equation (3) by $D_\ell(-i\partial/\partial u_\ell)Z_\ell(-i\partial/\partial u_\ell) = -(\square_{u_\ell} + m_\ell^2)$ is equivalent to the contraction of line ℓ to a point, which is achieved by replacing $\int_0^\infty d\alpha_\ell$ by $-\int_0^\infty d\alpha_\ell \partial/\partial \alpha_\ell$, i.e. $\mathcal{T}'_G = \mathcal{T}_{G/\ell}$, where \mathcal{T}'_G is the Feynman amplitude for G with an additional factor $D_\ell(-i\partial/\partial u_\ell)$ included. Therefore, it is sufficient to show

$$(\square_{u_\ell} + m_\ell^2 - ie) I_\ell(p, u, \alpha) = i(\partial/\partial \alpha_\ell) I_\ell(p, u, \alpha) \quad (18)$$

in the formal algebra of covariants. Up to trivial factors⁸ $I_\ell(p, u, \alpha)$ has the form

$$(\text{setting } \prod_{i \in \gamma_G} X_i(p_i, -i\partial/\partial u_\ell) \prod_{\ell \in \mathcal{L}_G} Z_\ell(-i\partial/\partial u_\ell) = Z(-i\partial/\partial u)),$$

$$I_\ell(p, u, \alpha) = Z(-i\partial/\partial u) d(\alpha)^{-n/2} \exp \{iV(p, u, \alpha) - i \sum_{\ell \in \mathcal{L}_G} \alpha_\ell (m_\ell^2 - ie)\}$$

⁸ Actually it is important that the only dependence of these factors on the number L of lines resp. M of vertices is through the number of loops $h = L - M + 1$, since the latter does not change by contracting a line

with

$$V(p, u, \alpha) = (p^+, u^+) M^{-1} \begin{pmatrix} p \\ u \end{pmatrix} \quad M = \begin{pmatrix} 0 & -2e^+ \\ -2e & -4\alpha \end{pmatrix}, d(\alpha) = \det M / (-4)^L.$$

Differentiation with respect to α_ℓ gives

$$-i \partial I_\varepsilon / \partial \alpha_\ell = i(2^{-1} n d^{-1} \partial d / \partial \alpha_\ell - i \partial V / \partial \alpha_\ell + i(m_\ell^2 - ie)) I_\varepsilon \quad (19)$$

whereas

$$-(\square_{u_\ell} + m_\ell^2 - ie) I_\varepsilon = i(-\square_{u_\ell} V - i(\partial V / \partial u_\ell) \cdot (\partial V / \partial u_\ell) + i(m_\ell^2 - ie)) I_\varepsilon. \quad (20)$$

Using

$$2^{-1} n d^{-1} \partial d / \partial \alpha_\ell = 2^{-1} n \text{Tr}((\partial M / \partial \alpha_\ell) M^{-1}) = -\square_{u_\ell} V \quad (21a)$$

(here $g_{\mu\mu} = n$ comes in !) and

$$\partial V / \partial \alpha_\ell = -(p^+ u^+) M^{-1} (\partial M / \partial \alpha_\ell) M^{-1} \begin{pmatrix} p \\ u \end{pmatrix} = (\partial V / \partial u_\ell) \cdot (\partial V / \partial u_\ell) \quad (21b)$$

we arrive at the desired equality (18).

Now we go over to the corresponding dimensionally regularized Feynman amplitude of both sides of Equation (18) as described above and obtain $\mathcal{T}'_G(p, n) = \mathcal{T}'_{G/\ell}(p, n)$ for $\text{Re } n \ll 0$. Analytic continuation and combinatorics will complete the proof of (13).

It remains to prove Equation (15)⁹. Again reducing it to the language of graphs it means

$$-i \partial \mathcal{T}_G / \partial m_\ell = \mathcal{T}'_G \quad (22)$$

where G' is obtained from G by insertion of the 2-vertex with vertex part $\partial D / \partial m_\ell$ into line ℓ of G .

From

$$\begin{aligned} D Z_\ell &= -(\square_{u_\ell} + m_\ell^2) \quad \text{we get} \quad (\partial D / \partial m_\ell) Z_\ell + D \partial Z_\ell / \partial m_\ell = -2m_\ell \\ \text{resp. } (\square_{u_\ell} + m_\ell^2) \partial Z_\ell / \partial m_\ell &= 2m_\ell Z_\ell + Z_\ell (\partial D / \partial m_\ell) Z_\ell. \end{aligned} \quad (23)$$

Using Equation (18) and $\int_0^\infty d\alpha_\ell I_G(p, u, \alpha) = -\int_0^\infty d\alpha_\ell \alpha_\ell \partial I_G / \partial \alpha_\ell$ we get¹⁰ for sufficiently small n

$$\begin{aligned} -i \partial \mathcal{T}_G / \partial m_\ell &\approx -i \int_0^\infty ((\partial Z_\ell / \partial m_\ell) Z_\ell^{-1} - 2i\alpha_\ell m_\ell) I_G(p, u, \alpha) d\alpha_\ell|_{u=0} \\ &= \int_0^\infty (i\alpha_\ell (\partial / \partial \alpha_\ell) (\partial Z_\ell / \partial m_\ell) Z_\ell^{-1} - 2\alpha_\ell m_\ell) I_G d\alpha_\ell|_{u=0} \\ &= \int_0^\infty \alpha_\ell Z_\ell (\partial D_\ell / \partial m_\ell) I_G d\alpha_\ell|_{u=0}. \end{aligned} \quad (24)$$

⁹ The only “nontrivial” case is, when \mathcal{L}_0 depends on λ . We consider only the case where λ is a mass, but the other possibilities can be treated in a similar way

¹⁰ Correct ordering of the Z -factors is understood here

On the other hand

$$\begin{aligned}\mathcal{T}_{G'} &\approx \int_0^\infty d\alpha_\ell d\alpha_{\ell'} Z_{\ell'} (-i \partial/\partial u_{\ell'}) (\partial D/\partial m_{\ell'}) I_G(p, u_\ell + u_{\ell'}, \alpha_\ell + \alpha_{\ell'})|_{u=0} \\ &= \int_0^\infty \alpha_\ell d\alpha_\ell Z_\ell (-i \partial/\partial u_\ell) (\partial D/\partial m_\ell) I_G(p, u, \alpha)|_{u=0}\end{aligned}\quad (25)$$

using

$$\int_0^\infty d\alpha_1 d\alpha_2 f(\alpha_1 + \alpha_2) = \int_0^\infty \alpha d\alpha f(\alpha).$$

Analytic continuation and some combinatorial considerations complete the proof of Equation (15).

II.4. Ward Identities, etc.

Choosing the infinitesimal transformations of some Lie-group as variations we may derive Ward identities from the action principle, Equation (13):

$$\delta_a \phi_i(x) = t_{ij}^a \phi_j(x) \delta e(x) \quad (26)$$

with real, antihermitean matrices t_{ij}^a . We get from Equation (13a)

$$\begin{aligned}\delta \mathcal{L}(x) &= (\delta \mathcal{L}/\delta \phi_i(x)) t_{ij}^a \phi_j(x) + (\delta \mathcal{L}/\delta \partial_\mu \phi_i(x)) t_{ij}^a \partial_\mu \phi_j(x) \\ &\quad - \partial_\mu (\delta \mathcal{L}/\delta \partial_\mu \phi_i(x)) t_{ij}^a \phi_j(x) \\ &= \delta_a \mathcal{L}(x) - \partial_\mu J_\mu^a(x)\end{aligned}\quad (27)$$

with the definition

$$J_\mu^a(x) = (\delta \mathcal{L}/\delta \partial_\mu \phi_i(x)) t_{ij}^a \phi_j(x).$$

Isolating the source term in $\mathcal{L} = \bar{\mathcal{L}} + \int J(x) \phi(x) dx$ we obtain from Equation (13) the Ward identity

$$\langle (\partial^\mu J_\mu^a(x) - \delta_a \bar{\mathcal{L}}(x) - J_i(x) t_{ij}^a \phi_j(x)) \exp i\hbar^{-1} \int \mathcal{L}_{\text{int}}(x) dx \rangle_{+,c}^0 = 0. \quad (28)$$

The generalization of this derivation to Lagrangians including gauge fields is straightforward¹¹. Using the so-called Slavnov transformation introduced in [3] one may in the same manner derive Slavnov-Taylor identities for non-abelian gauge theories.

A word of caution, however, has to be said about Lagrangians resp. variations involving γ_5 . Since γ_5 does not anticommute with γ_μ (see Prop. 1) certain simplifications usually performed in the evaluation of $\delta_a \mathcal{L}$ may not be possible in n dimensions. This can lead to anomalies in the process of renormalization. A typical example for that is the axial vector current in theories with fermions.

The same difficulty occurs for Ward identities in super symmetric theories which depend on the identity $\gamma_5 \psi (\bar{\psi} \gamma_5 \psi) + \psi (\bar{\psi} \psi) = 0$ for a Majorana field. For $n=4$ this follows from Fierz symmetry, a typical 4-dimensional property derived

¹¹ We want to recall, however, that we excluded theories with massless particles

from the completeness relation for the γ -algebra, which cannot directly be generalized to $n \neq 4$ and, therefore, may give rise to anomalies.

A similar situation arises in the derivation of trace identities from the variation $\delta\phi(x) = (d_\phi + x_\mu \partial^\mu)\phi(x)$. If the dimensions d_ϕ are chosen such that $\delta\mathcal{L}_0(x)$ is a divergence, $\delta\mathcal{L}_{\text{int}}$ will contain (apart from mass terms) a contribution proportional to $n-4$, which again will lead to anomalies in the process of renormalization. The problem of these anomalies is discussed in Section III.3.

As a second application of the action principle let us derive the renormalization group—resp. Callan-Symanzik equation in the ϕ^4 -theory. Our discussion follows closely Ref. [11].

We take the Lagrangian

$$\mathcal{L}(x) = ((1+b)/2) \partial_\mu \phi \partial^\mu \phi - ((m^2 - a)/2) \phi^2 - ((g - c)/4!) \phi^4 \quad (29)$$

where a , b , and c are chosen such that the following normalization conditions are fulfilled (Γ_N denotes the 1PI vertex functions with N arguments)

$$\Gamma_2(p^2 = m^2) = 0, \quad (30a)$$

$$\Gamma_2(p^2 = \mu^2) = i(\mu^2 - m^2), \quad (30b)$$

$$\Gamma_4(\text{s.p.}) = -ig, \text{ s.p. = symmetry point: } p_i p_j = (4\delta_{ij} - 1)\mu^2/3, \quad i, j = 1, \dots, 4. \quad (30c)$$

Applying the action principle Equation (15) we find

$$\partial\Gamma_N/\partial m^2 = ((\partial a/\partial m^2 - 1)\Delta_1 + (\partial b/\partial m^2)\Delta_2 + (\partial c/\partial m^2)\Delta_3)\Gamma_N, \quad (31a)$$

$$\partial\Gamma_N/\partial\mu^2 = ((\partial a/\partial\mu^2)\Delta_1 + (\partial b/\partial\mu^2)\Delta_2 + (\partial c/\partial\mu^2)\Delta_3)\Gamma_N, \quad (31b)$$

$$\partial\Gamma_N/\partial g = ((\partial a/\partial g)\Delta_1 + (\partial b/\partial g)\Delta_2 + (\partial c/\partial g - 1)\Delta_3)\Gamma_N, \quad (31c)$$

where the Δ_i signify the insertion of the extra vertices $\Delta_1 = i2^{-1} \int dx \phi^2(x)$, $\Delta_2 = i2^{-1} \int dx \partial_\mu \phi \partial^\mu \phi$ and $\Delta_3 = (i/4!) \int dx \phi^4(x)$ into the graphs for Γ_N . Putting $P = \phi$ in Equation (13) we get in addition

$$(-2(m^2 - a)\Delta_1 + 2(1+b)\Delta_2 - 4(c-g)\Delta_3 + N)\Gamma_N = 0. \quad (31d)$$

Eliminating the Δ_i from (31d) in favour of the diff. operators in (31a–c) it takes the form

$$[\mu^2 \partial/\partial\mu^2 + \varrho m^2 \partial/\partial m^2 + \sigma \partial/\partial g - \tau N]\Gamma_N = 0, \quad (32a)$$

$$[\mu^2 \partial/\partial\mu^2 + m^2 \partial/\partial m^2 + \beta \partial/\partial g - \gamma N]\Gamma_N = \alpha m^2 \Delta_1 \Gamma_N \quad (32b)$$

with some coefficients α depending on m , μ , g and β , γ , ϱ , σ , τ depending on m/μ and g . Applying Equation (32) to the normalization conditions Equation (30) we get

$$\varrho = 0, \quad \sigma = 4g\tau - i\mu^2 \partial\Gamma_4/\partial\mu^2 \text{ (s.p.)}, \quad \tau = \mu^2 (\partial\Gamma_2/\partial\mu^2)(p^2 = \mu^2)/2i(\mu^2 - m^2)$$

and similar for α , β , γ .

Equations (32) are the renormalization group resp. Callan-Symanzik equation. Note that there is no hard mass insertion like in Ref. [11] involved since Δ_1 is soft.

III. Dimensional Renormalization

III.1. Definition of the Renormalized Amplitudes

As we have emphasized in Section II, the dimensionally regularized Feynman amplitudes are meromorphic functions of the complex parameter n . The problem of renormalization consists in subtracting the poles of these functions at $n=4$ in a way which is compatible with additive renormalization. That is, the subtractions have to be performed in a way that they correspond to counterterms in the Lagrangian. What is to be subtracted for a graph G is the *singular part* of the amplitude for H inserted into the amplitude for the reduced graph G/H . We will symbolize this subtraction by the application of a subtraction operator C_H analogous to the Taylor series subtraction operator in the BPHZ scheme. The *singular part* of the amplitude for H is defined to be the singular part of the Laurant series expansion at $n=4$ of $\bar{\mathcal{T}}_H$, the amplitude resulting from \mathcal{T}_H after performance of the subtractions corresponding to all its genuine 1PI subgraphs.

Theorem 1. *The singular part of the dimensionally regularized amplitude for H as defined above consists of poles of order h_H or less and is a polynomial of degree ω_H in the external momenta of the graph. The singular part vanishes if H is superficially convergent. The amplitudes $\mathcal{R}_{G,\epsilon}(p, n)$, remaining after performance of the subtractions corresponding to all 1PI subgraphs, are analytic at $n=4$ in any order of perturbation theory. The $\lim_{\epsilon \rightarrow 0} \mathcal{R}_{G,\epsilon}(p, n)$ exists in $\mathcal{S}'(\mathbb{R}^{4M})$ and is again analytic $n=4$.*

This limit $\mathcal{R}_G(p, n)$ (or its value for $n=4$) is the renormalized amplitude.

Remark. Clearly the overall singularity at $n=4$ of the dimensionally regularized Feynman amplitude is not a polynomial in its external momenta. This becomes only true after combination with the counterterms corresponding to all its genuine 1PI subgraphs.

It is straightforward to show that the procedure described here differs by a finite renormalization from the BPHZ scheme. Therefore, the procedure defines a renormalization fulfilling the usual requirements of causality and unitarity [5]. As usual, the physical normalization conditions can be achieved by finite counterterms in the Lagrangian.

III.2. Proof of Theorem 1

Let us first discuss the limit $\epsilon \rightarrow 0$. Since the renormalized amplitude (for $n \neq 4$) is a finite sum of the unrenormalized, regularized amplitude + counterterms, each of which has a continuous $\lim_{\epsilon \rightarrow 0}$ in $\mathcal{S}'(\mathbb{R}^{4M})$, there is no question that also $\lim_{\epsilon \rightarrow 0} \mathcal{R}_{G,\epsilon}(p, n)$ exists in $\mathcal{S}'(\mathbb{R}^{4M})$ for $n \neq 4$. Since $\mathcal{R}_{G,\epsilon}(p, n)$ is analytic at $n=4$, $\lim_{\epsilon \rightarrow 0} \mathcal{R}_{G,\epsilon}(p, n)$ can be continued to $n=4$.

In order to prove that the subtractions actually have the required properties we have to exhibit the singularities connected with all the 1PI subgraphs of a given graph G and combine them with the corresponding counterterms. As usual, the complication due to overlapping divergences is resolved by the decomposition

of the integration region in α -space according to families of non-overlapping subgraphs. It is, however, to be remarked that in contrast to BPH-renormalization the subtraction corresponding to a particular subgraph H can be completely absorbed into the integrations over those sectors related to families containing H , leading to the simple representation Equation (3) below.

We use maximal forests as defined in Appendix A, since they correspond directly to counterterms in the Lagrangian. They differ from the s -families of Speer [9] mainly by a different definition of overlapping graphs.

According to Lemma 3 the regularized Feynman amplitude can be decomposed

$$\mathcal{T}_G = \lim_{\varepsilon \rightarrow 0} \mathcal{T}_{G,\varepsilon} = \lim_{\varepsilon \rightarrow 0} \int_0^\infty d\underline{\alpha} I_{G,\varepsilon}(p, \underline{\alpha}) = \lim_{\varepsilon \rightarrow 0} \sum_{(\mathcal{C}, \sigma)} \int_{\mathcal{D}(\mathcal{C}, \sigma)} d\underline{\alpha} I_{G,\varepsilon}(p, \underline{\alpha}) \quad (1)$$

where the sum extends over all labelled forests (\mathcal{C}, σ) for G . Resolving the recursive definition of the subtraction operators into a forest formula, anticipating that we know the action of C_H on I_G , the renormalized amplitude can be decomposed accordingly

$$\begin{aligned} \mathcal{R}_G &= \lim_{\varepsilon \rightarrow 0} \mathcal{R}_{G,\varepsilon} = \lim_{\varepsilon \rightarrow 0} \int_0^\infty d\underline{\alpha} \sum_{F \in \mathcal{F}} \prod_{H \in F} (-C_H) I_{G,\varepsilon}(p, \underline{\alpha}) \\ &= \lim_{\varepsilon \rightarrow 0} \sum_{(\mathcal{C}, \sigma)} \int_{\mathcal{D}(\mathcal{C}, \sigma)} d\underline{\alpha} \sum_{F \in \mathcal{F}} \prod_{H \in F} (-C_H) I_{G,\varepsilon}(p, \underline{\alpha}) \end{aligned} \quad (2)$$

where \mathcal{F} is the set of all forests of non-overlapping 1PI subgraphs of G . Since C_H will be defined such that $\int_{\mathcal{D}(\mathcal{C}, \sigma)} d\underline{\alpha} \prod_{H \in F} (-C_H) I_{G,\varepsilon} = 0$ unless $F \subset \mathcal{C}$, we can rewrite the renormalized amplitude

$$\begin{aligned} \mathcal{R}_{G,\varepsilon} &= \sum_{(\mathcal{C}, \sigma)} \int_{\mathcal{D}(\mathcal{C}, \sigma)} d\underline{\alpha} \sum_{F \subset \mathcal{C}} \prod_{H \in F} (-C_H) I_{G,\varepsilon}(p, \underline{\alpha}) \\ &= \sum_{(\mathcal{C}, \sigma)} \int_{\mathcal{D}(\mathcal{C}, \sigma)} d\underline{\alpha} \prod_{H \in \mathcal{C}} (1 - C_H) I_{G,\varepsilon}(p, \underline{\alpha}). \end{aligned} \quad (3)$$

It is easy to verify that the regularized amplitude for a 1PR graph is the product of the regularized amplitudes for its components and of the (momentum space) propagators for all lines connecting them. This factorization is obviously maintained for the decomposition of the renormalized amplitude as given by Equations (2, 3) [compare Lemma 3 (a)]. We will therefore assume for the remainder of this subsection (III.2) that G is 1PI in order to avoid complicated notation.

For each labelled forest (\mathcal{C}, σ) we introduce the familiar scaling variables $(t, \beta) = (t_H, H \in \mathcal{C}, \beta_\ell, \ell \in \mathcal{L}'_G = \mathcal{L}_G \setminus \sigma(\mathcal{C}))$ [8] as well as auxiliary variables ξ_H, ζ_H

$$\alpha_\ell = \begin{cases} \prod_{H \subset H' \in \mathcal{C}} t_{H'}^2 = t_H^2 \xi_H^2 = \zeta_H^2 & \text{if } \ell = \sigma(H), \quad H \in \mathcal{C}, \\ \beta_\ell \xi_H^2 & \text{if } \ell \in \mathcal{L}'_H = \mathcal{L}_H \setminus \sigma(H), \quad H \in \mathcal{C}, \end{cases}$$

and define $\beta_\ell = 1$ if $\ell \in \sigma(\mathcal{C})$ for convenience.

The image of $\mathcal{D}(\mathcal{C}, \sigma)$ under this substitution is the set

$$\{(\underline{t}, \beta) : 0 \leq t_G < \infty; 0 \leq t_H \leq 1 \text{ for } H \neq G; 0 \leq \beta_\ell \leq 1 \text{ for } \ell \in \mathcal{L}'_G\}.$$

Using Lemma 4 we may rewrite the contribution of $\mathcal{D}(\mathcal{C}, \sigma)$ to $\mathcal{T}_{G,\epsilon}$

$$\begin{aligned} \int_{\mathcal{D}(\mathcal{C}, \sigma)} d\underline{x} I_{G,\epsilon}(p, \underline{x}) &= \prod_{\ell \in \mathcal{L}'_G} \int_0^1 d\beta_\ell \prod_{H \in \mathcal{C}} \left\{ 2 \int_0^{\theta_H} (dt_H/t_H) t_H^{\omega_H - \omega_H} Z_H(-i \partial/\partial \tilde{u}) \right\} \\ &\quad \cdot g_\epsilon(\tilde{q}, \tilde{u}, \underline{t}, \beta, v) \\ &= \prod_{H \in \mathcal{C}} \left\{ \int d\mu_H \zeta^{v - \omega_H} Z_H(-i \partial/\partial \tilde{u}) \right\} g_\epsilon(\tilde{q}, \tilde{u}, \underline{t}, \beta, v) \end{aligned} \quad (4)$$

where $v = n - 4$, $\omega_{\bar{H}} = \omega_H - \sum_{H' \in \mathcal{M}(H)} \omega_{H'}$, $Z_H(-i \partial/\partial \tilde{u}) = \prod_{\ell \in \mathcal{L}_{\bar{H}}} \{\zeta_H^\ell Z_\ell(-i \partial/\partial u)\}$ and $\int d\mu_H = 2 \int_0^{\theta_H} dt_H/t_H \int_0^1 \prod_{\ell \in \mathcal{L}'_{\bar{H}}} d\beta_\ell$ with $\theta_G = \infty$ and $\theta_H = 1$ for $H \neq G$.

The Z_H are polynomials in \underline{t} and g_ϵ is C^∞ in (\underline{t}, β) , analytic in v and exponentially decreasing as $t_G \rightarrow \infty$. For the definition of $\bar{H}, \tilde{q}, \tilde{u}$ etc. see Appendix A.

Considered as a generalized function $t_{H,+}^\lambda$ is a meromorphic function of λ with simple poles at all negative integers

$$t_+^{\omega_H - \omega - 1} = (-)^\omega \delta^{(\omega)}(t)/\omega! v h + \text{regular at } v=0.$$

[We put $\delta^{(\omega)}(t)/\omega! = 0$ and $(d/dt)^\omega/\omega! = 0$ for $\omega < 0$.] This will enable us to define the action of the subtraction operator C on the Feynman integrand. Correspondingly we can decompose the t_H -integral in Equation (4) into

$$(\omega_H! v h_H)^{-1} (d/dt_H)^\omega g_\epsilon(\dots t_H \dots)|_{t_H=0} + \int_0^\infty (\dots)_{\text{reg.}} dt_H. \quad (5)$$

Due to the structure of the integrand I_ϵ the first part is a polynomial of degree ω_H in the external momenta of H . The second part is analytic at $n=4$ and may be represented by the following integral, convergent for $|\text{Re}(v h_H)| < 1$

$$\begin{aligned} \int_0^\infty (\dots)_{\text{reg.}} dt_H &= \int_0^\infty dt_H t_H^{\omega_H - \omega_H - 1} \left[g_\epsilon(\dots t_H \dots) - \sum_{k=0}^{\omega_H-1} (t_H^k/k!) g^{(k)}(\dots t_H=0 \dots) \right. \\ &\quad \left. - \theta(1-t_H)(t_H^{\omega_H}/\omega_H!) g^{(\omega_H)}(\dots t_H=0 \dots) \right]. \end{aligned} \quad (6)$$

Next we proceed to the definition of the subtraction operators C_H acting on the Feynman integrand $I_{G,\epsilon}$. Let us denote by \mathcal{A}_G the formal algebra of covariants generated by $\{p_{i,\mu}\}_{i \in \mathcal{V}_G}$ and $\{u_{\ell,\mu}\}_{\ell \in \mathcal{L}_G}$ (and by γ_μ 's etc. if required) with complex coefficients depending meromorphically on n .

On $A \in \mathcal{A}_G$ we define the operator C (not yet $C_H!$) by

$$C(A) = \text{sing. part at } n=4 \text{ of } \text{NF}(A|_{u=0}).$$

Notice that C is precisely the operator to be used in a recursive definition of the counterterms.

As a first step versus the construction of a suitable domain for the C_H we consider the set \mathcal{G}_G of triplets $\hat{X} = (X, F_X, (\mathcal{C}, \sigma)_X)$ with $X \subset G$, $F_X \in \hat{\mathcal{F}}'_X = \{\text{forests of pairwise disjoint, 1PI, proper subgraphs of } X\}$, $(\mathcal{C}, \sigma)_X$ a labelled forest for X/F_X .

Next we introduce a set \mathcal{E}_G of functions on \mathcal{G}_G with values in $\mathcal{A}_G : \mathcal{E}_G = \{f : \mathcal{G}_G \rightarrow \mathcal{A}_G, f(\hat{X}) \text{ depends only on the } (p, u) \text{ from } X/F_X\}$. As the domain for C_H we take the subset $\mathcal{D}_H = \{f \in \mathcal{E}_G : \text{for all } \hat{X} \text{ of the form } \hat{X} = (H, \emptyset, (\mathcal{C}, \sigma)_H) \quad C(f(\hat{X})) \text{ is a polynomial in } p\}$.

One particular such function I_G can be affiliated with the Feynman amplitude for G . $I_G : \hat{X} \rightarrow I_{X/F_X}$, represented in the form of Equation (4) in the sector $\mathcal{D}(\mathcal{C}, \sigma)_X$. The meromorphic dependence of the coefficients on n is achieved by interpreting the fractional powers of t 's in the distributional sense as displayed in Equations (5, 6).

On \mathcal{D}_H the operator C_H is defined by

$$(C_H f)(\hat{X}) = \begin{cases} f(\hat{X}) & \text{if either } X \cap H = \emptyset \text{ or } H \not\subset H' \text{ for some } H' \in F_X \\ C(f(\hat{X})) & \text{if } X = H \text{ and } F_X = \emptyset \\ U_H(C(f(\hat{H}_0))) f(\widehat{X:H}) & \text{if } H \notin F_X, F_X \cup \{H\} \in \mathcal{F}'_X \\ 0 & \text{else} \end{cases}$$

where $\hat{H}_0 = (H, \emptyset, (\mathcal{C}, \sigma)_H)$ and $\widehat{X:H} = (X, F_X \cup \{H\}, (\mathcal{C}, \sigma)_{X:H})$ with $(\mathcal{C}, \sigma)_H, (\mathcal{C}, \sigma)_{X:H}$ the pair of labelled forests for H resp. $X/F_X \cup \{H\}$ corresponding to $(\mathcal{C}, \sigma)_X$ (see Appendix A). U_H is defined in Appendix B.

Equation (3) is to be interpreted as $\sum_{(\mathcal{C}, \sigma)} \int d\xi d\beta \left[\prod_{H \in \mathcal{C}} (1 - C_H) I_G \right] (G, \emptyset, (\mathcal{C}, \sigma))|_{u=0}$ where the factors $(1 - C_H)$ are to be ordered such that those for smaller graphs act first.

The main problem in combining the singular part of the integral with the expression for the counterterm for H inserted into G/H comes from the fact that the coefficient of the pole at $v=0$ exhibited in Equation (4) depends on the number of loops of H . The subtractions corresponding to genuine subgraphs of H can be interpreted as counterterms inserted into reduced graphs and these reduced graphs will have fewer loops than H has. Therefore the term t^{v_h} in Equation (4) will be replaced by a sum of terms with different exponents and furthermore with coefficients which have poles at $v=0$. The following definition will be useful to trace such sums through the recursion.

Given a maximal forest \mathcal{C} for G , we define sets of functions J_H^K (for $0 \leq K < h_H$) and \tilde{J}_H^K (for $0 \leq K \leq h_H$) for all $H \in \mathcal{C}$

$$J_H^K = \left\{ f(\xi, v) : f(\xi, v) = \xi^v \prod_{H' \in \mathcal{M}(H)} g_{H'}(\xi, v) \text{ with } g_{H'} \in \tilde{J}_{H'}^{K_{H'}}, K = \sum_{H' \in \mathcal{M}(H)} K_{H'} \right\};$$

$$\tilde{J}_H^K = \left\{ g(\xi, v) : g \text{ or } \xi^v g \text{ with } g \in J_H^K \text{ or } g(\xi, v) = \int_1^\xi dx/x f(x, v) \text{ with } f \in J_H^{K-1} \right\}.$$

If $h_H = 1$ (i.e. $\mathcal{M}(H) = \emptyset$) there is only $K = 0$ and $J_H^0 = \{\xi^v\}$.

Lemma 2. Let $f(\xi, v)$ be any element of J_H^K , then

$$(a) \quad f(\xi, v) = v^{-K} \sum_{m=1}^{h_H} c_m \xi^{mv} \quad \text{with some constants } c_m; \quad (7)$$

$$(b) \quad f(\xi, 0) = c(\ln \xi)^K \quad \text{with some constant } c; \quad (8)$$

$$(c) \quad f(\xi t, v) = \sum_j f_{1j}(\xi, v) f_{2j}(t, v) \quad (9)$$

with $f_{ij} \in J_{H'}^{K_{ij}}$ such that $K_{1j} + K_{2j} = K$;

$$(d) \quad J_H^0 = \{\xi^{mv}, m=1, \dots, h_H\}. \quad (10)$$

Proof. All statements are true if $h_H = 1$. (a), (b) and (d) follow easily from the recursive definition of the sets J_H^K . Assume that (c) holds for all H' which are genuine subgraphs of H and let $h(x, v) \in J_{H'}^{K-1}$. Then

$$\begin{aligned} \int_1^{\xi t} (dx/x) h(x, v) &= \int_1^t (dx/x) h(x, v) + \int_1^{\xi} (dx/x) h(xt, v) \\ &= \int_1^t (dx/x) h(x, v) + \sum_j \int_1^{\xi} (dx/x) h_{1j}(x, v) h_{2j}(t, v) \end{aligned}$$

with $h_{ij} \in J_{H'}^{K_{ij}}$ such that $K_{1j} + K_{2j} = K - 1$. Insertion of this decomposition into the definition of J_H^K proves the statement (c) for H .

It should be noted that the pole of order K at $v=0$, indicated by the decomposition (7) is not really there. This is made explicit by Equation (8).

Now we are ready to prove

Proposition 3. Consider any $\mathcal{X}_0 \in \hat{\mathcal{P}}'_G$, $\mathcal{X}_0 \subset \mathcal{C}$ and define

$$\mathcal{X} = \{H' \in \mathcal{C} : H' \subset H \text{ for some } H \in \mathcal{X}_0\}.$$

After performance of all subtractions corresponding to subgraphs $H \in \mathcal{X}$, the contribution of (\mathcal{C}, σ) to $\mathcal{R}_{G, \varepsilon}$ is a sum of terms of the form

$$\prod_{H \in \mathcal{C} \setminus \mathcal{X}} \left\{ \int d\mu_H (1 - C_H) \zeta_H^{-\omega_H} Z_H(-i \partial/\partial \tilde{u}) \right\} \prod_{H \in \mathcal{X}_0} \left\{ \zeta_H^{-\omega_H} g_H(\zeta_H, v) \right\} \cdot g_{\mathcal{X}}(\tilde{q}, \tilde{u}, \underline{t}, \beta, v)|_{\tilde{u}=0} \quad (11)$$

where (\underline{t}, β) and \tilde{u} are scaling variables and scaled u 's for G/\mathcal{X}_0 (i.e. \underline{u}_H for $H \in \mathcal{X}$ are already set to zero) and \tilde{q} are scaled momenta appropriate for the family $\mathcal{C} \setminus \mathcal{X}$. $g_H \in \tilde{\mathcal{J}}_H^K$ for some K and $g_{\mathcal{X}}$ is some element of the abstract algebra of covariants with complex coefficients which are C^∞ in (\underline{t}, β) , analytic at $v=0$ and, due to $\varepsilon > 0$, exponentially decreasing as $t_G \rightarrow \infty$.

Proof (by induction with respect to $\sum_{H \in \mathcal{X}_0} h_H = |\mathcal{X}|$).

- i) The statement certainly holds for $\mathcal{X} = \emptyset$ [compare Eq. (4)].
- ii) Consider some $\mathcal{X} \subset \mathcal{C}$ ($\mathcal{X} \neq \emptyset$) $\mathcal{X}_0 \in \hat{\mathcal{P}}'_G$, choose any $H \in \mathcal{X}_0$ and assume that the statement is true for $\mathcal{X}'_0 = \mathcal{M}(H) \cup \mathcal{X}_0 \setminus \{H\}$ (i.e. $\mathcal{X}' = \mathcal{X} \setminus \{H\}$). If $\mathcal{X}_0 = \{G\}$ we will prove Theorem 1, otherwise we will show that the proposition holds for \mathcal{X}_0 as well.

The relevant part of Equation (11) is

$$\int d\mu_H (1 - C_H) \zeta_H^{-\omega_H} Z_H (-i \partial / \partial \tilde{u}) \prod_{H' \in \mathcal{M}(H)} (\zeta_{H'}^{-\omega_{H'}} g_{H'}(\zeta_{H'}, v)) g_{\mathcal{X}'}|_{\tilde{u}_H=0}. \quad (12)$$

First we notice that $\xi_{H'} = \zeta_H = t_H \xi_H$ for $H' \in \mathcal{M}(H)$. Evaluating the derivatives with respect to \tilde{u}_H , setting $\tilde{u}_H = 0$ and integrating over β 's we obtain

$$(12) = (1 - C_H) \int_0^\infty (dt_H/t_H) (\zeta_H t_H)^{-\omega_H} f(\zeta_H t_H, v) g'(t_H, \tilde{q}, \tilde{u}, \underline{t}, \underline{\beta}, v) \quad (13)$$

where $f \in J_H^K$ and we have replaced the variables $(\tilde{q}', \tilde{u}', \underline{t}', \underline{\beta}')$, appropriate for \mathcal{X}' , by those for \mathcal{X} . In particular $\tilde{q}_H = (\tilde{q}'_H/t_H, \tilde{q}'_{H'}, H' \in \mathcal{M}(H))$ and therefore g' is again C^∞ in $(t_H, \underline{t}, \underline{\beta})$ and analytic at $v=0$. Furthermore $g'=0$ for $t_H > 1$ unless $H=G$. Using Equations (5, 7) we get

$$(12) = (1 - C_H) \zeta_H^{-\omega_H} \left\{ v^{-K-1} \sum_{m=0}^{h_H} c_m m^{-1} \zeta_H^{\omega_H m} \left[\frac{1}{\omega_H!} (d/dt_H)^{\omega_H} g' \right]_{t_H=0} \right. \\ \left. + \int_0^\infty ((dt_H/t_H) t_H^{-\omega_H} f(\zeta_H t_H, v) g')_{\text{reg.}} \right\}. \quad (14)$$

In contrast to Equation (7) there is now really a pole of order $K+1 \leq h_H$ at $v=0$. If $H=G$ then $\zeta_H=1$, $\tilde{q}=q$ and there are no \tilde{u} 's left in g' . As shown in Appendix B

$$P(q) = [(\omega_H!)^{-1} (d/dt_H)^{\omega_H} g']_{t_H=0} \quad (15)$$

is a polynomial of degree ω_H in the external momenta q (a homogeneous polynomial in momenta, masses and $\sqrt{\varepsilon}$ if all Z_ℓ are homogeneous in these variables). We reduce $P(q)$ to NF in order to make its v -dependence explicit. We do, however, not interpret $P(q)$ as a 4-dimensional covariant at this point. Due to the exponential decrease of g' the regularized integral is absolutely convergent and analytic in v . Using the Taylor operator T_v^K of degree K in v we obtain

$$v^{-K-1} \sum_m c_m m^{-1} T_v^K P(q)$$

as one term in the contribution of (\mathcal{C}, σ) to the counterterm for H which is annihilated by $(1 - C_H)$. The contribution to $\mathcal{R}_{H,\varepsilon}$

$$(12) = \sum_m c_m m^{-1} v^{-K-1} (1 - T_v^K) P(q) + \int_0^\infty ((dt_H/t_H) t_H^{-\omega_H} f(t_H, v) g')_{\text{reg.}} \quad (16)$$

is analytic at $v=0$ ($n=4$) by construction. This proves Theorem 1, once we have completed the proof of Proposition 3.

If $H \neq G$ we use Lemma 5 to rewrite $[(\omega_H!)^{-1} (d/dt_H)^{\omega_H} g']_{t_H=0}$ as the amplitude for G/H with the polynomial P [Eq. (15)] inserted as vertex part:

$$[(\omega_H!)^{-1} (d/dt_H)^{\omega_H} g']_{t_H=0} = \tilde{P}(\tilde{q}_H, -i \partial / \partial \tilde{u}_{G/H}) g_{G/H}$$

where

$$\tilde{P}(\tilde{q}_H, -i \partial / \partial \tilde{u}_{G/H}) = \zeta_H^{\omega_H} P(q_H - i e_{GH}^+ \partial / \partial u_G^+)$$

is a polynomial in \tilde{q} , $\partial / \partial \tilde{u}$ and \underline{t} .

Let $((\mathcal{C}_1, \sigma_1), (\mathcal{C}_2, \sigma_2))$ be the pair of labelled forests for G/H and H corresponding to (\mathcal{C}, σ) . Due to Lemma 5 $g_{G/H}$ is the precise analogue of $g_{\mathcal{X}}$ in Equation (11) but for the contribution of $(\mathcal{C}_1, \sigma_1)$ to $\mathcal{R}_{G/H, \varepsilon}$ after subtracting all counterterms corresponding to subgraphs contained in $\mathcal{X} \cap \mathcal{C}_1$. The counterterm for H can now be inserted into the partially subtracted amplitude for G/H , i.e. $(1 - C_H)$ can be evaluated and we obtain

$$(12) = \xi_H^{-\omega_H} \left\{ v^{-K-1} \sum_m c_m m^{-1} [(\xi_H^{vm} - T_v^K) \tilde{P}] g_{G/H} + \int_0^\infty (\dots)_{\text{reg.}} \right\}. \quad (17)$$

Using Equation (7) we may rewrite the first part as

$$\begin{aligned} & v^{-K-1} \sum_m c_m m^{-1} [(\xi_H^{vm} - T_v^K) \tilde{P}] g_{G/H} \\ &= v^{-K-1} \sum_m c_m m^{-1} \{ [(\xi_H^{vm} - 1) T_v^K + \xi_H^{vm} (1 - T_v^K)] \tilde{P} \} g_{G/H} \\ &= \int_1^{\xi_H} (dt/t) f(t, v) (T_v^K \tilde{P}) g_{G/H} + \sum_m c_m m^{-1} \xi_H^{vm} [v^{-K-1} (1 - T_v^K) \tilde{P}] g_{G/H} \end{aligned} \quad (18)$$

and evaluate the derivatives with respect to \tilde{u} contained in \tilde{P} . For the regularized integral we use Equation (9) to get

$$\begin{aligned} \int_0^\infty (\dots)_{\text{reg.}} &= \sum_j f_{1j}(\xi_H, v) \left\{ \int_0^1 (dt/t) f_{2j}(t, v) t^{-\omega_H} (1 - T_t^{\omega_H}) g' \right. \\ &\quad \left. - \int_1^\infty (dt/t) f_{2j}(t, v) t^{-\omega_H} (T_t^{\omega_H-1} g') \right\} \end{aligned} \quad (19)$$

where both integrals are absolutely convergent and yield a function C^∞ in (\underline{t}, β) and analytic at $v=0$. Collecting all contributions of Equations (18, 19) we arrive at

$$(12) = \xi_H^{-\omega_H} \sum g_H(\xi_H, v) g_{\mathcal{X}}(\tilde{q}, \tilde{u}, \underline{t}, \beta, v) \quad (20)$$

where each g_H and $g_{\mathcal{X}}$ has all the properties required in Equation (11). Finally combining Equation (20) with all those terms not contained in Equation (12) we are led back to an expression of the form (11) but now for the set $\mathcal{X}_0 = \{H\} \cup \mathcal{X}'_0 \setminus \mathcal{M}(H)$. This completes the proof of both Proposition 3 and Theorem 1.

As a final remark it should be added that whenever we use the word counterterm it refers to the ones which are singular for $n=4$ and not to the additional finite counterterms which occur explicitly in the Lagrangian and serve to fix the physical normalization conditions.

III.3. The Renormalized Action Principle

Having defined the renormalized amplitudes in a satisfactory way we have to ask about the validity of the action principle in the renormalized theory, which is answered by

Theorem 2. *The action principle, Equations (II.13–15), remains valid without any change for the renormalized theory (for $n \neq 4$), if there is no explicit n -dependence involved. In particular, the classical equations of motion remain valid without any modification.*

The reservation made about an explicit n -dependence is to be understood in the following sense. The Action Principle leads to certain linear relations between regularized Feynman amplitudes reduced to their resp. NF. If these relations have n -independent coefficients they remain true for the renormalized amplitudes. In contrast, n -dependent coefficients are generally the source of anomalies, i.e. terms absent in the tree approximation. Such n -dependence can arise in two ways. Either the variation $\delta\phi$ depends explicitly on n , as for dilatations (the dimension d_ϕ of the field ϕ depends on n) or the n -dependence comes in during the reduction to NF in the course of eliminating $\hat{\mathcal{O}}$ -type covariants. A typical case for the latter situation is the Adler anomaly. $\{\gamma_\mu, \gamma_5\} = 2\hat{\gamma}_\mu\gamma_5$ can lead to terms containing $\hat{g}_{\mu\mu} = n - 4$. More generally one can say that operators like $(\bar{\psi}\{\gamma_\mu, \gamma_5\}\psi)^{\text{ren.}}$ or $((n-4)\phi^4)^{\text{ren.}}$ vanishing in the tree approximation are non-zero due to radiative corrections.

As was shown in Section II.3 the proof of the action principle is based essentially on the equality of two regularized amplitudes, one for a graph G with a factor $Z_\ell = i(\square + m^2)$ and the other one for a graph G/ℓ where the line ℓ is contracted to a point. Assume this equality still holds for $\tilde{\mathcal{T}}_G$, i.e. after performing all subtractions for genuine subgraphs, then uniqueness of the Laurant expansion guarantees that also the finite parts (i.e. the renormalized amplitudes) are equal and, therefore, satisfy the action principle. Here we have to use the convention that any parameter of the Lagrangian which might depend implicitly on n is *not* subject to the Laurant expansion.

If the maximal forests for the two graphs G and G/ℓ are in one to one correspondence we can immediately justify our assumption. Either a particular subgraph H does not contain the line ℓ : in this case the same singular part of the amplitude for H has to be inserted into the amplitudes for two reduced graphs which are equal. Or the line ℓ is contained in the subgraph H of G : then the subtractions corresponding to H and H/ℓ , being the singular part of two equal amplitudes, are equal and are inserted into the amplitude for the same reduced graph G/H . This again justifies recursively the above assumption, provided the maximal forests or equivalently the forests of non-overlapping 1PI subgraphs for G and G/ℓ are in one to one correspondence.

There will, however, in general be no such one to one correspondence for the following two reasons:

- i) There may exist 1PI subgraphs H of G which contain both vertices of ℓ but not the line ℓ itself. They do not have any corresponding subgraph in G/ℓ .
- ii) There may be 1PR subgraphs H of G such that H/ℓ becomes 1PI in G/ℓ .

Figures 1 and 2 exemplify these two possibilities. All other 1PI subgraphs of G are in one to one correspondence with 1PI subgraphs of G/ℓ .

The first case is very easily resolved for dimensional renormalization; subtractions for subgraphs like H in Figure 1 may just be omitted since they do not contribute to this particular amplitude. They do, however, contribute if there is




Fig. 1. Example of a subgraph $H \subset G$, which is 1PI, contains both endpoints of line ℓ , but not line ℓ'




Fig. 2. Example of a subgraph $H \subset G$, which is 1PR in G (a), but becomes 1PI in G/ℓ (b)

no factor $(\square + m^2 - i\varepsilon)$ for line ℓ . This is the reason why we do not use any normal ordering at all in the Gell-Mann-Low expansion (II.12). Since ℓ is a loop line in G/H , the contribution of C_H to \mathcal{R}_G would be proportional to

$$\begin{aligned} & \int_0^\infty d\alpha \alpha^{-n/2} Z'(\partial/\partial u)(\square_u + m^2 - i\varepsilon) \exp[-i(\alpha(m^2 - i\varepsilon) + u^2/4\alpha)]|_{u=0} \\ &= i \int_0^\infty d\alpha (\partial/\partial \alpha) \alpha^{-n/2} Z'(\partial/\partial u) \exp[-i(\alpha(m^2 - i\varepsilon) + u^2/4\alpha)]|_{u=0}. \end{aligned}$$

Assuming that Z' is a polynomial of degree $2k$ or less and performing the differentiations with respect to u , this gives

$$\begin{aligned} & \int_0^\infty d\alpha (\partial/\partial \alpha) \alpha^{-n/2-k} Z''(\alpha) \exp[-i\alpha(m^2 - i\varepsilon)] \\ &= \alpha^{-n/2-k} Z''(\alpha) \exp[-i\alpha(m^2 - i\varepsilon)]|_{\alpha=0} = 0. \end{aligned}$$

Here we have used that the integral is defined by analytic continuation in n and vanishes identical for $\text{Re}(n) < -2k$. The analogous problem has been discussed by Lam [8] for Zimmermann's renormalization scheme (normal product algorithm) [14, 15] whereas the problem of subtractions, related to subgraphs like the ones shown in Figure 2, has not been discussed in this context. It is shown in Appendix C that the classical equations of motion are valid also for this case without any modification. This is due to a special property of our subtraction operator C_H . Assume a graph $H = H_1 \cup H_2$ such that H_1 and H_2 overlap at a single vertex. The Feynman integrand for H factorizes $I_H = I_{H_1} \cdot I_{H_2}$ and the subtraction operators satisfy

$$(-C_H)(1 - C_{H_1} - C_{H_2}) = (-C_{H_1})(-C_{H_2}).$$

This is slightly stronger than the unitarity requirement which allows an additional finite counterterm. The proof in Appendix C shows, again in a recursive

way, that the whole sum (3) for $\mathcal{R}_{G/\ell}$ may be rewritten such that we finally get the analogous sum but over all forests of G . Therefore, the classical equations of motion are valid for the renormalized (amplitudes of) composite operators.

Acknowledgements. We are indebted to K. Symanzik for pointing out the relevance of Equation (II.32b) to us. Considerable improvements in the formulation of Section III and Appendix B are due to many discussions with E. R. Speer

Appendix A. Feynman Graphs

In this Appendix we have collected all relevant definitions and properties concerning Feynman graphs, following essentially the notation of Refs. [9] and [17].

A Feynman graph G is a collection of vertices $\{V_1, \dots, V_M\} = \mathcal{V}_G$ and of lines $\{\ell_1, \dots, \ell_L\} = \mathcal{L}_G$ together with a rule assigning to each line ℓ an initial vertex V_{i_ℓ} and a final vertex V_{f_ℓ} . The corresponding *incidence matrix* e_G is defined by

$$e_{\ell i} = \begin{cases} 1 & \text{if } V_i = V_{f_\ell} \neq V_{i_\ell}; \\ -1 & \text{if } V_i = V_{i_\ell} \neq V_{f_\ell}; \\ 0 & \text{otherwise.} \end{cases}$$

A subgraph $G' \subset G$ is a subset of lines and vertices of G which build again a Feynman graph in the above sense. A *generalized vertex* G' of G is a subgraph of G containing with any pair of vertices all lines in G joining them.

A Feynman graph G is called *connected* if $\text{rank}(e) = M - 1$, *irreducible* if it stays connected by removal of any one line or vertex, *one-particle irreducible* (1PI) if it is connected and stays so after removal of any one line, *one-particle reducible* (1PR) if it is not 1PI, *non-empty* if $\mathcal{V}_G \neq \emptyset$, *non-trivial* if $\mathcal{L}_G \neq \emptyset$. The maximal non-empty connected subgraphs of G are called its *connectivity components* (c -components).

Two subgraphs $G_1, G_2 \subset G$ are called *non-overlapping* ($G_1 \circledast G_2$) if either $G_1 \subset G_2$, $G_2 \subset G_1$ or $G_1 \cap G_2 = \emptyset$, otherwise they are overlapping ($G_1 \circ G_2$).

For any $G' \subset G$ the *reduced graph* G/G' is the graph obtained from G by removal of all lines of G' and replacement of all the vertices of each c -component of G' by one new vertex. For any set \mathcal{M} of pairwise disjoint subgraphs of G , G/\mathcal{M} is defined as the obvious extension of this definition.

In what follows G is a connected Feynman graph. The number $h_G = L - M + 1$ is called the *connectivity* (or number of independent loops). A *tree* T in G is a maximal subgraph of G with $h_T = 0$. A K -tree ($K \geq 2$) is a tree minus $K - 1$ lines. We call the maximal non-trivial 1PI subgraphs of G the *(1PI-) components* of G .

Let $\{\alpha_\ell\}$ be a set of L variables then we define a $L \times L$ matrix α by $\alpha_{\ell\ell'} = \alpha_\ell \delta_{\ell\ell'}$ and a $M \times M$ matrix A by

$$A_{ij} = (e^+ \alpha^{-1} e)_{ij} = \sum_{\ell=1}^L e_{\ell i} e_{\ell j} \alpha_\ell^{-1}.$$

The incidence matrix for a connected graph has rank $M - 1$, satisfying $\sum_{i=1}^M e_{\ell i} = 0$,

i.e. the vector $n = \begin{pmatrix} 1 \\ \vdots \\ 1 \end{pmatrix}$ is mapped to zero by e . Correspondingly A is singular

since $An=0$. Let us denote by $E = \left\{ (p) : \sum_{i=1}^M p_i = 0 \right\}$ the hyperplane orthogonal to n and by e_E resp. A_E the restriction of e resp. A to E . Then A_E is non-singular and we find $A_E = e_E^+ \alpha^{-1} e_E$. If we parametrize E by $\{p_1, \dots, \hat{p}_k, \dots, p_M\}$ —where \hat{p}_k means that p_k is missing—we find

$$(A_E^{-1})_{ij} = A(ki)/A(k) \quad \text{where} \quad A(i_1, \dots, i_r) \quad \text{is the corresponding}$$

minor of A .

The so-called *Symanzik functions* of G , $d(\alpha) = A(k) \prod_{\ell \in \mathcal{L}_G} \alpha_\ell$ and $D_{ij}^k(\alpha) = A(ki)$ · $\prod_{\ell \in \mathcal{L}_G} \alpha_\ell$, can be represented in the form $d(\alpha) = \sum_T \prod_{\ell \notin T} \alpha_\ell$ where the sum runs over all trees T in G , and $D_{ij}^k(\alpha) = \sum_{T_2} \prod_{\ell \notin T_2} \alpha_\ell$ where the sum is running over all 2-trees T_2 in G which separate V_k from V_i and V_j . Note that $d(\alpha)$ and the quadratic form $p^+ A_E^{-1} p = \sum_{i,j \neq k} (D_{ij}^k(\alpha)/d(\alpha)) p_i p_j$ are independent of k (for $p \in E$).

Let G be a connected graph with components G_1, \dots, G_n . A *maximal forest* for G is a maximal set of non-trivial non-overlapping 1PI subgraphs of G . For any maximal forest \mathcal{C} and any $H \in \mathcal{C}$ let $\mathcal{M}(H)$ be the set of all maximal elements of \mathcal{C} properly contained in H and let \bar{H} be $H/\mathcal{M}(H)$. A *labelled forest* is a pair (\mathcal{C}, σ) consisting of a maximal forest \mathcal{C} and a mapping $\sigma: \mathcal{C} \rightarrow \mathcal{L}_G$ such that $\sigma(H) \in \mathcal{L}_{\bar{H}}$ for all $H \in \mathcal{C}$. Note that $\mathcal{M}(H)$ is uniquely determined by H and $\sigma(H)$.

For any labelled forest (\mathcal{C}, σ) we define $\mathcal{D}(\mathcal{C}, \sigma)$ to be the subset of α -space given by

$$\mathcal{D}(\mathcal{C}, \sigma) = \{(\alpha_1, \dots, \alpha_L) : \alpha_\ell \geq 0 \text{ for all } \ell; \quad \alpha_\ell \leq \alpha_{\sigma(H)} \text{ for } \ell \in H \in \mathcal{C}\}.$$

Lemma 3. (a) Any maximal forest \mathcal{C} for G is a disjoint union $\mathcal{C} = \bigoplus_{i=1}^n \mathcal{C}_i$ of maximal forests \mathcal{C}_i for the components G_i of G ; $G_i \in \mathcal{C}_i$;

(b) any maximal forest \mathcal{C} for G can be labelled;

(c) for any 1PI subgraph H of G there is a natural one to one correspondence between labelled forests (\mathcal{C}, σ) for G such that $H \in \mathcal{C}$ and pairs $((\mathcal{C}_1, \sigma_1), (\mathcal{C}_2, \sigma_2))$ of labelled forests for G/H and H ;

(d) any maximal forest for G has exactly h_G elements;

(e) $G - \sigma(\mathcal{C})$ is a tree in G ;

(f) $\bigcup_{(\mathcal{C}, \sigma)} \mathcal{D}(\mathcal{C}, \sigma) = \{(\alpha_\ell) : \alpha_\ell \geq 0 \text{ for all } \ell\};$

(g) for $(\mathcal{C}, \sigma) \neq (\mathcal{C}', \sigma')$, $\mathcal{D}(\mathcal{C}, \sigma) \cap \mathcal{D}(\mathcal{C}', \sigma')$ is a set of Lebesgue measure zero (as subset of \mathbb{R}^L).

Proof. (a) Follows directly from the definition.

(b) Let $H \in \mathcal{C}$ and $\mathcal{M}(H) = \{H_1, \dots, H_m\}$. We first observe that \bar{H} is non-trivial. For $m \leq 1$ this follows directly from the definition of $\mathcal{M}(H)$. For $m \geq 2$ the H_i are pairwise disjoint (no common vertices and no common lines), therefore H contains lines which are in no one H_i . Choose any line of \bar{H} as $\sigma(H)$.

(c) For any (\mathcal{C}, σ) such that $H \in \mathcal{C}$ chose

$$\mathcal{C}_1 = \{H' : H' \in \mathcal{C} \text{ and } H' \cap H = \emptyset; \sigma_1(H') = \sigma(H')\} \quad \text{or}$$

$$H' = H''/H, H \subsetneq H'' \in \mathcal{C}; \sigma_1(H') = \sigma(H'')\} ;$$

$$\mathcal{C}_2 = \{H' \in \mathcal{C} : H' \subset H; \sigma_2(H') = \sigma(H')\} .$$

One easily verifies the one to one correspondence and, furthermore, if (\mathcal{C}, σ) is a labelled forest so are $(\mathcal{C}_1, \sigma_1), (\mathcal{C}_2, \sigma_2)$ and vice versa.

(d) Due to (a) it is sufficient to prove the statement for G 1PI. It trivially holds for $h_G < 2$. Let $h_G = n \geq 2$ and assume the statement holds for all 1PI graphs G' with $h_{G'} < n$. Any \mathcal{C} for G contains at least one genuine subgraph H of G (maximality). From (c) follows the one to one correspondence between \mathcal{C} and $(\mathcal{C}_1, \mathcal{C}_2)$. \mathcal{C}_1 resp. \mathcal{C}_2 have $h_{G/H} = h_G - h_H$ resp. h_H elements by assumption. Therefore, \mathcal{C} has h_G elements.

(e) Let G_1 be one component of G , then $G' = G - \{\sigma(G_1)\}$ is connected and $\mathcal{C} - \{G_1\}$ is a maximal forest for G' . Proceeding this way we finally have removed h_G lines and get a connected graph which must be a tree.

(f) Let $\underline{\alpha} = (\alpha_1, \dots, \alpha_L)$ be a point satisfying $\alpha_\ell \geq 0$ and let us relabel the lines of G such that $\alpha_1 \leq \alpha_2 \dots \leq \alpha_L$. Let G_j be the graph consisting of lines ℓ_1, \dots, ℓ_j and all their endpoints ($G_0 = \emptyset$). For each k ($1 \leq k \leq h_G$) there is a unique $j(k)$ with $h_{G_{j(k)}} = k$, $h_{G_{j(k)-1}} = k-1$. Let H_k be that component of $G_{j(k)}$ which contains line $\ell_{j(k)}$. Then (\mathcal{C}, σ) with $\mathcal{C} = \{H_k, k=1, \dots, h_G\}$ and $\sigma(H_k) = \ell_{j(k)}$ is a labelled forest and $(\alpha_1, \dots, \alpha_L) \in \mathcal{D}(\mathcal{C}, \sigma)$.

(g) Let $(\mathcal{C}, \sigma) \neq (\mathcal{C}', \sigma')$ and $\underline{\alpha} \in \mathcal{D}(\mathcal{C}, \sigma) \cap \mathcal{D}(\mathcal{C}', \sigma')$. We will show that $\alpha_\ell = \alpha_{\ell'}$ for $\ell \neq \ell'$. The set of such $\underline{\alpha}$ has Lebesgue measure zero. Since both \mathcal{C} and \mathcal{C}' contain all components G_i of G there must exist a subgraph $H \in \mathcal{C} \cap \mathcal{C}'$ for which $\sigma(H) \neq \sigma'(H)$ and hence $\alpha_{\sigma(H)} = \alpha_{\sigma'(H)}$.

Let G be a connected graph and \mathcal{C} a family of non-trivial non-overlapping subgraphs of G . The definition of $\mathcal{M}(H)$ and the concept of a labelled family (\mathcal{C}, σ) can obviously be extended to this situation. The elements $H \in \mathcal{C}$ need not be 1PI (not even connected) nor need \mathcal{C} be maximal. We require, however, that $G \in \mathcal{C}$ and that for every $H \in \mathcal{C}$ $H/\mathcal{M}(H)$ consists of a connected graph \tilde{H} plus possibly some trivial graphs not connected to \tilde{H} . Given any \mathcal{C} these requirements can always be satisfied by adding some suitable subgraphs to \mathcal{C} .

To any vertex V_a of \tilde{H} ($H \in \mathcal{C}$) we assign a momentum q_a which is the sum of all those p_i associated with vertices V_i which are mapped to V_a by $H \rightarrow H/\mathcal{M}(H)$. Let $q_H, H \in \mathcal{C}$ be the momenta associated with all but one of the vertices of \tilde{H} . The new momenta $\underline{q} = (q_H, H \in \mathcal{C})$ are obtained from $\underline{p} = (p_i, i=1, \dots, M-1)$ by a linear transformation $R: \underline{q} = R\underline{p}$. In order to express $\tilde{V}(p, \underline{q}, \underline{\alpha})$ in terms of \underline{q} , we decompose $\underline{u} = (u_\ell, \ell \in \mathcal{L}_{\tilde{H}})$ and the transformed incidence matrix eR^+ in a similar way: $\underline{u} = (\underline{u}_H = (u_\ell, \ell \in \mathcal{L}_H), H \in \mathcal{C})$, $eR^+ = (e_{HH'}, H, H' \in \mathcal{C})$. If we introduce in addition scaled quantities $\tilde{q} = (\tilde{q}_H = q_H \zeta_H, H \in \mathcal{C})$, $\tilde{u} = (\tilde{u}_H = u_H / \zeta_H, H \in \mathcal{C})$ and

$$\tilde{e} = (\tilde{e}_{HH'} = e_{HH'} \cdot \zeta_{H'}/\zeta_H, H, H' \in \mathcal{C})$$

we find $V(p, \underline{q}, \underline{\alpha}) = (\tilde{q}^+, \tilde{q}^-) \tilde{M}^{-1} \begin{pmatrix} \tilde{q} \\ \tilde{\underline{q}} \end{pmatrix}$ with $\tilde{M} = \begin{pmatrix} 0 & -2\tilde{\epsilon}^+ \\ -2\tilde{\epsilon} & -4\beta \end{pmatrix}$ where β is the diagonal matrix with elements $\beta_\ell = \alpha_\ell / \zeta_H^2$, $\ell \in \mathcal{L}_{\bar{H}}$, $H \in \mathcal{C}$.

Lemma 4. (a) $\det R = \pm 1$;

(b) eR^+ has the following block structure

$$e_{HH'} \begin{cases} = 0 \text{ if } H \nmid H'; \\ \text{is the incidence matrix } e_{\bar{H}} \text{ for } \bar{H} \text{ if } H = H'; \\ \text{indicates how } H' \text{ is contained in } H \text{ if } H \supsetneq H'; \end{cases}$$

(c) $d(\alpha) = \tilde{d} \prod_{H \in \mathcal{C}} \zeta_H^{2h_{\bar{H}}} = \tilde{d} \prod_{H \in \mathcal{C}} t_H^{2h_H}$ and $\tilde{d} = \det \tilde{M}/(-4)^L \geq \prod_{H \in \mathcal{C}} \tilde{d}_H$ as long as $\beta_\ell \geq 0$

for all $\ell \in \mathcal{L}_G$ where

$$\tilde{d}_H = \det \begin{pmatrix} 0 & -2\tilde{\epsilon}_{HH}^+ \\ -2\tilde{\epsilon}_{HH} & -4\beta_H \end{pmatrix} / (-4)^{|\mathcal{L}_{\bar{H}}|};$$

(d) considered as functions of (t, β) the matrix elements of \tilde{M} are independent of t_G and polynomials in the remaining variables;

(e) if \mathcal{C} is such that $h_{\bar{H}} \leq 1$ for all $H \in \mathcal{C}$, and ζ_H are scaling variables for (\mathcal{C}, σ) , then $\tilde{d} \geq 1$ and therefore \tilde{d}^{-1} and \tilde{M}^{-1} are C^∞ in (t, β) from the domain $\mathcal{D}(\mathcal{C}, \sigma)$. This is true in particular if G is 1PI and (\mathcal{C}, σ) is a labelled forest for G .

Proof. (a) Up to permutations R is a triangular matrix with all diagonal elements $= 1$;

(b) Introducing internal momenta $\underline{k} = (k_\ell, \ell \in \mathcal{L}_G)$ flowing through all lines, momentum conservation at each vertex is expressed by $p + e^+ \underline{k} = 0$. The block structure follows immediately from $q + \text{Re}^+ \underline{k} = 0$;

(c) The factorization follows from a), $|\mathcal{L}_{\bar{H}}| + 1 = |\mathcal{V}_{\bar{H}}| + h_{\bar{H}}$ and the relation between ζ and t . The inequality is due to the fact that every pair of trees for G/H and H corresponds to a unique tree for G and that the contribution of each tree to $d(\alpha)$ is non-negative;

(d) The matrix elements of $\tilde{\epsilon}$ are polynomials due to b) and clearly both $\tilde{\epsilon}$ and β are independent of t_G ;

(e) $\tilde{d}_H = 1$ if $h_{\bar{H}} = 0$ and $\tilde{d}_H = \sum_{\ell \in \mathcal{L}_{\bar{H}}} \beta_\ell$ if $h_{\bar{H}} = 1$. $\beta_{\sigma(H)} = 1$ and therefore $\tilde{d}_H \geq 1$.

In the domain $\mathcal{D}(\mathcal{C}, \sigma)$ $\beta_\ell \leq 1$ for all ℓ and $t_H \leq 1$ for $H \neq G$. The statements follow from these results and (d).

Remark. We may consider a family \mathcal{C}' obtained from \mathcal{C} by omission of one element $H_0 \in \mathcal{M}(H_1)$ and build momenta \underline{q}'_H , $H \in \mathcal{C}'$ for this new family. Obviously there is a choice for the \underline{q}_H and \underline{q}'_H (i.e. choice for the vertex whose momentum is omitted from \underline{q}_H) such that $\underline{q}'_{H_1} = (\underline{q}_{H_1}, \underline{q}_{H_0})$ and $\underline{q}'_H = \underline{q}_H$ for all other $H \in \mathcal{C}'$.

Appendix B

Let G be 1PI, (\mathcal{C}, σ) a labelled forest for G and $H \in \mathcal{C}$ ($H \neq G$). Corresponding to (\mathcal{C}, σ) there is a unique pair of labelled forests $((\mathcal{C}_1, \sigma_1), (\mathcal{C}_2, \sigma_2))$ for G/H and H [Lemma 3 (c)].

We want to relate the contributions of these three labelled forests to \mathcal{T}_G , $\mathcal{T}_{G/H}$ and \mathcal{T}_H respectively. We will prove that the singular part of the t_H -integral [as defined by Eq. (III.5)] in \mathcal{T}_H is a polynomial $P(g_H)$ of degree ω_H in the external momenta q_H of H (a homogeneous polynomial of degree ω_H in q_H , masses and $\sqrt{\epsilon}$ if all spin polynomials Z_ϵ are homogeneous in $-i\partial/\partial u$, masses and $\sqrt{\epsilon}$). On the other hand we will express the singular part of the t_H -integral in \mathcal{T}_G in terms of the same polynomial $P(g_H)$ inserted as vertex function into $\mathcal{T}_{G/H}$.

We will show all this for the regularized amplitudes \mathcal{T} . The results apply as well if we first perform subtractions for some subgraphs $H' \in \mathcal{C}$, $H' \not\supset H$, because the subtracted amplitudes are sums of regularized amplitudes for various reduced graphs, each of them with the above mentioned properties. In particular reducing a subgraph H' to a vertex with vertex function $P(g_{H'})$ which is a polynomial of degree $\omega_{H'}$ does not change the degree of divergence ω_H for any $H \supset H'$.

Let us first consider the t_G -integral in Equation (III.4)

$$\int_0^\infty (dt_G/t_G) t_G^{\omega_H - \omega_G} g_\epsilon(t, \beta, q).$$

Assuming that all Z_ϵ are homogeneous the entire dependence of g_ϵ on t_G , q , masses and ϵ comes from the combinations $t_G q$, $t_G m_\epsilon$ and $t_G \sqrt{\epsilon}$. First of all there is no singular part as long as $\omega_G < 0$. If $\omega_G \geq 0$ the singular part is proportional to $[(\omega_G !)^{-1} (d/dt_G)^{\omega_G} g_\epsilon]_{t_G=0}$ which certainly is a homogeneous polynomial of degree ω_G .

In order to study the singular part of the t_H -integrals we replace the maximal forest \mathcal{C} by a family $\mathcal{C} = \{G, H\}$ and (using the notation of Lemma 4) introduce momenta $q = (q_G, q_H)$ correspondingly. Inserting the polynomial $P(g_H)$ as vertex into the amplitude for G/H amounts to the substitution $U_H: q_H \rightarrow q_H - ie_{GH}^+ \partial/\partial u_G^+$ and acting with the differential operator $U_H(P(g_H))$ on the integrand $I_{G/H}$; the external momenta of H differ from the external momenta of G associated with vertices of H by linear combinations of internal momenta of G/H .

Apart from factors and integrals common to both amplitudes, the expressions for \mathcal{T}_G resp. $\mathcal{T}_{G/H}$ (with the insertion corresponding to H) are

$$\xi_H^{\omega_H} \int_0^1 (dt_H/t_H) t_H^{\omega_H - \omega_H} I_G|_{\tilde{q}_H=0}; I_G = Z_H (-i\partial/\partial \tilde{u}_H) (\det \tilde{M})^{-n/2} \exp iW$$

resp.

$$U_H(\xi_H^{\omega_H} P(g_H)) I_{G/H}; I_{G/H} = (\det \tilde{M}_{G/H})^{-n/2} \exp iW_{G/H}.$$

Using the homogeneity of Z_H and $P(g_H)$ we can compensate the difference in scaling variables used for \mathcal{T}_G and \mathcal{T}_H and find

$$\xi_H^{\omega_H} P(g_H) = \{(\omega_H !)^{-1} (d/dt_H)^{\omega_H} [Z_H (-i\partial/\partial \tilde{u}_H) (\det \tilde{M}_H)^{-n/2} \exp iW_H]_{\tilde{q}_H=0}\}_{t_H=0}$$

with (X stands for either G/H or H) $\tilde{M} = \hat{M} - 2t_H(\hat{e} + \hat{e}^+)$;

$$\hat{M} = \begin{pmatrix} \tilde{M}_{G/H} & 0 \\ 0 & \tilde{M}_H \end{pmatrix}; \quad \tilde{M}_X = \begin{pmatrix} 0 & -2e_X^+ \\ -2e_X^- & -4\beta_X \end{pmatrix}; \quad \hat{e} = \begin{pmatrix} 0 & 0 \\ e_{GH} & 0 \\ 0 & 0 \end{pmatrix};$$

$$W = Q^+ \tilde{M}^{-1} Q - m_{G/H}^2 - m_H^2; \quad W_X = Q_X^+ \tilde{M}_X^{-1} Q_X - m_X^2; \quad Q = \begin{pmatrix} Q_{G/H} \\ Q_H \end{pmatrix}; \quad Q_X = \begin{pmatrix} \tilde{q}_X \\ \tilde{u}_X \end{pmatrix};$$

and

$$m_X^2 = \sum_{\ell \in \mathcal{L}_X} \alpha_\ell (m_\ell^2 - i\varepsilon).$$

Lemma 5. $U_H [\xi_H^{\omega_H} P(q_H)] I_{G/H} = [(\omega_H!)^{-1} (d/dt_H)^{\omega_H} I_G]_{t_H = \tilde{q}_H = 0}.$

Remark. This establishes the required connection between the singular part of the t_H -integrals in \mathcal{T}_G and \mathcal{T}_H . The remaining difference, a factor $\xi_H^{vh_H}$, is the origin of $\sum_m c_m m^{-1} \xi_H^{vm}$ in Equation (III.14), each term ξ_H^{vm} arises from reduced graphs of H with m loops.

Proof. First we note that $(\det \tilde{M})|_{t_H=0} = (\det \tilde{M}_{G/H})(\det \tilde{M}_H)$, $W|_{t_H=0} = W_{G/H} + W_H|_{t_H=0}$ and $\tilde{q}_H = \xi_H t_H q_H$. It is therefore sufficient to show that

$$-i(d/dt_H) \{(\det \tilde{M})^{-n/2} \exp iW\} = U_H (dW_H/dt_H) \{(\det \tilde{M})^{-n/2} \exp iW\}.$$

where (d/dt_H) is taken either for fixed (q_H, u_H) or fixed (q_H, \tilde{u}_H) . This can be done, using $U_H(Q) = Q - it_H \hat{e}^+ \partial/\partial Q^+$ and some properties of the matrix \hat{e} : $\hat{e} \hat{M}^{-1} \hat{e} = 0$, $\text{Tr}(\hat{e} \hat{M}^{-1}) = 0$, $\hat{e}^+ \hat{M}^{-1} dQ/dt_H = 0$. Once again we need $\partial u_{\ell'}/\partial u_\ell = \delta_{\ell\ell'} g_{\mu\mu} = \delta_{\ell\ell'} n$ and it is therefore important that we interpret \underline{u} as an element of the abstract algebra of “ n -dimensional” covariants.

Appendix C. Completion of the Proof of Theorem 2

The problem to be considered is that there may be subgraphs H of G which have two components H_l and H_r , joined by line ℓ . These become 1PI by contraction of line ℓ . The forests used to renormalize \mathcal{T}_G may contain $\{H_1, H_r\}$ whereas the forests for $\mathcal{T}_{G/\ell}$ may contain $\{H\}$, $\{H, H_l\}$ or $\{H, H_r\}$. We shall recursively remove the latter forests from $\mathcal{T}_{G/\ell}$ and replace them by forests (for G) containing $\{H_l, H_r\}$. We show that $\Sigma\Pi(-C_H)$ does not change under these replacements due to the factorization $I_H = I_{H_l} I_{H_r}$ and the identity

$$(-C_H + C_H C_{H_l} + C_H C_{H_r}) = C_{H_l} C_{H_r}. \quad (1)$$

Define

$$\mathcal{H}_K = \{H \subset G : \ell \notin H; H = H_l \cup H_r; V_{i_\ell} \in \mathcal{V}_{H_l}; V_{f_\ell} \in \mathcal{V}_{H_r}; h_H = K; H_l \cap H_r = \emptyset; H_l, H_r \text{ are 1PI}\}.$$

Since there is a one to one correspondence between subgraphs of G/ℓ and those of G not containing line ℓ , we may consider these subgraphs also to be in G/ℓ .

Put $\mathcal{F} = \{\text{forests for } G\}$ which do not contain subgraphs of the type shown in Figure 1},

$$\mathcal{F}^\ell = \{\text{forests for } G/\ell\}, \quad \mathcal{F}_1 = \{F \in \mathcal{F}^\ell : F \ni H \text{ 1PI in } G/\ell\}.$$

\mathcal{F}_{K+1} is constructed in the following way from \mathcal{F}_K : For all $H \in \mathcal{H}_K$ replace all forests which contain either $\{H\}$ or $\{H, H_l\}$ or $\{H, H_r\}$ by corresponding forests which contain $\{H_l, H_r\}$. This procedure gives

$\mathcal{F}_K = \{F \in \mathcal{F} : \text{If } H \in F, \text{ then either } H \text{ is 1PI in } G \text{ or } H \text{ is 1PI in } G/\ell \text{ and } h_H > K; \text{ for any pair } \{H_1, H_2\} \in F \text{ either } H_1 \circ H_2 \text{ in } G/\ell \text{ or } H_1 \cup H_2 \in \bigcup_{K' \leq K} \mathcal{H}_{K'}\}.$

We find $\mathcal{F}_{h_G} = \{F \in \mathcal{F} : \exists H \text{ 1PI in } G\}.$

We prove now the

Proposition 4. *If (1) holds for all $H \in \bigoplus_K \mathcal{H}_K$, then*

$$\sum_{F \in \mathcal{F}_K} \prod_{H \in F} (-C_H) = \sum_{F \in \mathcal{F}_{K+1}} \prod_{H \in F} (-C_H) \quad \text{for } 1 \leq K < h_G. \quad (2)$$

Proof. We observe the following: Let $\{H^1, H^2\} \subset \mathcal{H}_{K+1}$, then

- i) $H^1 \circ H^2$ and, therefore, there is no $F \in \mathcal{F}_K$ with $\{H^1, H^2\} \in F$;
- ii) $H_l^1 \circ H_l^2$ or $H_r^1 \circ H_r^2$ or $\max\{h_{H_l^1 \cup H_l^2}, h_{H_r^1 \cup H_r^2}\} > K+1$ and, therefore, there is no $F \in \mathcal{F}_{K+1}$ with $\{H_l^1, H_r^1, H_l^2, H_r^2\} \subset F$.

That means that we can uniquely decompose ($\oplus = \text{disjoint union}$)

$$\begin{aligned} \mathcal{F}_K &= \bigoplus_{H \in \mathcal{H}_{K+1}} \bigoplus_{F \in \mathcal{F}^H} \{F \oplus \{H\}, F \oplus \{H, H_l\}, F \oplus \{H, H_r\}\} \oplus \hat{\mathcal{F}}_K; \\ \mathcal{F}_{K+1} &= \bigoplus_{H \in \mathcal{H}_{K+1}} \bigoplus_{F \in \mathcal{F}^H} \{F \oplus \{H_l, H_r\}\} \oplus \hat{\mathcal{F}}_K; \end{aligned}$$

where $\hat{\mathcal{F}}_K = \mathcal{F}_K \cap \mathcal{F}_{K+1}$, and

$$\mathcal{F}^H = \{F \in \hat{\mathcal{F}}_K : H' \in F \Rightarrow H' \text{ non-overlapping with } H, H_l, H_r\}.$$

Using this decomposition and Equation (1), Equation (2) is immediately proved. We get

$$\sum_{F \in \mathcal{F}_1} \prod_{H \in F} (-C_H) \mathcal{F}_{G/\ell} = \sum_{F \in \mathcal{F}_{h_G}} \prod_{H \in F} (-C_H) \mathcal{F}_{G/\ell}$$

and this completes the proof of Theorem 2.

References

1. t'Hooft, G., Veltman, M.: Nucl. Phys. B **44**, 189 (1972)
2. Bollini, C.G., Giambiagi, J.J.: Phys. Letters **40B**, 566 (1972)
Cicuta, G.M., Montaldi, E.: Nuovo Cimento Letters **4**, 329 (1972)
3. Becchi, C., Rouet, A., Stora, R.: Broken Symmetries: Perturbation theory of renormalizable models. Lectures given at the International School of Elementary Particle Physics, Basco Polje, 1974
4. t'Hooft, G., Veltman, M.: Diagrammar, CERN Report 73/9
5. Hepp, K.: Renormalization theory in statistical mechanics and quantum field theory (Les Houches 1970), (C. de Witt, R. Stora eds.). New York: Gordon and Breach 1971
6. Speer, E.R.: J. Math. Phys. **15**, 1 (1974)
7. Collins, J.C.: Nucl. Phys. B **92**, 477 (1975)
8. Lam, Y.M.P.: Phys. Rev. D **6**, 2145 (1972)
9. Speer, E.R.: Generalized Feynman amplitudes. Princeton: Princeton University Press 1969

10. Speer, E.R., Westwater, M.J.: Ann. Inst. H. Poincaré **14 A**, 1 (1971)
11. Lowenstein, J.H.: Commun. math. Phys. **24**, 1 (1971)
12. Gelfand, I.M., Shilov, G.E.: Generalized functions, Vol. I. New York: Academic Press 1964
13. Bollini, C.G., Giambiagi, J.J.: Acta Phys. Acustriaca **38**, 211 (1973)
14. Zimmermann, W.: Commun. math. Phys. **15**, 208 (1969)
15. Lowenstein, J.H.: Seminars on Renormalization Theory, Vol. I., Maryland Technical Report 73-068 (1972)
16. Gomez, M., Lowenstein, J.H.: Phys. Rev. D**7**, 550 (1973)
17. Nakanishi, N.: Graph theory and Feynman integrals. New York: Gordon and Breach 1971

Communicated by K. Symanzik

Received July 4, 1975; in revised form April 7, 1976