

Dinuclear Silver(I) and Copper(II) Complexes of Hexadentate Macrocyclic Ligands Containing *p*-Xylyl Spacers

Christine J. McKenzie,^{a,*} Lars Preuss Nielsen^a and Inger Søtofte^b

^aDepartment of Chemistry, Odense University, DK-5230 Odense M, Denmark and ^bDepartment of Chemistry, Technical University of Denmark, DK-2800 Lyngby, Denmark

McKenzie, C. J., Nielsen, L. P. and Søtofte, I., 1998. Dinuclear Silver(I) and Copper(II) Complexes of Hexadentate Macrocyclic Ligands Containing *p*-Xylyl Spacers. – Acta Chem. Scand. 52: 571–577. © Acta Chemica Scandinavica 1998.

The cyclocondensation of terephthalic aldehyde with *N,N*-bis(3-aminopropyl)-methylamine in the presence of silver(I) gives the dinuclear tetraimine Schiff base macrocyclic complex, $[\text{Ag}_2\text{L}^1](\text{NO}_3)_2$ ($\text{L}^1 = 7,22\text{-}N,N'$ -dimethyl-3,7,11,18,22,26-hexaazatricyclo[26.2.2^{1.18}.2^{13.16}]-tetracos-2, 11, 13, 15, 17, 26, 28,30,31,33-decaene). $[\text{Ag}_2\text{L}^1](\text{NO}_3)_2$ crystallizes in the monoclinic space group $P2_1/c$, with $a = 14.153(6)$, $b = 12.263(4)$, $c = 9.220(2)$ Å, $\beta = 97.52(3)$ Å and $Z = 2$. The silver ions are strongly coordinated at each end of the macrocycle by the two imine nitrogen atoms [2.177(3) and 2.182(3) Å] with close interatomic interactions to an oxygen atom of a nitrate ion and an amine nitrogen atom [2.580(2) and 2.690(2) Å]. The Ag...Ag distance is 6.892(3) Å. The free tetraimine macrocycle, L^1 , was obtained by treatment of $[\text{Ag}_2\text{L}^1](\text{NO}_3)_2$ with an excess of iodide, and the reduced hexaaza derivative 7,22-*N,N'*-dimethyl-3,7,11,18,22,26-hexaazatricyclo[26.2.2^{1.18}.2^{13.16}]-tetracos-13,15,28,30,31,33-decaene, L^2 , was prepared by reduction of $[\text{Ag}_2\text{L}^1](\text{NO}_3)_2$ with NaBH_4 . The copper(II) complexes of the reduced ligand, $[\text{Cu}_2\text{Cl}_2\text{L}^2]\text{Cl}_2 \cdot \text{CH}_3\text{OH}$, $[\text{Cu}_2\text{Cl}_2\text{L}^2](\text{PF}_6)_2$ and $[\text{Cu}_2(\text{CH}_3\text{CO}_2)_2\text{L}^2](\text{PF}_6)_2 \cdot 4\text{H}_2\text{O}$, have been prepared and characterized.

The construction of mono-, di- and polynuclear macrocyclic complexes by the cyclocondensation of diamines and dialdehydes using appropriate templating metal ions is a well established area of coordination chemistry.¹ In the case of systems containing more than one metal ion, interest has focused on the study of electronic interactions between the metal ions² or cooperative effects in terms of binding, and possibly even activation,³ of unusual species at the exogenous bridging sites created when two or more metal ions are bound by the macrocycle. An example of small molecule activation by the dicopper complex of a macrocyclic ligand has been reported by Martell and co-workers.⁴ This hexaaza hexadentate macrocycle ligand contains *m*-xylene spacers. In the presence of two equivalents of copper(I) and dioxygen, hydroxylation of one of the *m*-xylene spacers to give a phenolate group occurs. The resultant phenolato oxygen atom bridges between the two copper(II) ions in the product isolated from this reaction. A mechanism involving dioxygen activation by the dicopper(I) site was proposed.

Ligand modification influences the topology the macro-

cyclic complex and, often as a consequence of this, the reactivity of the metal ions. The present work concerns the preparation of the new hexadentate macrocycles related to the *m*-xylene-based ligands of Martell and co-workers,⁴ but instead the backbones contain *p*-xylene spacers. The new 30-membered hexadentate ligands, L^1 and L^2 , are shown below. The present paper describes the preparation of L^1 and L^2 and a preliminary investigation of their coordination chemistry with silver and copper.

The parent non-methylated macrocycle L^3 has been prepared previously by reduction of the product of the 2+2 cyclocondensation of terephthalic aldehyde with bis(3-aminopropyl)amine (Fig. 1).⁵ The initial product of this reaction contains two imine groups and two 1,3-aminal six-membered rings. Formation of the aminal rings is likely to be the driving force for this unusual reaction; the presence of metal templates or high dilution conditions are not necessary. Reduction of this product by LiAlH_4 results in aminal ring opening and Schiff base reduction to give the hexaaza macrocycle L^3 . A synthetic route similar to that in Fig. 1 but using *N,N*-bis(3-aminopropyl)methylamine is not possible for the preparation of L^2 , since formation of aminal rings is pre-

* To whom correspondence should be addressed.




Fig. 1. Synthetic route to the parent of L^2 , the hexaaza macrocycle L^3 ; (i) 0 °C, THF; (ii) 50 °C, $LiAlH_4$, THF, under Ar.

vented by the presence of the central tertiary amine group. The present paper describes an alternative approach of using silver to template the 2+2 cyclocondensation of *N,N*-bis(3-aminopropyl)methylamine and terephthalic aldehyde. This method is advantageous in that the tetraimine macrocycle is produced and can be isolated as a free ligand after silver ion removal. The tetraimine macrocycle, L^1 , is expected to be more suitable than the hexaaza ligands L^2 and L^3 for the study of dinuclear complexes of 'soft' transition metal ions, e.g. dicopper(I) complexes. The reactivity of macrocyclic dicopper(I) complexes towards molecular oxygen is of particular interest.

Experimental

Physical measurements. IR spectra were measured as KBr discs using a Hitachi 270–30 IR spectrometer. UV–visible absorption spectra were recorded on a Shimadzu UV-3100 spectrophotometer. FAB mass spectrometry was carried out on a Kratos MS50TC instrument. NMR

spectra were recorded on a Bruker AC 250. Elemental analyses were carried out by Atlantic Microlab Inc., GA. ESR measurements were carried out on powdered samples on a Bruker ESP-380EFT-EPR spectrometer at r.t.

(7,22-*N,N'*-Dimethyl-3,7,11,18,22,26-hexaazatri-cyclo[26.2.2^{1.18}.2^{13.16}]tetratricosa-2,11,13,15,17,26,28,30,31,33-decaene)disilver(I) dinitrate, $[Ag_2L^1](NO_3)_2$. Terephthalaldehyde (2.68 g, 20 mmol) and *N,N*-bis(3-aminopropyl)methylamine (2.90 g, 20 mmol) were added to a 60 °C solution of $AgNO_3$ (3.40 g, 20 mmol) in MeOH (500 ml). After 2 h the product precipitated as golden-brown crystals, which were isolated by filtration. Yield 5.8 g, 70.2%. *Anal.* Found C, 43.44; H, 5.02; N, 13.48. Calc. for $C_{30}H_{42}Ag_2N_8O_6$: C, 43.60; H, 5.12; N, 13.56. FABMS, m/z : 764 ($[Ag_2L^1(NO_3)]^+$, 40%), 593 ($[AgL^1]^+$, 100). 1H NMR (DMSO- d_6): δ = 1.86 (q, 8 H, $CH_2CH_2CH_2$), 2.21 (s, 6 H, CH_3), 2.50 [t, 8 H, $N(CH_3)CH_2$], 3.83 (t, 8 H, $NHCH_2$), 7.89 (s, 8 H, C_6H_4), 8.55 (s, 4 H, CH). ^{13}C NMR (DMSO- d_6): δ = 28.08 ($CH_2CH_2CH_2$), 40 [CH_3 (hidden in DMSO)], 55.26 [$N(CH_3)CH_2$], 59.14 ($NHCH_2$), 128.14 (phenylene CH), 137.36 (phenylene C), 163.63 (CH). Crystals for X-ray data collection were obtained by recrystallization from methanol/benzene (4:1 v/v).

7,22-*N,N'*-Dimethyl-3,7,11,18,22,26-hexaazatri-cyclo[26.2.2^{1.18}.2^{13.16}]tetratricosa-2,11,13,15,17,26,28,30,31,33-decaene, $L^1 \cdot CH_2Cl_2$. KI (260 mg, 1.56 mmol) was added to a solution of $[Ag_2L^1](NO_3)_2$ (400 mg, 0.484 mmol) in MeOH (200 ml). After 10 min AgI was removed by filtration and the solution was evaporated to dryness. The residue was resuspended in CH_2Cl_2 (30 ml), and the undissolved material was removed by filtration. The solution was evaporated to a brown oil, which was placed under vacuum for 24 h. Yield 114 mg, 48.4% of an oily yellow solid. *Anal.* Found C, 64.74; H, 7.48; N, 14.53. Calc. for $C_{31}H_{44}N_6Cl_2$: C, 65.14; H, 7.76; N, 14.70. FABMS, m/z : 487 (L^1H^+ , 100%). 1H NMR ($CDCl_3$): δ = 1.86 (m, 8 H, $CH_2CH_2CH_2$), 2.23 (s, 6 H, CH_3), 2.43 [m, 8 H, $N(CH_3)CH_2$], 3.63 (t, 8 H, $NHCH_2$), 7.56, 7.72 (2 × s, 8 H, C_6H_4), 8.19, 8.27 (2 × s, 4 H, CH). ^{13}C NMR ($CDCl_3$): δ = 28.31, 28.80 ($CH_2CH_2CH_2$), 42.48, 42.61 (CH_3), 54.49, 55.72 ($N(CH_3)CH_2$), 58.76, 60.06 ($NHCH_2CH_2$), 128.46, 128.55 (phenylene CH), 138.33, 138.43 (phenylene C), 160.83, 161.02 (CH).

7,22-*N,N'*-Dimethyl-3,7,11,18,22,26-hexaazatri-cyclo[26.2.2^{1.18}.2^{13.16}]tetratricosa-13,15,28,30,31,33-decaene, L^2 . $NaBH_4$ (4.5 g, 0.119 mol) was added to a suspension of $[Ag_2L^1](NO_3)_2$ (1.818 g, 2.2 mmol) in DMF (100 ml) and MeOH (50 ml) in three portions over a period of 1 h. The mixture was stirred at r.t. for a further 1 h, after which time the suspension was acidified to pH 4 with 4 M H_2SO_4 and evaporated to dryness. The solid was resuspended in MeOH (400 ml) and undissolved material was removed by filtration. The resultant solution was

evaporated to dryness, redissolved in H₂O (100 ml) and extracted with chloroform (3 × 20 ml). The combined chloroform extracts were dried over anhydrous Na₂SO₄ and evaporated to dryness. The solid was extracted by means of a Soxhlet apparatus with low boiling petroleum. Yield 1.09 g, 58%. *Anal.* Found C, 71.94; H, 10.16; N, 16.79. Calc. for C₃₀H₅₀N₆: C, 72.83; H, 10.19; N, 16.99. FABMS, *m/z*: 495 (L²H⁺, 100%). ¹H NMR (DMSO-*d*₆): δ = 1.63 (k, 8 H, CH₂CH₂CH₂), 2.18 (s, 6 H, CH₃), 2.35 [t, 8 H, N(CH₃)CH₂], 2.62 (t, 8 H, NHCH₂CH₂), 3.67 (s, 8 H, C₆H₄CH₂), 7.23 (s, 8 H, C₆H₄). ¹³C NMR (DMSO-*d*₆): δ = 27.58 (CH₂CH₂CH₂), 42.65 (CH₃), 48.56 (C₆H₄CH₂), 54.12 [N(CH₃)CH₂], 56.6 (NHCH₂CH₂), 128.51 (phenylene CH), 139.57 (phenylene C).

[(7,22-N,N'-Dimethyl-3,7,11,18,22,26-hexaazatricyclo-[26.2.2]^{1.18,2.13.16}]tetratricosa-13,15,28,30,31,33-decaene)-dicopper(II)dichloro]dichloride, [Cu₂Cl₂L²]Cl₂ · CH₃OH. CuCl₂ · 2H₂O (17.8 mg, 104.41 μmol) in MeOH (1 ml) was added to L² (25 mg, 50.6 μmol) in MeOH (2 ml). After 5 min the precipitated turquoise microcrystalline solid was isolated by filtration. Yield 31 mg, 77.3%. *Anal.* Found C, 46.76; H, 6.19; N, 10.72. Calc. for C₃₁H₅₄Cl₄Cu₂N₆O: C, 46.79; H, 6.84; N, 10.56. UV-Vis (DMSO) λ/nm (ε/dm³ mol⁻¹ cm⁻¹): 292 (9390), 703 (370). FABMS, *m/z*: 692 ([Cu₂Cl₂L²]⁺, 12%), 655 ([Cu₂ClL²]⁺, 31), 592 ([Cu₂ClL²]⁺, 68), 557 ([Cu₂L²]⁺, 100).

[(7,22-N,N'-Dimethyl-3,7,11,18,22,26-hexaazatricyclo-[26.2.2]^{1.18,2.13.16}]tetratricosa-13,15,28,30,31,33-decaene)-dicopper(II)dichloro]di-hexafluorophosphate, [Cu₂Cl₂L²](PF₆)₂. NH₄PF₆ (33.0 mg, 202 μmol) in MeOH (2 ml) was added to a solution of CuCl₂ · 2H₂O (17.2 mg, 101 μmol) and L² (25 mg, 51 μmol) in MeOH (20 ml). The turquoise microcrystals were isolated by filtration and washed with MeOH (1 ml). Yield 41 mg, 83%. *Anal.* Found C, 37.11; H, 5.22; N, 8.54. Calc. for C₃₀H₅₀Cl₂Cu₂F₁₂N₆P₂: C, 36.67; H, 5.13; N, 8.55. UV-Vis (DMSO) λ/nm (ε/dm³ mol⁻¹ cm⁻¹): 295 (8810), 700 (410). FABMS, *m/z*: 655 ([Cu₂ClL²]⁺, 30%), 592 ([Cu₂ClL²]⁺, 5), 557 ([Cu₂L²]⁺, 100).

[(7,22-N,N'-Dimethyl-3,7,11,18,22,26-hexaazatricyclo-[26.2.2]^{1.18,2.13.16}]tetratricosa-13,15,28,30,31,33-decaene)-dicopper(II)diacetato]di-hexafluorophosphate, [Cu₂(CH₃CO₂)₂L²](PF₆)₂ · 4H₂O. NH₄PF₆ (81 mg, 497 μmol) was added to a solution of L² (50 mg, 101 μmol) and Cu(CH₃CO₂)₂ · 2H₂O (55.4 mg, 277 μmol) in MeOH (4 ml) and water (0.5 ml). After standing overnight the blue crystals were isolated by filtration. Yield 74.7 mg, 67%. *Anal.* Found C, 37.16; H, 5.30; N, 7.48. Calc. for C₃₄H₆₄Cu₂F₁₂N₆O₈ P₂: C, 37.06; H, 5.85; N, 7.48. UV-Vis (DMSO) λ/nm (ε/dm³ mol⁻¹ cm⁻¹): 292 (8850), 699 (440). FABMS, *m/z*: 825 ([Cu₂(CH₃CO₂)₂L²(PF₆)₂]⁺, 8%), 765 ([Cu₂(CH₃CO₂)₂L²(PF₆)₂H₂]⁺, 100), 679 ([Cu₂(CH₃CO₂)₂L²]⁺, 5), 617 ([Cu₂(CH₃CO₂)₂L²H]⁺, 57), 557 ([Cu₂L²]⁺, 82).

X-Ray techniques. Crystal and experimental data for [Ag₂L¹](NO₃)₂ are listed in Table 1. The crystal was cooled to 120 K using the Cryostream nitrogen gas cooler system.⁶ Four standard reflections were measured for intensity and orientation control after every 4 h. No decay in intensities was observed. The intensities were corrected for Lorentz and polarization effects. Absorption correction was carried out by an empirical method,⁷ in which the crystal shape is approximated by an ellipsoid and the size (in units of μm⁻¹) and orientation are treated as parameters. The refinement of the parameters is based on ψ-scans on 4 reflections and symmetry equivalents, which amounts to 145 observations. The structure was solved by the Patterson method

Table 1. Crystallographic data and details of structural determination for [Ag₂L¹](NO₃)₂.

Formula	C ₃₀ H ₄₂ N ₆ O ₆ Ag ₂
Formula weight	826.46
Crystal system	Monoclinic
Space group	P2 ₁ /c
Unit-cell dimensions:	
<i>a</i> /Å	14.153(6)
<i>b</i> /Å	12.263(4)
<i>c</i> /Å	9.220(2)
β/°	97.52(3)
Unit-cell volume, <i>V</i> /Å ³	1586.4(9)
Formula units per unit cell, <i>Z</i>	2
<i>F</i> (000)	840
Calculated density <i>D</i> _x /g cm ⁻³	1.73
Radiation	Mo Kα
Wavelength, λ/Å	0.71073
Linear absorption coefficient/mm ⁻¹	1.29
Temperature, <i>T</i> /K	120
Crystal description	Golden-brown
Crystal size/mm	0.14 × 0.14 × 0.08
Diffractometer	Enraf-Nonius CAD-4F
Unit-cell determination:	
No. of reflections used	25
θ-Range/°	8.0–12.8
Intensity data collection:	
θ _{max} /°	30
Range of <i>h</i>	–19–0
Range of <i>k</i>	–17–0
Range of <i>l</i>	–12–12
Scan mode	ω
Scan range, Δω	2.15 + 0.35 tan θ
Total number of unique reflections	4614
No. of independent reflections, <i>I</i> > 2σ(<i>I</i>)	3646
Corrections	Lorentz-polarization absorption
Transmission factors	0.8471–0.7952
Structure refinement:	
Minimization of	Σ <i>w</i> (<i>F</i> _o ² – <i>F</i> _c ²) ²
Anisotropic thermal parameters	All non-hydrogen atoms
Isotropic thermal parameters	Hydrogen atoms
No. of refined parameters	292
Weighting scheme	[σ ² (<i>F</i> _o ²) + (0.0198 <i>P</i>) ² + 2.28 <i>P</i>] ⁻¹ , <i>P</i> = (<i>F</i> _o ² + 2 <i>F</i> _c ²)/3
<i>R</i> = Σ <i>F</i> _o – <i>F</i> _c /Σ <i>F</i> _o	0.0367 (3646 reflections)
<i>wR</i> 2 = [Σ <i>w</i> <i>F</i> _o ² – <i>F</i> _c ² ² /Σ <i>wF</i> _o ⁴] ^{1/2}	0.0699 (4614 reflections)
<i>S</i> = [Σ <i>w</i> (<i>F</i> _o ² – <i>F</i> _c ²) ² /(<i>N</i> _{obs} – <i>N</i> _{var})] ^{1/2}	1.10
Final (Δ/σ) _{max}	0.12
Final Δρ _{min} and Δρ _{max} /e Å ⁻³	–1.09 and 0.67

with partial structure expansion and refined by a full-matrix least-squares technique. All non-hydrogen atoms were refined anisotropically. The hydrogen atoms could all be located in the difference maps and were refined with fixed thermal parameters $U(\text{H}) = 1.2U$ for attached atoms. The crystallographic computations were performed with SHELXS86⁸ and SHELXL93.⁹ The atomic scattering factors were taken from the literature.¹⁰ The SHELXTL program¹¹ was used for illustration and PLATON¹² for molecular geometry calculations.

The final positional parameters are listed in Table 2. Anisotropic thermal parameters, positional parameters for the hydrogen atoms and a list of observed and calculated structure factors may be obtained from one of the authors (I.S.) on request.

Results and discussion

Syntheses. The 2+2 Schiff base cyclocondensation of 1,4-terephthalaldehyde and *N,N*-bis(3-aminopropyl)-methylamine in the absence of silver leads to intractable sticky products. The mass spectra of these materials show peaks due to several of the expected oligomeric products. By contrast, if the reaction is carried out in the presence of silver nitrate, a disilver complex of the tetra-Schiff base macrocycle L^1 , $[\text{Ag}_2L^1](\text{NO}_3)_2$, is isolated in a 70% yield. $[\text{Ag}_2L^1](\text{NO}_3)_2$ was used as starting material for preparation of the free ligand 7,22-*N,N'*-dimethyl-3,7,11,18,22,26-hexaazatricyclo[26.2.2^{1,18}.2^{13,16}]tetratricosa-2,11,13,15,17,26,28,30,31,33-decaene, L^1 , and the reduced macrocyclic ligand dimethyl-3,7,

11,18,22,26-hexaazatricyclo[26.2.2^{1,18}.2^{13,16}]tetratricosa-13,15,28,30,31,33-decaene, L^2 . The synthetic route to the disilver complex and the two macrocycles, L^1 and L^2 , derived from this complex is depicted in Fig. 2. Isolation of the free ligand was accomplished by the removal of the silver ions in $[\text{Ag}_2L^1](\text{NO}_3)_2$ using excess iodide ions. The doubling up of signals in the NMR spectrum of L^1 compared to the NMR spectrum of $[\text{Ag}_2L^1](\text{NO}_3)_2$ suggest the presence of at least two geometrical isomers of the free Schiff base, probably due to inversion about one or more of the imine nitrogen atoms. The reduced macrocycle, L^2 , was isolated after reduction of the silver complex by NaBH_4 in methanol. In this reaction the four imine groups and the silver(I) are reduced to amines and metallic silver, respectively. As a result the silver is effectively removed from the macrocycle. Dinuclear copper(II) complexes were prepared and characterised using L^2 , *vide infra*.

Silver complex. *X-Ray crystal structure of $[\text{Ag}_2L^1](\text{NO}_3)_2$.* Selected bond distances and angles are summarized in Table 3. The structure of $[\text{Ag}_2L^1](\text{NO}_3)_2$ is shown in Fig. 3. The cation contains a crystallographic centre of symmetry. The silver ions are coordinated to two imine nitrogen atoms at each end of the macrocycle, the Ag–N(1) and the Ag–N(3) bonds being 2.177(3) and 2.182(3) Å, respectively. There are close interactions of each silver ion to the oxygen atom of a nitrate anion [Ag–O(2) = 2.580(2) Å], and to the tertiary amine nitrogen atoms [Ag–N(2) = 2.690(2) Å]. The Ag atoms are situated 0.63 Å above the plane of the *p*-xylyl rings. The two planes of these rings are parallel with a separation of 1.9 Å between the planes. The shortest distance between the two xylyl rings is C(13)⋯C(14) [1–*x*, –*y*, –*z*] at a distance of 3.892(4) Å. The Ag⋯Ag separation is 6.892(3) Å.

It is interesting to compare the structure of the present complex $[\text{L}^1\text{Ag}_2](\text{NO}_3)_2$ with that of a dicopper complex of the non-methylated saturated macrocyclic analogue, L^3 .¹³ The interatomic metal–metal separation in $[\text{Ag}_2L^1](\text{NO}_3)_2$ is significantly smaller than that measured for $[\text{Cu}_2(\text{CH}_3\text{CO}_2)_2L^3](\text{ClO}_4)_2$ (8.40 Å).¹³ The most significant chemical difference in the two macrocycles is the presence of four secondary amine groups in L^3 as opposed to the four imine groups in L^1 . This apparently has a profound effect on the topology of the two complexes. π -Conjugation of the xylyl groups with the *para* imine groups imposes their coplanarity in the structure of $[\text{Ag}_2L^1](\text{NO}_3)_2$, whereas the presence of amine groups in the same position in the case of L^3 gives more flexibility due to the fact that rotation about the four aryl–CH₂ bonds is possible. Hence, in the structure of $[\text{Cu}_2(\text{CH}_3\text{CO}_2)_2L^3](\text{ClO}_4)_2$ a cleft is created in which the benzene rings are stacked with an interplanar distance of 3.8 Å on opposite sides of the cleft. This is an essentially 90° twist in the orientation of the benzene groups in the dicopper complex of L^3 compared to the disilver complex of L^1 . A schematic depiction of the

Table 2. Fractional atomic coordinates and equivalent isotropic thermal parameters (in Å²) for $[\text{Ag}_2L^1](\text{NO}_3)_2$.

Atom	<i>x</i>	<i>y</i>	<i>z</i>	U_{eq}^a
Ag	0.30655(2)	0.01152(2)	0.19086(2)	0.0153(1)
N(1)	0.32431(16)	–0.15071(19)	0.2864(3)	0.0135(6)
N(2)	0.12575(14)	0.0026(2)	0.2510(2)	0.0144(6)
N(3)	0.28444(16)	0.18673(19)	0.1630(3)	0.0139(6)
C(1)	0.2540(2)	–0.1691(2)	0.3896(3)	0.0157(7)
C(2)	0.1551(2)	–0.1952(2)	0.3106(3)	0.0178(8)
C(3)	0.1125(2)	–0.1104(2)	0.1990(3)	0.0171(8)
C(4)	0.0767(2)	0.0194(3)	0.3804(3)	0.0218(8)
C(5)	0.0880(2)	0.0776(2)	0.1329(3)	0.0159(8)
C(6)	0.1089(2)	0.1973(3)	0.1704(4)	0.0196(8)
C(7)	0.2090(2)	0.2218(2)	0.2488(3)	0.0176(8)
C(8)	0.32740(19)	0.2624(2)	0.1033(3)	0.0132(7)
C(9)	0.40315(19)	0.2475(2)	0.0104(3)	0.0126(7)
C(10)	0.4477(2)	0.3418(2)	–0.0328(3)	0.0151(7)
C(11)	0.5188(2)	0.3351(2)	–0.1226(3)	0.0146(7)
C(12)	0.54825(19)	0.2332(2)	–0.1689(3)	0.0133(7)
C(13)	0.50257(19)	0.1397(2)	–0.1273(3)	0.0140(7)
C(14)	0.43088(19)	0.1461(2)	–0.0393(3)	0.0140(7)
C(15)	0.37711(19)	–0.2324(2)	0.2656(3)	0.0138(7)
N	0.24436(17)	–0.02385(19)	–0.1922(3)	0.0176(7)
O(1)	0.29477(17)	–0.01533(18)	–0.2933(2)	0.0300(7)
O(2)	0.27965(17)	–0.06464(17)	–0.0722(2)	0.0242(6)
O(3)	0.16022	0.0077(2)	–0.2098(3)	0.0380(8)

$$^a U_{\text{eq}} = \frac{1}{3} \sum_i \sum_j U_{ij} a_i^* a_j^* a_i \cdot a_j.$$




Fig. 2. Synthetic route to the dinuclear silver complex $[Ag_2L^1](NO_3)_2$, the free Schiff base ligand L^1 and its reduced derivative L^2 . (i) MeOH, 60 °C, 2 h; (ii) KI, MeOH and (iii) $NaBH_4$, MeOH/DMF.

Table 3. Selected bond distances (in Å) and angles (in °) for $[Ag_2L^1](NO_3)_2$.

Ag–N(1)	2.177(2)	Ag–N(3)	2.182(2)
Ag···O(2)	2.580(2)	Ag···N(2)	2.691(2)
N(1)–C(15) ⁱ	1.280(3)	N(1)–C(1)	1.481(3)
N(2)–C(4)	1.471(3)	N(2)–C(3)	1.470(4)
N(3)–C(7)	1.474(3)	N(2)–C(5)	1.472(4)
C(1)–C(2)	1.526(4)	N(3)–C(8)	1.274(4)
C(5)–C(6)	1.528(4)	C(2)–C(3)	1.529(4)
C(8)–C(9)	1.468(4)	C(6)–C(7)	1.533(4)
C(9)–C(14)	1.398(4)	C(9)–C(10)	1.400(4)
C(11)–C(12)	1.402(4)	C(10)–C(11)	1.387(4)
C(12)–C(15)	1.468(4)	C(12)–C(13)	1.395(4)
O(1)–N	1.250(3)	C(13)–C(14)	1.382(4)
O(3)–N	1.242(3)	O(2)–N	1.257(3)
N(1)–Ag–N(3)	162.7(8)	Ag–N(1)–C(15) ⁱ	134.0(2)
Ag–N(1)–C(1)	110.3	C(3)–N(2)–C(4)	110.0(2)
C(1)–N(1)–C(15) ⁱ	115.6(2)	C(4)–N(2)–C(5)	110.4(2)
C(3)–N(2)–C(5)	109.3(2)	Ag–N(3)–C(8)	134.4(2)
Ag–N(3)–C(7)	109.1(2)	N(1)–C(1)–C(2)	112.1(2)
C(7)–N(3)–C(8)	116.1(2)	N(2)–C(3)–C(2)	113.5(2)
C(1)–C(2)–C(3)	115.8(2)	C(5)–C(6)–C(7)	115.8(3)
N(2)–C(5)–C(6)	113.1(2)	N(3)–C(8)–C(9)	126.0(2)
N(3)–C(7)–C(6)	112.3(2)	C(8)–C(9)–C(14)	124.1(2)
C(8)–C(9)–C(10)	116.9(2)	C(9)–C(10)–C(11)	120.7(2)
C(10)–C(9)–C(14)	119.0(2)	C(11)–C(12)–C(13)	118.8(2)
C(10)–C(11)–C(12)	120.1(2)	C(13)–C(12)–C(15)	124.0(2)
C(11)–C(12)–C(15)	117.1(2)	C(9)–C(14)–C(13)	120.2(2)
C(12)–C(13)–C(14)	121.1(2)	O(1)–N–O(2)	119.3(3)
C(12)–C(15)–N(1) ⁱ	125.2(2)	O(2)–N–O(3)	120.1(3)
O(1)–N–O(3)	120.5(3)		

Symmetry code: ⁱ1 – x, – y, – z.




Fig. 3. The structural arrangement in $[Ag_2L^1](NO_3)_2$ together with atomic numbering scheme. The interaction between silver ions and the closest nitrate oxygen atom is represented by the dashed lines.

orientations of the xyllyl groups in these two complexes is presented in Fig. 4.

Copper complexes. The reaction of the reduced macrocycle L^2 with copper chloride in the absence and presence of PF_6^- in methanolic solution gave the turquoise-coloured compounds $[Cu_2Cl_2L^2]Cl_2 \cdot CH_3OH$ and $[Cu_2Cl_2L^2](PF_6)_2$, respectively. Comparison of IR, UV–Vis, ESR and mass spectra indicates the cations in each of these salts are identical in solution. The blue compound $[Cu_2(CH_3CO_2)_2L^2](PF_6)_2 \cdot 4H_2O$ was obtained from the reaction of L^2 with copper acetate in methanol. The X-ray crystal structure of the related non-




Fig. 4. Schematic representations of the orientations of the two *p*-xylyl groups in (a) $[\text{Ag}_2\text{L}^1](\text{NO}_3)_2$ and (b) $[\text{Cu}_2(\text{CH}_3\text{CO}_2)_2\text{L}^3](\text{ClO}_4)_2$.¹³

methylated complex $[\text{Cu}_2(\text{CH}_3\text{CO}_2)_2\text{L}^3](\text{ClO}_4)_2$ shows that each acetate group was bound to one copper atom in a bidentate fashion.¹³ Since the IR spectra of $[\text{Cu}_2(\text{CH}_3\text{CO}_2)_2\text{L}^2](\text{PF}_6)_2 \cdot 4\text{H}_2\text{O}$ and $[\text{Cu}_2(\text{CH}_3\text{CO}_2)_2\text{L}^3](\text{PF}_6)_2$ are identical, apart from the expected differences due to different counter-anions and the methyl groups, we assume a similar bidentate acetate coordination with the acetate bands at $\nu_{\text{OCO}}(\text{asym})$, 1567 cm^{-1} , and $\nu_{\text{OCO}}(\text{sym})$, 1429 cm^{-1} , in the IR spectrum.

The ESR spectra of the copper(II) complexes reveal that there is no significant exchange coupling between the d^9 copper ions. The absence of magnetic interaction is not surprising in view of the lack of suitable exchange pathways,¹⁴ since the spectroscopic evidence suggests that neither the chloride ions, acetate ions or possible coordinated solvent act as bridging ligands between the two encapsulated copper ions; the only bridging element between the copper ions is the macrocycle framework which is not expected to provide a magnetic exchange pathway. The ESR spectra of solid samples of $[\text{Cu}_2\text{Cl}_2\text{L}^2]\text{Cl}_2 \cdot \text{CH}_3\text{OH}$ and $[\text{Cu}_2\text{Cl}_2\text{L}^2](\text{PF}_6)_2$ are different; $[\text{Cu}_2\text{Cl}_2\text{L}^2]\text{Cl}_2 \cdot \text{CH}_3\text{OH}$ affords an axial signal with $g_{\perp} = 2.208$ and $g_{\parallel} = 2.078$, while $[\text{Cu}_2\text{Cl}_2\text{L}^2](\text{PF}_6)_2$ shows a rhombic signal with $g_x = 2.046$, $g_y = 2.143$ and $g_z = 2.179$. The observation of a difference in the ESR spectra of these compounds which we propose contain identical cations is likely to be related to their solid state structure. An interaction of the chloride counter-anions with the copper ions may be present in the solid state, for example, by μ -chloro bridging between adjacent dinuclear cations and thus giving rise to pseudo-five-coordinated copper ions. The possibility of counter-anion interaction with the copper centres is removed in the case of the hexafluorophosphate complex. The acetate complex $[\text{Cu}_2(\text{CH}_3\text{CO}_2)_2\text{L}^2](\text{PF}_6)_2 \cdot 4\text{H}_2\text{O}$ gives rise to an axial signal with $g_{\perp} = 2.090$ and $g_{\parallel} = 2.191$.

Concluding remarks

The silver templated reaction reported here affords a simple one-pot preparation of the tetra Schiff base macrocycle L^1 . The isolation of the free L^1 by iodide assisted removal of silver represents a new development in the chemistry of Schiff base macrocycles and offers the

possibility of preparing other metal complexes of L^1 . We have carried out a preliminary investigation into the copper(I) coordination chemistry of L^1 ; however, as yet we have not characterized any products; the reaction of $[\text{Cu}(\text{CH}_3\text{CN})_4](\text{BF}_4)$ with L^1 in an inert atmosphere results in formation of yellow solutions from which we are unable to isolate solid products. On exposure to air these solutions turn green within seconds, indicating oxidation of the copper(I) to copper(II). Elaboration of this chemistry will be the subject of ongoing work. Apart from pursuing the study of copper(I) complexes of L^1 and their possible reactivity towards molecular oxygen, these macrocycles offer the possibility of further modification. Future work will involve the derivatization of the secondary nitrogen atoms of L^2 with metal binding groups to give bis-pentadentate macrocyclic ligands for the study of dinuclear complexes of transition metal ions with a preference for higher coordination numbers.

Acknowledgements. We thank ERASMUS student Valérie Gandubert for help in the initial stages of the synthetic work. Support from the Danish Natural Science Council (grant no. 9503162 to C.J.M.) is acknowledged. Jens Z. Pedersen is thanked for recording the ESR spectra.

References

1. L. F. Lindoy, *The Chemistry of Macrocyclic Ligand Complexes*. Cambridge University Press, Cambridge 1989. See, for example: (a) Harding, C., McDowell, D., Nelson, J., Raghunathan, S., Stevenson, C., Drew, M. G. B. and Yates, P. C. *J. Chem. Soc., Dalton Trans.* (1990) 2524; (b) Abid, K. K., Fenton, D. E., Casellato, U., Vigato, P. A. and Graziani, R. *J. Chem. Soc., Dalton Trans.* (1984) 351; (c) Hoskins, B. F., Robson, R. and Smith, P. *J. Chem. Soc., Chem Commun.* (1990) 488; (d) Tandon, S. S., Thompson, L. K. and Brisdon, J. N. *Inorg. Chem.* 34 (1995) 5507; (e) Brooker, S., McKee, V., Shepard, W. B. and Pannell, L. K. *J. Chem. Soc., Dalton Trans.* (1987) 2555; (f) Nanda, K. K., Venkatsubramanian, K., Majumdar, D. and Nag, K. *Inorg. Chem.* 33 (1994) 1581; (g) Adams, H., Bailey, N. A., Bertrand, P., Collinson, S. R., Fenton, D. E. and Kitchen, S. J. *Inorg. Chim. Acta*, 250 (1996) 139; (h) Collinson, R. and Fenton, D. E. *Coord. Chem. Rev.* 148 (1996) 19; (i) Salata, C. A., Youinou, M.-T. and Burrows, C. J. *J. Am. Chem. Soc.* 111 (1989) 9278. (j) Fenton, D. E., Matthews, R. W., McPartlin, M., Murphy, B. P., Scowen I. J. and Tasker, P. A. *J. Chem. Soc., Dalton Trans.* 16 (1996) 3421.
2. (a) Drew, M. G. B., Harding, C. J., Howarth, O. W., Lu, Q., Marrs, D. J., Morgan, G. G., McKee, V. and Nelson, J. *J. Chem. Soc., Dalton Trans.* (1996) 3021; (b) Drew, M. G. B., Yates, P. C., Esho, F. S., Trocha-Grimshaw, J., Lavary, A., McKillop, K. P. Nelson S. M. and Nelson, J., *J. Chem. Soc., Dalton Trans.* (1988) 2995; (c) Bailey, N. A., Fenton, D. E., Moody, R., Rodriguez de Barbarin, C. O., Sciambarella, I. N., Latour, J.-M., Limosin, D. and McKee, V. *J. Chem. Soc., Dalton Trans.* (1987) 2519.
3. (a) Nelson, S. M., Esho, F., Lavary, A. and Drew, M. G. B. *J. Am. Chem. Soc.* 105 (1983) 5693; (b) Sharma, P. and Vige, G. S. *Inorg. Chim. Acta* 88 (1984) 29; (c) Moore, K. and Vige, G. S. *Inorg. Chim. Acta* 66 (1982) 125; (d) Curtis, N. J., Hagen K. S. and Sargeson, A. M. *J. Chem. Soc.*,

- Chem. Commun.* (1984) 1571; (e) Agnus, Y., Gisselbrecht, J. P., Louis, R. and Metz, B. *J. Am. Chem. Soc.* 111 (1989) 1494; (f) Drew, M. G. B., Yates, P. C., Trocha-Grimshaw, J., McKillop K. P. and Nelson, S. M. *J. Chem. Soc., Chem Commun.* (1985) 262.
4. Menif, N., Martell, A. E., Squattrito, P. J. and Clearfield, A. *Inorg. Chem.* 29 (1990) 4723.
 5. Pietraszkiewicz, M. and Gasiorowski, R. *Chem. Ber.* 123 (1990) 405.
 6. Cosier, J. and Glazer, A. M. *J. Appl. Crystallogr.* 19 (1986) 105.
 7. Nielsen, K. *To be published.*
 8. Sheldrick, G. M. *Acta Crystallogr., Sect. A* 46 (1990) 467.
 9. Sheldrick, G. M. *SHELXL93. Program for Crystal Structure Refinement.* University of Göttingen, Germany (1993).
 10. *International Tables for X-Ray Crystallography.* Kynoch Press, Birmingham 1974, Vol. IV (Present distributor: Kluwer Academic Publishers, Dordrecht).
 11. Sheldrick, G. M. *SHELXTL95.* Siemens Analytical X-ray Instruments Inc., Madison, WI 1995.
 12. Spek, A. L. *Acta Crystallogr., Sect. A* 46 (1990) C-34.
 13. McKenzie, C. J., Toftlund, H., Pietraszkiewicz, M., Stojek, Zb. and Slowinski, K. *Inorg. Chim. Acta* 210 (1993) 143.
 14. D. N. Hendrickson. In Willett, R. D., Gatteschi, D. and Hahn, O., Eds., *Magneto-Structural Correlations in Exchange Coupled Systems*, Reidel, Dordrecht 1985, p. 523.

Received August 14, 1997.