

# Direct and indirect effects of alien insect herbivores on ecological processes and interactions in forests of eastern North America

Kamal J. K. Gandhi · Daniel A. Herms

Received: 28 January 2009 / Accepted: 20 August 2009 / Published online: 6 November 2009  
© Springer Science+Business Media B.V. 2009

**Abstract** Alien invasive insects such as gypsy moth, hemlock woolly adelgid, and emerald ash borer continue to disturb the mixed deciduous and hemlock forests of eastern North America by causing wide-scale defoliation, decline and/or mortality of their hosts. Some of the most devastating species are spreading in “defense free space”, causing extensive mortality of hosts that are inherently susceptible, perhaps due to their lack of coevolutionary history with the invader. These disturbances have altered the dynamics of canopy gaps, coarse woody debris, biogeochemical cycling, and ecological interactions among organisms in terrestrial and aquatic systems, with consequent effects on forest composition, structure, and function. Populations of indigenous species specialized to particular habitats and/or host trees are most likely to decrease, while some generalist and opportunistic species may increase in invaded forests, including exotic plants as their facilitation by alien insects sparks an “invasional meltdown”. Although poorly documented, alien insects may induce positive

feedback effects on ecological processes and interactions. For example, effects of herbivory on foliar chemistry may indirectly alter tri-trophic interactions of indigenous herbivores on their shared hosts, slow rates of terrestrial nutrient cycling, and decrease productivity of aquatic habitats based on allochthonous inputs. Tactics used to eradicate or suppress alien insects in forests such as insecticide applications, biological control, and silvicultural prescriptions can also have ecological impacts. As alien insects continue to establish and spread in forests of eastern North America, their already pervasive effects on ecological interactions and ecosystem processes will continue to magnify.

**Keywords** Invasive species · Invasional meltdown · Defense free space · Disturbance · Nutrient cycling · Gap ecology · Forest succession · Nontarget effects

## Introduction

During the last two centuries, the number of alien phytophagous insect species that have established in forests of North America has increased exponentially (Mattson et al. 1994; Liebhold et al. 1995; Niemelä and Mattson 1996). Some have been especially devastating, either alone or in conjunction with pathogenic symbionts, causing wide-spread mortality of their hosts that has substantially altered forest

---

K. J. K. Gandhi  
Daniel B. Warnell School of Forestry and Natural Resources, The University of Georgia, Athens, GA 30602, USA

D. A. Herms (✉)  
Department of Entomology, The Ohio State University, Ohio Agricultural Research and Development Center, 1680 Madison Avenue, Wooster, OH 44691, USA  
e-mail: herms.2@osu.edu

structure and ecosystem processes. For example, gypsy moth (*Lymantria dispar* L.) and beech scale (*Cryptococcus fagisuga* Lind.) with its pathogenic fungal associates have severely impacted oak (*Quercus* spp.) (Davidson et al. 1999) and beech (*Fagus grandifolia* Ehrh.) (Houston 1994; Morin et al. 2007) stands, respectively, throughout New England. Smaller European elm bark beetle (*Scolytus multistriatus* Marsh.) historically has been the most important vector of Dutch elm disease (Hanula and Berisford 1984), which decimated American elm (*Ulmus americana* L.) throughout North America (Barnes 1976; Karnosky 1979). Hemlock woolly adelgid (*Adelges tsugae* Annand) has caused wide-spread mortality of eastern [*Tsuga canadensis* (L.) Carr.] and Carolina hemlock (*Tsuga caroliniana* Engelm.), which are keystone species in riparian areas in the Appalachian forests (Orwig et al. 2002; Ellison et al. 2005; Small et al. 2005). Balsam woolly adelgid [*Adelges piceae* (Ratzeburg)] has devastated Fraser fir [*Abies fraseri* (Pursh) Poir.] in high altitude forests of the southern Appalachian Mountains (Pauley and Clebsch 1990; Hollingsworth and Hain 1991). More recent and still relatively localized introductions causing wide-spread mortality of their hosts include emerald ash borer (*Agrilus planipennis* Fairmaire) (Cappaert et al. 2005; Poland and McCullough 2006), viburnum leaf beetle [*Pyrrhalta viburni* (Paykull)] (Weston et al. 2007), and redbay ambrosia beetle (*Xyleborus glabratus* Eichhoff) and its fungal symbiont (*Raffaella* sp.) (Hanula et al. 2008). Other emerging threats that have the potential to impart severe impacts on forests of eastern North America include Asian longhorned beetle [*Anoplophora glabripennis* (Motschulsky)] (Townsend Peterson and Sachetti-Pereira 2004), and European woodwasp (*Sirex noctilio* Fabricius) (Ciesla 2003; Haugen and Hoebeke 2005).

Release from natural enemies has been considered a key factor in facilitating success of alien insects as they establish and spread readily in “enemy free space” (Liebhold et al. 1995). However, if indigenous host plants are highly susceptible to alien insects because their lack of coevolutionary history with the herbivore has left them without effective targeted defenses, then release from bottom-up regulation also may be critical in the establishment, spread, and ultimate impact of exotic herbivores. Wide-spread tree mortality can result as range expansion of alien insects occurs in “defense free space”. This appears

to be the case with some of the most devastating alien invaders, which threaten to functionally extirpate their host plants. For example, there is experimental evidence that North American hosts naïve to the alien insect are more susceptible than are their coevolved congeners in the case of balsam woolly adelgid and North American firs (Witter and Ragenovich 1986), beech scale and North American beeches (Houston 1987), hemlock woolly adelgid and eastern North American hemlocks (Havill et al. 2006), and emerald ash borer and North American ashes (*Fraxinus* spp.) (Rebek et al. 2008).

To date, more than 400 alien phytophagous insect species have established in North American forests, and nearly every genera of woody plant endemic to eastern North America has at least one alien insect and/or pathogen associated with it (Mattson et al. 1994, 2007). About 75% of the alien herbivorous insect fauna currently established in North American forests is endemic to Europe (Niemelä and Mattson 1996; Mattson et al. 2007). Pathways of introduction have been unintentional such as infested wood packing or soil and plant material (Work et al. 2005; Haack 2006; McCullough et al. 2006), and intentional as biocontrol agents (Elkinton and Liebhold 1990). As “biological pollutants” and anthropogenic disturbance agents, alien invasive phytophagous insects have caused widespread economic and ecological damage in urban and forested areas of North America (Wallner 1996). The economic impact has been estimated at billions of US dollars each year through loss of forest products, reforestation efforts, management efforts, removal and disposal of hazard trees in urban areas, and loss of livelihood of local communities (Pimentel et al. 2000; Sydnor et al. 2007).

Significant and diverse direct and indirect ecological impacts of alien insects have been documented (Lovett et al. 2006). For example, effects of wide-spread gypsy moth defoliation (Fig. 1) can range from relatively short-term alteration of the forest floor environment to longer-term effects of tree decline and mortality on successional trajectories. Other alien insects such as emerald ash borer selectively kill individual trees resulting in formation of small but widely distributed canopy gaps, ultimately removing their host taxa from invaded communities (Fig. 2). In turn, these disturbances clearly have the potential to unleash a cascade of indirect effects on ecological processes and




**Fig. 1** Wide scale defoliation by gypsy moth triggers short term changes in light, temperature, and moisture regimes on the forest floor, alters nutrient cycles, and exerts longer term effects on competitive interactions and successional trajectories



**Fig. 2** Ash mortality induced by emerald ash borer results in isolated but widely distributed canopy gaps, coupled with extirpation of ash species from invaded forests

interactions that reverberate throughout the forest ecosystem.

The last two decades have seen an increase in studies addressing the economic and ecological impacts of alien insects in forests, and the publication of several important reviews (e.g. Liebhold et al. 1995; Chornesky et al. 2005; Ellison et al. 2005; Lovett et al. 2006; Kenis et al. 2009). Perhaps the best documented ecological impacts of alien insects in eastern North American forests are those of gypsy moth and hemlock woolly adelgid on community composition and nutrient cycling (Lovett et al. 2006). A diverse potpourri of isolated studies has documented an array of additional ecological impacts, but few of these have been characterized comprehensively, and a full accounting



**Fig. 3** Generalized conceptual model depicting cascading direct and indirect effects of alien insect herbivores on ecological processes and interactions that ultimately impact community composition and successional trajectories in eastern North American forests

of impacts of alien insects on forest ecosystems remains elusive.

The objective of this review is to elucidate potential direct and indirect ecological impacts of alien herbivorous insects in the mixed deciduous and hemlock forests of eastern North America. To address the following questions, we employ the framework of Parker et al. (1999) for classifying impacts of invasive organisms, emphasizing effects on ecological processes (e.g. on disturbance regimes such as gap formation, and rates of resource availability and acquisition), as well as community and population level impacts. How can alien insects directly alter abiotic conditions and ecological processes? How can these changes cascade through forest ecosystems as indirect effects on ecological interactions, community structure, and successional trajectories? What are the direct and indirect impacts of management activities targeted at alien insects? A conceptual overview of our synthesis is outlined in Fig. 3. By necessity, much of this review is focused on illuminating knowledge gaps and hypotheses for testing.

### Impacts of alien insects on ecosystem processes

#### Formation of canopy gaps

Several alien insect species have caused extensive tree mortality in eastern forests, including gypsy

moth, hemlock woolly adelgid, and beech scale (Liebhold et al. 1995; Ellison et al. 2005; Lovett et al. 2006), and more recently emerald ash borer (Herms et al. 2004; Poland and McCullough 2006), which has caused nearly 100% mortality of ash near its point of introduction in southeastern Michigan (Gandhi et al. 2008). Gap formation resulting from decline and mortality of oak and beech, resulting from gypsy moth defoliation (Davidson et al. 1999; Jedlicka et al. 2004) and beech bark disease (Houston 1975; Morin et al. 2007; Runkle 2007), respectively, is particularly well documented. By inducing the decline and mortality of host trees, alien herbivorous insects cause the formation of canopy gaps (Rabenold et al. 1998; Runkle 2005), with their size and frequency being dependent upon host dominance, density, distribution, and mortality levels. The ecological significance of gap formation for community structure and successional trajectory is especially important in eastern forests where wildfire frequency is naturally low and has been further suppressed by management activities (Runkle 1982, 1990, 1998, 2000; Shang et al. 2007).

Gap dynamics induced by alien insects are likely to differ from those generated by gradual decline and senescence of canopy trees or abiotic disturbance events such as windthrow. In the mixed deciduous forests characteristic of eastern North America, natural decline and mortality of aging trees generates gradual but ongoing formation of small gaps, while storms can generate episodic formation of numerous gaps, which can vary in size depending on the intensity of the disturbance (Runkle 1990; Canham et al. 2001; Nagel and Svoboda 2008). Gap formation caused by alien insects is also likely to be episodic. However, decline and mortality of trees in response to insect infestation is more likely to be species-specific and generally slower, with dead trees remaining standing longer as snags (Krasny and DiGregorio 2001). In the Allegheny hardwood forests of New York impacted by beech bark disease and gypsy moth, the area covered by gaps increased from 20 to 32% over 6 years (Krasny and DiGregorio 2001).

Gap formation caused by alien insects (and other disturbance agents) alters the forest microenvironment including light, moisture, and temperature regimes (Twery 1990; Stadler et al. 2006), and increases edge-effects, patchiness, and habitat heterogeneity (Canham et al. 1990; Webb 1999; Gandhi

et al. 2007). For example, hemlock decline and mortality caused by hemlock woolly adelgid increased canopy gaps and levels of light reaching the forest floor (Orwig and Foster 1998), which decreased soil moisture within 3 years (Orwig et al. 2008). Such alterations in microhabitat conditions can alter understory vegetation dynamics (Orwig and Foster 1998; Rabenold et al. 1998; see section “Impacts on Native Understory Plants”).

#### Impacts on woody debris dynamics

As host trees experience greater mortality over time, dead standing trees (snags) eventually fall and become downed-woody debris (DWD), which provides important habitat for many animal and plant species (Gandhi et al. 2007; Owens et al. 2008). DWD, including fine—(small branches and roots) and coarse—(tree trunks and branches) woody debris, increases structural and chemical heterogeneity of the forest floor (Harmon et al. 1986).

Few studies have addressed rates and volume of woody debris generated by mortality caused by alien insects. The rate and manner of tree fall (e.g. uprooted trees disturb the adjacent soil environment to a greater degree than do trees that snap above ground) and decomposition will likely differ depending on tree species and rate of mortality (e.g. slower decline caused by defoliation vs. more rapid mortality caused by wood-borers), presence of decay agents such as fungi and other saproxylic insects, topography, and prevalence of wind- and ice-storms (Gandhi et al. 2007). For example, beech trees killed by beech bark disease fell at a rate that tripled over 6 years, while oaks killed by gypsy moth fell at a rate that remained constant over the same period (Krasny and DiGregorio 2001). Species-specific rates of tree-fall suggest that invasive insects can alter the dynamics of snags and downed woody debris. It remains to be seen, however, how increased habitat heterogeneity and other ecological impacts of woody debris generated by alien insects compare with those created by abiotic disturbances such as wind- and ice-storms (Gandhi et al. 2007).

#### Impacts on biogeochemical cycling

In a seminal paper, Mattson and Addy (1975) proposed that phytophagous insects can act as



regulators of primary productivity by influencing patterns of nutrient cycling. The boreal forest of North America has long been subjected to expansive outbreaks of indigenous defoliating insects including forest tent caterpillar (*Malacosoma disstria* Hübner) and spruce budworm [*Choristoneura fumiferana* (Clemens)] (Mattson et al. 1991). Eastern deciduous forests, however, did not experience episodic defoliation on a wide scale prior to introduction of gypsy moth (Mattson et al. 1991). Subsequent studies have confirmed the potential for gypsy moth defoliation to alter biogeochemistry of both terrestrial and aquatic ecosystems through increased inputs of DWD, green litter fall, frass, and their corpses (Lovett et al. 2002).

Expansive defoliation can transfer substantial quantities of foliar nitrogen in green leaf fragments and frass to the forest floor before it can be reallocated within the plant prior to leaf senescence (Schowalter et al. 1986; Lovett et al. 2002). Furthermore, Lovett and Ruesink (1995) found that highly labile carbon in gypsy moth frass stimulated microbial growth resulting in immobilization of more than 90% of nitrogen in soil samples within 10 days, which could potentially reduce availability of nitrogen to trees, at least in the short term. Similar results were found for nitrogen in frass of gypsy moth that had consumed oak foliage labeled with  $^{15}\text{N}$ . Nitrogen in the frass was mobilized more quickly than nitrogen in leaf litter, rendering it largely unavailable to plants (Christenson et al. 2002). In a Pennsylvania oak forest, gypsy moth defoliation increased quantities of nitrogen, potassium, and phosphorus in litter fall, and decreased quantities of calcium (Grace 1986). The seasonality of the nutrient pulse was also altered as most litter (including frass and leaf fragments) fell in early summer when larvae were feeding, rather than in early autumn during natural leaf abscission. In addition to effects imposed by defoliation, canopy gap formation resulting from tree mortality can also alter patterns of nitrogen and carbon cycling (Scharnbroch and Bockheim 2008a, b).

Effects of alien insects on biogeochemical cycles also have been detected at the level of the watershed (Swank et al. 1981; Ellison et al. 2005; Lewis and Likens 2007). Grady et al. (2007) found that 66 watersheds in western Virginia had reduced concentrations of dissolved silica from 1988 to 2003. Abiotic factors such as pH, precipitation, and water discharge were not correlated with changing amounts

of silica. However, the reduction in silica levels did coincide with initiation of gypsy moth outbreaks in that region. The authors hypothesized that defoliation increased light availability and nitrate concentrations in the streams, which, in turn, stimulated populations of benthic diatoms, leading to greater extraction of silica from the water. Webb et al. (1995) reported that gypsy moth defoliation affected the pH and buffering capacity of streams, and increased nitrate levels almost tenfold. This potentially may have consequences for aquatic animals, as high nitrate levels have been shown to adversely affect the behavior, morphology, and survival of amphibian larvae (Marco et al. 1999).

It can be hypothesized that the extensive defoliation characteristic of gypsy moth outbreaks may trigger positive feedbacks that decrease productivity of terrestrial and aquatic ecosystems for several years (Choudhury 1988; Schweitzer et al. 2005). For example, herbivory has been shown to decrease the rate of decomposition of litter derived from the damaged leaves (Findlay et al. 1996), as well as increase foliar concentrations of secondary metabolites and decrease foliar nutrient concentrations in the years following defoliation (Schultz and Baldwin 1982; Tuomi et al. 1984; Bryant et al. 1993). Higher secondary metabolite concentrations and carbon:nitrogen ratios in green foliage have been associated with decreased rates of litter decomposition and nitrogen mineralization (Horner et al. 1988; Hättenschwiler and Vitousek 2000), which can decrease nutrient availability and uptake by trees. In turn, soil nutrient limitation can elevate foliar concentrations of carbon-based secondary metabolites (Mattson 1980; Herms and Mattson 1992), which may further inhibit decomposition, thereby decelerating rates of nutrient cycling (Schweitzer et al. 2005). Such feedback effects should relax over time as delayed defoliation-induced effects on host quality subside over subsequent years (Haukioja 1990).

Indirect effects of herbivory on litter quality may also impact the productivity of allochthonous aquatic habitats by decreasing the quality of the leaf detritus upon which their food webs are founded, including streams (Irons et al. 1991) and tree-holes (Walker et al. 1997; Strand et al. 1999). In some cases, defoliation can also increase the quality of leaf litter. Following severe defoliation by gypsy moth during spring, oak trees generally refoliate during summer.

Hutchens and Benfield (2000) found that this second flush of foliage had higher ratios of protein to fiber than did the first flush leaves, resulting in their faster decomposition in streams.

Altered patterns of terrestrial and aquatic nutrient cycling have been well documented in hemlock stands impacted by hemlock woolly adelgid (Ellison et al. 2005). Hemlock mortality accelerated nitrogen mineralization and nitrification rates (Jenkins et al. 1999; Orwig et al. 2008), elevated nitrate and cation levels in soil water (Yorks et al. 1999), and increased dissolved organic carbon and nitrogen in through-fall (Stadler et al. 2006). Following adelgid-induced hemlock decline and mortality, Small et al. (2005) reported up to 140% increase in density of black birch (*Betula lenta* L.) seedlings and saplings, which have higher concentrations of foliar nitrogen and produce higher quality litter than does hemlock foliage (Cobb and Orwig 2002). Hemlock woolly adelgid may also exert ecologically significant impacts on hydrological cycles. For example, Ford and Vose (2007) concluded that wide-spread mortality of eastern hemlock will decrease annual stand-level transpiration by about 10%, and decrease winter and spring stand-level transpiration by about 30%, resulting in persistent increases in discharge by forest streams.

### Impacts of alien insects on biotic populations and communities

#### Impacts on canopy composition

Decline and mortality of indigenous tree species caused by alien phytophagous insects has altered forest community composition and successional trajectories (Campbell and Sloan 1977; Runkle 2005). Shifts in forest species composition following defoliation by gypsy moth have been well documented in eastern North America (Campbell and Sloan 1977; Davidson et al. 1999). Several studies found that growth of oaks declined in the year following defoliation, while the growth of less favored hosts such as maples (*Acer* spp.) and ashes increased, probably due to release from competitive suppression (Muzika and Liebhold 1999; Naidoo and Lechowicz 2001). Furthermore, there is evidence that oak mortality increased the importance of shade-tolerant

species that are less preferred by gypsy moth, such as maples, beech, and cherries (*Prunus* spp.) (Campbell and Sloan 1977; Nowacki and Abrams 1992; Muzika and Twery 1995; Fajvan and Wood 1996; Jedlicka et al. 2004). Beech bark disease has decreased the density of American beech in the overstory while, at least in some cases, increasing its density in the understory, probably via strong basal sprouting of infested mature trees (Houston 1975; Morin et al. 2007; Runkle 2007).

Forests infested with hemlock woolly adelgid have also experienced a shift in canopy composition, with black oak (*Q. velutina* Lam.), sassafras [*Sassafras albidum* (nutt.) Nees], red maple (*Acer rubrum* L.), black birch, yellow birch (*B. alleghaniensis* Britton), and American beech gaining in importance in the canopy layer as hemlock declined (Small et al. 2005). Successional responses were site-specific, with black oak gaining greater dominance on xeric ridges and ledges, and other species on mesic sites in ravines (Small et al. 2005).

Ironically, in several cases, endemic tree species that increased in dominance as alien insects killed their competitors have themselves in recent years become threatened by the next wave of alien invaders. For example, wide-scale mortality of American elm caused by Dutch elm disease led to increased dominance of ash in eastern North American forests (Barnes 1976), which are now threatened by emerald ash borer. It has been hypothesized that following the chestnut blight epidemic, American chestnut [*Castanea dentata* (Marsh.) Borkh.] was replaced in riparian areas of the Appalachians by hemlocks (Ellison et al. 2005), which are now threatened by the hemlock woolly adelgid. Oak trees increased in density in the eastern United States early in the twentieth century, in part due to the demise of American chestnut (Muzika and Liebhold 2001), only to become the preferred host of gypsy moth.

#### Impacts on native understory plants

The forest understory is also directly affected by defoliation and gap formation caused by alien insects, and the consequent increase in light penetration and temperature (e.g. Kasbohm et al. 1996). In New England, increased growth of black birch, red maple, and oak seedlings in the understory was observed in response to hemlock decline (Orwig 2002). Hemlock

mortality also increased cover of hay-scented fern [*Dennstaedtia punctilobula* (Michx.) T. Moore], which can form a dense herbaceous layer that inhibits tree regeneration following disturbance (Penrod and McCormick 1997). Populations of other bryophytes have also been observed to increase as hemlock mortality increased light availability in the understory (Cleavitt et al. 2008).

Following mortality of Fraser fir due to balsam woolly adelgid, the cover of woody shrubs such as blackberry (*Rubus canadensis* L.) increased, whereas the cover of various herbaceous species, including some bryophytes, decreased (DeSelm and Boner 1984). Similarly, in the Great Smoky Mountains, the density of blackberry increased, while that of herbaceous plants such as woodfern [*Dryopteris campyloptera* (Kunze) Clarkson] and oxalis (*Oxalis acetocella* Linnaeus) decreased, as did density of shade-adapted shrubs such as hobblebush (*Viburnum alnifolium* Marsh.) and mountain cranberry [*Vaccinium erythrocarpum* (Michx.)] (Rabenold et al. 1998).

The herbaceous plant layer can exert strong control over ecological processes in temperate forest ecosystems (Zak et al. 1990; Gilliam 2007). Hence, these studies suggest that effects of alien insects on canopy species will invariably cascade to the forest floor, leading to reorganization of the understory community with important implications for forest successional trajectories.

#### Impacts on invasive plants

It has been hypothesized that alien forest insects, including hemlock woolly adelgid (Orwig and Foster 1998) and emerald ash borer (Herms et al. 2008), can facilitate the establishment and spread of invasive plants by creating canopy gaps that increase light availability while relaxing interspecific competition for space and resources, thereby igniting an “invasional meltdown” (Simberloff and Von Holle 1999). Many invasive plant species are not highly tolerant of shade and are more common at forest edges where light is not as limiting (Knapp and Canham 2000; Parendes and Jones 2000; Hunter and Mattice 2002). Furthermore, disturbances that open the canopy have been shown to facilitate establishment of alien plants in the forest interior (Horvitz et al. 1998; Snitzer et al. 2005; Webster et al. 2005; Glasgow and Matlack

2007; Eschtruth and Battles 2009). McEwan et al. (2009) speculated that gypsy moth may also facilitate the spread of invasive shrubs in North America by selectively feeding on native species.

Although few studies have directly investigated facilitation of invasive plants by alien herbivores, increased incidence and cover of invasive plants has been documented in stands impacted by hemlock woolly adelgid (Orwig 2002; Eschtruth et al. 2006), with interactions between canopy disturbance and propagule pressure being the most important factors affecting the distribution of exotic plants (Eschtruth and Battles 2009). Increased abundance of exotic plants may prevent recruitment and establishment of indigenous species, and thus alter successional trajectories within invaded forests (Levine et al. 2003), while exerting feedback effects that further alter community composition (e.g., Gómez-Aparicio and Canham 2008). While not yet well documented, alien invasive insects clearly have the potential to exert pervasive indirect effects that cascade through forest communities through facilitative effects on invasive plants.

#### Impacts on native fauna

Alien herbivorous insects may impact indigenous animals by decreasing (or, in some cases, increasing) the distribution and abundance of plants on which they depend for at least part of their life-cycle (Koh et al. 2004). The effect may be especially strong for herbivorous insects that specialize on host plants strongly impacted by alien herbivores. Wagner (2007) catalogued at least 21 species of monophagous Lepidoptera that specialize on North American ash species that will be threatened by the demise of North American ash due to emerald ash borer. Specialist herbivores of eastern hemlock and Fraser fir are similarly threatened by hemlock woolly adelgid and balsam woolly adelgid, respectively. To date, there has been little synthesis of the natural history of the community of organisms associated with tree species threatened by alien insects and pathogens. Such research is urgently needed before these trees become rare or locally extirpated, resulting in biotic impoverishment.

Alien insects may also decrease the abundance of native phytophagous insects through competitive interactions (Scriber 2004). Work and McCullough

(2000) assessed the response of forest Lepidoptera to defoliation by gypsy moth in Michigan. While most native species appeared to be resilient to the gypsy moth outbreak, the populations of a subset of species associated with oak (a favored gypsy moth host) were negatively affected during the year of the outbreak, possibly because of interspecific competition for food. Gypsy moth defoliation also temporarily decreased populations of native Lepidoptera in West Virginia; however, the effects were small relative to annual fluctuations in caterpillar density due to factors such as weather (Sample et al. 1996).

Alien herbivorous insects may exert indirect effects on the population and community structure of indigenous fauna by altering their habitat and food supply. For example, balsam woolly adelgid reduced the dominance of endemic fir trees by about 50% and drastically opened the canopy in the Appalachian forests. In response, populations of birds that forage in the canopy and sub-canopy of the fir forests decreased, whereas populations of bird species more characteristic of disturbed habitats increased (Rabenold et al. 1998). Similarly, populations of birds that are habitat specialists of hemlock forests, such as Acadian flycatcher [*Empidonax virescens* (Vieillot)], have been adversely affected by extensive adelgid-induced hemlock mortality (Tingley et al. 2002; Becker et al. 2008). In contrast, populations of other birds increased in infested hemlock stands, including the hooded warbler [*Wilsonia citrina* (Boddaert)], a species of regional concern that benefited from a dense black birch understory created by hemlock mortality (Tingley et al. 2002), and opportunistic species such as the wood thrush [*Hylocichla mustelina* (J.F. Gmelin)] that benefited from increased numbers of dead trees and canopy gaps (Becker et al. 2008).

During outbreaks, gypsy moth larvae represent a super-abundant food source for predators such as native cuckoos (*Coccyzus* spp.) that specialize on hairy caterpillars, and Barber et al. (2008) observed a shift in cuckoo distributions during outbreaks, with more cuckoos in areas where gypsy moth was epidemic. As predators respond spatially and temporally to populations of alien insects, predation pressure on indigenous species may change in complex ways.

Stress imposed on oaks by severe gypsy moth defoliation decreases or eliminates acorn production (Kasbohm et al. 1996), which is an important food source for many species of wildlife (Ostfield et al.

1996; McShea 2000; McShea et al. 2007). Effects of gypsy moth defoliation on acorn production have been shown to reduce the reproductive success of small mammals including white-footed mice (*Peromyscus leucopus* Rafinesque) and eastern chipmunks (*Tamias striatus* L.), alter the foraging patterns of white-tailed deer [*Odocoileus virginianus* (Zimmerman)] (Ostfield et al. 1996), and shift the dietary choices of black bear (*Ursus americanus* Pallas) (Kasbohm et al. 1996). Gypsy moth defoliation also increased the rate of nest predation of forest birds, possibly by increasing nest visibility (Thurber et al. 1994). Conversely, longer-term habitat changes including increased availability of snags and opened canopy may create more nesting habitat for many bird species (Bell and Whitmore 2000).

Gypsy moth density may be indirectly tied to acorn production through effects of acorn availability on mice populations, which are key predators of gypsy moth. In a fascinating experiment, Jones et al. (1998) manipulated the availability of acorns in the forest by removing them from some plots and distributing the collected acorns in other plots. Where the acorn supply was experimentally increased, mouse populations increased dramatically, as did predation of gypsy moth pupae by mice. Where acorn density was experimentally lowered, mouse populations decreased and gypsy moth predation declined. Hence, it is possible that gypsy moth outbreaks could be prolonged if defoliation-induced reductions in acorn production have positive feedback effects on gypsy moth survival through effects on mice.

Indirect negative interactions between alien and endemic insect herbivores mediated by host trees and natural enemies have also been documented. In Michigan, Redman and Scriber (2000) found that gypsy moth defoliation decreased growth and survival of native swallowtail butterflies (*Papilio canadensis* Rothschild & Jordan) on quaking aspen (*Populus tremuloides* Michx.) by inducing reductions in host quality. Further, tiger swallowtail larvae experienced higher rates of parasitism in the vicinity of high gypsy moth populations. The generalist parasitic wasp was likely attracted in greater numbers by the presence of gypsy moth. This study suggests that alien herbivorous insects can indirectly exert both top-down and bottom-up impacts on native fauna. Preisser and Elkinton (2008) documented competitive interactions between two alien insect



specialists of hemlock: hemlock woolly adelgid and elongate hemlock scale (*Fiorinia externa* Ferris). The densities of both species were reduced by 30% in mixed-species relative to single-species trials. Interestingly, foliage growth was greater in the mixed-species treatment, indicating that the presence of elongate hemlock scale (a less damaging species on hemlock trees) can mitigate some of the adverse effects of hemlock woolly adelgid.

Not all interactions between alien and native insects are negative. For example, Hammons et al. (2009) found that feeding on ripening fruits by the exotic scarab Japanese beetle (*Popillia japonica* Newman) facilitated subsequent feeding by native scarab beetles. Gypsy moth defoliation has been shown to facilitate colonization of oaks by twolined chestnut borer [*Agilus bilineatus* (Weber)], probably by compromising tree defenses to this phloem-feeding wood borer (Dunbar and Stephens 1975; Wargo 1977).

### Ecological impacts of management practices for controlling alien insects

#### Impacts of chemical control

Direct and indirect effects of alien insects on forest ecosystems can be considered to extend to the tactics used to manage them. For example, aerial applications of the biopesticide *Bacillus thuringiensis kurstaki* (*Btk*) used to eradicate and suppress gypsy moth infestations can have nontarget effects on indigenous Lepidoptera (Herms 2003; Scriber 2004). In West Virginia, a single aerial application of *Btk* to suppress gypsy moth decreased the diversity and abundance of endemic butterflies and moths for one season (Sample et al. 1996). Similar effects were observed in Virginia (Rastall et al. 2003). In Oregon, where three applications of *Btk* were made during the same season to eradicate small populations of gypsy moth, caterpillar diversity was lower 3 years after application, although overall numbers of caterpillars rebounded within 1 year (Miller 1990). The short residual activity of *Btk* in the field (considered to be only a few days) is thought to minimize effects on nontarget Lepidoptera. However, Johnson et al. (1995) found that foliage treated with *Btk* applied with ground equipment at a high labeled rate remained toxic to

swallowtail butterfly (*P. glaucus* L.) larvae for 30 days following application. Conversely, nucleopolyhedrosis virus (NPV) applied to suppress gypsy moth in West Virginia had no detectable adverse effects on populations of native Lepidoptera (Rastall et al. 2003), or in laboratory studies on mammalian predators of gypsy moth such as white-footed mouse, short-tailed shrew [*Blarina brevicauda* (Say)], and Virginia opossum (*Didelphis marsupialis* L.) (Lautenschlager et al. 1977).

Diflubenzuron is a growth regulating insecticide that interferes with exoskeleton formation of immature insects during the molting process. It is considered more effective than *Btk* for suppressing gypsy moth, but is not used as widely because it impacts a much broader diversity of insects with effects that persist much longer (Eisler 1992). Aquatic arthropods, including insects and crustaceans, are especially sensitive, but aerial applications of diflubenzuron also have been shown to decrease the abundance and diversity of numerous taxa of terrestrial forest insects, with effects persisting beyond the year of application (Martinat et al. 1988; Butler et al. 1997). Residues can persist on foliage throughout the season, and soil and aquatic arthropods have been impacted by senesced leaves that abscise in autumn (Griffith et al. 2000). In a study on litter- and soil-dwelling arthropods in a West Virginia forest, spiders (Araneae) and mites (Acari) were not affected by application of diflubenzuron, but populations of springtails (Collembola) were decreased (Perry et al. 1997).

Effects of aerial insecticide applications on nontarget insects may also extend to higher trophic levels by decreasing their food supply. For example, in a study in which *Btk* applications caused significant declines in Lepidoptera larvae, populations of 18 of 27 insectivorous birds also declined (Strazanac and Butler 2005). However, effects on most bird species were temporary, with all but two recovering fully within 3 years following the treatments (Strazanac and Butler 2005). Another study found that when caterpillar populations were decreased by aerial application of diflubenzuron, birds consumed fewer caterpillars and spent more time foraging for food (Cooper et al. 1990).

#### Impacts of biological control

Classical biological control, the introduction of natural enemies from the endemic range of the alien

herbivore, is considered a corner-stone of integrated pest management. However, there have been cases when these “enemies of the enemy” have turned against nontarget indigenous fauna with unintended consequences (Hawkins and Marino 1997). For example, high rates of parasitism by the alien tachinid fly, *Compsilura concinnata* Meig, which was introduced to eastern North America more than 100 years ago to control gypsy moth, has been implicated in the decline of North American silk moths (Saturniidae) in New England, including *Hyalophora cecropia* (Linnaeus) and *Callosamia promethea* (Drury) (Boettner et al. 2000). Kellogg et al. (2003) reported similar effects of this parasitoid on experimental populations of luna moth (*Actias luna* L.). As might be expected, these impacts on North American silk moth populations appear to vary regionally (Selfridge et al. 2007).

There is some evidence that introduced biocontrol agents can competitively exclude indigenous parasitoids. For example, comparison of surveys of parasitoids of endemic Lepidoptera conducted between 1915–1929 and 1999–2005 in Massachusetts revealed that the tachinid fly, *Lespesia frenchii* (Williston), was a common parasitoid of the silk moth *C. promethea* and eastern tent caterpillar (*Malacosoma americanum* Fabricius) early in the twentieth century. However, this indigenous parasitoid species was not recovered in the latter survey when parasitism by the alien *C. concinnata* was common (Parry 2009). Interestingly, in New York, where *C. concinnata* parasitism rates are much lower, *L. frenchii* was reared from its native Lepidoptera hosts in similar frequencies in 1915–1929 and 1999–2005 (Parry 2009).

#### Impacts of silvicultural treatments used to manage alien insects

Host trees are sometimes removed from forested landscapes and destroyed in an attempt to eradicate alien species, as was the case for the emerald ash borer eradication program (Herms et al. 2004; Poland and McCullough 2006), or to salvage logs from trees killed by alien species (Orwig and Kittredge 2005), which can alter microhabitat conditions and successional trajectories. For example, logging of hemlock woolly adelgid-infested trees in New England increased soil compaction, pH, nitrification rates, scarification, and

biomass of woody debris, while increasing dominance of black birch, brambles (*Rubus* spp.), sedges (Cyperaceae), and hay-scented fern (Kizlinski et al. 2002; Orwig and Kizlinski 2002). Hausman et al. (2008) reported that removal of ash trees from a natural area to eradicate emerald ash borer increased soil compaction, possibly facilitating colonization of the newly formed gaps by invasive plants.

Thinning and pre-salvage harvesting of living trees have also been evaluated as tactics for preemptively decreasing forest susceptibility to invasion by alien insects. One study assessed the impact on salamanders of selective pre-salvage harvesting of green hemlock trees in anticipation of hemlock woolly adelgid invasion (Brooks 2001). Although populations of eastern redback salamander (*Plethodon cinereus* Green) decreased in harvested areas, the effects were ephemeral as populations rebounded within a few years (Brooks 2001). Muzika et al. (2004) assessed the effects of thinning of oaks to reduce vulnerability of forests to gypsy moth on small mammal populations in West Virginia. Mice (*Peromyscus* spp.) were found in higher numbers in thinned than in unthinned control stands, while populations of masked shrew (*Sorex cinereus* Baird) and redback vole [*Clethrionomys gapperi* (Vigors)] were not affected. Comparable studies on other taxa (invertebrates and vertebrates) are notably absent.

#### Synthesis and conclusions

Alien insect herbivores, and the programs to manage them, clearly have the potential to unleash a diverse cascade of direct and indirect effects on ecosystem processes and ecological interactions that can alter community composition and successional trajectories of eastern North American forests (Fig. 3). Direct effects can range from widespread defoliation (e.g. gypsy moth) to formation of isolated but widely distributed gaps caused by species-specific tree mortality (e.g. emerald ash borer). Effects of gypsy moth and hemlock woolly adelgid on the abiotic environment, nutrient cycling, and community composition are the most comprehensively documented (Lovett et al. 2006), but a diverse array of additional direct, indirect, and feedback effects have also been documented or can be hypothesized (Kenis et al. 2009).

The impact of some of the most devastating alien species that cause wide-spread tree mortality, such as hemlock woolly adelgid (Havill et al. 2006) and emerald ash borer (Rebek et al. 2008), is magnified by the low level of resistance in their hosts, possibly because their lack of coevolutionary history has rendered them effectively defenseless. These alien invaders threaten to decimate their hosts on a continental scale as they continue to spread in “defense free space.” More studies are required to document the effects of extensive host tree mortality on the native fauna that depend on them for food and shelter. Such studies would generate a better understanding of the threat of host extirpation on co-extinction of their specialized fauna.

Tree mortality creates canopy gaps and increases woody debris (both standing and fallen), which in turn affects the physical environment and biogeochemical cycling (Stadler et al. 2006), and ultimately community composition (Runkle 2005). Canopy gaps caused by alien insects may exert ecological impacts that differ qualitatively from those caused by natural tree decline and senescence, which tend to form more slowly, or those caused by disturbances such as wind and ice storms, which do not selectively remove individual taxa from the community (Gandhi et al. 2007). The dynamics and effects of gaps caused by extensive species-specific tree mortality should be more thoroughly investigated.

Defoliation and tree mortality induced by alien insects may generate a cascade of indirect effects on ecological interactions. For example, gypsy moth defoliation can substantially decrease acorn production by oak, which in turn can reduce small mammal populations by decreasing their food supply (Ostfield et al. 1996; McShea et al. 2007). Gypsy moth defoliation has also been shown to increase the rate of nest predation of forest birds, possibly by increasing nest visibility (Thurber et al. 1994). Although poorly documented, feedback effects of herbivory and tree mortality caused by alien insects may be pervasive. For example, changes in foliage and litter quality induced by defoliation (e.g. Findlay et al. 1996) may trigger positive feedbacks on nutrient cycles that could decrease the productivity of terrestrial and aquatic ecosystems over several years (e.g. Choudhury 1988; Schweitzer et al. 2005), as well as alter competitive (Redman and Scriber 2000) and facilitative (Hammons et al.

2009) interactions between endemic and alien herbivores. Some studies also suggest widespread potential for alien insects to facilitate the establishment and spread of exotic plants by altering resource availability and competitive interactions (e.g. Eschtruth et al. 2006; Eschtruth and Battles 2009), thereby unleashing an “invasional meltdown” (Simberloff and Von Holle 1999).

An increasing body of evidence suggests that tactics used to eradicate and suppress alien species also can have direct and indirect effects on forest communities. Aerial insecticide applications have been shown to temporarily suppress endemic nontarget fauna (Sample et al. 1996), with effects that can extend to higher trophic levels (Strazanac and Butler 2005). There is evidence that some generalist parasitoids released as biological control agents have exerted long-term negative effects on indigenous fauna (Boettner et al. 2000; Parry 2009). Tree removal associated with eradication programs (Hausman et al. 2008) and salvage logging of stands impacted by invasive insects (Kizlinski et al. 2002; Orwig and Kizlinski 2002) also has the potential to alter forest communities and ecosystem processes. Additional research is required to fully characterize the costs and benefits of programs to manage invasive forest insects.

The mixed deciduous and hemlock forests of eastern North America have experienced large-scale disturbance during the last two centuries, resulting in habitat destruction, isolation, and fragmentation (Russell and Davis 2001). Combined with ongoing climate change, the addition of alien herbivorous insects to the “disturbance potpourri” will almost certainly continue to increase the strain on the stability and resilience of North American forests (Mattson 1997; Dale et al. 2009). However, major knowledge gaps remain regarding short and long-term direct and indirect effects of alien invasive insects on structure and function of forest ecosystems. Future research may focus on longer-term assessments of impacts of alien herbivorous insects across various abiotic conditions, taxa, and ecosystems; provide models for predicting and mitigating some of these adverse impacts at the landscape level (e.g. Dale et al. 2009); and document the benefits (and costs) of quarantine, eradication, and containment measures for preventing the establishment and spread of alien invasive insects.

**Acknowledgments** We thank Catherine P. Herms (The Ohio State University) and two anonymous reviewers for their helpful comments on earlier drafts of this manuscript, and Lee A. Odgen (The University of Georgia) for formatting the manuscript. Financial support was provided by the USDA Forest Service, Northern Research Station, Morgantown, West Virginia, and Delaware, Ohio, and by state and federal funds appropriated to the Daniel B. Warnell School of Forestry and Natural Resources, The University of Georgia, Athens, and to the Ohio Agricultural Research and Development Center, The Ohio State University, Wooster.

## References

- Barber NA, Marquis RJ, Tori WP (2008) Invasive prey impacts the abundance and distribution of native predators. *Ecology* 89:2678–2683
- Barnes BV (1976) Succession in deciduous swamp communities of southern Michigan formerly dominated by American elm. *Can J Bot* 54:19–24
- Becker DA, Brittingham MC, Goguen CB (2008) Effects of hemlock woolly adelgid on breeding birds at Fort Indiantown Gap, Pennsylvania. *Northeast Nat* 15:227–240
- Bell JL, Whitmore RC (2000) Bird nesting ecology in a forest defoliated by gypsy moths. *Wilson Bull* 112:524–531
- Boettner GH, Elkinton JS, Boettner CJ (2000) Effects of a biological control introduction on three nontarget native species of saturniid moths. *Conserv Biol* 14:1798–1806
- Brooks RT (2001) Effects of the removal of overstory hemlock from hemlock-dominated forests on eastern redback salamanders. *For Ecol Manage* 2001:197–204
- Bryant JP, Reichardt JB, Clausen TP, Werner RA (1993) Effects of mineral nutrition on delayed inducible resistance in Alaska paper birch. *Ecology* 74:2072–2084
- Butler L, Chrislip GA, Kondo VA, Townsend EC (1997) Effect of diflubenzuron on nontarget canopy arthropods in closed, deciduous watersheds in a central Appalachian forest. *J Econ Entomol* 90:784–794
- Campbell RW, Sloan RJ (1977) Forest stand responses to defoliation by the gypsy moth. *For Sci Monogr* 19:1–34
- Canham CD, Denslow JS, Platt WJ, Runkle JR, Spies TA, White PS (1990) Light regimes beneath closed canopies and tree-fall gaps in temperate and tropical forests. *Can J For Res* 20:620–631
- Canham CD, Papaik MJ, Latty EF (2001) Interspecific variation in susceptibility to windthrow as a function of tree size and storm severity for northern temperate tree species. *Can J For Res* 31:1–10
- Cappaert D, McCullough DG, Poland TM, Siegert NW (2005) Emerald ash borer in North America: a research and regulatory challenge. *Am Entomol* 51:152–165
- Chornesky EA, Bartuska AM, Aplet GH, Britton KO, Cummings-Carlson J, Davis FW, Eskow J, Gordon DR, Gottschalk KW, Haack RA, Hansen AJ, Mack RN, Rahel FJ, Shannon MA, Wainger LA, Wigley TB (2005) Science priorities for reducing the threat of invasive species to sustainable forestry. *Bioscience* 55:335–348
- Choudhury D (1988) Herbivore induced changes in leaf-litter resource quality: a neglected aspect of herbivory in ecosystem nutrient dynamics. *Oikos* 51:389–393
- Christenson LM, Lovett GM, Mitchell MJ, Groffman PM (2002) The fate of nitrogen in gypsy moth frass deposited to an oak forest floor. *Oecologia* 131:444–452
- Ciesla WM (2003) European woodwasp: a potential threat to North America's conifer forests. *J For* 101:18–23
- Cleavitt NL, Eschtruth AK, Battles JJ, Fahey TJ (2008) Bryophyte response to eastern hemlock decline caused by hemlock woolly adelgid infestation. *J Torr Bot Soc* 135:12–25
- Cobb RC, Orwig DA (2002) Impacts of hemlock woolly adelgid infestation on decomposition: an overview. In: Onken B, Reardon R (eds) Third symposium on hemlock woolly adelgid in the eastern United States. Asheville, NC. FHET-2005-01, pp 317–323
- Cooper RJ, Dodge KM, Martinat PJ, Donahoe SB, Whitmore RC (1990) Effect of diflubenzuron application on eastern deciduous forest birds. *J Wildl Manage* 54:486–493
- Dale VH, Lannom KO, Tharp ML, Hodges DG, Fogel J (2009) Effects of climate change, land-use change, and invasive species on the ecology of the Cumberland forests. *Can J For Res* 39:467–480
- Davidson CB, Gottschalk KW, Johnson JE (1999) Tree mortality following defoliation by the European gypsy moth (*Lymantria dispar* L.) in the United States: a review. *For Sci* 45:74–84
- DeSelm HR, Boner RR (1984) Understory changes in spruce-fir during the first 16–20 years following the death of fir. In: White PS (ed) The southern Appalachian spruce-fir ecosystem: its biology and threats. USDA National Park Service, Atlanta, GA, pp 51–69
- Dunbar DM, Stephens GR (1975) Association of twolined chestnut borer and shoestring fungus with mortality of defoliated oak in Connecticut. *For Sci* 21:169–174
- Eisler R (1992) Diflubenzuron hazards to fish, wildlife, and invertebrates: a synoptic review. U.S. Department of Interior Fish and Wildlife Service, Contaminant Hazard Reviews 25, Biological Report 4
- Elkinton JS, Liebhold AM (1990) Population dynamics of gypsy moth in North America. *Annu Rev Entomol* 35:571–596
- Ellison AM, Bank MS, Clinton BD, Colburn EA, Elliott K, Ford CR, Foster DR, Kloeppel BD, Knoepp JD, Lovett GM, Mohan J, Orwig DA, Rodenhouse NL, Sobczak WV, Stinson KA, Stone JK, Swan CM, Thompson J, Von Holle B, Webster JR (2005) Loss of foundation species: consequences for the structure and dynamics of forested ecosystems. *Front Ecol* 3:479–486
- Eschtruth AK, Battles JJ (2009) Assessing the relative importance of disturbance, herbivory, diversity, and propagule pressure in exotic plant invasions. *Ecol Monogr* 79:265–280
- Eschtruth AK, Cleavitt NL, Battles JJ, Evans RA, Fahey TJ (2006) Vegetation dynamics in declining eastern hemlock stands: 9 years of forest response to hemlock woolly adelgid infestation. *Can J For Res* 36:1435–1450
- Fajvan MA, Wood JM (1996) Stand structure and development after gypsy moth defoliation in the Appalachian plateau. *For Ecol Manage* 89:79–88



- Findlay S, Carreiro M, Krischik V, Jones CG (1996) Effects of damage to living plants on leaf litter quality. *Ecol Appl* 6:269–275
- Ford CR, Vose JM (2007) *Tsuga canadensis* (L.) Carr. mortality will impact hydrologic processes in southern Appalachian forest ecosystems. *Ecol Appl* 17:1156–1167
- Gandhi KJK, Gilmore DW, Katovich SA, Mattson WJ, Spence JR, Seybold SJ (2007) Physical effects of weather disturbances on the abundance and diversity of forest insects in North American forests. *Environ Rev* 15:113–152
- Gandhi KJK, Smith A, Long RA, Taylor RAJ, Herms DA (2008) Three-year progression of emerald ash borer induced ash decline and mortality in the forests of southeastern Michigan. In: Mastro V, Lance D, Reardon R, Parra G (eds) Proceedings of the emerald ash borer and Asian longhorned beetle research and technology development meeting, Pittsburgh, Pennsylvania, 23–24 Oct 2007. USDA Forest Health Technology Enterprise Team, FHTET-2008-07, p 27
- Gilliam FS (2007) The ecological significance of the herbaceous layer in temperate forest ecosystems. *Bioscience* 57:845–858
- Glasgow LS, Matlack GR (2007) The effects of prescribed burning and canopy openness on establishment of two non-native plant species in a deciduous forest, southeast Ohio, USA. *For Ecol Manage* 238:31–329
- Gómez-Aparicio L, Canham CD (2008) Neighborhood models of the effects of invasive tree species on ecosystem processes. *Ecol Monogr* 78:69–86
- Grace JR (1986) The influence of gypsy moth on the composition and nutrient content of litter fall in a Pennsylvania oak forest. *For Sci* 32:855–870
- Grady AE, Scanlon TM, Galloway JN (2007) Declines in dissolved silica concentrations in western Virginia streams (1988–2003): gypsy moth defoliation stimulates diatoms? *J Geophys Res* 112:1–11
- Griffith MB, Barrows EM, Perry SA (2000) Effect of diflubenzuron on flight of adult aquatic insects (Plecoptera, Trichoptera) following emergence during the second year after aerial application. *J Econ Entomol* 93:1695–1700
- Haack RA (2006) Exotic bark- and wood-boring Coleoptera in the United States: recent establishments and interceptions. *Can J For Res* 36:269–288
- Hammons DL, Kurtural SK, Newman MC, Potter DA (2009) Invasive Japanese beetles facilitate aggregation and injury by a native scarab pest of ripening fruits. *Proc Nat Acad Sci USA* 106:3686–3691
- Hanula JL, Berisford CW (1984) Seasonal flight activity of the smaller European elm bark beetle, *Scolytus multistriatus* (Coleoptera: Scolytidae), and associates in Georgia. *Can Entomol* 116:1251–1258
- Hanula JL, Mayfield AE, Fraedrich SW, Rabaglia RJ (2008) Biology and host associations of redbay ambrosia beetle (Coleoptera: Curculionidae: Scolytinae), exotic vector of laurel wilt killing redbay trees in the southeastern United States. *J Econ Entomol* 101:1276–1286
- Harmon ME, Franklin JF, Swanson FJ, Sollins P, Gregory SV, Lattin JD, Anderson NH, Cline SP, Aumen NG, Sedell JR, Lienkaemper GW, Cromack K Jr, Cummins KW (1986) Ecology of coarse woody debris in temperate ecosystems. *Adv Ecol Res* 15:133–302
- Hättenschwiler S, Vitousek PM (2000) The role of polyphenols in terrestrial ecosystem nutrient cycling. *Trends Ecol Evol* 15:238–243
- Haugen DA, Hoebeke ER (2005) *Sirex* woodwasp—*Sirex noctilio* F. (Hymenoptera: Siricidae). USDA-Forest Service, Pest Alert NA-PR-07-05
- Haukioja E (1990) Induction of defenses in trees. *Annu Rev Entomol* 36:25–42
- Hausman CE, Rocha OJ, Jaeger JF (2008) How emerald ash borer facilitates a secondary spread of invasive plant species: impacts of emerald ash borer eradication and tree mortality. In: Mastro V, Lance D, Reardon R, Parra G (eds) Proceedings of the emerald ash borer and Asian longhorned beetle research and technology development meeting, Pittsburgh, Pennsylvania, Oct 23–24 2007. USDA Forest Health Technology Enterprise Team, FHTET-2008-07, pp 14–15
- Havill NP, Montgomery ME, Yu G, Shiyake S, Caccone A (2006) Mitochondrial DNA from hemlock woolly adelgid (Hemiptera: Adelgidae) suggests cryptic speciation and pinpoints the source of the introduction to eastern North America. *Ann Entomol Soc Am* 99:195–203
- Hawkins BA, Marino PC (1997) The colonization of native phytophagous insects in North America by exotic parasitoids. *Oecologia* 112:566–571
- Herms DA (2003) Assessing management options for gypsy moth. *Pestic Outlook* 14:14–18
- Herms DA, Mattson WJ (1992) The dilemma of plants: to grow or defend. *Quart Rev Biol* 67:283–335
- Herms DA, Stone AK, Chatfield JA (2004) Emerald ash borer: the beginning of the end of ash in North America? In: Chatfield JA, Draper EA, Mathers HM, Dyke DE, Bennett PJ, Boggs JF (eds) Ornamental plants: annual reports and research reviews 2003, OARDC/OSU Extension Special Circular 193, pp 62–71
- Herms DA, Gandhi KJK, Cardina J, Long RP, Knight KS, Smith A, McCullough DG (2008) Impacts of emerald ash borer-induced gap formation on forest communities. In: Mastro V, Lance D, Reardon R, Parra G (eds) Proceedings of the emerald ash borer and Asian longhorned beetle research and technology development meeting, Pittsburgh, Pennsylvania, 23–24 Oct 2007, p 10
- Hollingsworth RG, Hain FP (1991) Balsam woolly adelgid (Homoptera: Adelgidae) and spruce-fir decline in the southern Appalachians: assessing pest relevance in a damaged ecosystem. *Fla Entomol* 74:179–187
- Horner JD, Gosz JR, Cates RG (1988) The role of carbon-based plant secondary metabolites in decomposition in terrestrial ecosystems. *Am Nat* 132:869–883
- Horvitz CC, Pascarella JB, McMann S, Freedman A, Hofstetter RH (1998) Functional roles of invasive non-indigenous plants in hurricane-affected subtropical hardwood forests. *Ecol Appl* 8:947–974
- Houston DR (1975) Beech bark disease: the aftermath forests are structured for a new outbreak. *J For* 73:660–663
- Houston DR (1987) Forest tree declines of past and present: current understanding. *Can J Plant Pathol* 9:349–360
- Houston DR (1994) Major new tree disease epidemics: beech bark disease. *Annu Rev Phytopathol* 32:75–87
- Hunter JC, Mattice JA (2002) The spread of woody exotics into the forests of a northeastern landscape, 1938–1999. *J Torr Bot Soc* 129:220–227

- Hutchens JJ Jr, Benfield EF (2000) Effects of forest defoliation by the gypsy moth on detritus processing in southern Appalachian streams. *Am Midl Nat* 143:397–404
- Irons JG III, Bryant JP, Oswood MW (1991) Effects of moose browsing on decomposition rates of birch leaf litter in a subarctic stream. *Can J Fish Aquat Sci* 48:442–444
- Jedlicka J, Vandermeer J, Aviles-Vazquez K, Barros O, Perfecto I (2004) Gypsy moth defoliation of oak trees and a positive response of red maple and black cherry: an example of indirect interaction. *Am Midl Nat* 152:231–236
- Jenkins JC, Aber JD, Canham CD (1999) Hemlock woolly adelgid impacts on community structure and N cycling rates in eastern hemlock forests. *Can J For Res* 29:630–645
- Johnson KS, Scriber JM, Nitao JK, Smitley DR (1995) Toxicity of *Bacillus thuringiensis* var. *kurstaki* to three non-target Lepidoptera in field studies. *Environ Entomol* 24:288–297
- Jones CG, Ostfeld RS, Richard MP, Schaubert EM, Wolff JO (1998) Chain reactions linking acorns to gypsy moth outbreaks and Lyme disease risk. *Science* 279:1023–1026
- Karnosky DF (1979) Dutch elm disease: a review of the history, environmental implications, control, and research needs. *Environ Conserv* 6:311–322
- Kasbohm JW, Vaughan MR, Kraus JG (1996) Effects of gypsy moth infestation on black bear reproduction and survival. *J Wildl Manage* 60:408–416
- Kellogg SK, Fink LS, Brower LP (2003) Parasitism of native luna moths, *Actias luna* (L.) (Lepidoptera: Saturniidae) by the introduced *Compsilura concinnata* (Meigen) (Diptera: Tachnidae) in central Virginia, their hyperparasitism by trigonalid wasp (Hymenoptera: Trigonalidae). *Environ Entomol* 32:1019–1027
- Kenis M, Auger-Rozenberg M, Roques A, Timms L, Péré C, Cock MJW, Settele J, Augustin S, Lopez-Caamonde C (2009) Ecological effects of invasive alien insects. *Biol Invasions* 11:21–45
- Kizlinski ML, Orwig DA, Cobb RC, Foster DR (2002) Direct and indirect ecosystem consequences of an invasive pest on forests dominated by eastern hemlock. *J Biogeogr* 29:1489–1503
- Knapp LB, Canham CD (2000) Invasion of old-growth forests by *Ailanthus altissima*: sapling growth and recruitment in canopy gaps. *J Torr Bot Soc* 127:307–315
- Koh LP, Dunn RR, Sodhi NS, Colwell RK, Proctor HC, Smith VS (2004) Species coextinctions and the biodiversity crisis. *Science* 305:1632–1634
- Krasny ME, DiGregorio LM (2001) Gap dynamics in Allegheny northern hardwood forests in the presence of beech bark disease and gypsy moth disturbances. *For Ecol Manage* 144:265–274
- Lautenschlager RA, Kircher CH, Podgwaite JD (1977) Effect of nucleopolyhedrosis virus on selected mammalian predators of the gypsy moth. USDA Forest Service, Northeastern Research Station, Res Pap NE-377
- Levine JM, Vila M, D'Antonio CM, Dukes JS, Grigulis K, Lavorel S (2003) Mechanisms underlying the impacts of exotic plant invasions. *Proc R Soc Lond B* 270:775–781
- Lewis GP, Likens GE (2007) Changes in stream chemistry associated with insect defoliation in a Pennsylvania hemlock-hardwoods forest. *For Ecol Manage* 238:199–211
- Liebold AM, MacDonald WL, Bergdahl D, Mastro VC (1995) Invasion by exotic forest pests: a threat to forest ecosystems. *For Sci Monogr* 30:1–49
- Lovett GM, Ruesink AE (1995) Carbon and nitrogen mineralization from decomposing gypsy moth frass. *Oecologia* 104:133–138
- Lovett GM, Christenson LM, Groffman PM, Jones CG, Hart JE, Mitchell MJ (2002) Insect defoliation and nitrogen cycling in forests. *Bioscience* 52:335–341
- Lovett GM, Canham CD, Arthur MA, Weathers KC, Fitzhugh RD (2006) Forest ecosystem responses to exotic pests and pathogens in eastern North America. *Bioscience* 56:395–405
- Marco A, Quilchano C, Blaustein AR (1999) Sensitivity to nitrate and nitrite in pond-breeding amphibians from the Pacific Northwest, USA. *Environ Toxicol Chem* 18:2836–2839
- Martinat PJ, Coffman CC, Dodge K, Cooper RJ, Whitmore RC (1988) Effect of diflubenzuron on the canopy arthropod community in a central Appalachian forest. *J Econ Entomol* 81:261–267
- Mattson WJ (1980) Herbivory in relation to plant nitrogen content. *Annu Rev Ecol Syst* 11:119–161
- Mattson WJ (1997) Exotic insects in North American forests: ecological systems forever altered. In: Britton KO (ed) Exotic pests of eastern forests, conference proceedings, Nashville, TN, 8–10 April 1997. USDA Forest Service and UN Exotic Pest Council
- Mattson WJ, Addy ND (1975) Phytophagous insects as regulators of forest primary production. *Science* 190:515–522
- Mattson WJ, Herms DA, Witter JA, Allen DA (1991) Woody plant grazing systems: North American outbreak folivores and their host plants. In: Baranchikov YO, Mattson WJ, Hain F, Payne TL (eds) Forest insect guilds: patterns of interaction with host trees. USDA General Technical Report NE-153, pp 53–84
- Mattson WJ, Niemelä P, Millers I, Inguanzo Y (1994) Immigrant phytophagous insects on woody plants in the United States and Canada: an annotated list. USDA Forest Service, General Technical Report NC-169
- Mattson WJ, Vanhanen H, Veteli T, Sivonen S, Niemelä P (2007) Few immigrant phytophagous insects on woody plants in Europe: legacy of the European crucible? *Biol Invasions* 9:957–974
- McCullough DG, Work TT, Cavey JF, Liebold AM, Marshall D (2006) Interceptions at nonindigenous plant pests at US ports of entry and border crossings over a 17-year period. *Biol Invasions* 8:611–630
- McEwan RW, Rieske LK, Arthur MA (2009) Potential interactions between invasive woody shrubs and the gypsy moth (*Lymantria dispar*), an invasive insect herbivore. *Biol Invasions* 11:1053–1058
- McShea WJ (2000) The influence of acorn production on annual variation in rodent and bird populations. *Ecology* 81:228–238
- McShea WJ, Healy WM, Devers P, Fearer T, Koch FH, Stauffer D, Waldon J (2007) Forestry matters: decline of oaks will impact wildlife in hardwood forests. *J Wildl Manage* 71:1717–1728

- Miller JC (1990) Field assessment of the effects of a microbial pest control agent on nontarget Lepidoptera. *Am Entomol* 36:135–139
- Morin RS, Liebhold AM, Tobin PC, Gottschalk KW, Luzader E (2007) Spread of beech bark disease in the eastern United States and its relationship to regional forest composition. *Can J For Res* 37:726–736
- Muzika RM, Liebhold AM (1999) Changes in radial increment of host and non-host tree species with gypsy moth defoliation. *Can J For Res* 29:1365–1373
- Muzika RM, Liebhold AM (2001) A critique of silvicultural approaches to managing defoliating insects in North America. *Agric For Entomol* 2:97–105
- Muzika RM, Twery MJ (1995) Regeneration in defoliated and thinned hardwood stands of north-central West Virginia. In: Gottschalk KW, Fosbroke SLC (eds) Proceedings, 10th Central Hardwood Forest Conference, Radnor, Pennsylvania, General Technical Report NE-197, U.S. Department of Agriculture, Forest Service, Northeastern Forest Experiment Station, pp 326–340
- Muzika RM, Grushecky ST, Liebhold AM, Smith RL (2004) Using thinning as a management tool for gypsy moth: the influence on small mammal abundance. *For Ecol Manage* 192:349–359
- Nagel TA, Svoboda M (2008) Gap disturbance regime in an old-growth *Fagus-Abies* forest in Dinaric Mountains, Bosnia-Herzegovina. *Can J For Res* 38:2728–2737
- Naidoo R, Lechowicz MJ (2001) Effects of gypsy moth on radial growth of deciduous trees. *For Sci* 47:338–348
- Niemelä P, Mattson WJ (1996) Invasion of North American forests by European phytophagous insects. *Bioscience* 46:741–753
- Nowacki GJ, Abrams MD (1992) Community, edaphic, and historical analysis of mixed oak forests of the Ridge and Valley Province in central Pennsylvania. *Can J For Res* 22:790–800
- Orwig DA (2002) Stand dynamics associated with chronic hemlock woolly adelgid infestations in southern New England. In: Reardon RC, Onken BP, Lashomb J (eds) Symposium on the Hemlock Woolly adelgid in eastern North America. New Jersey Agricultural Experiment Station Publication, New Brunswick, New Jersey, pp 36–47
- Orwig DA, Foster DR (1998) Forest response to the introduced hemlock woolly adelgid in southern New England, USA. *J Torr Bot Soc* 125:60–73
- Orwig DA, Kittredge D (2005) Silvicultural options for managing hemlock forests threatened by hemlock woolly adelgid. In: Onken B, Reardon R (eds) Third symposium on Hemlock Woolly adelgid in the eastern United States. Asheville, North Carolina, FHTET-2005-01, pp 106–118
- Orwig DA, Kizlinski ML (2002) Vegetation response following hemlock woolly adelgid infestation, hemlock decline, and hemlock salvage logging. In: Reardon RC, Onken BP, Lashomb J (eds) Proceedings: symposium on Hemlock Woolly adelgid in the eastern United States. New Jersey Agricultural Experiment Station Publication, New Brunswick, New Jersey, pp 106–117
- Orwig DA, Foster DR, Mausel DL (2002) Landscape patterns of hemlock decline in New England due to the introduced hemlock woolly adelgid. *J Biogeogr* 29:1475–1487
- Orwig DA, Cobb RC, D'Amato AW, Kizlinski ML, Foster DR (2008) Multi-year ecosystem response to hemlock woolly adelgid infestation in southern New England. *Can J For Res* 38:834–843
- Ostfield RS, Jones CG, Wolff JO (1996) Of mice and mast. *Bioscience* 46:323–330
- Owens AK, Moseley KR, McCay TS, Castleberry SB, Kilgo JC, Ford WM (2008) Amphibian and reptile community response to coarse woody debris manipulations in upland loblolly pine (*Pinus taeda*) forests. *For Ecol Manage* 256:2078–2083
- Parendes LA, Jones JA (2000) Role of light availability and dispersal in exotic plant invasion along roads and streams in the H. J. Andrews Experimental Forest, Oregon. *Conserv Biol* 14:64–75
- Parker IM, Simberloff D, Lonsdale WM, Goodell K, Wonham M, Kareiva PM, Williamson MH, Von Holle B, Moyle PB, Byers JE, Goldwasser L (1999) Impact: toward a framework for understanding the ecological effects of invaders. *Biol Invasions* 1:3–19
- Parry D (2009) Beyond Pandora's box: quantitatively evaluating non-target effects of parasitoids in classical biological control. *Biol Invasions* 11:47–58
- Pauley EF, Clebsch EEC (1990) Patterns of *Abies fraseri* regeneration in a Great Smokey Mountains spruce-fir forest. *Bull Torr Bot Club* 117:375–381
- Penrod KA, McCormick LH (1997) Hay-scented fern spore production following clearcutting. In: Pallardy SG, Cecich RA, Garrett HG, Johnson PS (eds) Proceedings of the 11th Central Hardwood Forest Conference, General Technical Report NC-188, U.S. Department of Agriculture, Forest Service, North Central Forest Experimental Station, St. Paul, Minnesota, pp 401
- Perry WB, Christiansen TA, Perry SA (1997) Response of soil and leaf litter microarthropods to forest application of diflubenzuron. *Ecotoxicology* 6:87–99
- Pimentel D, Lach L, Zuniga R, Morrison D (2000) Environmental and economic costs associated with non-indigenous species in the United States. *Bioscience* 50:53–65
- Poland T, McCullough D (2006) Emerald ash borer: invasion of the urban forest and the threat to North America's ash resource. *J For* 104:118–124
- Preisser EL, Elkinton JS (2008) Exploitative competition between invasive herbivores benefits a native host plant. *Ecology* 89:2671–2677
- Rabenold KN, Fauth PT, Goodner BW, Sadowski JA, Parker PG (1998) Response of avian communities to disturbance by an exotic insect in spruce-fir forests of the Southern Appalachians. *Conserv Biol* 12:177–189
- Rastall K, Kondo V, Strazanac JS, Butler L (2003) Lethal effects of biological insecticide applications on non-target lepidopterans in two Appalachian forests. *Environ Entomol* 32:1364–1369
- Rebek EJ, Herms DA, Smitley DR (2008) Interspecific variation in resistance to emerald ash borer (Coleoptera: Buprestidae) among North American and Asian ash (*Fraxinus* spp.). *Environ Entomol* 37:242–246
- Redman AM, Scriber JM (2000) Competition between the gypsy moth, *Lymantria dispar*, and the northern tiger swallowtail, *Papilio canadensis*: interactions mediated by

- host plant chemistry, pathogens, and parasitoids. *Oecologia* 125:218–228
- Runkle JR (1982) Patterns of disturbance in some old-growth mesic forests of eastern North America. *Ecology* 63:1533–1546
- Runkle JR (1990) Gap dynamics in an Ohio *Acer-Fagus* forest and speculation on the geography of disturbance. *Can J For Res* 20:632–641
- Runkle JR (1998) Changes in southern Appalachian canopy tree gaps sampled thrice. *Ecology* 79:1768–1780
- Runkle JR (2000) Canopy tree turnover in old-growth mesic forests of eastern North America. *Ecology* 81:554–567
- Runkle JR (2005) Twenty-four years of change in an old *Tsuga canadensis* woods affected by beech bark disease. *J Torr Bot Soc* 132:483–491
- Runkle JR (2007) Impacts of beech bark disease and deer browsing on the old-growth forest. *Am Midl Nat* 157:241–249
- Russell EWB, Davis RB (2001) Five centuries of changing forest vegetation in the northeastern United States. *Plant Ecol* 155:1–13
- Sample BE, Butler L, Zivkovich C, Whitmore RC, Reardon R (1996) Effects of *Bacillus thuringiensis* Berliner var. *kurstaki* and defoliation by the gypsy moth (*Lymantria dispar* (L.), Lepidoptera: Lymantriidae) on native arthropods in West Virginia. *Can Entomol* 128:573–592
- Scharenbroch BC, Bockheim JG (2008a) The effects of gap disturbance on nitrogen cycling and retention in late-successional northern hardwood-hemlock forests. *Biogeochemistry* 87:231–248
- Scharenbroch BC, Bockheim JG (2008b) Gaps and soil C dynamics in old growth northern hardwood-hemlock forests. *Ecosystems* 11:426–441
- Schowalter TD, Hargrove WW, Crossley DA Jr (1986) Herbivory in forested ecosystems. *Annu Rev Entomol* 31:177–196
- Schultz JC, Baldwin IT (1982) Oak leaf quality declines in response to defoliation by gypsy moth larvae. *Science* 217:149–151
- Schweitzer JA, Bailey JK, Hart SC, Wimp GM, Chapman SK, Whitham TG (2005) The interaction of plant genotype and herbivory decelerate leaf litter decomposition and alter nutrient dynamics. *Oikos* 110:133–145
- Scriber JM (2004) Non-target impacts of forest defoliator management options: decision for no spraying may have worse impacts on non-target Lepidoptera than *Bacillus thuringiensis* insecticides. *J Insect Conserv* 8:241–261
- Selfridge JA, Parry D, Boettner GH (2007) Parasitism of barrens buck moth *Hemileuca maia* Drury in early and late successional pine barrens habitats. *J Lepid Soc* 61:213–221
- Shang Z, He HS, Lytle DE, Shifley SR, Crow TR (2007) Modeling the long-term effects of fire suppression on central hardwood forests in Missouri Ozarks, using LANDIS. *For Ecol Manage* 242:776–790
- Simberloff D, Von Holle B (1999) Positive interactions of nonindigenous species: invasional meltdown? *Biol Invasions* 1:21–32
- Small MJ, Small CJ, Dreyer GD (2005) Changes in a hemlock-dominated forest following woolly adelgid infestation in southern New England. *J Torr Bot Soc* 132:458–470
- Snitzer JL, Boucher DH, Kyde KL (2005) Response of exotic invasive plant species to forest damage caused by Hurricane Isabel. In: Sellner KG (ed) Hurricane Isabel in perspective. Chesapeake Research Consortium, CRC Publication 05-160, Edgewater, Maryland, pp 209–214
- Stadler B, Müller T, Orwig D (2006) The ecology of energy and nutrient fluxes in hemlock forests invaded by hemlock woolly adelgid. *Ecology* 87:1792–1804
- Strand RM, Herms DA, Ayres MP, Kubiske ME, Kaufman MG, Walker ED, Pregitzer PS, Merritt RW (1999) Effects of atmospheric CO<sub>2</sub>, light availability, and tree species on the quality of leaf detritus as a resource for treehole mosquitoes. *Oikos* 84:277–283
- Strazanac JS, Butler L (eds) (2005) Long-term evaluation of the effects of *Bacillus thuringiensis kurstaki*, gypsy moth nucleohedrosis virus product Gypchek, and *Entomophaga maimaiga* on nontarget organisms in mixed broadleaf-pine forests in the central Appalachians. USDA Forest Service Forest Health Technology Publication FHTET-2004-14
- Swank WT, Waide JB, Crossley DA Jr, Todd RL (1981) Insect defoliation enhances nitrate export from forest ecosystems. *Oecologia* 51:297–299
- Sydnor TD, Bumgardner M, Todd A (2007) The potential economic impacts of emerald ash borer (*Agrilus planipennis*) on Ohio, U.S., communities. *Arbor Urban For* 33:48–54
- Thurber DK, McLain WR, Whitmore RC (1994) Indirect effects of gypsy moth defoliation on nest predation. *J Wildl Manage* 58:493–500
- Tingley MW, Orwig DW, Field R, Motzkin G (2002) Avian response to removal of a forest dominant: consequences of hemlock woolly adelgid infestations. *J Biogeogr* 29:1505–1516
- Townsend Peterson A, Sachetti-Pereira R (2004) Potential geographic distribution of *Anoplophora glabripennis* (Coleoptera: Cerambycidae) in North America. *Am Midl Nat* 151:170–178
- Tuomi J, Niemelä P, Haukioja E, Sirén S, Neuvonen S (1984) Nutrient stress: an explanation for plant anti-herbivore responses to defoliation. *Oecologia* 61:208–210
- Twery MJ (1990). Effects of defoliation by gypsy moth. In: Gottschalk KW, Twery MJ, Smith SI (eds) USDA gypsy moth research review, USDA Forest Service, Northeast Forest Experiment Station General Technical Report NE-146, pp 27–39
- Wagner DL (2007) Emerald ash borer threatens ash-feeding Lepidoptera. *News Lepid Soc* 49:10–11
- Walker ED, Kaufman MG, Ayres MP, Riedel MH, Merritt RW (1997) Effects of variation in quality of leaf detritus on growth of the eastern tree-hole mosquito, *Aedes triseriatus* (Diptera: Culicidae). *Can J Zool* 75:706–718
- Wallner WE (1996) Invasive pests ('biological pollutants') and US forests: whose problem, who pays? *Bull OEPP/EPPO Bull* 26:167–180
- Wargo PM (1977) *Armillariella mellea* and *Agrilus bilineatus* and mortality of defoliated oak trees. *For Sci* 23:485–492
- Webb SL (1999) Disturbance by wind in temperate-zone forests. In: Walker LR (ed) *Ecosystems of disturbed ground. Ecosystems of the world 16*. Elsevier, Amsterdam, The Netherlands, pp 187–222



- Webb JR, Cosby BJ, Deviney FA, Eshleman KN, Galloway JN (1995) Change in the acid–base status of an Appalachian mountain catchment following forest defoliation by the gypsy moth. *Water Air Soil Pollut* 85:535–540
- Webster CR, Nelson K, Wangen SR (2005) Stand dynamics of an insular population of an invasive tree, *Acer platanoides*. *For Ecol Manage* 208:85–99
- Weston PA, Desurmont G, Hoebeke ER (2007) *Viburnum* leaf beetle (Coleoptera: Chrysomelidae): biology, invasion history in North America, and management options. *Am Entomol* 53:96–101
- Witter JA, Ragenovich IR (1986) Regeneration of Fraser fir at Mt. Mitchell, North Carolina, after depredations by the balsam woolly adelgid. *For Sci* 32:585–594
- Work TT, McCullough DG (2000) Lepidopteran communities in two forest ecosystems during the first gypsy moth outbreaks in northern Michigan. *Environ Entomol* 29:884–900
- Work TT, McCullough DG, Cavey JG, Komsa R (2005) Arrival rate of nonindigenous insect species into the United States through foreign trade. *Biol Invasions* 7:323–332
- Yorks TE, Jenkins JC, Leopold DJ, Raynal DJ, Orwig DA (1999) Influences of eastern hemlock mortality on nutrient cycling. In: McManus KA, Shields KS, Souto DR (eds) *Proceedings: symposium on sustainable management of hemlock ecosystems in eastern North America*. 22–23 June, 100, Durham, NH, USDA Forest Service, General Technical Report NE-267, pp 126–133
- Zak DR, Groffman PM, Pregitzer KS, Christensen S, Tiedje JM (1990) The vernal dam: plant-microbe competition for nitrogen in northern hardwood forests. *Ecology* 71:651–656