

Direct observation of DNA knots using a solid-state nanopore

Plesa, Calin; Verschueren, Daniel; Pud, Sergii; van der Torre, Jaco; Ruitenberg, Justus W.; Witteveen, Menno J.; Jonsson, Magnus P.; Grosberg, Alexander Y.; Rabin, Yitzhak; Dekker, Cees

DOI

[10.1038/nnano.2016.153](https://doi.org/10.1038/nnano.2016.153)

Publication date

2016

Document Version

Accepted author manuscript

Published in

Nature Nanotechnology

Citation (APA)

Plesa, C., Verschueren, D., Pud, S., van der Torre, J., Ruitenberg, J. W., Witteveen, M. J., Jonsson, M. P., Grosberg, A. Y., Rabin, Y., & Dekker, C. (2016). Direct observation of DNA knots using a solid-state nanopore. *Nature Nanotechnology*, *11*(12), 1093-1097. <https://doi.org/10.1038/nnano.2016.153>

Important note

To cite this publication, please use the final published version (if applicable).
Please check the document version above.

Copyright

Other than for strictly personal use, it is not permitted to download, forward or distribute the text or part of it, without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license such as Creative Commons.

Takedown policy

Please contact us and provide details if you believe this document breaches copyrights.
We will remove access to the work immediately and investigate your claim.

Direct observation of DNA knots using a solid-state nanopore

Calin Plesa¹, Daniel Verschueren¹, Sergii Pud¹, Jaco van der Torre¹, Justus W. Ruitenberg¹, Menno J. Witteveen¹, Magnus P. Jonsson^{1,†}, Alexander Y. Grosberg², Yitzhak Rabin³, Cees Dekker^{1}*

¹Department of Bionanoscience, Kavli Institute of Nanoscience, Delft University of Technology, Lorentzweg 1, 2628 CJ Delft, The Netherlands.

²Department of Physics and Center for Soft Matter Research, New York University, 4 Washington Place, New York, NY 10003, USA.

³Department of Physics and Institute for Nanotechnology and Advanced Materials, Bar Ilan University, Ramat Gan 52900, Israel.

***Corresponding author.** E-mail: c.dekker@tudelft.nl

[†]Current address: Department of Science and Technology, Campus Norrköping, Linköping University, SE-60174 Norrköping, Sweden.

KEYWORDS: DNA, knots, nanopore

ABSTRACT: Long DNA molecules can self-entangle into knots. Experimental techniques to observe such DNA knots (primarily gel electrophoresis) are limited to bulk methods and circular molecules below 10 kilobase pairs in length. Here, we show that solid-state nanopores can be used to directly observe individual knots in both linear and circular single DNA molecules of arbitrary length. The DNA knots are observed as short spikes in the nanopore current traces of traversing DNA molecules and their detection is dependent on sufficiently high measurement resolution, which can be achieved using high-concentration LiCl buffers. We study the percentage of molecules with knots for DNA molecules of up to 166 kilobase pairs in length, and find that the knotting occurrence rises with the length of the DNA molecule, consistent with a constant knotting probability per unit length. Our experimental data compare favorably with previous simulation-based predictions for long polymers. From the translocation time of the knot through the nanopore, we estimate that the majority of the DNA knots are tight, with remarkably small sizes below 100 nm. In the case of linear molecules, we also observe that knots are able to slide out upon applying high driving forces (voltage).

It is well established that long polymers are subject to increasing self-entanglement as their length increases. DNA knots have been the focus of significant study in polymer physics^{1, 2} and, although they are ubiquitous in nature³, their exact role in biological processes is still under investigation^{4, 5, 6}. Despite their interest, knotting remains among the least understood properties of polymers due to a lack of both experimental techniques to observe them as well as rigorous theoretical approaches to describe and characterize them. Many open questions remain⁷, such as what determines the characteristic chain length beyond which knots become prevalent upon cyclization, knot localization^{8, 9}, the existence of metastable tight knots¹⁰, and many others.

A number of experimental techniques have been developed to study knots, particularly in DNA. Knots have been induced with high-electric fields¹¹, optical tweezers^{12, 13}, topoisomerase enzymes^{14, 15, 16}, DNA recombinases¹⁷, and through the cyclization of linear DNA molecules^{18, 19}. Electron microscopy¹⁴ and atomic force microscopy²⁰ have been used to image knots with excellent resolution but these techniques are limited to small molecules and low statistics. Optical techniques^{11, 12} have been used to introduce and study the behaviour of knots in DNA strands. Gel electrophoresis^{15, 16, 18, 19, 21}, the dominant tool used in knot studies, is a bulk technique where knots are trapped in circular molecules which are limited to lengths in the range of 10 kbp or lower.

Solid-state nanopores have emerged as an important tool with a large number of potential applications^{22, 23} at the crossroads of physics, biology and chemistry. Nanopores have provided a method to investigate various concepts in polymer physics such as polymer translocation through pores^{24, 25}, the Zimm relaxation time^{25, 26}, and the polymer capture process^{27, 28}. In nanopore sensing, biomolecules in an aqueous solution are placed into one of two reservoirs separated by a 20 nm thick membrane containing a nanometer scale pore (Figure 1). Subsequent application of

voltage using electrodes bathed in each reservoir produces an electric field and a resulting electrophoretic force on charged molecules such as DNA, causing them to be pulled through the nanopore. The presence of a molecule in the pore causes a blockade in the ionic current that (in high salt conditions) is proportional to the volume of the segment of the molecule in the pore. For long polymers, the translocation process is much faster than the typical relaxation time²⁹, which allows us to investigate the polymers before they have a chance to relax. This enables us to probe long DNA molecules for the presence of topological structures such as knots. Several theoretical papers have addressed some of the issues arising from the presence of knots in molecules translocating through nanopores^{30, 31, 32}. Here we report the direct experimental observation of knots in measurements on long dsDNA molecules that translocate through solid-state nanopores.

Ionic current signatures of DNA knots

Knots produce distinctive signatures in the current blockade of translocating DNA molecules. When a linear unknotted DNA molecule passes through the pore, it causes a blockade with magnitude I_l in the current level as shown in Figure 2a. If the molecule enters the pore in a folded configuration this produces a blockade level twice as high ($2I_l$) for the duration of the fold, as shown in Figure 2b. These folds occur primarily at the start of the translocation process and their probability of occurrence has been shown to increase as the capture point gets closer to the end of the molecule²⁸. High-resolution investigation of DNA translocation events reveals the presence of additional sharp blockage spikes with a high amplitude and very short duration occurring within a fraction of the events. Examples of such events are shown in Figure 2c-d,f. We observe such events in both linear DNA (Fig 2cd) and circular dsDNA (Fig 2f, Supplementary Fig 18cd). These spikes occur only within the DNA translocation events and

have an amplitude (on top of the DNA blockade level I_1) which is an integer multiple of $2I_1$. Such spikes can only be associated to two types of molecular configurations: an internal fold within the molecule (Supplementary Fig. 5ii), or a topologically constrained DNA knot (Supplementary Fig. 5i). The former configuration of local folds has extremely low rates of occurrence due to the nature of the translocation process, which can be seen as follows. The nanopore can be thought of as applying a point force to the polymer at the location of the nanopore, since the electric field strength is highest in the pore and strongly decreases as $1/r^2$ away from the pore³³. A long DNA molecule coils up into a polymer blob that has a large size (many hundreds of nm) compared to the nm-size nanopore. For translocation, a DNA end is pulled through the pore, thus disentangling the blob. The tension in the DNA strand resulting from the pulling action towards the pore propagates along the strand outwards from the nanopore, which pulls out any internal folds (since the velocity of the leading (captured) DNA is much higher than that of the lagging DNA (behind the fold)). Knots, on the other hand, cannot unfold due to the topological constraints imposed and are pulled towards and translocated through the nanopore. This argument that the observed blockades are due to knots as opposed to folds is strongly supported by the small size observed for the majority of knots, the higher-order $4I_1, 6I_1, \dots$ current blockades observed, the observation of knot sliding, and the differences in occurrence observed between linear and circular molecules. A detailed overview of all arguments supporting this conclusion can be found in Supplementary Section 2.

The observed current signatures allow us to probe the knotting probability, knot size, and knot position. The $2I_1$ additional magnitude of the spikes provides strong evidence for the occurrence of DNA knots, indicating the presence of 3 double-stranded DNA segments in total within the nanopore. Although the knot amplitude can be used to determine the number of dsDNA strands

inside the pore, the current signatures for the knots do not provide information about the crossing number of the knot. Different knot types (trefoil, figure-eight,...) have the same number of dsDNA strands when linearly stretched out (without changing the topology of the knot) and accordingly can have the same amount of dsDNA strands simultaneously inside the pore during translocation. As a consequence, our experimental results lead to statements encompassing many knot types. Most spikes that we observe consistently have an amplitude of $2I_1$ beyond the normal DNA blockage level. Occasionally, however, we observe events during which the additional current blockade has an amplitude of only I_1 (beyond the normal DNA blockage level), which may be due to the presence of replication forks, that occur due to the nature of the plasmid replication process, or due to simultaneous co-translocation of two different molecules. This latter phenomenon is only significant if either a high concentration of DNA is used or if there is a significant amount of smaller DNA fragments due to handling, and such events can easily be distinguished from the blockades produced by knots (Supplementary Section 2). For circular DNA molecules, DNA catenanes may be possible as well. Higher blockades ($4I_1$, $6I_1$, $8I_1$,...) as shown in Supplementary Figure 18, are observed in a small fraction of events and can be attributed to more complex knots, since these higher-order knot structures bring each time two additional dsDNA strands simultaneously into the pore. The fraction of these complex knots increases as a function of increasing polymer length³⁴.

In common nanopore experiments, knots are hard to observe since their small size (< 100 nm; see below) makes their observation very sensitive to the resolution of the solid-state nanopore measurement. Most nanopore experiments are carried out in 1M KCl^{25, 26, 27, 28}, where most observed knots have a duration (τ) of 15 μ s at 100 mV, indicating that their detection is at the edge of what is resolvable (Supplementary Section 4). For this reason the majority of the

experiments presented in this study are carried out in solutions of 2M or 4M LiCl, which as we have previously shown, can increase the translocation time of DNA by a factor of 7x or 10x respectively, relative to 1M KCl³⁵. Furthermore, the conductance blockades are also higher in 2M and 4M LiCl, which leads to a higher signal-to-noise-ratio (SNR) and allows for higher measurement bandwidths. Indeed, for 48.5 kbp DNA molecules at the same conditions, we observe fewer than 50% of the knots in 1M KCl relative to measurements in 2M LiCl solution, as further discussed below.

Knotting probability as a function of DNA length

We measured molecules of different length to study how the knotting probability scales as a function of DNA length, and we compare the observed rates of knot occurrence to previous simulation-based predictions. Four different DNA lengths were used: 2686 bp pUC19 linearized with XmnI, a 20678 bp linearized plasmid, 48502 bp lambda phage DNA, and 165648 bp phage T4 GT7 DNA. Figure 3 shows the measured knotting probabilities and their standard deviation for linear molecules as a function of DNA molecule length (Supplementary Section 1). The observed scaling can be fit with a simple model, described below, which is linear below ~40 kbp. For 2686 bp molecules in 4M LiCl we find a knotting probability of 1.8%, while for the 20678 bp molecules we observe $13.8 \pm 1.0\%$ of molecules with knots. For 48502 bp molecules we find $26.8 \pm 3.4\%$ of molecules to contain knots, while similar measurements carried out in 2M LiCl find a $24.4 \pm 4.6\%$ knotting occurrence (Supplementary Section 8). Since these molecules are linear, the knotting probabilities may be reduced due to the effect of knots slipping out, as discussed in detail further on. We fit the observed scaling of knotting occurrence with length using a model of the form $P = 1 - \exp(-N/N_0)$, where N is the length in bp, and N_0 is the random knotting length (details in Supplementary Section 3). We find a fitted value of 143 ± 5 kbp for

N_0 . As mentioned above, data taken in 1M KCl showed significantly lower knotting levels, an effect attributed to the limited resolution in these measurements. Nevertheless, it can provide a lower bound for the knot probability. These results provide the first experimental validation of the knotting abundances in long polymers.

The observed rates of knotting occurrence are co-plotted in Figure 3 alongside the simulation-based predictions of Deguchi and Tsurusaki³⁴ and Rybenkov et al¹⁸ (Supplementary Section 9). Neither of these simulations include complex knots and thus can only serve as a lower bound since the amount of more complex knots increases significantly as a function of the polymer length. For 2686, 20678, 48502, and 165648 bp DNA molecules these simulations predict knotting rates of 0.2%, 14%, 35%, and 80% respectively. Qualitatively we thus clearly see that the simulation-based predictions show the same overall trend as the experiments. Since our measurements are carried out in very high ionic strength conditions where the electrostatic screening is very strong, we expect the effective diameter to be very close to the physical diameter of DNA. This leads to higher knotting probabilities relative to those expected in the case of lower ionic strength regimes, such as those found at physiological conditions¹⁸.

DNA knot position, sliding, and slipping out

Since these measurements are carried out on linear molecules, we investigated the possibility of knots slipping out before being captured. Once a knot reaches the pore, translocation can only occur after the knotted strand has been bent to a size set by the pore diameter, i.e. on scales below the persistence length of dsDNA. If this process does not occur quickly enough, the low-friction DNA-DNA interactions could allow a knot to slip out^{12, 30, 36} by remaining at the pore entrance while the non-knotted DNA strand translocates through (Supplementary Fig. 6a). In

circular DNA molecules, however, the closed curve topology prevents such slipping out and knots are intrinsically trapped. We investigated the position of the observed knots in 20.7 kbp relaxed circular molecules where any knots contained inside would not be able to escape, although within a translocation event they might slip until the end. Figure 4 shows the normalized center position of the knots (τ_p/τ_{DNA}) in a 20 nm pore at 4M LiCl for applied voltages of 100 mV and 200 mV. We observe that knots occur at random positions for 100 mV, whereas at 200 mV they indeed occur with a strong preference for the end of the molecule that translocates through the pore last. We thus see a clear indication that the higher voltage causes knots to slide towards the end of the molecule. As a consequence, the numbers for the knot occurrence in linear molecules (Figure 3) provide a reasonable value at low applied voltage (100 mV), but should be treated as a lower bound to the equilibrium knotting levels at higher voltages, where a significant amount of knots may slip out. In order to quantify the amount of knots that slip out, we carried out experiments using lambda phage DNA (48.5 kbp). This molecule has a 12 bp complementary overhang allowing it to exist in both linear and circular forms depending on temperature and salt conditions³⁷. We heated solutions of lambda DNA and quickly cooled them while in 2M LiCl, to form mixed populations of linear and circular DNA from which to compare the knotting occurrence. Previously Rybenkov et al.¹⁸ observed no temperature dependence for the knotting probability, allowing us to attribute any differences between the knotting occurrence in the two populations to knots slipping out. We observed a 55% higher knotting occurrence in the circular molecules compared to the linear ones (Supplementary Section 8). Furthermore, we have tentative evidence suggesting that the knot sliding can sometimes be directly observed within the current trace of a translocation event, as shown in Supplementary Fig. 19. This effect is due to a slight increase in the current blockade caused by

the knot sitting against the pore mouth before being translocated, and is consistent with similar effects reported in literature for DNA docked onto the pore mouth^{38, 39, 40}. These results indicate that knots are able to slip out of linear DNA molecules during their translocation through nanopores.

DNA knot size

We estimated the size of knots from the time required for the knots to translocate through the nanopore. This estimate uses the average velocity of the DNA translocation, which is determined by dividing the known length of the molecules by their total translocation time. The total contour length of DNA within the knot can be determined by multiplying the knot size by the total number of strands in the knot (which is three for simple knots such as trefoil, figure-eight,...). Figure 5 shows the measured time duration of knots on a log scale for linear 20.7 kbp molecules at 100 mV. The vertical dashed line indicates the filtering distortion point ($2T_f$) while the solid line is a fit to the model described below. We estimate the size of the knots to be in the tens of nm at both 100 mV and 200 mV (Supplementary Section 6) applied voltage, assuming mean translocation velocities (Supplementary Section 7) of 872 nm/ms and 1937 nm/ms respectively, as shown on the top x-axis of Figure 5. Note that such a small knot size indicates that the DNA knots in these long DNA polymers are remarkably tight. The numbers may underestimate the size of the knots somewhat, especially for those which occur at the end of the translocation process where we know that the velocity is higher than average^{41, 42}. Measurements on circular versions of the same molecules in the same conditions reveal similar distributions of very tight knots (Supplementary Section 6). From all our data, we conclude that the majority of DNA knots have a size below 100 nm. The observation of these tight knots provides evidence for the occurrence of metastable tight knots which have been predicted to exist at equilibrium in long

polymers^{10, 43}. Given the nature of the translocation process, it remains to be determined how much these knots are being tightened relative to their equilibrium sizes although the similar numbers for the data at 100 vs 200 mV and linear vs circular DNA suggest that the tight knots are intrinsic.

The knot translocation duration distribution $P(\tau)$ is consistent with a model based on the previously proposed¹⁰ free energy penalty for knot formation, i.e., $P(\tau) = A \cdot \exp\left(-\frac{c_1}{\tau} - c_2\tau^{\frac{1}{3}}\right)$, where A is the normalization factor and the two terms in the exponential reflect the bending energy and confinement entropy contributions, respectively. Although this distribution was originally derived for the knot length and knot size, we adopt it here to characterize the knot translocation duration which is assumed to scale linearly with knot size. Fits were made to the data of 20678 bp DNA in 4M LiCl at 100 mV in 20 nm pores, see Figure 5. This yields $c_2 = 0.61 \pm 0.09 \mu\text{s}^{-1/3}$, while it is hard to obtain reliable estimates for c_1 , as fitted values range from 0.3 to 18 μs . Physically, coefficient c_1 controls the decay of probability for very small times τ , i.e., for very tight knots, and the large uncertainty in its magnitude is due to knots slipping out as well as the limited temporal resolution of the measurements⁴⁴. The tail of the distribution is consistent with the functional form $\exp(-c_2\tau^{1/3})$ which theoretically derives from DNA undulations inside the effective tube formed by the knot.

As an important control, we investigated biologically knotted circular DNA plasmids that were treated with Topoisomerase IV (Topo), an enzyme that is capable of removing knots from plasmids⁴⁵. As expected, we found that incubation with Topo led to a reduction in the amount of current spikes observed (Supplementary Section 12). Circular DNA (20.7kbp) was incubated with 4.1 $\mu\text{U}/\mu\text{L}$ Topo for 30 min, nicked, re-purified, and measured with a nanopore (4M LiCl,

20nm, 100mV). Analysis of the current spikes that correspond to simple knots ($2I_1$) revealed a decrease of 31-37% in the amount of events containing these spikes for the Topo-incubated DNA relative to the controls. Because the presence of DNA-bound Topo proteins (which causes additional high-current spikes, see SI) counteracts the unknotting effects of Topo, this number represents a lower-bound estimate for the unknotting activity of Topo in our conditions. This assay provides further and compelling evidence that the current spikes are due to knots, and shows that it is possible to assess enzymatic unknotting with this approach.

We have demonstrated that it is possible to detect knots in DNA molecules with solid-state nanopores and we have used this approach to estimate the knotting occurrence for a variety of DNA molecules with lengths far longer than previously possible. The single-molecule nature of the technique allows us to observe and analyze individual knots while still being able to generate population statistics. The measurements reveal that knots are capable of slipping out of linear molecules during translocation at high driving voltage, so care must be taken when comparing observed knotting rates to those predicted for molecules at equilibrium. Additionally, the knot translocation duration can be used to estimate the knot size, which is observed to be less than 100 nm for the majority of knots, although a further understanding of the translocation process is required to accurately relate this to the equilibrium knot size. From a nanopore applications perspective, efforts to sequence or detect DNA-bound proteins with nanopores will have to take into account the presence and effects of these knots⁴⁶. These results present a major step towards the ability to directly interrogate polymer knots and thus increase our understanding of this ubiquitous phenomenon.

REFERENCES

1. Sumners DW, Whittington SG. Knots in self-avoiding walks. *J Phys A: Math Gen* 1988, **21**(7): 1689.
2. Kawauchi A. *Survey on knot theory*. Springer, 1996.
3. Meluzzi D, Smith DE, Arya G. Biophysics of Knotting. *Annual Review of Biophysics* 2010, **39**(1): 349-366.
4. Staczek P, Higgins NP. Gyrase and Topo IV modulate chromosome domain size in vivo. *Mol Microbiol* 1998, **29**(6): 1435-1448.
5. Rodríguez-Campos A. DNA Knotting Abolishes in Vitro Chromatin Assembly. *J Biol Chem* 1996, **271**(24): 14150-14155.
6. Portugal J, Rodríguez-Campos A. T7 RNA Polymerase Cannot Transcribe Through a Highly Knotted DNA Template. *Nucleic Acids Res* 1996, **24**(24): 4890-4894.
7. Grosberg AY. A few notes about polymer knots. *Polym Sci Ser A* 2009, **51**(1): 70-79.
8. Metzler R, Hanke A, Dommersnes PG, Kantor Y, Kardar M. Equilibrium Shapes of Flat Knots. *Phys Rev Lett* 2002, **88**(18): 188101.
9. Orlandini E, Stella AL, Vanderzande C. The size of knots in polymers. *Physical Biology* 2009, **6**(2): 025012.
10. Grosberg AY, Rabin Y. Metastable Tight Knots in a Wormlike Polymer. *Phys Rev Lett* 2007, **99**(21): 217801.
11. Tang J, Du N, Doyle PS. Compression and self-entanglement of single DNA molecules under uniform electric field. *Proceedings of the National Academy of Sciences* 2011, **108**(39): 16153-16158.
12. Bao XR, Lee HJ, Quake SR. Behavior of Complex Knots in Single DNA Molecules. *Phys Rev Lett* 2003, **91**(26): 265506.

13. Arai Y *et al.* Tying a molecular knot with optical tweezers. *Nature* 1999, **399**(6735): 446-448.
14. Krasnow MA *et al.* Determination of the absolute handedness of knots and catenanes of DNA. *Nature* 1983, **304**(5926): 559-560.
15. Liu LF, Davis JL, Calendar R. Novel topologically knotted DNA from bacteriophage P4 capsids: studies with DNA topoisomerases. *Nucleic Acids Res* 1981, **9**(16): 3979-3989.
16. Trigueros S, Arsuaga J, Vazquez ME, Summers DW, Roca J. Novel display of knotted DNA molecules by two-dimensional gel electrophoresis. *Nucleic Acids Res* 2001, **29**(13): e67-e67.
17. Wasserman SA, Dungan JM, Cozzarelli NR. Discovery of a predicted DNA knot substantiates a model for site-specific recombination. *Science* 1985, **229**(4709): 171-174.
18. Rybenkov VV, Cozzarelli NR, Vologodskii AV. Probability of DNA knotting and the effective diameter of the DNA double helix. *Proceedings of the National Academy of Sciences* 1993, **90**(11): 5307-5311.
19. Shaw SY, Wang JC. Knotting of a DNA chain during ring closure. *Science* 1993, **260**(5107): 533-536.
20. Ercolini E *et al.* Fractal Dimension and Localization of DNA Knots. *Phys Rev Lett* 2007, **98**(5): 058102.
21. Wasserman SA, Cozzarelli NR. Biochemical topology: applications to DNA recombination and replication. *Science* 1986, **232**(4753): 951-960.
22. Haque F, Li J, Wu H-C, Liang X-J, Guo P. Solid-state and biological nanopore for real-time sensing of single chemical and sequencing of DNA. *Nano Today* 2013, **8**(1): 56-74.
23. Wanunu M. Nanopores: A journey towards DNA sequencing. *Physics of Life Reviews* 2012, **9**(2): 125-158.
24. Muthukumar M. Mechanism of DNA Transport Through Pores. *Annu Rev Biophys Biomol Struct* 2007, **36**(1): 435-450.

25. Storm AJ *et al.* Fast DNA Translocation through a Solid-State Nanopore. *Nano Lett* 2005, **5**(7): 1193-1197.
26. Plesa C, Cornelissen L, Tuijtel MW, Dekker C. Non-equilibrium folding of individual DNA molecules recaptured up to 1000 times in a solid state nanopore. *Nanotechnology* 2013, **24**(47): 475101.
27. Gershow M, Golovchenko JA. Recapturing and trapping single molecules with a solid-state nanopore. *Nat Nano* 2007, **2**(12): 775-779.
28. Mihovilovic M, Hagerty N, Stein D. Statistics of DNA Capture by a Solid-State Nanopore. *Phys Rev Lett* 2013, **110**(2): 028102.
29. Kantor Y, Kardar M. Anomalous dynamics of forced translocation. *Physical Review E* 2004, **69**(2): 021806.
30. Rosa A, Di Ventra M, Micheletti C. Topological Jamming of Spontaneously Knotted Polyelectrolyte Chains Driven Through a Nanopore. *Phys Rev Lett* 2012, **109**(11): 118301.
31. Huang L, Makarov DE. Translocation of a knotted polypeptide through a pore. *The Journal of Chemical Physics* 2008, **129**(12): 121107.
32. Suma A, Rosa A, Micheletti C. Pore Translocation of Knotted Polymer Chains: How Friction Depends on Knot Complexity. *ACS Macro Letters* 2015, **4**(12): 1420-1424.
33. Ando G, Hyun C, Li J, Mitsui T. Directly Observing the Motion of DNA Molecules near Solid-State Nanopores. *ACS Nano* 2012, **6**(11): 10090-10097.
34. Deguchi T, Tsurusaki K. A Statistical Study of Random Knotting Using the Vassiliev Invariants. *Journal of Knot Theory and Its Ramifications* 1994, **03**(03): 321-353.
35. Kowalczyk SW, Wells DB, Aksimentiev A, Dekker C. Slowing down DNA Translocation through a Nanopore in Lithium Chloride. *Nano Lett* 2012, **12**(2): 1038-1044.
36. Vologodskii A. Brownian Dynamics Simulation of Knot Diffusion along a Stretched DNA Molecule. *Biophys J* 2006, **90**(5): 1594-1597.

37. Wang JC, Davidson N. Thermodynamic and kinetic studies on the interconversion between the linear and circular forms of phage lambda DNA. *J Mol Biol* 1966, **15**(1): 111-123.
38. Carlsen AT, Zahid OK, Ruzicka J, Taylor EW, Hall AR. Interpreting the Conductance Blockades of DNA Translocations through Solid-State Nanopores. *ACS Nano* 2014, **8**(5): 4754-4760.
39. Rosenstein JK, Wanunu M, Merchant CA, Drndic M, Shepard KL. Integrated nanopore sensing platform with sub-microsecond temporal resolution. *Nat Meth* 2012, **9**(5): 487-492.
40. Kowalczyk SW, Dekker C. Measurement of the Docking Time of a DNA Molecule onto a Solid-State Nanopore. *Nano Lett* 2012, **12**(8): 4159-4163.
41. Plesa C, van Loo N, Ketterer P, Dietz H, Dekker C. Velocity of DNA during translocation through a solid state nanopore. *Nano Lett* 2015, **15**(1): 732-737.
42. Lu B, Albertorio F, Hoogerheide DP, Golovchenko JA. Origins and consequences of velocity fluctuations during DNA passage through a nanopore. *Biophys J* 2011, **101**(1): 70-79.
43. Dai L, Renner CB, Doyle PS. Metastable Tight Knots in Semiflexible Chains. *Macromolecules* 2014, **47**(17): 6135-6140.
44. Plesa C *et al.* Fast Translocation of Proteins through Solid State Nanopores. *Nano Lett* 2013, **13**(2): 658-663.
45. Deibler RW, Rahmati S, Zechiedrich EL. Topoisomerase IV, alone, unknots DNA in *E. coli*. *Genes Dev* 2001, **15**(6): 748-761.
46. Plesa C, Ruitenbergh JW, Witteveen MJ, Dekker C. Detection of Individual Proteins Bound along DNA Using Solid-State Nanopores. *Nano Lett* 2015, **15**(5): 3153-3158.

ACKNOWLEDGMENT

The authors would like to thank Cristian Micheletti, Marco Di Stefano, and Peter Virnau for discussions, Meng-Yue Wu for TEM drilling of nanopores, and Rini Joseph and Stefan W. Kowalczyk for early experiments. This work was supported by the Netherlands Organisation for Scientific Research (NWO/OCW), as part of the Frontiers of Nanoscience program, and by the European Research Council under research grant NanoforBio (no. 247072) and SynDiv (no. 669598), the Koninklijke Nederlandse Akademie van Wetenschappen (KNAW) Academy Assistants Program, and by the Wenner-Gren Foundations. Y.R. and A.Y.G. would like to acknowledge support from the US-Israel Binational Science foundation.

AUTHOR CONTRIBUTIONS

C.P., D.V., S.P., J.v.d.T, J.W.R, M.J.W., M.P.J. carried out the measurements; C.P. and D.V. analyzed experimental data; A.Y.G. and Y.R. provided theoretical interpretation; all authors discussed and interpreted results; C.P. and C.D. wrote the manuscript with input from all of the co-authors;

ADDITIONAL INFORMATION

Supplementary information is available in the online version of the paper. Reprints and permissions information is available online at www.nature.com/reprints.

Correspondence and requests for materials should be addressed to C.D.

COMPETING FINANCIAL INTERESTS

The authors declare no competing financial interests.




Figure 1 – A schematic illustration of a DNA molecule with a trefoil (3_1) knot translocating through a solid-state nanopore. This figure depicts a simplified schematic of the process for illustrative purposes. In reality, the knot may be located anywhere within the DNA blob, and most strands of the blob will be closer to the nanopore opening.




Figure 2 – Six example events for lambda DNA translocating through a 10 nm pore at 200 mV in 2M LiCl at 30kHz bandwidth. The molecular configuration attributed to each type of event are shown above each current trace in red. **a)** Current trace of an unfolded event. **b)** Current trace of an event with a fold at the start. **c)** Current trace of an event with an internal blockade which can be associated with a knot. The blockade has an additional amplitude of $2I_1$ on top of the I_1 blockade from the linear molecule, consistent with a total of 3 dsDNA strands that occupy the pore at the moment that the knot is translocating. **d)** Another event similar to *c* but with an

additional fold at the start. **e)** Current trace of an unfolded circular molecule translocating. **f)** Current trace of a circular molecule containing a knot. The blockade has an additional amplitude of $2I_1$ on top of the $2I_1$ blockade from the circular molecule, consistent with a total of 4 dsDNA strands in the pore while the knot is translocating.




Figure 3 – Percentage of events with knots as a function of DNA length for measurements carried out in 4M LiCl (red-squares) and 1M KCl (blue-circles) for linear dsDNA molecules in 20 nm pores. Errors are standard deviation, except for the data point for 166 kbp in 4M LiCl (Supplementary Section 3). The 1M KCl data provide a lower limit of the knot occurrence due to its low resolution. The dashed lines represent knot occurrence predictions based on simulations by Deguchi and Tsurusaki³⁴ (green) and Rybenkov et al.¹⁸ (magenta) (Supplementary Section 9). The solid line is a fit of $1 - \exp(-N/N_0)$ viewing localized knots as a Poisson process, with $N_0 = 143 \pm 5$ kbp (Supplementary Section 3).




Figure 4 – Normalized center position of knots observed in 20.7 kbp relaxed circular molecules in 4M LiCl at 100 mV (top) and 200 mV (bottom). $\tau_p/\tau_{DNA} = 0$ denotes the start of the translocation and $\tau_p/\tau_{DNA} = 1$ the end of the translocation event. As the voltage is increased, we see knots sliding towards the end of the molecule.




Figure 5 – Translocation duration (τ) of knots observed in 20.7 kbp linear molecules in 4M LiCl at 100 mV, plotted on a logarithmic time scale. Solid line is a fit to the model described in the main text, while the vertical dashed line indicates the filtering distortion point ($2T_r$). The top x-axis provides an estimated knot size scale based on the mean translocation velocity of each population. The majority of the knots have very short durations and thus small sizes, indicating that we are observing primarily tight knots.

Methods

Solid-state nanopores were fabricated and used as described previously⁴⁷. All buffers were pH 8 with 10 mM Tris and 1mM EDTA. Data analysis was carried out using custom Matlab scripts described in detail elsewhere⁴⁸. Lambda phage DNA and T4 GT7 DNA were purchased from Promega (Madison, WI) and Nippon Gene (Toyama, Japan) respectively. A 20.7 kbp plasmid was grown in XL10-Gold E.coli cells and midipreped. This plasmid was subsequently either linearized with BamHI or relaxed using the nt.BbvCI nickase. The resulting products were purified with phenol/chloroform and concentrated using ethanol precipitation. Mixed populations of both linear and circular Lambda DNA were formed by heating the DNA in 2M LiCl to 65°C for 5 min and then placing the solution on ice until measuring. The linear 2686 bp DNA molecules were pUC19 plasmids minipreped from E.coli cells and linearized with XmnI. Further methods can be found in the Supplementary Information.

REFERENCES for Methods

47. Janssen XJA *et al.* Rapid manufacturing of low-noise membranes for nanopore sensors by trans -chip illumination lithography. *Nanotechnology* 2012, **23**(47): 475302.
48. Plesa C, Dekker C. Data analysis methods for solid-state nanopores. *Nanotechnology* 2015, **26**(8): 084003.