

Directional Monotone Comparative Statics*

By

Anne-Christine Barthel

Department of Economics and Business
Randolph College
Lynchburg, VA, 24503, USA
abarthel@randolphcollege.edu

Tarun Sabarwal

Department of Economics
University of Kansas
Lawrence KS, 66045, USA
sabarwal@ku.edu

Abstract

Many questions of interest can be stated in terms of monotone comparative statics: if a parameter of a constrained optimization problem “increases,” when does its solution “increase” as well. This paper studies monotone comparative statics in different directions in finite-dimensional Euclidean space. The conditions on the objective function are ordinal and retain the same flavor as their counterparts in the standard theory. They can be naturally specialized to cardinal conditions, and to differential conditions using directional derivatives. Conditions on both the objective function and the constraint set do not require new binary relations or convex domains. The results allow flexibility to explore comparative statics with respect to the constraint set, with respect to parameters in the objective function, or both. Results from Quah (2007) are included as a special case. Several examples highlight applications of the results.

JEL Numbers: C60, C61, D10, D20, D40

Keywords: monotone comparative statics, directional single-crossing property, directional set order

First Draft: November 2012

This Version: April 27, 2015

*Part of this paper was written when Sabarwal was visiting Université Paris 1 Panthéon-Sorbonne as an invited professor. He is grateful for their warm welcome and hospitality.

1 Introduction

In economics, we are frequently interested in how solutions to a constrained optimization problem change when the economic environment changes. In many cases, the question of interest can be stated in terms of “monotone” comparative statics: if a parameter of the constrained optimization problem increases, when does its solution increase as well. Both the constraint set and the objective function may include parameters. For example, let X be a set, $f : X \rightarrow \mathbb{R}$, and A, B be subsets of X ordered by some relation, $A \sqsubseteq B$. When is it true that $A \sqsubseteq B \Rightarrow \arg \max_A f \sqsubseteq \arg \max_B f$? Intuitively, when is $\arg \max_A f$ increasing in A ? Or, more generally, $f : X \times T \rightarrow \mathbb{R}$, where T is a partially ordered set. When is it true that $A \sqsubseteq B$ and $t \preceq t' \Rightarrow \arg \max_A f(\cdot, t) \sqsubseteq \arg \max_B f(\cdot, t')$? Intuitively, when is $\arg \max_A f(\cdot, t)$ increasing in (A, t) ?

Milgrom and Shannon (1994) show that when X is a lattice¹ and \sqsubseteq is the standard lattice set order, denoted \sqsubseteq^{lso} , $\arg \max_A f(\cdot, t)$ is increasing in (A, t) in the standard lattice set order, if, and only if, for every $t \in T$, $f(\cdot, t)$ is quasisupermodular on X and f satisfies single crossing property on $X \times T$.² There are several appealing features of such lattice-theoretic monotone comparative statics results. For example, the sets X and A are not required to be convex and can be finite, the objective function f is not required to be differentiable or continuous, and the results apply even when there are multiple solutions to the optimization problem. Moreover, the notion of quasisupermodularity has a nice economic intuition in terms of complementarities: when X is a product space, when one component variable increases, the “marginal” benefit of another component variable goes up. Some of this standard theory is developed in Topkis (1978), Topkis (1979), LiCalzi and

¹Recall that a lattice is a partially ordered set in which every two points have a supremum and an infimum. For example, \mathbb{R}^N is a lattice, with the standard product partial order.

²Recall: $A \sqsubseteq^{lso} B$, if for every $a \in A, b \in B$, $a \wedge b \in A$ and $a \vee b \in B$. Moreover, $f : X \rightarrow \mathbb{R}$ is quasisupermodular, if for every $a, b \in X$, $f(a) \geq (>) f(a \wedge b) \implies f(a \vee b) \geq (>) f(b)$, and $f : X \times T \rightarrow \mathbb{R}$ satisfies single-crossing property on $X \times T$, if for every $a, b \in X$ with $a \succeq b$ and for every $t, t' \in T$ with $t' \succeq t$, $f(a, t) \geq (>) f(b, t) \implies f(a, t') \geq (>) f(b, t')$.

Veinott (1992), Veinott (1992), and Milgrom and Roberts (1994). These ideas have had many applications in developing the theory of supermodular games, submodular games, aggregative games, and comparing equilibria.³

A limitation of these results is that they do not apply to some basic economic problems in which constraint sets are not ordered in the standard lattice set order. Consider, for example, the standard budget set in consumer theory: $B(p, w) = \{x \in \mathbb{R}_+^N \mid p \cdot x \leq w\}$, where $p \in \mathbb{R}^N$, $p \gg 0$ is a price system, and wealth is $w > 0$. As is well-known, for $w < w'$, $B(p, w) \not\subseteq^{lso} B(p, w')$, and therefore, the standard lattice-based monotone comparative statics results cannot be applied directly to the theory of demand.

Quah (2007) develops monotone comparative statics results to include such problems.⁴ He considers $f : X \rightarrow \mathbb{R}$, where X is a convex sublattice of \mathbb{R}^N , and $i \in \{1, \dots, N\}$ is a direction in \mathbb{R}^N . His techniques include new binary relations, denoted $\Delta_i^\lambda, \nabla_i^\lambda$, a new set order, termed \mathcal{C}_i -flexible set order, and a new notion of \mathcal{C}_i -quasisupermodular function.⁵ In particular, if $w < w'$, then $B(p, w)$ is lower than $B(p, w')$ in the \mathcal{C}_i -flexible set order. A main result is: *arg max_A f is increasing in A in the \mathcal{C}_i -flexible set order, if, and only if, f is \mathcal{C}_i -quasisupermodular.* Moreover, a sufficient condition for f to be \mathcal{C}_i -quasisupermodular is that f is supermodular and i -concave.⁶

Quah (2007) uses some assumptions that are less typical in the standard theory of monotone comparative statics. The domain, X , of the objective function is assumed to be convex. This rules out discrete spaces; in particular, finite games and cases where

³Some of this can be seen in Bulow, Geanakoplos, and Klemperer (1985), Vives (1990), Milgrom and Roberts (1990), Milgrom and Roberts (1994), Zhou (1994), Amir (1996), Amir and Lambson (2000), Echenique (2002), Echenique (2004), Zimmer (2007), Roy and Sabarwal (2008), Roy and Sabarwal (2010), Roy and Sabarwal (2012), Acemoglu and Jensen (2013), Acemoglu and Jensen (2015), Monaco and Sabarwal (2015), and others.

⁴For additional development and generalizations, confer Quah and Strulovici (2009) and Quah and Strulovici (2012).

⁵Formal definitions are presented in appendix A.

⁶Intuitively, i -concave requires concavity in every direction u , where u is a vector with $u_i = 0$.

consumption of some goods is more naturally modeled as discrete, for example, automobiles and homes. Moreover, the notion of \mathcal{C}_i -quasisupermodular function uses the binary relations $\Delta_i^\lambda, \nabla_i^\lambda$ and convexity of domain in important ways, and it is less transparent than standard assumptions of quasisupermodularity and single-crossing property. Furthermore, the binary relations $\Delta_i^\lambda, \nabla_i^\lambda$ have some counter-intuitive properties; they are non-commutative, and their outcomes are not necessarily comparable in the underlying order in \mathbb{R}^N . Finally, the framework does not include parameterized objective functions, which rules out cases involving the effect of actions of others on a given agent's payoff, for example, cases with public goods, externalities from other consumers or producers, and strategic effects based on actions of other players.

This paper presents an extension of the theory of monotone comparative statics. The basic framework is as follows. Consider a sublattice X of \mathbb{R}^N , T a partially ordered set, $f : X \times T \rightarrow \mathbb{R}$, and $i \in \{1, \dots, N\}$. A main result is: *arg max $_A f(\cdot, t)$ is increasing in (A, t) in the i -directional set order, if, and only if, for every $t \in T$, $f(\cdot, t)$ is i -quasisupermodular and satisfies i -single crossing property on X , and f satisfies basic i -single crossing property on $X \times T$.* These terms are defined more concretely in the next section, but intuitively, increase in the i -directional set order formalizes the idea of increase in the i -th direction in \mathbb{R}^N . In this characterization, X is not required to be convex and there is no use of the binary relations $\Delta_i^\lambda, \nabla_i^\lambda$. The framework allows for parameter effects in the objective function. The new properties i -quasisupermodular, i -single crossing, and basic i -single crossing retain the same flavor as their counterparts in the standard theory of monotone comparative statics. The i -directional set order is a reformulation of \mathcal{C}_i -flexible set order to align more closely with the spirit of monotone methods; on convex sublattices, it coincides with \mathcal{C}_i -flexible set order and this helps subsume results in Quah (2007).⁷

The main result is explored in several directions. It is extended to apply to all di-

⁷Notably, this strand of the literature does not address comparison of budget sets with respect to price effects. That turns out to be a more complex problem. Antoniadu (2007) and Mirman and Ruble (2008) present some results for that case.

rections i , it is specialized to consider comparative statics with respect to A only or to t only, and the ordinal nature of the properties allows for increasing transformations of the objective function to also respect the same characterization. Sufficient conditions are explored as well; in particular, Quah’s sufficient conditions of supermodular and i -concave remain sufficient here. Furthermore, the characterization here has a natural formulation in terms of cardinal assumptions – i -supermodular and i -increasing differences, and in turn, this has a natural formulation in terms of differential conditions using directional derivatives. Several examples highlight applications of these results.

The paper proceeds as follows. Section 2 formalizes the constrained optimization problem, the set orders, and properties on objective function. The main results on directional monotone comparative statics are presented next. The main results are explored further in subsections formalizing sufficient conditions and differential conditions. Section 3 presents several applications of the main result. Appendix A presents some connections to Quah (2007) and appendix B includes details of some proofs.

2 Constrained Optimization

Recall that a lattice⁸ is a partially ordered set in which every two elements, a and b , have a supremum in the set, denoted $a \vee b$, and an infimum in the set, denoted $a \wedge b$. The supremum and infimum operations are with respect to the partial order. In this paper, we work with finite-dimensional Euclidean space, represented by \mathbb{R}^N . This is a lattice in the standard product order on \mathbb{R}^N , denoted, as usual, by \leq ,⁹ and in this order, for $a, b \in \mathbb{R}^N$, $a \wedge b = (\min \{a_1, b_1\}, \dots, \min \{a_N, b_N\})$ and $a \vee b = (\max \{a_1, b_1\}, \dots, \max \{a_N, b_N\})$. A subset X of a lattice is a sublattice, if for every a and b in X , their supremum in the overall lattice, $a \vee b$, is in X , and their infimum in the overall lattice, $a \wedge b$, is in X .

Let X be a sublattice of \mathbb{R}^N , (T, \preceq) be a partially ordered set, $f : X \times T \rightarrow \mathbb{R}$, A

⁸This paper uses standard lattice terminology. See, for example, Topkis (1998).

⁹For $a, b \in \mathbb{R}^N$, $a \leq b$ means that for every $i = 1, \dots, N$, $a_i \leq b_i$.

be a subset of X , and consider the constrained maximization problem $\max_A f(\cdot, t)$. We are interested in how $\arg \max_A f(\cdot, t)$ changes with (A, t) . As the set of maximizers is not necessarily a singleton, this involves a comparison of sets.

2.1 Set Orders

There are several set orders on subsets of a lattice (confer Topkis (1998)). Two of the more common ones are as follows. Consider a sublattice X of \mathbb{R}^N , and subsets A and B of X . A **is lower than B in the standard lattice set order**, denoted $A \sqsubseteq^{lso} B$, if for every $a \in A, b \in B$, it follows that $a \wedge b \in A$ and $a \vee b \in B$. A **is lower than B in the weak lattice set order**, denoted $A \sqsubseteq^{wlsO} B$, if for every $a \in A$, there is $b \in B$ such that $a \leq b$, and for every $b \in B$, there is $a \in A$ such that $a \leq b$.¹⁰ Moreover, another set order is of interest when we are considering increases in a particular component of vectors: for $i \in \{1, 2, \dots, N\}$, A **is lower than B in the i -weak lattice set order**, denoted $A \sqsubseteq_i^{wlsO} B$, if for every $a \in A$, there is $b \in B$ such that $a_i \leq b_i$, and for every $b \in B$, there is $a \in A$ such that $a_i \leq b_i$. As is well-known and easy to check: $A \sqsubseteq^{lso} B \implies A \sqsubseteq^{wlsO} B \implies A \sqsubseteq_i^{wlsO} B$.

The standard results in monotone comparative statics typically use the standard lattice set order, but that order cannot compare some of the constraint sets of interest here, and therefore, to expand comparability of sets, we work with the following weakenings of the standard lattice set order. Let X be a sublattice of \mathbb{R}^N , A and B be subsets of X , and $i \in \{1, 2, \dots, N\}$. A **is lower than B in the i -directional set order**, denoted, $A \sqsubseteq_i^{dso} B$, if for every $a \in A$ and $b \in B$ with $a_i > b_i$, there is $v = s(b - a \wedge b)$ for some $s \in [0, 1]$ such that $a + v \in B$ and $b - v \in A$.¹¹ In this definition, notice that the vector

¹⁰In all the set orders considered here, when convenient, we may say A is lower than B equivalently as B is higher than A .

¹¹The i -directional set order is a reformulation of the \mathcal{C}_i -flexible set order in Quah (2007). The definition here retains the spirit of monotone methods, does not require X to be convex, and there is no use of the operators $\Delta_i^\lambda, \nabla_i^\lambda$. Some comparisons to Quah (2007) are presented in appendix A.

v satisfies $v \geq 0$, and therefore, $a \leq a + v$ and $b - v \leq b$. Moreover, when $a \geq b$, this condition is satisfied trivially, and therefore, a non-trivial application of this order is when $a_i > b_i$ and $a \not\leq b$. Figure 1 shows this idea graphically. For intuition, we can consider the two-good discretized consumption space, and budget-type sets given by the green and the purple lines. For these sets to be ranked in the 1-directional set order, for each a in the lower set and b in the higher set with $a_1 > b_1$, there is $v = s(b - a \wedge b)$ such that $a + v$ is in the higher set and $b - v$ is in the lower set. Similarly, say that A is lower than B in




Figure 1: i -Directional Set Order

the directional set order, denoted $A \sqsubseteq^{dso} B$, if for every $i \in \{1, 2, \dots, N\}$, A is lower than B in the i -directional set order.

Proposition 1. *Let X be a sublattice of \mathbb{R}^N and A, B be non-empty subsets of X .*

- (1) $A \sqsubseteq^{lso} B \implies A \sqsubseteq_i^{dso} B \implies A \sqsubseteq_i^{wlsso} B$, for each $i \in \{1, 2, \dots, N\}$, and
- (2) $A \sqsubseteq^{lso} B \implies A \sqsubseteq^{dso} B \implies A \sqsubseteq^{wlsso} B$.

Proof. The proof of (1) is similar to that of (2). To prove (2), suppose first that $A \sqsubseteq^{lso} B$. Fix $i \in \{1, 2, \dots, N\}$, $a \in A$, and $b \in B$ with $a_i > b_i$. Let $s = 1$. Then $b - v = b - 1(b - a \wedge b) = a \wedge b \in A$ and $a + v = a + 1(a \vee b - a) = a \vee b \in B$. Thus, for every $i \in \{1, 2, \dots, N\}$, $A \sqsubseteq_i^{dso} B$, whence $A \sqsubseteq^{dso} B$. Now suppose $A \sqsubseteq^{dso} B$. Fix $a \in A$. As B is non-empty, let $b \in B$. If $a \leq b$, then we are done. Otherwise, there is i such that $a_i > b_i$. In this case, there is $v = s(b - a \wedge b)$ for some $s \in [0, 1]$ such that $a + v \in B$.

Moreover, $v \geq 0$ implies $a \leq a + v$. The proof is similar for the other case: $b \in B$ implies there is $a \in A$ such that $a \leq b$. ■

As shown in this proposition, the i -directional set order is weaker than the standard lattice set order and stronger than the i -weak lattice set order. Similarly, the directional set order is weaker than the standard lattice set order and stronger than the weak lattice set order. One benefit of the i -directional set order is that it can order budget sets for different levels of wealth, whereas the standard lattice set order cannot.

Example 1-1 (Walrasian budget sets). Let $X = \mathbb{R}_+^N$, $N \geq 2$, $p \gg 0$, and $w > 0$. The Walrasian budget set at (p, w) is given by $B(p, w) = \{x \in \mathbb{R}_+^N \mid p \cdot x \leq w\}$. We know that in the standard lattice set order when $w < w'$, $B(p, w) \not\sqsubseteq^{lso} B(p, w')$, but these budget sets are comparable in the directional set order: $w < w' \implies B(p, w) \sqsubseteq^{dso} B(p, w')$, as follows. Fix $i \in \{1, 2, \dots, N\}$, $a \in B(p, w)$ and $b \in B(p, w')$ with $a_i > b_i$. If $p \cdot (a \vee b) \leq w'$, let $s = 1$, and therefore, $v = b - a \wedge b$. In this case, $b - v = a \wedge b \in B(p, w)$, and $a + v = a \vee b \in B(p, w')$. Moreover, if $p \cdot b \leq w$, let $s = 0$, and so, $v = 0$. In this case, $b - v = b \in B(p, w)$, and $a + v = a \in B(p, w')$. In the other cases, let $s \in \left[\frac{p \cdot b - w}{p \cdot (b - a \wedge b)}, \frac{w' - p \cdot a}{p \cdot (b - a \wedge b)} \right] \subset [0, 1]$, and therefore, $v = s(b - a \wedge b)$. In this case, $p \cdot (b - v) \leq w$ and $p \cdot (a + v) \leq w'$. Consequently, $a + v \in B(p, w')$ and $b - v \in B(p, w)$, as desired.

Example 1-2 (Two-good discretized Walrasian budget sets). In the two-good case, the directional set order can be used to order budget sets with discrete consumption. Consider two goods, each consumed in integer amounts. Let $X = \mathbb{Z}_+^2$, $p = (p_1, p_2) \gg 0$, and $w > 0$. The (discretized) Walrasian budget set at (p, w) is given by $B(p, w) = \{x \in \mathbb{Z}_+^2 \mid p \cdot x \leq w\}$. Consider $w < w'$ and suppose p_1 divides $w' - w$ and p_2 divides $w' - w$. In this case, $w < w' \implies B(p, w) \sqsubseteq^{dso} B(p, w')$, as follows. Fix $i = 1$. Let $a \in B(p, w)$ and $b \in B(p, w')$ with $a_1 > b_1$. As above, if $p \cdot (a \vee b) \leq w'$, let $s = 1$, and if $p \cdot b \leq w$, let $s = 0$. Notice that these cases include the case where $a \geq b$. So suppose $a_1 > b_1$ and $a_2 \not\geq b_2$. Then $b - a \wedge b = (0, b_2 - a_2) > 0$, and $p \cdot (b - a \wedge b) = p_2(b_2 - a_2)$. Let $s = \frac{w' - w}{p \cdot (b - a \wedge b)} = \frac{w' - w}{p_2(b_2 - a_2)}$ and $v = s(b - a \wedge b)$. Notice

that $b - v = (b_1, b_2 - s(b_2 - a_2)) = (b_1, b_2 - \frac{w' - w}{p_2}) \in \mathbb{Z}_+^2$, because p_2 divides $w' - w$. Thus $B(p, w) \sqsubseteq_1^{dso} B(p, w')$. Similarly, $B(p, w) \sqsubseteq_2^{dso} B(p, w')$, whence $B(p, w) \sqsubseteq^{dso} B(p, w')$.

When there are three or more discrete goods, the discretized Walrasian budget set is not necessarily comparable in the directional set order. Consider $X = \mathbb{Z}_+^3$, $p = (1, 1, 1)$, $w = 1$, $w' = 2$, and $B(p, w) = \{x \in \mathbb{Z}_+^3 \mid p \cdot x \leq 1\}$ and $B(p, w') = \{x \in \mathbb{Z}_+^3 \mid p \cdot x \leq 2\}$. Let $i = 1$, $a = (1, 0, 0) \in B(p, w)$, and $b = (0, 1, 1) \in B(p, w')$. Then $a_1 > b_1$, and for $s \in [0, 1]$ consider $v = s(b - a \wedge b)$. It is easy to check that for $s = 0$, $b - v \notin B(p, w)$, for $s = 1$, $a + v \notin B(p, w')$, and for $s \in (0, 1)$, $b - v \notin \mathbb{Z}_+^3$. Thus, $B(p, w) \not\sqsubseteq_1^{dso} B(p, w')$.

This does not imply that other sets in higher dimensions are not comparable in the directional set order. For example, consider $A = \{(0, 0, 0), (1, 0, 0), (0, 1, 0), (0, 0, 1)\}$ and $B = \{(0, 2, 0), (1, 1, 0), (0, 1, 1), (1, 1, 1)\}$. In this case, $A \not\sqsubseteq^{lso} B$, because for $a = (1, 0, 0)$ and $b = (0, 2, 0)$, $a \vee b = (1, 2, 0) \notin B$. But it is easy to check that for $i = 1, 2, 3$, $A \sqsubseteq_i^{dso} B$, and therefore, $A \sqsubseteq^{dso} B$.

Additional classes of sets comparable in the i -directional set order can be derived in a manner analogous to Quah (2007). One such class is presented in appendix B.

2.2 Objective Function

Let X be a sublattice of \mathbb{R}^N , $f : X \rightarrow \mathbb{R}$, and $i \in \{1, 2, \dots, N\}$. The function f is **i -quasisupermodular on X** , if for every $a, b \in X$ with $a_i > b_i$, $f(a) \geq (>) f(a \wedge b) \implies f(a \vee b) \geq (>) f(b)$. In this definition, notice that when $a \geq b$, these conditions are satisfied trivially. Therefore, non-trivial application of this definition is when $a_i > b_i$ and $a \not\geq b$. The graphical intuition is the same as in the standard notion of a quasisupermodular function, as shown in figure 2. In other words, if the tradeoff between a and $a \wedge b$ is favorable (in the sense that $f(a) \geq f(a \wedge b)$ or $f(a) > f(a \wedge b)$), then the tradeoff remains favorable at $a \vee b$ and b , in the same sense. Indeed, recall the definition of a quasisupermodular function: f is **quasisupermodular on X** , if for every $a, b \in X$




Figure 2: i -Quasisupermodular on X

$f(a) \geq (>) f(a \wedge b) \implies f(a \vee b) \geq (>) f(b)$. It is easy to check that for every i , f is i -quasisupermodular on X , if, and only if, f is quasisupermodular on X .

Another useful property is the following. Let X be a sublattice of \mathbb{R}^N , $f : X \rightarrow \mathbb{R}$, and $i \in \{1, 2, \dots, N\}$. The function f satisfies **i -single crossing property on X** , if for every $a, b \in X$ with $a_i > b_i$, and for every $v \in \{s(b - a \wedge b) \mid s \in \mathbb{R}, s \geq 0\}$ such that $a + v, b + v \in X$, $f(a) \geq (>) f(b) \implies f(a + v) \geq (>) f(b + v)$. In this definition, notice that $v \geq 0$, and $v_i = 0$. Moreover, when $a \geq b$, these conditions are satisfied trivially. Therefore, non-trivial application of this property is when $a_i > b_i$ and $a \not\geq b$. Figure 3 presents a graphical idea.




Figure 3: i -Single Crossing Property on X

Notice that the black arrow is $(b - a \wedge b)$ and the red arrow is (translated) $s(b - a \wedge b)$. Intuitively, this property says that if the tradeoff between a and b is initially favorable (in the sense that $f(a) \geq f(b)$ or $f(a) > f(b)$), then it remains favorable when we move in the direction $b - a \wedge b$. This intuition is similar to that of the standard single crossing property. In particular, as $v = s(b - a \wedge b)$ satisfies $v \geq 0$ and $v_i = 0$, we may reformulate i -single crossing property as follows: for every $a, b \in X$ with $a_i > b_i$, and for every $v \in \{s(b - a \wedge b) \mid s \geq 0\}$ such that $a + v, b + v \in X$, $f(a_i, a_{-i}) \geq (>) f(b_i, b_{-i}) \implies f(a_i, a_{-i} + v_{-i}) \geq (>) f(b_i, b_{-i} + v_{-i})$. This reformulation captures the flavor of the standard single crossing property as follows. For a, b with $a_i > b_i$, if $f(a_i, a_{-i}) \geq (>) f(b_i, b_{-i})$, then when we increase a_{-i} and b_{-i} by a non-negative $v_{-i} = [s(b - a \wedge b)]_{-i}$, the tradeoff remains favorable. Similarly, f satisfies **directional single crossing property on X** , if for every $i \in \{1, 2, \dots, N\}$, f satisfies i -single crossing property on X .

In order to consider parameterized objective functions, let X be a sublattice of \mathbb{R}^N , (T, \preceq) be a partially ordered set, $f : X \times T \rightarrow \mathbb{R}$, and $i \in \{1, 2, \dots, N\}$. The function f satisfies **basic i -single crossing property on $X \times T$** , if for every $a, b \in X$ with $a_i > b_i$, and for every $t, t' \in T$ with $t' \succeq t$, $f(a, t) \geq (>) f(b, t) \implies f(a, t') \geq (>) f(b, t')$. The function f satisfies **basic directional single crossing property on $X \times T$** , if for every $i \in \{1, 2, \dots, N\}$, f satisfies basic i -single crossing property on $X \times T$. For convenience of reference, the word “basic” is used in basic i -single crossing property on $X \times T$ to distinguish this definition from that for i -single crossing property on X . It is easy to check that *if f satisfies basic directional single crossing property on $X \times T$, then f satisfies (standard) single crossing property in $(x; t)$.*¹²

¹²For every $a, b \in X$ with $a \geq b$ and for every $t, t' \in T$ with $t' \succeq t$, $f(a, t) \geq (>) f(b, t) \implies f(a, t') \geq (>) f(b, t')$.

2.3 Directional Monotone Comparative Statics

Some of the main results in this paper concern conditions on f that yield monotone comparative statics. This is formalized as follows. Let X be a sublattice of \mathbb{R}^N , (T, \preceq) be a partially ordered set, $f : X \times T \rightarrow \mathbb{R}$, and $i \in \{1, 2, \dots, N\}$. The function f satisfies ***i -directional monotone comparative statics on $X \times T$*** , if for every A, B subset of X , and for every t, t' in T , $A \sqsubseteq_i^{dso} B$ and $t \preceq t' \implies \arg \max_A f(\cdot, t) \sqsubseteq_i^{dso} \arg \max_B f(\cdot, t')$. In other words, f satisfies *i -directional monotone comparative statics* formalizes the idea that $\arg \max_A f(\cdot, t)$ is increasing in (A, t) in the i -directional set order. Similarly, f satisfies ***i -directional monotone comparative statics on $X \times T$*** , if for every $i \in \{1, 2, \dots, N\}$, f satisfies *i -directional monotone comparative statics on $X \times T$* .

Theorem 1. *Let X be a sublattice of \mathbb{R}^N , (T, \preceq) be a partially ordered set, $f : X \times T \rightarrow \mathbb{R}$, and $i \in \{1, 2, \dots, N\}$. The following are equivalent.*

- (1) *f satisfies i -directional monotone comparative statics on $X \times T$.*
- (2) *For every $t \in T$, $f(\cdot, t)$ is i -quasisupermodular and satisfies i -single crossing property on X , and f satisfies basic i -single crossing property on $X \times T$.*

Proof. Suppose first that (2) holds. Let $A \sqsubseteq_i^{dso} B$ and $t \preceq t'$. Let $a \in \arg \max_A f(\cdot, t)$, $b \in \arg \max_B f(\cdot, t')$, and $a_i > b_i$. Then there is $v = s(b - a \wedge b)$ for some $s \in [0, 1]$ such that $a + v \in B$ and $b - v \in A$.

As case 1, suppose $s = 1$. Then $a \wedge b = b - b + a \wedge b = b - v \in A$, and $a \vee b = a + s(a \vee b - a) = a + v \in B$. As $a \in \arg \max_A f(\cdot, t)$, it follows that $f(a, t) \geq f(a \wedge b, t)$, and then i -quasisupermodularity on X implies $f(a \vee b, t) \geq f(b, t)$, and then basic i -single crossing property on $X \times T$ implies $f(a \vee b, t') \geq f(b, t')$. As $b \in \arg \max_B f(\cdot, t')$, it follows that $a + v = a \vee b \in \arg \max_B f(\cdot, t')$. Therefore, $f(a \vee b, t') = f(b, t')$. In particular, $f(a \vee b, t') \not\geq f(b, t')$, and again i -quasisupermodularity implies $f(a, t') \not\geq f(a \wedge b, t')$, and then basic i -single crossing property on $X \times T$ implies $f(a, t) \not\geq f(a \wedge b, t)$. Consequently, $f(a, t) \leq f(a \wedge b, t)$, and it follows that $b - v = a \wedge b \in \arg \max_A f(\cdot, t)$.

As case 2, suppose $s < 1$. Then $a \in \arg \max_A f(\cdot, t)$ and $b - v \in A$ imply $f(a, t) \geq f(b - v, t)$. Moreover, when looking at the i -th component, $a_i > b_i \geq (b - v)_i$, because $v = s(b - a \wedge b) \geq 0$. Applying i -single crossing property on X to a and $b - v$, with the directional vector $w = \frac{s}{1-s} [(b - v) - a \wedge (b - v)]$ implies $f(a + w, t) \geq f(b - v + w, t)$. Notice that $v = s(b - a \wedge b) = s[(b - v) - a \wedge b] + sv = s[(b - v) - a \wedge (b - v)] + sv$, and therefore, $v = \frac{s}{1-s} [(b - v) - a \wedge (b - v)] = w$. In other words, $f(a + v, t) \geq f(b, t)$, and then basic i -single crossing property on $X \times T$ implies $f(a + v, t') \geq f(b, t')$, whence $a + v \in \arg \max_B f(\cdot, t')$. Thus, $f(a + v, t') = f(b, t')$, whence $f(a + v, t') \not\geq f(b, t')$, or equivalently, $f(a + w, t') \not\geq f(b - v + w, t')$ and then using i -single crossing property on X , $f(a, t') \not\geq f(b - v, t')$, and then using basic i -single crossing property on $X \times T$, $f(a, t) \not\geq f(b - v, t)$. Thus, $b - v \in \arg \max_A f(\cdot, t)$, as desired.

In the other direction, suppose f satisfies i -directional monotone comparative statics on $X \times T$. Let's first see that for every t , $f(\cdot, t)$ is i -quasisupermodular on X . Fix t , and a, b with $a_i > b_i$. Form the sets $A = \{a, a \wedge b\}$ and $B = \{b, a \vee b\}$. Notice that $A \sqsubseteq_i^{dso} B$. (Consider $a \in A$ and $b \in B$. Let $v = b - a \wedge b$. Then $a + v = a \vee b \in B$ and $b - v = a \wedge b \in A$. The other cases are satisfied vacuously, because in those cases the i -th component of the element from A is not greater than the i -th component of the element from B .)

Suppose $f(a, t) \geq f(a \wedge b, t)$. Then $a \in \arg \max_A f(\cdot, t)$. Suppose to the contrary that $f(a \vee b, t) < f(b, t)$. Then $\arg \max_B f(\cdot, t) = \{b\}$. Applying f satisfies i -directional monotone comparative statics to (A, t) and (B, t) , there is $s \in [0, 1]$ such that $a + s(a \vee b - a) \in \arg \max_B f(\cdot, t) = \{b\}$. But the i -th component of $a + s(a \vee b - a)$ is a_i which is strictly greater than b_i , a contradiction. Therefore, $f(a \vee b, t) \geq f(b, t)$, as desired.

Now suppose $f(a, t) > f(a \wedge b, t)$. Then $\{a\} = \arg \max_A f(\cdot, t)$. Suppose to the contrary that $f(a \vee b, t) \leq f(b, t)$. Then $b \in \arg \max_B f(\cdot, t)$. By i -directional monotone comparative statics, there is $s \in [0, 1]$ such that $b - s(b - a \wedge b) \in \arg \max_A f(\cdot, t) = \{a\}$. But the i -th component of $b - s(b - a \wedge b)$ is b_i which is strictly less than a_i , a contradiction.

Therefore, $f(a \vee b, t) > f(b, t)$, as desired.

Let's now check that for every t , $f(\cdot, t)$ satisfies i -single crossing property on X . Fix t , and $a, b \in X$ with $a_i > b_i$. Fix $v = s(b - a \wedge b)$ with $s \geq 0$ such that $a + v, b + v \in X$. Before we proceed further, consider the following calculations. Let $y = b + v$, and let $u = y - a \wedge y = a \vee y - a$. Notice that $u = y - a \wedge y = y - a \wedge b = (1 + s)(b - a \wedge b)$. This implies that $v = s(b - a \wedge b) = \frac{s}{1+s}u$. Let $s' = \frac{s}{1+s} \in [0, 1]$ and write $v = s'u$. In particular, $y - s'(y - a \wedge y) = y - v$, and $a + s'(a \vee y - a) = a + v$. Now let $A = \{a, y - v\}$ and $B = \{y, a + v\}$. Then $A \sqsubseteq_i^{dso} B$, because for $a \in A$, and $y \in B$, there is $s' \in [0, 1]$, as above such that $a + s'(a \vee y - a) = a + v \in B$ and $y - s'(y - a \wedge y) = y - v \in A$. The other comparisons are vacuously true, because when considering the i -th components, $(y - v)_i \leq y_i = b_i < a_i \leq (a + v)_i$.

Suppose $f(a, t) \geq f(b, t) = f(y - v, t)$. Then $a \in \arg \max_A f(\cdot, t)$. Suppose to the contrary that $f(a + v, t) < f(b + v, t) = f(y, t)$. Then $\{y\} = \arg \max_B f(\cdot, t)$. As f satisfies i -directional monotone comparative statics on $X \times T$, there is $\hat{s} \in [0, 1]$ such that $a + \hat{s}(a \vee y - a) \in \arg \max_B f(\cdot, t) = \{y\}$. But considering the i -th components, $(a + \hat{s}(a \vee y - a))_i = a_i > b_i = y_i$, a contradiction. Thus $f(a + v, t) \geq f(b + v, t)$, as desired.

Now suppose $f(a, t) > f(b, t) = f(y - v, t)$. Then $\{a\} = \arg \max_A f(\cdot, t)$. Suppose to the contrary that $f(a + v, t) \leq f(b + v, t) = f(y, t)$. Then $y \in \arg \max_B f(\cdot, t)$. As f satisfies i -directional monotone comparative statics, there is $\hat{s} \in [0, 1]$ such that $y - \hat{s}(y - a \wedge y) \in \arg \max_A f(\cdot, t) = \{a\}$. But considering the i -th components, $(y - \hat{s}(y - a \wedge y))_i = y_i = b_i < a_i$, a contradiction. Thus $f(a + v, t) > f(b + v, t)$, as desired.

Finally, let's check that f satisfies basic i -single crossing property in $X \times T$. Fix a, b with $a_i > b_i$, and fix $t' \succeq t$. Let $A = \{a, b\}$. Then $A \sqsubseteq_i^{dso} A$. Suppose $f(a, t) \geq f(b, t)$. Then $a \in \arg \max_A f(\cdot, t)$. As f satisfies i -directional monotone comparative statics on $X \times T$, there is $s \in [0, 1]$ such that $a + v = a + s(b - a \wedge b) \in \arg \max_A f(\cdot, t')$. Notice that $(a + s(b - a \wedge b))_i = a_i > b_i$, and therefore, $a + v = a$, whence $f(a, t') \geq f(b, t')$.

Now suppose $f(a, t) > f(b, t)$. Then $\{a\} = \arg \max_A f(\cdot, t)$. Suppose to the contrary that $f(a, t') \leq f(b, t')$. Then $b \in \arg \max_A f(\cdot, t')$. By i -directional monotone comparative statics, there is $s \in [0, 1]$ such that $b - v = b - s(b - a \wedge b) \in \arg \max_A f(\cdot, t) = \{a\}$, a contradiction. Thus, $f(a, t') > f(b, t')$. ■

Some implications of this theorem are formalized in the following corollaries.

Corollary 1. *Let X be a sublattice of \mathbb{R}^N , (T, \preceq) be a partially ordered set, and $f : X \times T \rightarrow \mathbb{R}$. The following are equivalent.*

- (1) *f satisfies directional monotone comparative statics on $X \times T$.*
- (2) *For every $t \in T$, $f(\cdot, t)$ is quasisupermodular and satisfies directional single crossing property on X , and f satisfies basic directional single crossing property on $X \times T$.*

Proof. For this equivalence, notice that f satisfies directional monotone comparative statics on $X \times T$ means that for every $i \in \{1, 2, \dots, N\}$, f satisfies i -directional monotone comparative statics on $X \times T$, which is equivalent to, for every $i \in \{1, 2, \dots, N\}$, for every $t \in T$, $f(\cdot, t)$ is i -quasisupermodular and satisfies i -single crossing property on X , and f satisfies basic i -single crossing property on $X \times T$, and this is equivalent to (2). ■

Corollary 2. *Let X be a sublattice of \mathbb{R}^N , (T, \preceq) be a partially ordered set, $f : X \times T \rightarrow \mathbb{R}$, and $i \in \{1, \dots, N\}$.*

- (1) *If f satisfies i -directional monotone comparative statics on $X \times T$, then*

$$A \sqsubseteq_i^{dso} B \text{ and } t \preceq t' \implies \arg \max_A f(\cdot, t) \sqsubseteq_i^{wls} \arg \max_B f(\cdot, t').$$

- (2) *If f satisfies directional monotone comparative statics on $X \times T$, then*

$$A \sqsubseteq^{dso} B \text{ and } t \preceq t' \implies \arg \max_A f(\cdot, t) \sqsubseteq^{wls} \arg \max_B f(\cdot, t').$$

Proof. Statement (1) follows from relations between i -directional set order and i -weak lattice set order (proposition 1). For statement (2), suppose f satisfies directional monotone comparative statics on $X \times T$. Consider $A \sqsubseteq^{dso} B$ and $t \preceq t'$. Then for every $i \in \{1, 2, \dots, N\}$, $A \sqsubseteq_i^{dso} B$, and by the theorem, for every $i \in \{1, 2, \dots, N\}$, $\arg \max_A f(\cdot, t) \sqsubseteq_i^{dso} \arg \max_B f(\cdot, t')$, whence $\arg \max_A f(\cdot, t) \sqsubseteq^{dso} \arg \max_B f(\cdot, t')$, and consequently, $\arg \max_A f(\cdot, t) \sqsubseteq^{wls} \arg \max_B f(\cdot, t')$. ■

In other words, under (1), f satisfies i -directional monotone comparative statics on $X \times T$ implies that when $A \sqsubseteq_i^{dso} B$ and $t \preceq t'$, then no matter which maximizer of $f(\cdot, t)$ we take from A , we can find a maximizer of $f(\cdot, t')$ from B that is larger in the i -th component, and symmetrically, no matter which maximizer of $f(\cdot, t')$ we take from B , we can find a maximizer of $f(\cdot, t)$ from A that is smaller in the i -th component. In particular, when the set of maximizers is a singleton, we conclude that the solution to the optimization problem is increasing in the i -th component, in the standard order in the real numbers.

Similarly, f satisfies directional monotone comparative statics on $X \times T$ implies that when $A \sqsubseteq^{dso} B$ and $t \preceq t'$, then no matter which maximizer of $f(\cdot, t)$ we take from A , we can find a larger maximizer of $f(\cdot, t')$ from B , and symmetrically, no matter which maximizer of $f(\cdot, t')$ we take from B , we can find a smaller maximizer of $f(\cdot, t)$ from A . In particular, when the set of maximizers is a singleton, we conclude that the solution to the optimization problem is increasing in the standard vector order in \mathbb{R}^N .

The framework in theorem 1 can be specialized naturally to the case of non-parameterized objective functions. Let X be a sublattice of \mathbb{R}^N , $f : X \rightarrow \mathbb{R}$, and $i \in \{1, 2, \dots, N\}$. The function f satisfies ***i -directional monotone comparative statics on X*** , if for every A, B subset of X , $A \sqsubseteq_i^{dso} B \implies \arg \max_A f \sqsubseteq_i^{dso} \arg \max_B f$. In other words, f satisfies i -directional monotone comparative statics on X formalizes the idea that $\arg \max_A f(\cdot)$ is increasing in A in the i -directional set order.

Corollary 3. *Let X be a sublattice of \mathbb{R}^N , $f : X \rightarrow \mathbb{R}$, and $i \in \{1, \dots, N\}$.*

The following are equivalent.

- (1) *f satisfies i -directional monotone comparative statics on X*
- (2) *f is i -quasisupermodular and satisfies i -single crossing property on X*

Proof. Apply theorem with singleton $T = \{t\}$. ■

Similarly, say that f satisfies ***directional monotone comparative statics on X*** , if for every $i \in \{1, 2, \dots, N\}$, f satisfies i -directional monotone comparative statics on X .

It follows immediately that f satisfies directional monotone comparative statics on X , if, and only if, f is quasisupermodular and satisfies directional single crossing property on X .

Theorem 1 can be used to inquire separately about comparative statics with respect to the parameter in the objective function, holding fixed the constraint set. In this case, the condition i -single crossing property on X may be dropped, as follows.

Corollary 4. *Let X be a sublattice of \mathbb{R}^N , A be a subset of X , (T, \preceq) be a partially ordered set, $f : X \times T \rightarrow \mathbb{R}$, and $i \in \{1, 2, \dots, N\}$.*

If f is i -quasisupermodular on X and satisfies basic i -single crossing property on $X \times T$, then $t \preceq t' \implies \arg \max_A f(\cdot, t) \sqsubseteq_i^{dso} \arg \max_A f(\cdot, t')$.

Proof. Follow the proof in the corresponding direction in theorem 1, setting $s = 0$ and note that i -directional set order is reflexive. ■

In this corollary, A is an arbitrary subset of X . Therefore, under the conditions in this corollary, for an arbitrary constraint set A , as long as the set of maximizers is nonempty, i -directional monotone comparative statics holds with respect to the parameter. (Of course, if the set of maximizers is empty, i -directional monotone comparative statics holds trivially.)

Finally, the ordinal nature of the conditions in theorem 1 implies that i -directional (and directional) monotone comparative statics property is preserved under increasing transformations of the objective function. This is useful in applications.

Corollary 5. *Let X be a sublattice of \mathbb{R}^N , (T, \preceq) be a partially ordered set, $f, g : X \times T \rightarrow \mathbb{R}$, and $i \in \{1, 2, \dots, N\}$. Suppose g is a strictly increasing transformation of f .*

f satisfies i -directional (respectively, directional) monotone comparative statics on $X \times T$, if, and only if, g satisfies i -directional (respectively, directional) monotone comparative statics on $X \times T$.

Proof. If f satisfies i -directional monotone comparative statics on $X \times T$, then f is

i -quasisupermodular and satisfies i -single crossing property on X , and satisfies basic i -single crossing property on $X \times T$. As these properties are ordinal, g satisfies these as well, and another application of the theorem yields the result. The other direction is similar. Moreover, the proof for directional monotone comparative statics is similar. ■

2.4 Sufficient Conditions

Quah (2007) shows that when X is a convex sublattice (a sublattice that is also a convex set) of \mathbb{R}^N , if $f : X \rightarrow \mathbb{R}$ is supermodular and i -concave, then $\arg \max_A f$ is increasing in A in the \mathcal{C}_i -flexible set order. In particular, if f is supermodular and concave, then this condition is satisfied for every i . This is useful, because supermodular and concave are conditions that are easy to check.

We show that these conditions are also sufficient for f to satisfy i -single crossing property on X . Therefore, we can use the same conditions here, apply them to some additional potentially discrete problems, and extend them naturally to include parameterized objective functions, as follows.

Let X be a sublattice of \mathbb{R}^N , $f : X \rightarrow \mathbb{R}$, and $u \in \mathbb{R}^N, u \neq 0$. The function f **is (relatively) concave in direction** u , if for every $a \in X$, the function $f(a + su)$, when viewed as a real-valued function of a real variable s , is a concave function relative to its domain in the real numbers. It is easy to check that f is (relatively) concave on X ,¹³ if, and only if, for every $u \in \mathbb{R}^N, u \neq 0$, f is (relatively) concave in direction u .

For $i \in \{1, 2, \dots, N\}$, f is **(relatively) i -concave on** X , if for every $u \in \mathbb{R}^N \setminus \{0\}$ with $u_i = 0$, f is (relatively) concave in direction u , and f is **(relatively) directionally concave on** X , if for every $i \in \{1, 2, \dots, N\}$, f is (relatively) i -concave on X . Notice that if f is (relatively) concave on X , then f is directionally concave on X .

¹³With the standard definition, $f(\alpha x + (1 - \alpha)y) \geq \alpha f(x) + (1 - \alpha)f(y)$, with $\alpha \in [0, 1]$ and with the quantifier “relative” applied to mean the points are in the domain of f , as usual.

Theorem 2. Let X be a sublattice of \mathbb{R}^N , (T, \preceq) be a partially ordered set, $f : X \times T \rightarrow \mathbb{R}$, and $i \in \{1, 2, \dots, N\}$.

If for every $t \in T$, $f(\cdot, t)$ is i -supermodular and (relatively) i -concave on X , and f satisfies basic i -single crossing property on $X \times T$, then f satisfies i -directional monotone comparative statics on $X \times T$.

Proof. Suppose for every $t \in T$, $f(\cdot, t)$ is i -supermodular and (relatively) i -concave on X , and f satisfies basic i -single crossing property on $X \times T$. It is sufficient to show that for every $t \in T$, $f(\cdot, t)$ satisfies i -single crossing property on X and then invoke theorem 1. To do so, fix $t \in T$, $a, b \in X$ with $a_i > b_i$, and $v = s(b - a \wedge b)$ with $s \geq 0$ such that $a + v, b + v \in X$.

Consider the following computations. Let $b' = b + v$, $a' = a + v$ and $u = a \vee b' - a'$. It is easy to check that $(a \vee b') - v = a \vee (b + v) - v = a \vee b$, and therefore, $u = a \vee b - a = b - a \wedge b$. Consequently, $v = su$. Moreover, notice that $u_i = 0$ and $a \vee b' = a' + u = a + (1 + s)u$.

Now, i -concavity in direction u implies that $f(a', t) - f(a \vee b', t) = f(a \vee b' - u, t) - f(a \vee b', t) \geq f(a \vee b' - u - su, t) - f(a \vee b' - su, t) = f(a, t) - f(a \vee b, t)$, and i -supermodularity implies $f(a \vee b', t) - f(b', t) \geq f(a \vee b, t) - f(b, t)$. Consequently, $f(a', t) - f(b', t) = f(a', t) - f(a \vee b', t) + f(a \vee b', t) - f(b', t) \geq f(a, t) - f(a \vee b, t) + f(a \vee b, t) - f(b, t) = f(a, t) - f(b, t)$. It follows that $f(a, t) \geq (>) f(b, t) \Rightarrow f(a', t) \geq (>) f(b', t)$, as desired. ■

The following corollaries follow immediately.

Corollary 6. Let X be a sublattice of \mathbb{R}^N , (T, \preceq) be a partially ordered set, and $f : X \times T \rightarrow \mathbb{R}$.

If for every $t \in T$, $f(\cdot, t)$ is supermodular and (relatively) directionally concave on X , and f satisfies basic directional single crossing property on $X \times T$, then f satisfies directional monotone comparative statics on $X \times T$.

Proof. The hypothesis implies that for every $i \in \{1, \dots, N\}$, for every $t \in T$, $f(\cdot, t)$ is i -supermodular and (relatively) i -concave on X , and f satisfies basic i -single crossing

property on $X \times T$, and the theorem then shows that for every $i \in \{1, \dots, N\}$, f satisfies i -directional monotone comparative statics on $X \times T$, as desired. ■

Corollary 7. *Let X be a sublattice of \mathbb{R}^N and $f : X \rightarrow \mathbb{R}$.*

(1) *If f is i -supermodular and (relatively) i -concave on X , then f satisfies i -directional monotone comparative statics on X .*

(2) *If f is supermodular and (relatively) directionally concave on X , then f satisfies directional monotone comparative statics on X .*

(3) *If f is supermodular and (relatively) concave on X , then f satisfies directional monotone comparative statics on X .*

Proof. Apply the previous theorem with singleton $T = \{t\}$. ■

Moreover, corollary 5 implies that in each of these sufficient conditions, if g is a strictly increasing transformation of f , then g also satisfies the corresponding i -directional (or directional) monotone comparative statics.

2.5 Differential Conditions

An appealing feature of the different single crossing properties defined here is that they are closely aligned to their counterparts in the standard theory. In particular, they possess natural extensions to cardinal properties and can also be formulated in terms of differential conditions in a manner similar to the standard case.

Consider the following cardinal property naturally suggested by the i -single crossing property on X . Let X be a sublattice of \mathbb{R}^N , $f : X \rightarrow \mathbb{R}$, and $i \in \{1, 2, \dots, N\}$. The function f satisfies ***i -increasing differences on X*** , if for every $a, b \in X$ with $a_i > b_i$, and for every $v \in \{s(b - a \wedge b) \mid s \geq 0\}$ such that $a + v, b + v \in X$, $f(a) - f(b) \leq f(a + v) - f(b + v)$. As earlier, when $a \geq b$, $v = 0$, and this condition is satisfied trivially. Nontrivial application of this definition is when $a_i > b_i$ and $a \not\geq b$. Similarly, f satisfies ***directional increasing differences on X*** , if for every $i \in \{1, 2, \dots, N\}$, f satisfies

i -increasing differences on X . It is easy to check that if f satisfies i -increasing differences on X , then f satisfies i -single crossing property on X , and it follows immediately that if f satisfies directional increasing differences on X , then f satisfies directional single crossing property on X .

Recall that in the standard theory, f satisfies (standard) increasing differences on \mathbb{R}^N , if, and only if, f satisfies increasing differences for every pair of component indices i, j with $i \neq j$. Thus, f satisfies increasing differences on \mathbb{R}^N , if, and only if, f is supermodular. Moreover, assuming differentiability, f is supermodular, if, and only if, every pair of cross-partials is nonnegative (for every $i \neq j$, $\frac{\partial^2 f}{\partial x_i \partial x_j} \geq 0$). The notion of i -increasing differences can be characterized similarly, using directional derivatives, as follows.

Notice that for $u \in \mathbb{R}^N$, if we let $a = b + u$, then $b - a \wedge b = (b - a)_+ = (-u)_+$. Say that a function $f : X \rightarrow \mathbb{R}$ satisfies i -increasing differences (u) on X , if for every $b \in X$, $u \in \mathbb{R}^N$ with $u_i > 0$, for every $s \geq 0$, such that $b+u, b+s(-u)_+, b+u+s(-u)_+ \in X$, $f(b+u) - f(b) \leq f(b+u+s(-u)_+) - f(b+s(-u)_+)$. Notice that for $u \geq 0$, $(-u)_+ = 0$, and this condition is satisfied trivially. Therefore, nontrivial application of this definition is when $u_i > 0$ and $u \not\geq 0$. It is easy to check that f satisfies i -increasing differences on X , if, and only if, f satisfies i -increasing differences (u) on X . This recasts i -increasing differences in terms of differences in f based on changes in direction u (where $u_i > 0$). Figure 4 presents the graphical intuition.




Figure 4: Cross Partial Directional Derivatives

The graphical intuition suggests a potential “cross partial” characterization based on directions u and $(-u)_+$. This is achieved as follows. Let X be a sublattice of \mathbb{R}^N , $f : X \rightarrow \mathbb{R}$, and $i \in \{1, 2, \dots, N\}$. Say that f satisfies *i -increasing differences (*) on X* , if for every $b \in X$, $u \in \mathbb{R}^N$ with $u_i > 0$, and for every $\sigma \geq 0$, $f(b + \sigma u + s(-u)_+) - f(b + s(-u)_+)$ is (weakly) increasing in s . As earlier, we consider only points $b + \sigma u + s(-u)_+, b + s(-u)_+ \in X$. As shown in appendix B, *f satisfies i -increasing differences (u) on X , if, and only if, f satisfies i -increasing differences (*) on X .*

These formulations show that *i -increasing differences on X is equivalent to i -increasing differences (*) on X* . A benefit of this equivalence is that the condition *i -increasing differences (*) on X* has the same mathematical structure as the one used to show that a supermodular function can be characterized by the sign of its cross-partials (confer Topkis (1978)). The only difference is that this definition uses a more general vector u whereas supermodularity uses the basis vectors. This connection can be seen more clearly as follows.

Recall the definition of a directional derivative. Let X be an open set in \mathbb{R}^N , $b \in X$ and $u \in \mathbb{R}^N$, and suppose $f : X \rightarrow \mathbb{R}$ is continuously differentiable. The *directional derivative of f at b in the direction u* is $D_u f(b) = \lim_{\sigma \rightarrow 0} \frac{f(b + \sigma u) - f(b)}{\sigma}$. Recall from the standard theory of supermodular functions (confer Topkis (1978), page 310, for the submodular case) that if u^i is the i -th basis vector, then a function f is supermodular on X (assuming X is an open set and a sublattice in \mathbb{R}^N , and f is twice continuously differentiable), if, and only if, for all $b \in X$, for all $i, j \in \{1, 2, \dots, N\}$ with $i \neq j$, and for all $\sigma \geq 0$, $f(b + \sigma u^i) - f(b)$ is (weakly) increasing in the j -th component (that is, in direction u^j). This is equivalent to: for all $b \in X$, for all $j \neq i$, $D_{u^i} f(b)$ is (weakly) increasing in the j -th component (that is, in direction u^j), which is further equivalent to: for all $b \in X$, for all $j \neq i$, $D_{u^j} D_{u^i} f(b) \geq 0$. Using the same logic yields the following result.

Proposition 2. *Let X be an open set and a sublattice of \mathbb{R}^N , $f : X \rightarrow \mathbb{R}$ be twice continuously differentiable, and $i \in \{1, 2, \dots, N\}$. The following are equivalent.*

- (1) f satisfies i -increasing differences on X .
- (2) For every $b \in X, u \in \mathbb{R}^N$ with $u_i > 0$, $D_{(-u)_+} D_u f(b) \geq 0$.

Proof. We know that f satisfies i -increasing differences on $X \iff f$ satisfies i -increasing differences (*) on X . In other words, (1) is equivalent to: for every $b \in X, u \in \mathbb{R}^N$ with $u_i > 0$, and for every $\sigma \geq 0$, $f(b + \sigma u + s(-u)_+) - f(b + s(-u)_+)$ is (weakly) increasing in s (that is, in the direction $(-u)_+$). Using the fundamental theorem of calculus, this is equivalent to: $\forall b \in X, \forall u \in \mathbb{R}^N$ with $u_i > 0$, $D_u f(b + s(-u)_+)$ is (weakly) increasing in s (that is, in direction $(-u)_+$). This, in turn, is equivalent to $\forall b \in X, \forall u \in \mathbb{R}^N$ with $u_i > 0$, $D_{(-u)_+} D_u f(b) \geq 0$. ■

The second statement can be given a convenient name in terms of nonnegative cross derivatives, as follows. Let X be an open set and a sublattice of \mathbb{R}^N , $f : X \rightarrow \mathbb{R}$ be twice continuously differentiable, and $i \in \{1, 2, \dots, N\}$. The function f **has nonnegative i -cross derivative property on X** , if for every $b \in X, u \in \mathbb{R}^N$ with $u_i > 0$, $D_{(-u)_+} D_u f(b) \geq 0$, and f **has nonnegative directional cross derivative property on X** , if for every $i \in \{1, 2, \dots, N\}$, f has nonnegative i -cross derivative property on X . This proposition shows that *i -increasing differences on X is equivalent to nonnegative i -cross derivative property on X* , and it follows immediately that *directional increasing differences on X is equivalent to nonnegative directional cross derivative property on X* .

Similarly, consider the following cardinal property naturally suggested by the basic i -single crossing property on $X \times T$. Let X be a sublattice of \mathbb{R}^N , (T, \preceq) be a partially ordered set, $f : X \times T \rightarrow \mathbb{R}$, and $i \in \{1, 2, \dots, N\}$. The function f satisfies **basic i -increasing differences on $X \times T$** , if for every $a, b \in X$ with $a_i > b_i$, and for every $t, t' \in T$ with $t \preceq t'$, $f(a, t) - f(b, t) \leq f(a, t') - f(b, t')$. The function f satisfies **basic directional increasing differences on $X \times T$** , if for every $i \in \{1, 2, \dots, N\}$, f satisfies basic i -increasing differences on $X \times T$. As earlier, it is easy to check that *if f satisfies basic i -increasing differences on $X \times T$, then f satisfies basic i -single crossing property on $X \times T$* , and it follows immediately that *if f satisfies basic directional increasing differences*

on $X \times T$, then f satisfies basic directional single crossing property on $X \times T$. The following result now obtains easily.

Proposition 3. *Let X be an open set and a sublattice of \mathbb{R}^N , T be an open subset of \mathbb{R}^M , $f : X \times T \rightarrow \mathbb{R}$ be twice continuously differentiable, and $i \in \{1, 2, \dots, N\}$.*

The following are equivalent.

- (1) *f satisfies basic i -increasing differences on $X \times T$.*
- (2) *For every $b \in X, u \in \mathbb{R}^N$ with $u_i > 0$, $D_t D_u f(b, t) \geq 0$.*

Proof. It is easy to check that f satisfies basic i -increasing differences on $X \times T \iff$ for every $b \in X, u \in \mathbb{R}^N$ with $u_i > 0$, $f(b + u, t) - f(b, t)$ is (weakly) increasing in t . This is equivalent to: $\forall b \in X, \forall u \in \mathbb{R}^N$ with $u_i > 0$, $D_u f(b, t)$ is (weakly) increasing in t , and this is further equivalent to: $\forall b \in X, \forall u \in \mathbb{R}^N$ with $u_i > 0$, $D_t D_u f(b, t) \geq 0$. ■

The second statement can be given a convenient name in terms of nonnegative cross derivatives, as follows. Let X be an open set and a sublattice of \mathbb{R}^N , T be an open subset of \mathbb{R}^M , $f : X \times T \rightarrow \mathbb{R}$ be twice continuously differentiable, and $i \in \{1, 2, \dots, N\}$. The function f **has nonnegative basic i -cross derivative property on $X \times T$** , if for every $b \in X, u \in \mathbb{R}^N$ with $u_i > 0$, $D_t D_u f(b, t) \geq 0$, and f **has nonnegative basic directional cross derivative property on $X \times T$** , if for every $i \in \{1, 2, \dots, N\}$, f has nonnegative basic i -cross derivative property on X . The above proposition shows that *basic i -increasing differences on $X \times T$ is equivalent to nonnegative basic i -cross derivative property on $X \times T$* , and it follows immediately that *basic directional increasing differences on $X \times T$ is equivalent to basic nonnegative directional cross derivative property on $X \times T$* . We have the following result.

Theorem 3. *Let X be an open set and a sublattice of \mathbb{R}^N , T be an open subset of \mathbb{R}^M , $f : X \times T \rightarrow \mathbb{R}$ be twice continuously differentiable, and $i \in \{1, 2, \dots, N\}$.*

If for every $t \in T$, $f(\cdot, t)$ is i -supermodular¹⁴ and has nonnegative i -cross derivative

¹⁴For every $a, b \in X$ with $a_i > b_i$, $f(a, t) - f(a \wedge b, t) \leq f(a \vee b, t) - f(b, t)$.

property on X , and f has nonnegative basic i -cross derivative property on $X \times T$, then f satisfies i -directional monotone comparative statics on $X \times T$.

Proof. The hypothesis in this statement, combined with the propositions above, implies that f satisfies basic i -single crossing property on $X \times T$, and for every $t \in T$, $f(\cdot, t)$ is i -quasisupermodular and satisfies i -single crossing property on X , and the conclusion follows from an application of theorem 1. ■

It follows immediately that *if for every $t \in T$, $f(\cdot, t)$ is supermodular and has nonnegative directional cross derivative property on X , and f has nonnegative basic directional cross derivative property on $X \times T$, then f satisfies directional monotone comparative statics on $X \times T$.*

The following corollaries help specialize this theorem to the case of comparative statics with respect to A or to t separately.

Corollary 8. *Let X be an open set and a sublattice of \mathbb{R}^N , $f : X \rightarrow \mathbb{R}$ is twice continuously differentiable, and $i \in \{1, 2, \dots, N\}$.*

If f is i -supermodular and has nonnegative i -cross derivative property on X , then f satisfies i -directional monotone comparative statics on X .

Proof. Apply the theorem with singleton $T = \{t\}$. ■

This corollary implies immediately that *if f is supermodular and has nonnegative directional cross derivative property on X , then f satisfies directional monotone comparative statics on X .*

In order to understand more concretely the nonnegative i -cross derivative property on X , let's compute $D_{(-u)_+} D_u f(b)$. For convenience, we use subscripts for partial derivatives. Notice that $D_u f(b) = \sum_{j=1}^N f_j(b) u_j$, where $f_j(b) \equiv \frac{\partial f}{\partial x_j}(b)$ and u_j is the j -th component of u . Therefore,

$$D_{(-u)_+} D_u f(b) = \sum_{k=1}^N \sum_{j=1}^N f_{k,j}(b) u_j (-u)_{+,k}.$$

Here $f_{k,j}(b)$ is the k, j -th cross-partial of f evaluated at b , u_j is the j -th component of u , and $(-u)_{+,k}$ is the k -th component of $(-u)_+$. This is easier to understand if we let $L = \{\ell \mid u_\ell < 0\}$. In this case,

$$(-u)_{+,k} = \begin{cases} -u_k & \text{if } k \in L, \text{ and} \\ 0 & \text{if } k \notin L, \end{cases}$$

and therefore,

$$\begin{aligned} D_{(-u)_+} D_u f(b) &= \sum_{k=1}^N \sum_{j=1}^N f_{k,j}(b) u_j (-u)_{+,k} \\ &= \sum_{k \in L} \sum_{j=1}^N f_{k,j}(b) u_j (-u_k) \\ &= \sum_{k \in L} \sum_{j \notin L} f_{k,j}(b) u_j (-u_k) + \sum_{k \in L} \sum_{j \in L} f_{k,j}(b) u_j (-u_k) \\ &= \sum_{k \in L} \sum_{j \notin L} f_{k,j}(b) u_j (-u_k) - \sum_{k \in L} \sum_{j \in L} f_{k,j}(b) (-u_j) (-u_k) \\ &= \sum_{k \in L} \sum_{j \notin L} f_{k,j}(b) u_j (-u_k) - [w'_L D^2 f_L(b) w_L], \end{aligned}$$

where f_L is the restriction of f to the components in L , $D^2 f_L(b)$ is the second derivative of f_L evaluated at b , w_L is the restriction of $(-u)_+$ to L , and w'_L is the transpose of w_L .

Notice that for $k \in L$, $-u_k > 0$ and for $j \notin L$, $u_j \geq 0$. In this case, the sign of the term $f_{k,j}(b) u_j (-u_k)$ is determined by the sign of the cross-partial $f_{k,j}(b)$. Similarly, for $k \in L$, $-u_k > 0$ and for $j \in L$, $u_j < 0$. In this case, the sign of the term $f_{k,j}(b) u_j (-u_k)$ is determined by the sign of $-f_{k,j}(b)$. In particular, if f is supermodular, then the first term, $\sum_{k \in L} \sum_{j \notin L} f_{k,j}(b) u_j (-u_k) \geq 0$. Moreover, if f is concave in direction $(-u)_+$, then the matrix of second derivatives is negative semidefinite, and therefore, the second term, $-[w'_L D^2 f_L(b) w_L] \geq 0$.

Corollary 9. *Let X be an open set and a sublattice of \mathbb{R}^N , T be an open subset of \mathbb{R}^M , $f : X \times T \rightarrow \mathbb{R}$ be twice continuously differentiable, and $i \in \{1, 2, \dots, N\}$.*

If for every $t \in T$, $f(\cdot, t)$ is i -supermodular on X , and f has nonnegative basic i -cross derivative property on $X \times T$, then f satisfies i -directional monotone comparative statics on $X \times T$.

Proof. The conditions in this statement imply that for every $t \in T$, $f(\cdot, t)$ is i -quasisupermodular on X , and f satisfies basic i -single crossing property on $X \times T$, and the conclusion follows from an application of corollary 4. ■

In order to understand more concretely the nonnegative basic i -cross derivative property on $X \times T$, let's compute $D_t D_u f(b, t)$. Recall that $D_u f(b, t) = \sum_{j=1}^N f_{x_j}(b, t) u_j$, where $f_{x_j}(b, t) \equiv \frac{\partial f}{\partial x_j}(b, t)$ and u_j is the j -th component of u . Therefore,

$$D_t D_u f(b, t) = \left[\sum_{j=1}^N f_{t_1, x_j}(b, t) u_j, \dots, \sum_{j=1}^N f_{t_M, x_j}(b, t) u_j \right],$$

where $f_{t_m, x_n}(b, t) \equiv \frac{\partial^2 f}{\partial t_m \partial x_n}(b, t)$, for $m = 1, \dots, M$, $n = 1, \dots, N$. This may be written in standard matrix form as

$$D_t D_u f(b, t) = \begin{bmatrix} f_{t_1, x_1}(b, t) & \cdots & f_{t_1, x_N}(b, t) \\ \vdots & & \vdots \\ f_{t_M, x_1}(b, t) & \cdots & f_{t_M, x_N}(b, t) \end{bmatrix} \begin{bmatrix} u_1 \\ \vdots \\ u_N \end{bmatrix}.$$

A useful sufficient condition for nonnegative basic i -cross derivative property on $X \times T$ is the following. Let X be an open set and a sublattice of \mathbb{R}^N , T be an open subset of \mathbb{R}^M , $f : X \times T \rightarrow \mathbb{R}$ be twice continuously differentiable, and $i \in \{1, 2, \dots, N\}$. If for some subset M' of $\{1, \dots, M\}$, $f_{t_m, x_i}(b, t) \geq 0$ for $m \in M'$, and $f_{t_m, x_j}(b, t) = 0$ otherwise, then f has nonnegative basic i -cross derivative property on $X \times T$. To see that this is true, fix $u \in \mathbb{R}^N$ with $u_i > 0$, and notice that the m -th component of $D_t D_u f(b, t)$ is $f_{t_m, x_i}(b, t) u_i \geq 0$ for $m \in M'$ and zero otherwise. This condition retains the flavor of standard increasing differences in $(x; t)$ by working with nonnegative cross-partial, and it is useful in applications, as detailed in the next section.

3 Examples

Example 2 (Consumer Demand). Consider a consumption space X that is a sublattice of \mathbb{R}_+^L , a partially ordered parameter space (T, \preceq) , a utility function $u : X \times T \rightarrow \mathbb{R}$

and a subset A of X , and consider the utility maximization problem, $\max_A u(\cdot, t)$. When utility is continuous on X and A is a nonempty compact set, this problem has a solution termed consumer demand. Let's denote it by $D(A, t) = \arg \max_A u(\cdot, t)$. Theorem 1 provides conditions characterizing when $D(A, t)$ is increasing in (A, t) in the i -directional set order and in the directional set order. Some special cases are notable.

Consider Walrasian demand, that is, let the consumption space be $X = \mathbb{R}_+^L$ or $X = \mathbb{R}_{++}^L$, a price vector $p \in \mathbb{R}_{++}^L$, wealth $w > 0$, and let $B(p, w) = \{x \in X \mid p \cdot x \leq w\}$ be the Walrasian budget set and let $D(p, w) = \arg \max_{B(p, w)} u(\cdot, t)$ be Walrasian demand. We know that $w \leq w' \Rightarrow (\forall i) B(p, w) \sqsubseteq_i^{dso} B(p, w')$. Say that demand for good i is *normal*, if $w \leq w' \Rightarrow D(p, w) \sqsubseteq_i^{wls} D(p, w')$. In this setting, the result on sufficient conditions implies that *if u is i -supermodular and i -concave, then Walrasian demand for good i is normal*, and *if u is supermodular and directionally concave, then Walrasian demand for all goods is normal*. Moreover, corollary 5 implies that strictly increasing transformations of u yield the same conclusions. This implies, for example, that some of the standard cases such as general Cobb-Douglas preferences (with no restriction that exponential parameters add to 1), constant elasticity of substitution, and taking logarithms of standard preferences are all admissible. Furthermore, in the case of two goods, these results hold with discrete consumption sets as well, when conditions in example 1-2 hold.

We can also consider comparative statics with respect to parameter t . Corollary 4 implies that if u is i -quasisupermodular and satisfies basic i -single crossing property on $X \times T$, then $t \preceq t' \implies \arg \max_A u(\cdot, t) \sqsubseteq_i^{dso} \arg \max_A u(\cdot, t')$. Notably, A can be an arbitrary subset of \mathbb{R}^L . This can be seen concretely with Stone-Geary utility.

Example 3 (Stone-Geary utility) Consider consumption space $X = \mathbb{R}_+^L$ or $X = \mathbb{R}_{++}^L$, a bundle $b \in \mathbb{R}_+^L$, and utility given by $u(x, b) = \prod_{j=1}^L (x_j + b_j)^{\alpha_j}$, where $\alpha_j > 0$ for all j . The bundle b is sometimes viewed as a survival bundle available as an outside option, perhaps through a government program, or through a soup kitchen, or through a charity,

and so on, although other interpretations are available.¹⁵ Theoretically, it is a parameter in the utility function. Notice that for each b , $u(\cdot, b)$ is quasisupermodular and quasiconcave. Moreover, when $b = 0$, Stone-Geary specializes to Cobb-Douglas preferences. There is no restriction that α_j add up to 1.

In order to use derivatives, let $a \in \mathbb{R}_{--}^L$, and write $u(x, a) = \prod_{j=1}^L (x_j - a_j)^{\alpha_j}$, where $\alpha_j > 0$ for all j , and consider the monotonic transformation, $v(x, a) = \sum_{j=1}^L \alpha_j \log(x_j - a_j)$. Then for each $a \in \mathbb{R}_{--}^L$, $v(\cdot, a)$ is supermodular and concave on X . Moreover, for fixed $i \in \{1, \dots, L\}$, and for every $u \in \mathbb{R}^L$ with $u_i > 0$, $D_u v(x, a) = \sum_{j=1}^L \frac{\alpha_j}{x_j - a_j} u_j$ and therefore, $D_{a_i} D_u v(x, a) = \frac{\alpha_i}{(x_i - a_i)^2} u_i > 0$. Consequently, v satisfies basic i single crossing property on $X \times \mathbb{R}_{--}$, where \mathbb{R}_{--} indexes a_i . By theorem 3, v (and u) satisfies i -directional monotone comparative statics on $X \times \mathbb{R}_{--}$. In particular, when a_i goes up, (and as long as the corresponding budget sets (weakly) increase in the i -directional set order,) demand for good i goes up.

In terms of the original problem with nonnegative b , this implies that when a component of the survival bundle is increased, a consumer's optimal response is to decrease the same component of her demand. This is consistent with results in public economics on the effect of more generous social welfare options. It follows here from a minimal calculation on the objective function and is valid for an arbitrarily fixed compact budget set.

Example 4 (Multi-market competition). Consider a firm that is competing in two markets; market 1 is imperfectly competitive, say, an oligopoly, and market 2 is perfectly competitive. (For example, the firm might produce a generic product for the competitive market and a differentiated version to have some market power.) Suppose the firm's profits are given by $\Pi(x_1, y; x_2, \dots, x_M) = \Pi_1(x_1, x_2, \dots, x_M) + p \cdot y - c(y)$, where Π_1 is the firm's profit in market 1, which has $M \geq 2$ firms, and $p \cdot y - c(y)$ is its profit in market 2. The firm's choice variables are $(x_1, y) \in \mathbb{R}_+^2$. Suppose the actions of

¹⁵For example, in models of charitable giving, b may be viewed as a consumer or donor's intrinsic benefit from donation, as in Harbaugh (1998).

other firms in market 1 are complementary to the actions of the firm; that is, $\frac{\partial^2 \Pi_1}{\partial x_j \partial x_1} \geq 0$, for $j = 2, \dots, M$. (For example, this follows if market 1 competition is of a standard differentiated Bertrand variety. It could also follow if there are production externalities or network externalities in market 1.) The firm's problem is to $\max_A \Pi(x_1, y; x_2, \dots, x_M)$. Notice that Π is supermodular in (x_1, y) , because of additivity. To check that Π has the basic 1-single crossing property in $(x_1, y; x_2, \dots, x_M)$, observe that for u with $u_1 > 0$, $D_{x_{-1}} D_u \Pi(x_1, y; x_{-1}) = [\frac{\partial^2 \Pi_1}{\partial x_2 \partial x_1} u_1, \dots, \frac{\partial^2 \Pi_1}{\partial x_M \partial x_1} u_1] \geq 0$. Therefore, when competitor action goes up, firm 1's best response in market 1 goes up as well. Notably, this result holds for arbitrary constraint set A .

We can also inquire about comparative statics with respect to A . For motivation, suppose market 1 is subject to production or network externalities. (For example, when other firms produce more, a given firm's marginal cost goes down, either because of a direct upstream or downstream production externality or an indirect one, perhaps through the availability of more skilled labor, more efficient supply chains, and so on.) In other words, suppose x_1 is the firm's production in market 1, and suppose again that $\frac{\partial^2 \Pi_1}{\partial x_j \partial x_1} \geq 0$ for $j = 2, \dots, M$. The firm faces a capacity constraint for producing both outputs, $A(k) = \{(x_1, y) \mid x_1 + y \leq k\}$. This can be generalized somewhat by considering $A(k) = \{(x_1, y) \mid \alpha_1 x_1 + \alpha_2 y \leq k\}$, where α_1 and α_2 are arbitrary positive constants, perhaps indexing different production requirements. (For example, it may be that some more factory space is needed to produce a unit of the differentiated good, as compared to the competitive good, and the "weighted" output is constrained by the "base" plant size of k units.) Consider the firm's problem: $\max_{A(k)} \Pi(x_1, y; x_2, \dots, x_M)$. For fixed $A(k)$, we still have the earlier result that the firm's supply in market 1 is (weakly) increasing with respect to supply of other firms. But now we can also inquire about the effects of an increase in plant capacity simultaneously with an increase in other firm actions. To do so, we need to check for x_1 -concavity of Π . This holds when the cost function in the competitive market, $c(y)$ is convex (which follows from concave production technology). In this case,

$$(k, x_{-1}) \leq (k', x'_{-1}) \Rightarrow \arg \max_{A(k)} \Pi(x_1, y; x_{-1}) \sqsubseteq_1^{dso} \arg \max_{A(k')} \Pi(x_1, y; x'_{-1}).^{16}$$

We can also inquire about monotone comparative statics for the competitive market. Notice that $\frac{\partial^2 \Pi}{\partial p \partial y} = 1$, and therefore, an increase in the competitive price increases output in the competitive market, regardless of the constraint set. Moreover, y -concavity of Π follows from convex cost in market 1 (again, following from concave production technology) and concave total revenue function in market 1 (which follows if demand is linear, and also if demand has constant elasticity less than or equal to 1, which includes Cobb-Douglas preferences).

A more general example can be formulated as follows. Consider a firm producing outputs in $N \geq 2$ markets, with profit in market i given by $\Pi_i(x_i, t_i)$, where $x_i \in \mathbb{R}_+$ is the firm's output in market i , and $t_i \in \mathbb{R}^{M_i}$ is a vector of parameters for market i . Total profit of the firm is $\Pi(x; t) = \sum_{i=1}^N \Pi_i(x_i, t_i)$ and the firm's problem is to $\max_A \Pi(x; t)$ for x in some constraint set A . In this case, Π is supermodular in (x_1, \dots, x_N) , due to additivity. If we assume that each $\Pi_i(x_i, t_i)$ is concave in x_i and satisfies basic i -single crossing property in (x_i, t_i) , (for example, if $\frac{\partial^2 \Pi_i}{\partial t_{i,j} \partial x_i} \geq 0$ for $j = 1, \dots, M_i$) then we may conclude that for each i , $A \sqsubseteq_i^{dso} A'$ and $t_i \leq t'_i$ implies $\arg \max_A \Pi(x; t_i, t_{-i}) \sqsubseteq_i^{dso} \arg \max_{A'} \Pi(x; t'_i, t_{-i})$.

Example 5 (Emissions Standards). Consider the emissions standards model in Montero (2002) and Bruneau (2004). A firm is producing an output $q \geq 0$ that causes pollution. It is subject to an emissions ceiling $e > 0$ and can produce produce more by engaging in costly abatement $a \geq 0$. The firm's payoff is given by $\pi(q, a; k) = p \cdot q - c(q) - kc(a)$. Here, revenue is $p \cdot q$, cost of output, $c(q)$ is assumed to be increasing and convex, as is cost of abatement, $c(a)$. The firm can consider technological progress $k \leq 1$, measured as a decrease in abatement cost to $kc(a)$. It is easy to check that Π is supermodular in (q, a) , Π is concave, and technological innovation (decrease in k , or increase in $-k$) satisfies $\frac{\partial^2 \Pi}{\partial (-k) \partial q} = 0$ and $\frac{\partial^2 \Pi}{\partial (-k) \partial a} \geq 0$. Therefore, an increase in technological in-

¹⁶The same result holds for minimum production quotas; constraints sets of the form $A(k) = \{(x_1, y) \mid \alpha_1 x_1 + \alpha_2 y \geq k\}$, where α_1 and α_2 are arbitrary positive constants. In this case as well, $k \leq k' \Rightarrow A(k) \sqsubseteq^{dso} A(k')$.

novation, say $(-k) \leq (-k')$ implies that $\arg \max_A \Pi(q, a; k) \sqsubseteq_a^{dso} \arg \max_A \Pi(q, a; k')$, for arbitrary constraint set A . In particular, it holds for the emissions constraint set, $A(e) = \{(q, a) \mid q - a = e\}$. Moreover, at optimum, an increase in a leads to an increase in q , and therefore, *increase in technological innovation increases both abatement and output*. This main result follows from an easy calculation on the objective function.

Example 6 (Discrete labor supply). Recent models of labor supply frequently incorporate a discrete choice model, for example, Aaberge, Gagsvik, and Strøm (1995), van Soest (1995), and Hoynes (1996). In order to work with integer data, these models consider integer work-leisure choices. Let h denote hours worked and l denote hours of leisure. Given total hours available T , the constraint set is $B(T) = \{(h, l) \in \mathbb{Z}_+ \times \mathbb{Z}_+ \mid h + l \leq T\}$. Using example 1-2, it is easy to check that $T \leq T' \Rightarrow B(T) \sqsubseteq^{dso} B(T')$. Preferences are given by $u(wh + I, l)$ where w is wage rate and I is non-labor income, both exogenously specified. Using our results, when preferences are supermodular and concave, both hours worked and leisure hours are increasing in the time constraint T , even in a discrete choice framework. In particular, for standard preferences such as Cobb-Douglas, CES, and their increasing transformations, this result holds. Moreover, consider the utility from Hoynes (1996), $u(wh + I, l) = \alpha_1 \ln(wh + I) + \alpha_2 \ln(l)$ with $\alpha_1, \alpha_2 > 0$. In this case, the basic h -single crossing property holds for parameters $(w, -I)$, because $\frac{\partial^2 u}{\partial(-I)\partial h} = \frac{\alpha_1 w}{(wh+I)^2} \geq 0$, $\frac{\partial^2 u}{\partial w \partial h} = \frac{\alpha_1 I}{(wh+I)^2} \geq 0$, $\frac{\partial^2 u}{\partial(-I)\partial l} = 0$, and $\frac{\partial^2 u}{\partial w \partial l} = 0$. Consequently, optimal labor supply increases when either wage rate goes up or non-labor income goes down. Notably, this result holds for discrete choices and for an arbitrary compact constraint set.

Example 7 (Auctions with budget constraints). Auctions with budget constraints are common in practice (for example ad auctions run by Google and Yahoo, Treasury auctions, and spectrum or electricity auctions), but less widely studied in the auction theory literature (for some examples, confer Rothkopf (1977), Palfrey (1980), and Dobzinski, Lavi, and Nisan (2012)). Consider an auction of $N \geq 2$ indivisible items. A bidder has exogenously specified valuation $v = (v_1, \dots, v_N)$ for the N objects, and can bid $b = (b_1, \dots, b_N)$ subject to a resource constraint $b_1 + \dots + b_N \leq T$. The probability

of winning object i with bid b_i is given by $F_i(b_i)$, where $\frac{\partial F_i}{\partial b_i} \geq 0$. The expected payoff from winning object i is $u_i(b_i, v_i)F_i(b_i)$ and the expected payoff from losing is normalized to 0. A bidder maximizes expected payoff $u(b, v) = \sum_{i=1}^N u_i(b_i, v_i)F_i(b_i)$ subject to her resource constraint. Suppose marginal utility for object i is increasing in v_i ($\frac{\partial u_i}{\partial v_i} \geq 0$), and bids and valuations are complementary ($\frac{\partial^2 u_i}{\partial v_i \partial b_i} \geq 0$). To inquire into monotone comparative statics, suppose $u_i(b_i, v_i)F_i(b_i)$ is concave in b_i . In this case, expected payoff is supermodular and concave in b , and therefore, the payoff-maximizing bid profile b is (weakly) increasing in total resources T . In the two-item case, this holds for discrete choice as well. Moreover, for an arbitrarily fixed item i , it is easy to compute that $\frac{\partial^2 u_i(b_i, v_i)F_i(b_i)}{\partial v_i \partial b_i} = \frac{\partial^2 u_i}{\partial v_i \partial b_i} F_i(b_i) + \frac{\partial u_i}{\partial v_i} \frac{\partial F_i}{\partial b_i} \geq 0$. Consequently, an increase in valuation of item i increases its optimal bid (for an otherwise arbitrary constraint set).

4 Conclusion

This paper presents an extension of the theory of monotone comparative statics in different directions in finite-dimensional Euclidean space. The new notions of i -single crossing property and basic i -single crossing property are similar in spirit to the single-crossing property in the standard theory of monotone comparative statics, both are ordinal properties, and both can be naturally specialized to related cardinal and differential properties. The results here use more standard assumptions on the objective function, include parameters in the objective function, do not require the use of new binary relations or convex domains, and include results in Quah (2007) as a special case. The results allow flexibility to explore comparative statics with respect to the constraint set, with respect to parameters in the objective function, or both. Moreover, the formulation here is easy to apply in many applications.

References

- AABERGE, R., J. K. GAGSVIK, AND S. STRØM (1995): “Labor Supply Responses and Welfare Effects of Tax Reforms,” *The Scandinavian Journal of Economics*, 97(4), 635–659.
- ACEMOGLU, D., AND M. K. JENSEN (2013): “Aggregate comparative statics,” *Games and Economic Behavior*, 81, 27–49.
- (2015): “Robust comparative statics in large dynamic economies,” *Journal of Political Economy*, forthcoming.
- AMIR, R. (1996): “Cournot Oligopoly and the Theory of Supermodular Games,” *Games and Economic Behavior*, 15, 132–148.
- AMIR, R., AND V. E. LAMBSON (2000): “On the Effects of Entry in Cournot Markets,” *The Review of Economic Studies*, 67(2), 235–254.
- ANTONIADU, E. (2007): “Comparative statics for the consumer problem,” *Economic Theory*, 31(1), 189–203.
- BRUNEAU, J. F. (2004): “A note on permits, standards, and technological innovation,” *Journal of Environmental Economics and Management*, 48(3), 1192–1199.
- BULOW, J. I., J. D. GEANAKOPOLOS, AND P. D. KLEMPERER (1985): “Multimarket Oligopoly: Strategic Substitutes and Complements,” *Journal of Political Economy*, 93(3), 488–511.
- DOBZINSKI, S., R. LAVI, AND N. NISAN (2012): “Multi-unit auctions with budget limits,” *Games and Economic Behavior*, 74(2), 486–503.
- ECHENIQUE, F. (2002): “Comparative statics by adaptive dynamics and the correspondence principle,” *Econometrica*, 70(2), 257–289.
- (2004): “A characterization of strategic complementarities,” *Games and Economic Behavior*, 46(2), 325–347.
- HARBAUGH, W. T. (1998): “The Prestige Motive for Making Charitable Transfers,” *American Economic Review*, 88(2), 277–282.
- HOYNES, H. H. (1996): “Welfare Transfers in Two-Parent Families: Labor Supply and Welfare Participation Under AFDC-UP,” *Econometrica*, 64(2), 295–332.
- LICALZI, M., AND A. F. VEINOTT, JR (1992): “Subextremal Functions and Lattice Programming,” Working paper.

- MILGROM, P., AND J. ROBERTS (1990): “Rationalizability, learning, and equilibrium in games with strategic complementarities,” *Econometrica*, 58(6), 1255–1277.
- (1994): “Comparing Equilibria,” *American Economic Review*, 84(3), 441–459.
- MILGROM, P., AND C. SHANNON (1994): “Monotone Comparative Statics,” *Econometrica*, 62(1), 157–180.
- MIRMAN, L. J., AND R. RUBLE (2008): “Lattice Theory and the Consumer’s Problem,” *Mathematics of Operations Research*, 33(2), 301–314.
- MONACO, A. J., AND T. SABARWAL (2015): “Games with Strategic Complements and Substitutes,” *Economic Theory*, forthcoming.
- MONTERO, J.-P. (2002): “Permits, Standards, and Technology Innovation,” *Journal of Environmental Economics and Management*, 44(1), 23–44.
- PALFREY, T. R. (1980): “Multiple-object, discriminatory auctions with bidding constraints: A game-theoretic analysis,” *Management Science*, 26(9), 935–946.
- QUAH, J. K.-H. (2007): “The Comparative Statics of Constrained Optimization Problems,” *Econometrica*, 75(2), 401–431.
- QUAH, J. K.-H., AND B. STRULOVICI (2009): “Comparative statics, informativeness, and the interval dominance order,” *Econometrica*, 77(6), 1949–1992.
- (2012): “Aggregating the single crossing property,” *Econometrica*, 80(5), 2333–2348.
- ROTHKOPF, M. H. (1977): “Bidding in simultaneous auctions with a constraint on exposure,” *Operations Research*, 25(4), 620–629.
- ROY, S., AND T. SABARWAL (2008): “On the (Non-)Lattice Structure of the Equilibrium Set in Games With Strategic Substitutes,” *Economic Theory*, 37(1), 161–169.
- (2010): “Monotone comparative statics for games with strategic substitutes,” *Journal of Mathematical Economics*, 46(5), 793–806.
- (2012): “Characterizing stability properties in games with strategic substitutes,” *Games and Economic Behavior*, 75(1), 337–353.
- TOPKIS, D. (1978): “Minimizing a submodular function on a lattice,” *Operations Research*, 26, 305–321.
- (1979): “Equilibrium points in nonzero-sum n -person submodular games,” *SIAM Journal on Control and Optimization*, 17(6), 773–787.
- (1998): *Supermodularity and Complementarity*. Princeton University Press.

- VAN SOEST, A. (1995): “Structural Models of Family Labor Supply: A Discrete Choice Approach,” *The Journal of Human Resources*, 30(1), 63–88.
- VEINOTT, JR, A. F. (1992): “Lattice programming: qualitative optimization and equilibria,” Working paper.
- VIVES, X. (1990): “Nash Equilibrium with Strategic Complementarities,” *Journal of Mathematical Economics*, 19(3), 305–321.
- ZHOU, L. (1994): “The Set of Nash Equilibria of a Supermodular Game is a Complete Lattice,” *Games and Economic Behavior*, 7(2), 295–300.
- ZIMPER, A. (2007): “A fixed point characterization of the dominance-solvability of lattice games with strategic substitutes,” *International Journal of Game Theory*, 36(1), 107–117.

Appendix A. Relation to Quah (2007)

Quah (2007) uses different techniques based on new binary relations, denoted ∇_i^λ and Δ_i^λ , and convex sets. Using these binary relations, he defines a new set order, termed \mathcal{C}_i -flexible set order, and a new notion of \mathcal{C}_i -quasisupermodular function. Some connections to these ideas are explored here.

Let X be a convex sublattice of \mathbb{R}^N (that is, X is a sublattice that is also a convex set), and $i \in \{1, 2, \dots, N\}$. For $a, b \in X$ and $\lambda \in [0, 1]$, let

$$a\Delta_i^\lambda b = \begin{cases} a & \text{if } a_i \leq b_i \\ \lambda b + (1 - \lambda)a \wedge b & \text{if } a_i > b_i, \end{cases} \quad \text{and} \quad a\nabla_i^\lambda b = \begin{cases} b & \text{if } a_i \leq b_i \\ \lambda a + (1 - \lambda)a \vee b & \text{if } a_i > b_i. \end{cases}$$

Figure 5 shows the graphical intuition.




Figure 5: \mathcal{C}_i -flexible Set Order

When $a_i > b_i$, the set $\{a, a\Delta_i^\lambda b, a\nabla_i^\lambda b, b\}$ forms a “backward-bending” parallelogram, as compared to the standard lattice theory rectangle formed by the set $\{a, a \wedge b, a \vee b, b\}$. The shape of this parallelogram varies with λ , ranging from the standard lattice theory rectangle when $\lambda = 0$ to the degenerate line segment formed by $\{a, b\}$ when $\lambda = 1$.

The binary operations $\Delta_i^\lambda, \nabla_i^\lambda$ have some counter-intuitive properties when compared to the standard lattice operations \wedge, \vee . For example, *the relations $\Delta_i^\lambda, \nabla_i^\lambda$ are non-commutative*: suppose $N = 2, i = 1$, consider $a = (1, 0)$, $b = (0, 1)$, and $\lambda = \frac{1}{2}$. Then $a\Delta_i^\lambda b = \frac{1}{2}b \neq b = b\Delta_i^\lambda a$, and $a\nabla_i^\lambda b = (1, \frac{1}{2}) \neq a = b\nabla_i^\lambda a$. Moreover, *$a\Delta_i^\lambda b$ and $a\nabla_i^\lambda b$ are not necessarily comparable in the underlying lattice order*: suppose $N = 2, i = 1$, and consider $a = (1, 1)$ and $b = (2, 0)$. Then for every $\lambda \in [0, 1]$, $a\Delta_i^\lambda b = a \not\leq b = a\nabla_i^\lambda b$. It is easy to see that additional classes of examples of these instances can be provided as well.

The binary relations $\Delta_i^\lambda, \nabla_i^\lambda$ are used to define the \mathcal{C}_i -flexible set order and the notion of a \mathcal{C}_i -quasisupermodular function, as follows.

Let X be a convex sublattice of \mathbb{R}^N and $i \in \{1, 2, \dots, N\}$. For subsets A, B of X , A is **lower than B in the \mathcal{C}_i -flexible set order**, denoted $A \sqsubseteq_i^{\mathcal{C}} B$, if for every $a \in A, b \in B$,

there is $\lambda \in [0, 1]$ such that $a\Delta_i^\lambda b \in A$ and $a\nabla_i^\lambda b \in B$. The \mathcal{C}_i -flexible set order is flexible in the sense that the choice of λ may vary for each $a \in A$ and $b \in B$, and therefore, the “backward bendedness” of the parallelogram may vary for each $a \in A$ and $b \in B$. On convex sublattices, the \mathcal{C}_i -flexible set order is the same as the i -directional set order, as shown next.

Proposition 4. *Let X be a convex sublattice of \mathbb{R}^N , $i \in \{1, 2, \dots, N\}$, and A, B be subsets of X . The following are equivalent.*

- (1) *A is lower than B in the \mathcal{C}_i -flexible set order ($A \sqsubseteq_i^{\mathcal{C}} B$).*
- (2) *A is lower than B in the i -directional set order ($A \sqsubseteq_i^{dso} B$).*

Proof. Suppose $A \sqsubseteq_i^{\mathcal{C}} B$. Fix $a \in A, b \in B$, and suppose $a_i > b_i$. Let $\lambda \in [0, 1]$ be such that $a\Delta_i^\lambda b \in A$ and $a\nabla_i^{1-\lambda} b \in B$. Let $t = 1 - \lambda \in [0, 1]$. Then $b - v = b - t(b - a \wedge b) = (1-t)b + t(a \wedge b) = a\Delta_i^{1-t} b \in A$, and $a + v = a + t(a \vee b - a) = (1-t)a + t(a \vee b) = a\nabla_i^{1-t} b \in B$, as desired.

In the other direction, suppose $A \sqsubseteq_i^{dso} B$. Fix $a \in A, b \in B$. Suppose $a_i \leq b_i$. Then $a\Delta_i^{1-t} b = a \in A$ and $a\nabla_i^{1-t} b = b \in B$, as desired. Suppose $a_i > b_i$. Let $t \in [0, 1]$ be such that $v = t(b - a \wedge b) = t(a \vee b - a)$ satisfies $b - v \in A$ and $a + v \in B$. Then for $\lambda = 1 - t$, $a\Delta_i^\lambda b = (1-t)b + t(a \wedge b) = b - t(b - a \wedge b) = b - v \in A$, and $a\nabla_i^\lambda b = (1-t)a + t(a \vee b) = a + t(a \vee b - a) = a + v \in B$, as desired. ■

The i -directional set order may be viewed as reformulating the \mathcal{C}_i -flexible set order to work more closely with monotone methods. In particular, i -directional set order does not invoke the binary relations $\Delta_i^\lambda, \nabla_i^\lambda$, it does not require convex sets, and it uses the standard properties of order and direction in \mathbb{R}^N .

Let X be a convex sublattice of \mathbb{R}^N , $f : X \rightarrow \mathbb{R}$, and $i \in \{1, 2, \dots, N\}$. The function f is **\mathcal{C}_i -quasisupermodular**, if for every $a, b \in X$ and for every $\lambda \in [0, 1]$, $f(a) \geq (>) f(a\Delta_i^\lambda b) \Rightarrow f(a\nabla_i^\lambda b) \geq (>) f(b)$. One of the main results in Quah (2007) is the following: *for every $i \in \{1, \dots, N\}$, $\arg \max_A f$ is increasing in A in the \mathcal{C}_i -flexible set order, if, and only if, f is \mathcal{C}_i -quasisupermodular.*

Notice that the property \mathcal{C}_i -quasisupermodular is symbolically similar to the notion of a quasisupermodular function. Its interpretation is more complex for two reasons: first, the use of the quantifier “for every $\lambda \in [0, 1]$ ” in the definition forces consideration of the whole line segment joining a and $a \vee b$ and the whole line segment joining $a \wedge b$ and b , and essentially forces consideration of convex sets, and second, the interpretive issues with using $\Delta_i^\lambda, \nabla_i^\lambda$ carry over to this definition.

The use of the quantifier “for every $\lambda \in [0, 1]$ ” in this definition is required by the \mathcal{C}_i -flexible set order. This can be seen as follows. Suppose we consider weakening the definition of f is \mathcal{C}_i -quasisupermodular by requiring it to hold for only some collection of $\lambda \in [0, 1]$, as follows. Let X be a convex sublattice of \mathbb{R}^N , $f : X \rightarrow \mathbb{R}$, $i \in \{1, 2, \dots, N\}$, and Λ be a nonempty subset of $[0, 1]$. The function f is **(i, Λ) -quasisupermodular**, if

for every x, y in X , and every $\lambda \in \Lambda$, $f(x) \geq (>) f(x\Delta_i^\lambda y) \Rightarrow f(x\Delta_i^\lambda y) \geq (>) f(y)$. Notice that f is \mathcal{C}_i -quasisupermodular is a special case of this definition, when $\Lambda = [0, 1]$.

In order to characterize the type of monotone comparative statics possible with (i, Λ) -quasisupermodular functions, consider the following set order. Let X be a convex sublattice of \mathbb{R}^N , $i \in \{1, 2, \dots, N\}$, and Λ be a nonempty subset of $[0, 1]$. For subsets A, B of X , A is (i, Λ) -**lower than** B , denoted $A \sqsubseteq_i^\Lambda B$, if for every $a \in A$, for every $b \in B$, there is $\lambda \in \Lambda$ such that $a\Delta_i^\lambda b \in A$ and $a\nabla_i^\lambda b \in B$. Notice that A is lower than B in the \mathcal{C}_i -flexible set order is a special case of this definition, when $\Lambda = [0, 1]$. Say that a function $f : X \rightarrow \mathbb{R}$ has (i, Λ) -**increasing property**, if for every A, B subset of X , $A \sqsubseteq_i^\Lambda B \Rightarrow \arg \max_A f \sqsubseteq_i^\Lambda \arg \max_B f$. We can prove the following result.

Proposition 5. *Let X be a convex sublattice of \mathbb{R}^N , $f : X \rightarrow \mathbb{R}$, $i \in \{1, 2, \dots, N\}$, and Λ be a nonempty subset of $[0, 1]$.*

f is (i, Λ) -quasisupermodular, if, and only if, f has (i, Λ) -increasing property.

Proof. (\Rightarrow) Suppose f is (i, Λ) -quasisupermodular. Fix $A \sqsubseteq_i^\Lambda B$. Let $a \in \arg \max_A f$ and $b \in \arg \max_B f$. Notice that $A \sqsubseteq_i^\Lambda B$ implies that there is $\lambda \in \Lambda$ such that $a\Delta_i^\lambda b \in A$ and $a\nabla_i^\lambda b \in B$. Fix this λ . Thus $a \in \arg \max_A f \Rightarrow f(a) \geq f(a\Delta_i^\lambda b) \Rightarrow f(a\nabla_i^\lambda b) \geq f(b)$, where the last implication follows from (i, Λ) -quasisupermodularity of f . Moreover, as $b \in \arg \max_B f$, it follows that $f(a\nabla_i^\lambda b) = f(b)$, whence $a\nabla_i^\lambda b \in \arg \max_B f$. Furthermore, $f(a\nabla_i^\lambda b) = f(b) \Rightarrow f(a\nabla_i^\lambda b) \not> f(b) \Rightarrow f(a) \leq f(a\Delta_i^\lambda b)$, where the last implication follows from (i, Λ) -quasisupermodularity of f . As $a \in \arg \max_A f$, it follows that $f(a) = f(a\Delta_i^\lambda b)$, whence $a\Delta_i^\lambda b \in \arg \max_A f$, as desired.

(\Leftarrow) Considering the contrapositive, suppose f is not (i, Λ) -quasisupermodular. Then there exists $\lambda \in \Lambda$, and there exist a, b in X , such that either (1) $f(a) \geq f(a\Delta_i^\lambda b)$ and $f(a\nabla_i^\lambda b) < f(b)$, or (2) $f(a) > f(a\Delta_i^\lambda b)$ and $f(a\nabla_i^\lambda b) \leq f(b)$. Notice that in either case, it must be that $a_i > b_i$. Therefore, $a\nabla_i^\lambda b \neq b$, $a\Delta_i^\lambda b \neq a$, and $a\Delta_i^\lambda b \neq a\nabla_i^\lambda b$. Let $C = \{a, a\Delta_i^\lambda b\}$ and $C' = \{b, a\nabla_i^\lambda b\}$. Then $C \sqsubseteq_i^\Lambda C'$. Suppose (1) is true. Then $a \in \arg \max_C f$ and $y = \arg \max_{C'} f$, but for every $\lambda' \in \Lambda$, $a\nabla_i^{\lambda'} b \notin \arg \max_{C'} f$, because $a_i > b_i$ implies that for every $\lambda' \in [0, 1]$, $a\nabla_i^{\lambda'} b \neq b$. Therefore, f does not have (i, Λ) -increasing property (for $C \sqsubseteq_i^\Lambda C'$). Suppose (2) is true. Then $a = \arg \max_C f$ and $b \in \arg \max_{C'} f$, but for every $\lambda' \in \Lambda$, $a\Delta_i^{\lambda'} b \notin \arg \max_C f$, because $a_i > b_i$ implies that for every $\lambda' \in [0, 1]$, $a\Delta_i^{\lambda'} b \neq a$. Again, f does not have (i, Λ) -increasing property. ■

The result in Quah (2007) is a special case of this result, when $\Lambda = [0, 1]$. The result here shows that if we want to weaken the notion of a \mathcal{C}_i -quasisupermodular function by requiring the condition to hold for fewer λ , then we must make the comparability of the set order more restrictive (that is, fewer sets can be ordered) by requiring less flexibility in the choice of λ as well. To say this differently, if we want a monotone comparative statics result applicable to a larger collection of constraint sets, we can expand the collection of sets that can be ordered by allowing the greatest flexibility in choosing λ , by setting $\Lambda = [0, 1]$. (This gives us the \mathcal{C}_i -flexible set order.) In this case, characterizing monotone

comparative static requires imposing the strictest conditions on the objective function by requiring $\Lambda = [0, 1]$. In particular, for every a and b , we are forced to consider the whole line segment joining a and $a \vee b$ and the whole line segment joining $a \wedge b$ and b , and we are essentially forced to consider convex sets.

One way to think about theorem 1 is that it presents monotone comparative statics in the \mathcal{C}_i -flexible set order but with new conditions on objective functions that do not force a restriction to convex sets. Moreover, these new conditions on the objective functions (i -quasisupermodular and i -single crossing property) are closer in flavor to the standard assumptions in the theory of monotone comparative statics, they do not use the relations $\Delta_i^\lambda, \nabla_i^\lambda$, and they work with the natural order and direction in \mathbb{R}^N . Furthermore, they can be naturally specialized to cardinal and differential conditions.

Appendix B. Some Proofs

One set of conditions under which sets can be ordered in the i -directional set order is as follows. Let X be a sublattice of \mathbb{R}^N , $f : X \rightarrow \mathbb{R}$, and $i \in \{1, 2, \dots, N\}$. The function f is *i -quasisubmodular on X* , if for every $a, b \in X$ with $a_i > b_i$, $f(a) \leq (<) f(a \wedge b) \implies f(a \vee b) \leq (<) f(b)$. The function f satisfies *dual i -single crossing property on X* , if for every $a, b \in X$ with $a_i > b_i$, and for every $v \in \{s(b - a \wedge b) \mid s \in \mathbb{R}, s \geq 0\}$ such that $a + v, b + v \in X$, $f(a) \leq (<) f(b) \implies f(a + v) \leq (<) f(b + v)$.

Proposition 6. *Let X be a convex sublattice of \mathbb{R}^N , $f : X \rightarrow \mathbb{R}$, and $i \in \{1, 2, \dots, N\}$. If f is continuous, (weakly) increasing, i -quasisubmodular, and satisfies dual i -single crossing on X , then $\tau \leq \tau' \implies \{x \mid f(x) \leq \tau\} \sqsubseteq_i^{dso} \{x \mid f(x) \leq \tau'\}$.*

Proof. Let $\tau \leq \tau'$, $A = \{x \mid f(x) \leq \tau\}$, $B = \{x \mid f(x) \leq \tau'\}$, and suppose $a \in A$, $b \in B$ with $a_i > b_i$. As case 1, suppose $f(b) \leq \tau$. In this case, let $v = 0$. Then $b - v = b \in A$ and $f(a) \leq \tau \leq \tau'$ implies that $a + v = a \in B$. As case 2, suppose $f(b) > \tau$. Then f is (weakly) increasing implies that $f(a \wedge b) \leq f(a) \leq \tau$. For $s \in [0, 1]$, consider $v(s) = s(b - a \wedge b)$. Then $s = 0$ implies $f(b - v(s)) > \tau$ and $s = 1$ implies $f(b - v(s)) \leq \tau$. By continuity, there is $\hat{s} \in (0, 1]$ such that $f(b - v(\hat{s})) = \tau$. Set $\hat{v} = \hat{s}(b - a \wedge b)$. Then $b - \hat{v} \in A$ and $f(b - \hat{v}) \geq f(a)$. As subcase 1, suppose $\hat{s} = 1$. Then $f(a \wedge b) = f(b - \hat{v}) \geq f(a)$ and i -quasisubmodularity implies that $f(b) \geq f(a \vee b)$, whence $a + 1(a \vee b - a) \in B$. As subcase 2, suppose $\hat{s} \in (0, 1)$. Applying dual i -single crossing to vectors to a and $b - \hat{v}$, with the directional vector $w = \frac{\hat{s}}{1-\hat{s}} [(b - \hat{v}) - a \wedge (b - \hat{v})]$ implies $f(b - \hat{v} + w) \geq f(a + w)$. But notice that $\hat{v} = \hat{s}(b - a \wedge b) = \hat{s} [(b - \hat{v}) - a \wedge b] + \hat{s} \hat{v} = \hat{s} [(b - \hat{v}) - a \wedge (b - \hat{v})] + \hat{s} \hat{v}$, and therefore, $\hat{v} = \frac{\hat{s}}{1-\hat{s}} [(b - \hat{v}) - a \wedge (b - \hat{v})] = w$. In other words, $f(b) \geq f(a + \hat{v})$, whence $a + \hat{v} \in B$. ■

It is easy to check that for given prices $p \gg 0$, the function $\phi : X \rightarrow \mathbb{R}$, $\phi(x) = p \cdot x$ satisfies these conditions. This provides another proof that with respect to wealth w , Walrasian budgets sets are ordered in the i -directional set order.

To show the equivalence of i -increasing differences (u) on X and i -increasing differences (*) on X , consider first the following slight modification of i -increasing differences (u) on X . Let X be a sublattice of \mathbb{R}^N , $f : X \rightarrow \mathbb{R}$, and $i \in \{1, 2, \dots, N\}$. f satisfies *i -increasing differences (σu)*, if for every $b \in X, u \in \mathbb{R}^N$ with $u_i > 0$, for every $\sigma, s \geq 0$, such that $b + \sigma u, b + s(-u)_+, b + \sigma u + s(-u)_+ \in X$, $f(b + \sigma u) - f(b) \leq f(b + \sigma u + s(-u)_+) - f(b + s(-u)_+)$. Consider the following equivalence.

Lemma 1. *Let X be a sublattice of \mathbb{R}^N , $f : X \rightarrow \mathbb{R}$, and $i \in \{1, 2, \dots, N\}$. f satisfies i -increasing differences (u) on X , if, and only if, f satisfies i -increasing differences (σu) on X .*

Proof. For sufficiency, fix $b \in X$, $u \in \mathbb{R}^N$ with $u_i > 0$, and fix $\sigma, s \geq 0$. If $\sigma = 0$, we are done, because left-hand side and right-hand side of the condition are both zero. Suppose

$\sigma > 0$. Let $\hat{u} = \sigma u$ and $\hat{s} = \frac{s}{\sigma} \geq 0$. Then $\hat{u}_i > 0$ and $\hat{s}(-\hat{u}_+) = s(-u)_+$, and therefore,

$$\begin{aligned} f(b + \sigma u) - f(b) &= f(b + \hat{u}) - f(b) \\ &\leq f(b + \hat{u} + \hat{s}(-\hat{u})_+) - f(b + \hat{s}(-\hat{u})_+) \\ &= f(b + \sigma u + s(-u)_+) - f(b + s(-u)_+), \end{aligned}$$

as desired. For necessity, let $\sigma = 1$. ■

Now recall that f satisfies i -increasing differences (*) on X , if for every $b \in X, u \in \mathbb{R}^N$ with $u_i > 0$, for every $\sigma \geq 0$, $f(b + \sigma u + s(-u)_+) - f(b + s(-u)_+)$ is (weakly) increasing in s , (where we consider only points $b + \sigma u + s(-u)_+, b + s(-u)_+ \in X$). Consider the following equivalence.

Lemma 2. *Let X be a sublattice of \mathbb{R}^N , $f : X \rightarrow \mathbb{R}$, and $i \in \{1, 2, \dots, N\}$. f satisfies i -increasing differences (σu) on X , if, and only if, f satisfies i -increasing differences (*) on X .*

Proof. Suppose f satisfies i -increasing differences (σu) on X . To check for i -increasing differences (*) on X , fix $b \in X, u \in \mathbb{R}^N$ with $u_i > 0$, and $\sigma \geq 0$. Fix $s_1 \leq s_2$. If $\sigma = 0$, we are done, because the expression is 0 for all s . Suppose $\sigma > 0$. Let $\hat{b} = b + s_1(-u)_+$ and $\hat{s} = s_2 - s_1 \geq 0$. Then

$$\begin{aligned} f(b + \sigma u + s_1(-u)_+) - f(b + s_1(-u)_+) &= f(\hat{b} + \sigma u) - f(\hat{b}) \\ &\leq f(\hat{b} + \sigma u + \hat{s}(-u)_+) - f(\hat{b} + \hat{s}(-u)_+) \\ &= f(b + \sigma u + s_1(-u)_+ + (s_2 - s_1)(-u)_+) - f(b + s_1(-u)_+ + (s_2 - s_1)(-u)_+) \\ &= f(b + \sigma u + s_2(-u)_+) - f(b + s_2(-u)_+), \end{aligned}$$

as desired.

Suppose f satisfies i -increasing differences (*) on X . To check that f satisfies i -increasing differences (σu) on X , fix $b \in X, u \in \mathbb{R}^N$ with $u_i > 0$, and fix $\sigma, s \geq 0$. Let $s_1 = 0$. Then $s_1 \leq s$, and therefore, $f(b + \sigma u) - f(b) = f(b + \sigma u + s_1(-u)_+) - f(b + s_1(-u)_+) \leq f(b + \sigma u + s(-u)_+) - f(b + s(-u)_+)$, as desired. ■

These lemmas imply the equivalence of i -increasing differences (u) on X and i -increasing differences (*) on X , as desired.