

DISABILITY, PAIN, PSYCHOLOGICAL FACTORS AND PHYSICAL PERFORMANCE IN HEALTHY CONTROLS, PATIENTS WITH SUB-ACUTE AND CHRONIC LOW BACK PAIN: A CASE-CONTROL STUDY

Jens Ivar Brox,¹ Kjersti Storheim,² Inger Holm,³ Astrid Friis³ and Olav Reikerås¹

From the ¹Department of Orthopaedic Surgery, Section for Back Surgery and Physical Medicine and Rehabilitation, National Hospital, Oslo, ²Norwegian University of Sports and Physical Education, Oslo and ³Department of Physiotherapy, National Hospital, Oslo, Norway

Objective: To compare measures of disability, psychological factors, pain and physical performance in healthy controls, and patients with sub-acute and chronic low back pain. To evaluate the concept of the deconditioning syndrome and to explore factors that may contribute to chronicity.

Design: Case-control study.

Subjects: Three age- and gender-matched groups were included in the study; healthy controls ($n = 45$), patients sick-listed 8–12 weeks ($n = 46$) and patients with chronic low back pain on a waiting list for lumbar instrumented fusion ($n = 45$).

Methods: Measures of disability, pain, psychological factors, and physical performance were obtained from the 3 groups using validated measures.

Results: Gender, age, body weight and height were not significantly different between the groups. Comparable scores were found for self-rated working ability, fear-avoidance beliefs for physical activity and aerobic capacity in the 2 patient groups. Oswestry Disability Index, pain, emotional distress, abdominal and back muscle endurance were significantly different between the 3 groups. Self-efficacy for pain and fear-avoidance beliefs for work was significantly different between the 2 patient groups.

Conclusion: The results suggest a stepwise deterioration of impairment and disability from healthy controls to patients with chronic low back pain. Most variables distinguished between healthy controls and patients with sub-acute or chronic low back pain. Deconditioning was more related to psychophysical measures of abdominal and back muscle endurance than to cardiovascular fitness. Comparable scores for fear-avoidance and working ability in the 2 patient categories suggest that these factors appear at an early stage and contribute to the transition from acute to chronic low back pain.

Key words: low back pain, healthy controls, sub-acute, chronic, fear-avoidance beliefs, physical performance.

J Rehabil Med 2005; 37: 95–99

Correspondence address: Jens Ivar Brox, Department of Orthopaedics, Section for Back Surgery and Physical Medicine and Rehabilitation, National Hospital, 0027 Oslo, Norway. E-mail: jens.ivar.brox@rikshospitalet.no

Submitted July 14, 2003; accepted June 1, 2004

© 2005 Taylor & Francis. ISSN 1650–1977
DOI 10.1080/16501970410017738

INTRODUCTION

Low back pain is a common complaint and its 1-year prevalence was 53% in a Norwegian epidemiological study (1). A Danish study reported that the lifetime, 1-year and point-prevalence of low back pain in the adult population was reduced in persons who are physically active at least 3 hours a week, compared with those who are less active (2).

Deconditioning is thought to be both a cause and a consequence of low back pain. The hypothesis of the deconditioning syndrome as a factor contributing to the chronicity of low back pain forms the basis for intensive physical rehabilitation of patients with chronic low back pain (CLBP). A recent study reported comparable levels of aerobic fitness in patients with CLBP, and therefore questioned the deconditioning myth. Moreover, they reported that there is no association between pain intensity and aerobic fitness. They concluded that deconditioning, defined as a lack of cardiovascular fitness levels normal for age and gender, therefore does not contribute to pain intensity in patients with CLBP (3).

Recent evidence suggests that psychosocial factors are important in predicting patients who will progress from an acute to a chronic stage. Patients with chronic pain often demonstrate somatization, anxiety and depression (4). Fear-avoidance beliefs have been hypothesized as the most important psychosocial factor in predicting disability among patients with CLBP. The cognitive-behaviour concept of how a chronic pain problem develops suggests that fear-avoidance behaviour appears at an early stage. A model for exaggerated pain perception was developed by Lethem et al. to explain why some individuals with acute pain develop chronic pain while others recover (5). This model proposes that the patients fear of pain, and subsequent avoidance behaviour, are determined by the relation between sensory and emotional components of pain. The cognitive model hypothesizes a vicious circle in which the patients beliefs and fears concerning symptoms and activity lead to unhelpful ways of managing symptoms, including avoidance behaviours, activity restriction and depression, that in turn lead to deconditioning which reflects a state of no condition or not feeling well.

The aim of the present study was to compare measures of disability, psychological factors, pain and physical performance in healthy controls, and patients with sub-acute and chronic low back pain. In addition, we wanted to evaluate the concept of

the deconditioning syndrome and to explore factors that may contribute to chronicity.

METHODS

Three age- and gender matched groups were included in the study; healthy controls ($n = 45$), patients sick-listed for low back pain 8–12 weeks ($n = 46$) and patients with CLBP on a waiting list for lumbar instrumented fusion ($n = 45$). Patients were matched for gender and age using 4 age cohorts (18–35 years, 36–45 years, 46–55 years and 55–67 years).

The control group included health workers who had not visited a physician, chiropractor or physiotherapist for back pain within the last 3 months. This group included physicians, nurses, nurse aids, secretaries, maintenance workers, cleaning and kitchen staff.

Patients in the sub-acute group had been sick-listed for 8–12 weeks due to non-specific LBP before inclusion. They were excluded if they had previous back surgery and had participated in regular physical exercise more than 3 times per week for the last 6 months. A more detailed description of these patients has been given previously (6).

Fifty-four percent ($n = 24$) of the patients on the waiting list had previous back surgery, while none of the participants in the other groups had been operated upon. The patients on the waiting list were labelled with the following diagnoses: facet joint syndrome (5 patients), spondylolisthesis (4 patients), spinal stenosis (7 patients), postlaminectomy or failed back surgery syndrome (24 patients) and degenerative disc disease (5 patients). Eighty percent ($n = 36$) were sick-listed or on disability pension (7).

Measures

Demographics. Background variables, including age, gender, height, weight, diagnosis, sick leave and physical activity, were recorded.

Pain. The participants scored their pain intensity on activity and at night on 9-point scales where 1 = no pain and 9 = worst pain (8).

Psychological factors. Fear-avoidance beliefs. Waddell's Fear-Avoidance Belief Questionnaire (FABQ) was used, higher numbers indicating increased levels of fear-avoidance beliefs (9). The questionnaire is divided into subscales for physical activity (FABQ-PA) and work (FABQ-W).

Self-efficacy beliefs for pain. Beliefs were assessed using 4 questions about how the patient believed he/she could manage pain, i.e. "Do you believe you can reduce your symptoms by your-self" (10). A high score indicates that the patient believed he/she was able to manage pain.

Emotional distress. Symptoms were rated by the short version of the Hopkins Symptom Check List (HSCL-25) (11). Patients rate 25 symptoms from 1 (not at all) to 4 (extremely). A mean symptom score of 1.75 or more was found in 20% of women and 9% of men in a large Norwegian epidemiological study (12). A score > 1.75 is a high predictor of current help-seeking, but seems to reflect illness or non-specific distress more than psychiatric diagnoses (13).

Life satisfaction was estimated by Cantrils Ladder Scale, a 10-point vertical numerical rating scale where 1 = very dissatisfied and 10 = very satisfied (14).

Disability. A Norwegian version of the original Oswestry Disability Index (ODI) (version 1.0) was used to evaluate condition-specific disability and pain (15, 16). This score has 10 questions about pain and pain-related disability in activities of daily life and social participation. Each question has 6 different response alternatives. The sum is calculated and presented as a percentage, where 0% represents no pain and disability and 100% represents the worst possible pain and disability.

Self-rated work capacity. Because most of the patients were out of work, a single question was used to assess the perceived capacity for work including paid work and housework. The response alternatives was graded: 1 = no limitations; 2 = some limitations, but able to do my ordinary work; 3 = periods not able to do my ordinary work; 4 = not able to do my ordinary work; and 5 = hardly self-reliant.

Physical performance. Cardiovascular fitness. Estimated by a sub-maximal bicycle ergometer test according to the method described by Aastrand (17).

Abdominal muscle endurance. Dynamic strength endurance of the

Table I. Mean (SD) characteristics of the participants

	Control ($n = 45$)	Sub-acute ($n = 46$)	Chronic ($n = 45$)
Age (years)	45.7 (8.1)	44.9 (9.2)	47.2 (10.2)
Gender (men/women)	21/24	22/26	21/24
Body weight (kg)	74.5 (10.8)	77.7 (16.2)	75.6 (15.7)
Height (cm)	173.8 (10.8)	173.0 (16.2)	172.0 (15.7)

abdominal muscles was measured with the patient in the supine position, knees bent at a 90° angle and with unsupported feet flat at the surface. With straight arms, a slow curl-up was performed until the fingertips just touched the proximal border of each patella. A metronome was set to 80 beats per minute and subjects took 2 beats to curl up and 2 beats to curl down (i.e. 20 curl-ups per minute). Performance with jerks or heels rise from the surface was not accepted. A maximum limit of 120 curl-ups was set (18, 19).

Back muscle endurance. Isometric endurance of trunk extensors was evaluated by measuring how many seconds (maximum 240 seconds) the patient was able to maintain the unsupported upper part of the body horizontal, when placed prone on a couch with legs fixed and the arms along the trunk. This modification of the Biering-Sørensen test is used in a former study of patients sick-listed for sub-acute LBP (20).

Statistical analysis

The non-parametric Kruskal-Wallis one-way analysis of variance rank test and the Mann-Whitney rank sum test were applied to compare the groups. To protect against Type 1 errors while performing a lot of different comparisons, the level of significance was set to 0.01 (21). Multiple linear regression was used to adjust for the influence of pain and psychological impairment on physical performance (22, 23).

RESULTS

Characteristics of the patients are given in Table I. Gender, age, body weight and height were not significantly different between the groups. Self-rated working capacity ($p = 0.3$) and fear-avoidance (FABQ-PA) ($p = 0.7$) were not significantly different between patients with sub-acute and chronic pain (Table II). When adjusted for pain and emotional distress, group explained less than 1% of the variance in aerobic capacity and the difference between patient groups was not significant ($p = 0.60$). Entering fear-avoidance (FABQ-PA) into the model and excluding the healthy controls from the model did not change results. Pain was not significantly associated with cardiovascular endurance. When adjusted for pain and emotional distress, group explained 27% of the variance in back muscle endurance and 21% of the variance in abdominal muscle endurance and differences between groups were significant ($p < 0.001$). All other outcome measures were significantly different between the 3 groups, demonstrating a stepwise increase in psychological impairment and disability, and a decrease in physical performance (Table II). Figure 1 shows separate box-plots in men and women for all outcome variables in the 3 groups.

DISCUSSION

We found a stepwise deterioration of pain, disability, psychological factors and physical performance from healthy controls to patients with CLBP. All the measures used except aerobic

Table II. Mean (SD) results of the participants

	Control (n = 45)	Sub-acute (n = 46)	Chronic (n = 45)	p-value for differences
Oswestry Disability Index	1.6 (3.5)	25.2 (12.2)	43.5 (13.3)	Co-S: <0.001 Co-Chr: <0.001 S-Chr: <0.001
Pain on activity	1.1 (0.4)	4.9 (1.9)	6.3 (1.9)	Co-S: <0.001 Co-Chr: <0.001 S-Chr: 0.002
Pain at rest	1.0 (0.4)	2.9 (1.9)	5.2 (2.2)	Co-S: <0.001 Co-Chr: <0.001 S-Chr: <0.001
Self-efficacy for pain Emotional distress	1.2 (0.2)	4.4 (1.4) 1.4 (0.4)	3.5 (1.5) 1.8 (0.6)	S-Chr: 0.004 Co-S: <0.001 Co-Chr: <0.001 S-Chr: <0.001
Life satisfaction	8.3 (1.7)	6.8 (2.0)	5.0 (1.9)	Co-S: <0.001 Co-Chr: <0.001 S-Chr: <0.001
Fear-avoidance physical activity		3.4 (1.2)	3.3 (1.5)	S-Chr: 0.73
Fear-avoidance work		3.4 (1.3)	4.4 (1.6)	S-Chr: <0.001
Self-rated work capacity	1.0 (0.3)	2.5 (0.9)	2.7 (1.0)	S-Chr: 0.30
Aerobic endurance (VO ₂ /mlkg)	34.7 (9.0)	30.7 (5.8)	31.8 (8.6)	Co-S: 0.004 Co-Chr: 0.06 S-Chr: 0.63
Abdominal muscle endurance (number of sit-ups)	35 (12)	27 (16)	19 (11)	Co-S: <0.001 Co-Chr: 0.002 S-Chr: 0.008
Back muscle endurance (seconds)	110 (49)	77 (40)	46 (42)	Co-S: 0.002 Co-Chr: <0.001 S-Chr: <0.001

endurance, fear-avoidance beliefs for physical activity and self-reported disability were able to distinguish between groups.

Cardiovascular fitness was not significantly different between healthy controls and patients with CLBP. Scores for fear-avoidance beliefs for physical activity and self-reported disability were comparable in patients with sub-acute and CLBP. Our results suggest that physical performance is just one component of the deconditioning syndrome, and that the concept may be misleading if interpreted as a reduction in cardiovascular fitness.

The hypothesis of the deconditioning syndrome as a factor contributing to chronicity forms the basis of intensive physical rehabilitation for CLBP (24). Deconditioning is thought to be both a cause and a consequence of back pain. In agreement with Wittink et al. we found no association between pain and aerobic endurance (3). In addition, the differences between groups are within measurement error for the Astrand test (25). The small differences between groups observed for cardiovascular fitness does not mean that exercises and physical fitness training are unhelpful for patients with CLBP. Physical exercise was ranked as the most important element in a comprehensive treatment programme in patients with depression, but non-aerobic forms were as effective as aerobic forms of exercise in the treatment of anxiety disorders and depression (26, 27). The reduction in psychological symptoms following exercises observed in these studies, and in recently published studies in patients with sub-acute and CLBP, are most probably mediated by psychological mechanisms, such as distraction and mastering (6, 27, 28).

A recent British study suggested that patients who have back pain for more than 3 months may require referral to psychological services (29). In keeping with previous studies we found considerably elevated scores for emotional distress in CLBP (4, 13, 30). In the sub-acute group mean scores for emotional distress were moderately elevated. The most important psychological finding in this group was the high scores on fear-avoidance for physical activity.

A large British survey suggested that the rises in outpatient attendance and sickness absence for low back pain are not explained by a greater incidence of severe back disease (31). They observed a change in back pain over the last decade that does not greatly impair function and suggested that a cultural shift may have rendered back pain more acceptable as a reason for absence attributed to sickness. The results from the British survey suggest that the rises in outpatient attendance and sickness absence may be attributed advice given by physicians, physiotherapists and others. Thus, the high scores for fear-avoidance of patients with sub-acute pain in the present study may reflect advice given in the acute phase and suggest that they believe that they should restrict physical activity to protect themselves from injury and chronic back pain. Although fear-avoidance is a natural psychological reaction in patients with acute back pain, prolonged avoidance is an example of maladaptive behaviour.

Fear-avoidance beliefs were not assessed in the healthy controls. According to current models personality traits and experiences pre-dating the onset of pain largely determine fear of pain and avoidance behaviour (32). Nevertheless, we considered that measures of fear-avoidance in the absence of




Fig. 1. Box-and-whisker plots of outcome variables for men and women in the 3 groups. Boxes show the median, 25th and 75th percentiles, whiskers show the values that are not outliers.

pain would be doubtful and did not use these questions in the healthy controls. Recent studies have reported elevated scores in patients with acute low back pain, and that fear-avoidance

believes predicts future disability (33, 34). If fear-avoidance beliefs are present from the initial experience of LBP, and represent detriment to recovery, interventions in the acute or

sub-acute stage may be useful in reducing fear-avoidance and promoting return to normal activity. A recently published clinical trial in patients with acute low back pain showed promising results in patients assigned to receive fear-avoidance based physiotherapy (35). In previously published clinical trials we have reported that cognitive intervention and exercises effectively reduced FABQ (PA) in patients with subacute and CLBP (6, 28).

A limitation of the present study is that it is small and has a cross-sectional design that makes it prone to selection bias. It is difficult to make a causal inference. We did not assess the level of physical activity in all participants, but patients with sub-acute pain were not included if they performed physical exercise more than 3 times a week, which may have contributed to significantly lower aerobic endurance in this group.

In conclusion the findings of the present study highlight the importance of fear-avoidance for physical activity in patients with sub-acute low back pain and question the concept of the deconditioning syndrome of patients with low back pain.

REFERENCES

- Natvig B, Nessjøy I, Bruusgaard D, Rutle O. Musculoskeletal symptoms in a local community. *Eur J Gen Practice* 1995; 1: 25–28.
- Harreby M, Hesselsø G, Kjær J, Neergaard K. Low back pain and physical exercise in leisure time in 38-year old men and women: a 25-year prospective cohort study of 640 school children. *Eur Spine J* 1997; 6: 181–186.
- Wittink H, Michel TH, Sukiennik A, Gascon C, Rogers W. The association of pain with aerobic fitness in patients with chronic low back pain. *Arch Phys Med Rehabil* 2002; 83: 1467–1471.
- Gatchel RJ, Polatin PB, Mayer T, Gatchel R. Psychopathology and the rehabilitation of patients with chronic low back pain disability. *Arch Phys Med Rehabil* 1994; 75: 666–700.
- Lethem J, Slade PD, Troup JD, Bentley G. Outline of a fear-avoidance model of exaggerated pain perception – I. *Behav Res Ther* 1983; 21: 401–408.
- Storheim K, Brox JI, Holm I, Koller AK, Bø K. Intensive group training compared with cognitive intervention in patients sick-listed for sub-acute low back pain: a single blind randomized controlled trial. *J Rehabil Med* 2003; 35: 132–140.
- Brox JI, Friis A, Holm I, Grundnes O, Sørensen R, Lange JE, et al. Patients with chronic degenerative spinal disease – can conservative treatment reduce the waiting list for surgery? *Tidsskr Nor Lægeforen* 1999; 119: 1784–1787.
- Brox JI, Staff PH, Ljunggren AE, Brevik JI. Arthroscopic surgery compared with supervised exercises in patients with rotator cuff disease. *BMJ* 1993; 307: 899–903.
- Waddell G, Newton M, Henderson I, Somerville D, Main CJ. A fear-avoidance beliefs questionnaire (FABQ) and the role of fear-avoidance beliefs in chronic low back pain and disability. *Pain* 1993; 52: 157–168.
- Lorig K, Chastain RL, Ung E, Shoor S, Holman HR. Development and evaluation of a scale to measure perceived self-efficacy in people with arthritis. *Arthritis Rheum* 1989; 32: 37–44.
- Derogatis LR, Lipman RS, Rickels K, Uhlenuth EH, Cori L. The Hopkins Symptom Checklist (HSCL): a self-report inventory. *Behav Sci* 1974; 19: 1–15.
- Sandanger I, Nygård JF, Ingebrigtsen G, Sørensen T, Dalgard OS. Prevalence, incidence and age at onset of psychiatric disorders in Norway. *Soc Psychiatry Psychiatr Epidemiol* 1999; 34: 570–579.
- Sandanger I, Moum T, Ingebrigtsen G, Sørensen T, Dalgard OS, Bruusgaard D. The meaning and significance of caseness: The Hopkins Symptom Checklist – 25 and the composite international diagnostic interview. II. *Soc Psychiatry Psychiatr Epidemiol* 1999; 34: 53–59.
- Andrews FM, Robinson JP. Measures of subjective well-being. In: Robinson JP, Shaver PR, Wrightsman LS, eds. *Measures of personality and social psychological attitudes*. San Diego: Academic Press; 1991.
- Fairbank JCT, Davis J, Couper J, O'Brian JP. Oswestry disability questionnaire. *Physiotherapy* 1980; 66: 271–273.
- Roland M, Fairbank J. The Roland-Morris Disability Questionnaire and the Oswestry Disability Questionnaire. *Spine* 2000; 25: 3115–3124.
- Astrand PO. Quantification of exercise capability and evaluation of physical capacity in man. *Prog Cardiovasc Dis* 1976; 19: 51–67.
- Faulkner RA, Sprigings EJ, McQuarrie A, Bell RD. A partial curl-up protocol for adults based on an analysis of two procedures. *Can J Sport Sci* 1989; 14: 135–141.
- Hyytiäinen K, Salminen JJ, Suviola T, Wickström, Pentti J. Reproducibility of nine tests to measure spinal mobility and trunk muscle strength. *Scand J Rehabil Med* 1991; 23: 3–10.
- Lindström I, Öhlund C, Eek C, Wallin L, Peterson L-E, Nachemson A. Mobility, strength, and fitness after a graded activity program for patients with subacute low back pain. *Spine* 1992; 17: 641–649.
- Tabachnick BG, Fidell LS. *Using multivariate statistics*. 3rd edn. New York: Harper Collins; 1996.
- Brox JI, Brevik JI, Ljunggren AE, Staff PH. Influence of anthropometric and psychological variables pain and disability on isometric endurance of shoulder abduction in patients with rotator tendinosis of the shoulder. *Scand J Rehabil Med* 1996; 28: 193–200.
- Keller A, Johansen JG, Hellesnes J, Brox JI. Predictors of isokinetic back muscle strength in patients with low back pain. *Spine* 1999; 24: 275–280.
- Mayer TG, Gatchel RJ, Mayer H, Kishino ND, Keeley J, Mooney V. A prospective two-year study of functional restoration in industrial low back injury. An objective assessment procedure. *JAMA* 1987; 258: 1763–1767.
- Keller A, Hellesnes J, Brox JI. Reliability of the isokinetic trunk extensor test, Biering-Sorensen test, and Aastrand bicycle test: assessment of intraclass correlation coefficient and critical difference in patients with chronic low back pain and healthy individuals. *Spine* 2001; 26: 771–777.
- Martinsen EW, Medhus A, Sandvig L. Effects of aerobic exercise on depression: a controlled study. *BMJ* 1985; 291: 109.
- Martinsen EW, Hoffart A, Solberg Ø. Aerobic and non-aerobic forms of exercise in the treatment of anxiety disorders. *Stress Medicine* 1989; 5: 115–120.
- Brox JI, Sørensen R, Friis A, Nygaard Ø, Indahl A, Keller A, et al. Lumbar instrumental fusion compared with cognitive intervention and exercises of patients with chronic low back pain and disc degeneration. *Spine* 2003; 28: 1913–1921.
- Waxman R, Tennant A, Helliwe P. Community survey of factors associated with consultation. *BMJ* 1998; 317: 1564–1567.
- Rush AJ, Polatin P, Gatchel R. Depression and chronic low back pain. *Spine* 2000; 20: 2566–2571.
- Palmer KT, Walsh K, Bendall H, Cooper C, Coggon D. Back pain in Britain: comparison of two prevalence surveys at an interval of 10 years. *BMJ* 2000; 320: 1577–1578.
- Vlaeyen JW, Linton SJ. Fear-avoidance and its consequences in chronic musculoskeletal pain: a state of the art. *Pain* 2000; 85: 317–332.
- Fritz JM, George SZ, Delitto A. The role of fear-avoidance beliefs in acute low back pain: relationships with current and future disability and work status. *Pain* 2001; 94: 7–15.
- Crombez G, Vlaeyen JW, Heuts PH, Lysens R. Pain-related fear is more disabling than fear itself: evidence and role of pain-related fear in chronic back pain disability. *Pain* 1999; 80: 329–339.
- George SZ, Fritz JM, Bialosky JE, Donald DA. The effect of fear-avoidance-based physical therapy intervention for patients with acute low back pain: results of a randomized trial. *Spine* 2003; 28: 2551–2560.