

Erratum to: Discontinuation of adalimumab treatment in rheumatoid arthritis patients after achieving low disease activity

Masayoshi Harigai · Tsutomu Takeuchi ·
Yoshiya Tanaka · Tsukasa Matsubara ·
Hisashi Yamanaka · Nobuyuki Miyasaka

Published online: 14 March 2012
© Japan College of Rheumatology 2012

Erratum to: Mod Rheumatol DOI 10.1007/s10165-011-0586-5

The authors regret that the names of some members of the BRIGHT Study Investigators Group were missing or misspelled in the Appendix of this article. The corrected Appendix is provided here.

Appendix

The following rheumatologists are the members of the BRIGHT Study Investigators Group: T. Koike, T. Atsumi, T. Odani (Hokkaido University); H. Takahashi (Sapporo Medical University); M. Mukai (Sapporo City General Hospital); S. Ohta (Taga General Hospital); T. Sumida,

For the BRIGHT Study Investigators Group.

The members of the BRIGHT Study Investigators Group are listed in the “[Appendix](#).”

The online version of the original article can be found under doi:[10.1007/s10165-011-0586-5](https://doi.org/10.1007/s10165-011-0586-5).

M. Harigai (✉)
Department of Pharmacovigilance, Graduate School of Medical and Dental Sciences, Tokyo Medical and Dental University, 1-5-45 Yushima, Bunkyo-ku, Tokyo 113-8519, Japan
e-mail: mharigai.mpha@tmd.ac.jp

M. Harigai · N. Miyasaka
Department of Medicine and Rheumatology,
Graduate School of Medical and Dental Sciences,
Tokyo Medical and Dental University, Tokyo, Japan

T. Takeuchi
Division of Rheumatology and Clinical Immunology,
Department of Internal Medicine, Faculty of Medicine,
Keio University, Tokyo, Japan

T. Hayashi (Tsukuba University); H. Nakajima, N. Watanabe, K. Ikeda (Chiba University); R. Matsumura (NHO Chiba East Hospital); T. Mimura, Y. Akiyama (Saitama Medical University); T. Hanyu (Nagaoka Red Cross Hospital); Y. Kaneko, N. Kimura (Keio University); T. Kasama (Showa University); H. Yamagata (NHO Murayama Medical Center); H. Yamada (St. Marianna University School of Medicine); S. Nagaoka (Yokohama Minami Kyou Sai Hospital); S. Tohma, T. Matsui (NHO Sagamihara National Hospital); S. Hirohata, T. Nagai (Kitasato University); H. Taki (Toyama University); K. Sugimoto (Fukui General Hospital); T. Tsuji (Fukui Onsen Hospital); N. Ishiguro, T. Kojima, D. Katoh (Nagoya University); Y. Eto, A. Kaneko, D. Kida, T. Sato, M. Tsukamoto (NHO Nagoya Medical Center); T. Fujii (Kyoto University); Y. Kawahito, A. Yamamoto (Kyoto Prefectural University); S. Kumagai, A. Morinobu (Kobe University); H. Makino, H. Wakabayashi (Okayama University); E. Suematsu, H. Miyahara (NHO Kyusyu Medical Center); T. Fukuda, S. Honda, M. Matsuoka, M. Yano (Kurume University); K. Saito, S. Hirata (University of Occupational and Environmental Health).

Y. Tanaka
The First Department of Internal Medicine,
University of Occupational and Environmental Health,
Kitakyushu, Japan

T. Matsubara
Matsubara Mayflower Hospital, Hyogo, Japan

H. Yamanaka
Institute of Rheumatology,
Tokyo Women's Medical University, Tokyo, Japan

N. Miyasaka
Global Center of Excellence Program, International Research
Center for Molecular Science in Tooth and Bone Diseases,
Tokyo, Japan