## Discontinuous Galerkin Methods for Solving Elliptic and Parabolic Equations. Theory and Implementation B. Riviere List of typos June 2012

Here is a list of misprints and clarifications. I would like to thank the readers for helping find the typos.

- page 3 line -3: replace j by n.
- page 29 line -10: the variable  $\epsilon$  is misplaced. The correct formula is:

$$L(v) = \int_{\Omega} fv + \sum_{e \in \Gamma_{\mathrm{D}}} \int_{e} \left( \epsilon \mathbf{K} \nabla v \cdot \mathbf{n}_{e} + \frac{\sigma_{e}^{0}}{|e|^{\beta_{0}}} v \right) g_{\mathrm{D}} + \sum_{e \in \Gamma_{\mathrm{N}}} \int_{e} v g_{\mathrm{N}}.$$

- page 31 line 5: clarification. The space  $\mathcal{D}(E)$  is the space of  $\mathcal{C}^{\infty}$  functions with compact support in E.
- page 52 line -5: the sign for the first term in the formula for  $m_e^{21}$  is wrong. The correct formula is:

$$m_e^{21} = \frac{1}{2} \int_e \mathbf{K} \nabla P_{h,1} \cdot \mathbf{n}_e v_2 + \frac{\epsilon}{2} \int_e \mathbf{K} \nabla v_2 \cdot \mathbf{n}_e P_{h,1} - \frac{\sigma_e^0}{|e|^{\beta_0}} \int_e P_{h,1} v_2.$$

• page 53 line 2: the sign for the first term in the formula for  $\mathbf{M}_{e}^{21}$  is wrong. The correct formula is:

$$(\mathbf{M}_{e}^{21})_{ij} = \frac{1}{2} \int_{e} \mathbf{K} \nabla \phi_{j,E_{e}^{1}} \cdot \mathbf{n}_{e} \phi_{i,E_{e}^{2}} + \frac{\epsilon}{2} \int_{e} \mathbf{K} \nabla \phi_{i,E_{e}^{2}} \cdot \mathbf{n}_{e} \phi_{j,E_{e}^{1}} - \frac{\sigma_{e}^{0}}{|e|^{\beta_{0}}} \int_{e} \phi_{j,E_{e}^{1}} \phi_{i,E_{e}^{2}} \cdot \mathbf{n}_{e} \phi_{j,E_{e}^{1}} \cdot \mathbf{n}_{e} \phi_{j,E_{e}^{1}} + \frac{\epsilon}{2} \int_{e} \mathbf{K} \nabla \phi_{i,E_{e}^{2}} \cdot \mathbf{n}_{e} \phi_{j,E_{e}^{1}} \cdot \mathbf{$$

• page 53 line 7: the correct formula for  $(\mathbf{b}_e)_i$  is:

$$(\mathbf{b}_e)_i = \int_e \left( \epsilon \mathbf{K} \nabla \phi_{i, E_e^1} \cdot \mathbf{n}_e + \frac{\sigma_e^0}{|e|^{\beta_0}} \phi_{i, E_e^1} \right) g_{\mathrm{D}}.$$

• page 55 line 4: the correct formula for  $\mathbf{M}_{e}^{21}(i,j)$  is:

$$\mathbf{M}_{e}^{21}(i,j) = \mathbf{M}_{e}^{21}(i,j) - \sigma_{e}^{0}w(k)\phi_{i,E_{e}^{2}}(s(k))\phi_{j,E_{e}^{1}}(s(k))$$

- page 55 line 5 of algorithm 2.3: the correct sentence is: for i = 1 to  $N_{loc}$  do.
- page 73 line -6:  $H_1^0(\Omega)$  should read  $H_0^1(\Omega)$ .
- page 74 line 10: in the definition of the energy norm, for the second term,  $||v||^2_{L^2(e)}$  should read  $||[v]||^2_{L^2(e)}$ .
- page 84 line 13: the term  $(\mathbf{M} + \Delta t\mathbf{A})$  should read  $(\mathbf{M} \Delta t\mathbf{A})$
- page 127 lines -1, -2: the terms  $h^k |\boldsymbol{u}|_{H^{k+1}(\Omega)}$  should read  $h^{2k} |\boldsymbol{u}|_{H^{k+1}(\Omega)}^2$ .
- page 129 line 6: the variable U should read  $U_h$ .
- page 129 last line of Theorem 6.12: the line should read where  $\delta = 1$  for SIPG and  $\delta = 0$  for IIPG and NIPG.
- page 166 line -4: the third argument for Aloc should be removed: Aloc(idofs,jdofs). The same comment holds for the third argument of variables Bloc11, Bloc22, Bloc12, Bloc21 on pages 170 and 171.