



UvA-DARE (Digital Academic Repository)

Discovery of circularly polarized radio emission from SS 433

Fender, R.P.; Rayner, D.; Norris, R.; Sault, R.; Pooley, G.G.

DOI

[10.1086/312481](https://doi.org/10.1086/312481)

Publication date

2000

Published in

Astrophysical Journal

[Link to publication](#)

Citation for published version (APA):

Fender, R. P., Rayner, D., Norris, R., Sault, R., & Pooley, G. G. (2000). Discovery of circularly polarized radio emission from SS 433. *Astrophysical Journal*, 530, L29. <https://doi.org/10.1086/312481>

General rights

It is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), other than for strictly personal, individual use, unless the work is under an open content license (like Creative Commons).

Disclaimer/Complaints regulations

If you believe that digital publication of certain material infringes any of your rights or (privacy) interests, please let the Library know, stating your reasons. In case of a legitimate complaint, the Library will make the material inaccessible and/or remove it from the website. Please Ask the Library: <https://uba.uva.nl/en/contact>, or a letter to: Library of the University of Amsterdam, Secretariat, Singel 425, 1012 WP Amsterdam, The Netherlands. You will be contacted as soon as possible.

DISCOVERY OF CIRCULARLY POLARIZED RADIO EMISSION FROM SS 433

ROBERT FENDER,¹ DAVID RAYNER,² RAY NORRIS,³ R. J. SAULT,³ AND GUY POOLEY⁴

Received 1999 October 26; accepted 1999 December 3; published 2000 January 10

ABSTRACT

We report the discovery of circularly polarized radio emission from the radio-jet X-ray binary SS 433 with the Australia Telescope Compact Array. The flux density spectrum of the circular polarization, clearly detected at four frequencies between 1 and 9 GHz, is of the form $V \propto \nu^{-0.9 \pm 0.1}$. Multiple components in the source and a lack of very high spatial resolution do not allow a unique determination of the origin of the circular polarization or of the spectrum of fractional polarization. However, we argue that the emission is likely to arise in the inner regions of the binary, possibly via propagation-induced conversion of linear to circular polarization, and the fractional circular polarization of these regions may be as high as 10%. Observations such as these have the potential to help us investigate the composition, whether pairs or baryonic, of the ejecta from X-ray binaries.

Subject headings: accretion, accretion disks — ISM: jets and outflows — radio continuum: stars — stars: individual (SS 433)

1. INTRODUCTION

High-velocity synchrotron-emitting jets are commonly observed in active galactic nuclei (AGNs; e.g., Ostrowski et al. 1997; Shields 1999) and X-ray binary (XRB) systems, both of which contain black holes and neutron stars (e.g., Hjellming & Han 1995; Fender 2000 and references therein). The composition of the jet plasma, whether electron-proton (e^-p^+) or electron-positron (e^-e^+), remains a fundamental yet unanswered question in nearly all cases.

SS 433 is one of the most celebrated of Galactic objects. The source is an XRB system consisting, most probably, of a mass-losing star in a 13 day orbit with a stellar-mass black hole or neutron star. The system produces bright quasi-continuous radio jets that precess with a period of ~ 162.5 days (Vermeulen 1989, 1993; Brinkmann 1998). Moving optical emission lines (Margon 1984 and references therein) indicate a jet velocity of $\beta = v/c = 0.26$, confirmed by both VLA and VLBI radio observations (Vermeulen 1993 and references therein). These optical lines and their X-ray counterparts (Kotani et al. 1996 and references therein) are the only direct evidence for the existence of baryonic material (i.e., e^-p^+) in a jet from any X-ray binary.

Progress toward determining the composition of the plasma in jets from AGNs has been made recently by the detection and modeling of a circularly polarized radio component from the quasar 3C 279 (Wardle et al. 1998). Wilson & Weiler (1997) argue that radio circular polarization upper limits for the Crab supernova remnant come close to determining the positron content of the nebula. In addition Bower, Falcke, & Backer (1999) and Sault & Macquart (1999) have recently detected circularly polarized radio emission from Sagittarius A* at the Galactic center.

In this Letter, we report the detection of circularly polarized radio emission from SS 433, the first from any XRB system,

at four radio frequencies. This observation has the potential to be the benchmark against which other jets from other XRB systems may be compared in an effort to determine whether they produce e^-p^+ or e^-e^+ jets.

2. OBSERVATIONS AND RESULTS

The Australia Telescope Compact Array (ATCA) consists of six 22 m altitude-azimuth antennas near Narrabri, New South Wales (Frater, Brooks, & Whiteoak 1992). Each ATCA antenna is equipped with two wide-band feed horns, and each feed horn is equipped with two pairs of orthogonal-linear probes. This allows both orthogonal polarizations at two separate frequencies to be observed with each feed horn at the same time. Observations were made using the ATCA “continuum mode,” which gives bandwidths of 128 MHz simultaneously at each of two frequencies and four correlation products (XX, YY, XY, and YX). The observations on 1999 May 10 were centered on 4.80 and 8.64 GHz. On 1999 May 20, observations at 1.38 and 2.50 GHz were alternated with those at 4.80 and 8.64 GHz, with a cycle time of ~ 25 minutes. Calibration sources, selected from the 1997 ATCA Calibrator Source Catalogue by J. E. Reynolds,⁵ were observed every cycle. The ATCA primary calibrator PKS 1934–638 was used to calibrate the bandpass and to set the absolute flux scale (Reynolds 1994).

Data reduction was performed with the MIRIAD package (Sault, Teuben, & Wright 1995). Prior to gain calibration, the XY-phase correction measured by a noise-diode system was applied, and the data were corrected for a small (otherwise unmodeled) field rotation due to the antenna’s pointing model (Kesteven 1997). The main calibration step involves simultaneously solving for time-dependent complex gains, a time-independent residual XY-phase, and time-independent polarization leakages for each feed as well as the linear and circular polarization of the calibration source. A good parallactic-angle coverage is required in order that the leakages and calibrator linear polarization can be decoupled in the solution process through the relative rotation of the feeds and parallactic angle (Conway & Kronberg 1969).

Circular polarization data require calibration using the “strongly polarized” equations (Sault, Killeen, & Kesteven 1991), which include terms in the leakage times the Stokes

¹ Astronomical Institute “Anton Pannekoek,” University of Amsterdam, and Center for High Energy Astrophysics, Kruislaan 403, Amsterdam, SJ NL-1098, Netherlands.

² School of Physics and Mathematics, University of Tasmania, GPO Box 252-21, Hobart, TAS 7001, Australia.

³ Australia Telescope National Facility, CSIRO, P.O. Box 76, Epping, NSW 1710, Australia.

⁴ Mullard Radio Astronomy Observatory, Cavendish Laboratory, Madingley Road, Cambridge, CB3 0HE, England, UK.

⁵ See <ftp://ftp.atnf.csiro.au/pub/atnfdocs/guides/at.cat>.

parameters Q and U . To obtain such a solution requires a calibration source that has a few percent of linear polarization (for this experiment, PKS 1908–202) so that there is sufficient signal in the second-order terms. The accuracy of the leakage solution was estimated by repeating the entire calibration procedure for the 1999 May 20 observation, for all four frequencies, with a different calibrator (either PKS 1947+079 or PKS 2029+121). For SS 433, the differences in the resultant circular polarization are of the same order as the errors expected from system noise alone; these differences have been incorporated into our circular polarization error estimates.

All calibrators were imaged in circular polarization as a consistency check, and the results for 1999 May 20 are shown in Figure 1. The leakage calibration of these calibrators was constrained so that the results gave zero circular polarization for PKS 1934–638 (an absolute circular polarization calibrator is needed for the interferometer; see Sault, Hamaker, & Bregman 1996). A bias in the observed circular polarization of compact steep-spectrum sources observed at the ATCA (Rayner, Norris, & Sault 2000) suggests that PKS 1934–638 is in fact circularly polarized with $V/I \sim 2.5 \times 10^{-4} \pm 0.5 \times 10^{-4}$ at 4.8 GHz. An error of this order in the absolute circular polarization flux scale is much smaller than the observed circular polarization of SS 433 and does not affect the conclusions of this Letter. The sign convention of V follows the IAU convention (transactions of the IAU, vol. 15B, 166 [1973]), which conforms to the IEEE definition (1969, Standard Definitions of Terms for Radio-Wave Propagation, IEEE Trans. AP-17, 270).

SS 433 was slightly resolved (i.e., $\geq 1''$) in the east-west direction in both total intensity and linear polarization, consistent with the observations of Hjellming & Johnston (1981). However, the source is unresolved in Stokes V , consistent with all the circularly polarized flux arising in an unresolved point source.

On 1999 May 10, we also observed the radio-jet X-ray binary GRS 1915+105; at the time, the source had a flat spectrum between 4.80 and 8.64 GHz, at a flux density of ~ 20 mJy. We can place 3σ limits of 1.2% on both the fractional linear and circular polarization of the emission from this source.

In addition to our ATCA observations, we also utilize data from the Green Bank Interferometer (GBI) variable source monitoring program. These data provide a long-term view of the state of activity of the radio source with daily flux density measurements at 2.3 and 8.3 GHz (see, e.g., Waltman et al. 1991).

3. EVOLUTION OF THE PROPAGATING EJECTA

Both the GBI monitoring data and our ATCA observations are presented in Figure 2. Note that the lower two panels show the flux density measured in linear and circular polarization, respectively (i.e., not the *fractional* spectra). A best fit to the observed flux density spectrum of the circular polarization has a spectral index of $\alpha_\nu = -0.9 \pm 0.1$ (where $V \propto \nu^{\alpha_\nu}$). Between the two epochs, it is clear that the total radio flux dropped by $\geq 40\%$ and was dominated by the decaying stages of the two major flare events that peaked at 2 GHz on \sim MJD 51,304 and \sim MJD 51,312, respectively. Over the same period, the linearly polarized flux may have increased slightly, and the circularly polarized flux appears to have remained constant.

Our current understanding of the evolution of external and internal opacity effects in the outflow from SS 433 is summarized in Figure 3 (based on Hjellming & Johnston 1981, Vermeulen 1993, and Paragi et al. 1999). Also indicated are our best estimates of the physical locations of the components




FIG. 1.—Fractional circular polarization for SS 433 and calibrators for 1999 May 20 observations. The scale is set by assuming that PKS 1934–638 has Stokes $V = 0$ at all frequencies (see text).

corresponding to the two flare events, at the times of our observations on 1999 May 10 and 20. We can presume that on our first epoch, we only observed linear polarization from the first ejection, which contributed $\leq 1/3$ of the total flux density at this time. By our second epoch of observations, both components contributed to the linearly polarized flux. Qualitatively, this can explain the increase in the linearly polarized flux density between the two epochs, even though the total flux density decreased. Qualitatively, similar behavior is seen in the evolution of linear polarization in ejections from GRS 1915+105 (Fender et al. 1999), where the core also remains persistently (linearly) depolarized.

4. THE ORIGIN OF THE CIRCULAR POLARIZATION

Circular polarization may be produced in a synchrotron-emitting plasma either directly as a result of the synchrotron process or via conversion of linear to circular polarization (Kennett & Melrose 1998 and references therein). Below we briefly discuss the possible interpretations of the circularly polarized radio emission from SS 433 in the context of these models.

4.1. Intrinsicly Circularly Polarized Synchrotron Emission

An electron of Lorentz factor γ can be considered to radiate synchrotron emission primarily at a frequency $\nu = 4.2B_\perp\gamma^2$ MHz, where B_\perp is the component of the magnetic field perpendicular to the line of sight, measured in units of gauss. The fractional circular polarization $m_c (=|V|/I)$ produced intrinsically by synchrotron radiation is of order $1/\gamma$ (Legg & Westfold 1968), and hence the observed circular polarization spectrum should follow the relation $m_c \propto \nu^{-1/2}$. For an observed $m_c \sim 0.005$ at 8640 MHz, we can estimate a magnetic field strength of ~ 50 mG (corresponding to $\gamma \sim 230$).

This estimate is in order-of-magnitude agreement with magnetic field estimates for major ejections from SS 433 and other X-ray binaries (e.g., Hjellming & Han 1995; Fender et al. 1999). Note, in addition, that “Faraday depolarization,” which can severely reduce the observed linear polarization, will have no effect on circular polarization. It is therefore possible to have an optically thin synchrotron source that displays a large ratio of circular to linear polarization, as is observed in this case. However, reversals in the line-of-sight component of the magnetic field that are likely to occur within the source will require significantly higher magnetic fields. Similarly, a sig-




Fig. 2.—Observations of SS 433 in 1999 May. The top panel illustrates GBI (quasi-)daily monitoring at 2.3 and 8.3 GHz; our best estimates of the ejection dates, corresponding to the two major flares, are indicated by arrows. The lower panels indicate mean flux densities in total intensity, linear polarization, and circular polarization for the two epochs of ATCA observations.

nificant e^+ population within the source will also cause a reduction in the observed V and require a significantly higher magnetic field. In addition, the approximately constant circularly polarized flux density at 8640 MHz during a drop by $\geq 40\%$ in the total flux density does not seem compatible with an origin for the circular polarization in the optically thin ejecta that correspond to the two major flares.

4.2. Propagation-induced Circular Polarization

Linearly polarized radiation can be converted to circularly polarized radiation during propagation through a plasma with elliptical (or linear) propagation modes (Pacholczyk 1973; Kennett & Melrose 1998 and references therein). In the event of the admixture of a small amount of relativistic plasma to a thermal plasma, propagation modes through the plasma will become slightly elliptical, and a spectrum of the form $m_c \propto \nu^{-1}$ is predicted (Pacholczyk 1973). In the event of plasma that is dominated by highly relativistic particles, the propagation modes may approach linear, and a much steeper spectrum of the form $m_c \propto \nu^{-3}$ is predicted (Kennett & Melrose 1998).

Wardle et al. (1998) argued that the circular polarization observed from 3C 279 arose because of such propagation-induced “repolarization.” They concluded that the low-energy spectrum of the relativistic particles must extend to $\gamma \ll 100$, and therefore the jet must be composed of an e^+e^- plasma (if there were protons accompanying each emitting electron, the kinetic energy of the jet would be several orders of magnitude greater than that which is seen to be dissipated at the head of the jet). We note that analogous considerations may also be applicable to SS 433, where X-ray and radio hot spots are observed within the W50 radio nebula, presumably at the site

of the jet–interstellar medium interaction. Indeed, the kinetic energy in the jets of SS 433, if they are composed of a e^-p^+ plasma, is $\geq 10^{40}$ ergs s^{-1} (Brinkmann et al. 1988), which is much greater than that which is directly observed to be dissipated on larger scales within W50 (this is one of the arguments against an e^-p^+ plasma that is put forward by Kundt 1998).

4.3. Alternative Mechanisms?

Given the complexity of the SS 433 system and the high densities and magnetic field strengths likely to be present near the base of the jet, alternative origins for the circularly polarized emission cannot be ruled out. These include gyrosynchrotron emission from low-energy electrons and cyclotron maser emission (Dulk 1985). We note that if the circularly polarized emission is associated with a region on the scale of one of the binary components (e.g., $< 10^{12}$ cm), then the brightness temperature is $\geq 10^{10}$ K at 8.64 GHz and $\geq 10^{12}$ K at 1.4 GHz.

5. DISCUSSION

It is of great importance to determine in which of the various emitting regions the circularly polarized flux density originated, in order to determine both the spectrum and the relative strength of the emission. Our ATCA observations rule out an extension with the optically thin jets on scales of $\geq 1''$, and the lack of correlated variability in Stokes I and V argues against an association with the two major ejection events. The underlying radio spectrum of SS 433 is typically around 250 mJy at 8 GHz, with an optically thin spectral index of about -0.7 , corresponding to a quasi-continuous flow of matter into the




FIG. 3.—Progression of internal and external absorption and depolarization effects as ejecta move away from SS 433 (Hjellming & Johnston 1981; Vermeulen 1993; Paragi et al. 1999). The gray circles at the top and bottom of the schematic indicate our best estimates of the physical locations of the components responsible for the radio flares that dominate the total flux density, on 1999 May 10 and 20, respectively.

jets. If associated with this component, then the spectrum of the relative circular polarization could be as flat as the $m_c \propto \nu^{-1/2}$ predicted for intrinsic synchrotron emission. However, if associated with this component, then why not with the two flares, which are presumably just enhancements of the same flow? Alternatively, Paragi et al. (1999) show that the innermost regions (≤ 50 mas) of the jets have a flat/inverted spectrum between 1 and 15 GHz. The core region has a peak flux density typically of ≤ 100 mJy on VLBI scales. If associated with this region, the fractional circular polarization may be as high as 10%, and the spectrum may steepen to the $m_c \propto \nu^{-1}$ predicted for a mildly relativistic plasma (it seems unlikely that the $m_c \propto \nu^{-3}$ spectrum can be recovered, unless the emission arises right in the binary core of the system, which has the most inverted radio spectrum). If associated with the inner regions, this implies a very large ratio of circular to linear polarizations, as found for Sgr A* by Bower et al. (1999).

Further precessional-phase-resolved monitoring and the spatial resolution of the regions responsible for the circular polarization are essential to investigate this discovery further. If, as seems likely, the circular polarization is associated with the synchrotron-emitting ejecta, comparison with circular polarization measurements of other X-ray binaries has the potential to reveal, finally, the composition of the relativistic plasmas.

R. F. would like to acknowledge useful discussions with Ralph Spencer and Al Stirling and to thank Mark Walker for the original suggestion to look for circular polarization in X-ray binaries. The Australia Telescope is funded by the Commonwealth of Australia for operation as a National Facility managed by CSIRO. R. F. was funded during the period of this research by a EC Marie Curie Fellowship (ERBFMBICT 972436).

REFERENCES

- Bower, G. C., Falcke, H., & Backer, D. C. 1999, *ApJ*, 523, L29
 Brinkmann, W. 1998, *NewA Rev.*, 42, 589
 Brinkmann, W., Fink, H. H., Massaglia, S., Bodo, G., & Ferrari, A. 1988, *A&A*, 196, 313
 Conway, R. G., & Kronberg, P. P. 1969, *MNRAS*, 142, 11
 Dulk, G. A. 1985, *ARA&A*, 23, 169
 Fender, R. P. 2000, in *Astrophysics and Cosmology: A Collection of Critical Thoughts*, ed. W. Kundt & C. van de Bruck (Lecture Notes in Physics; New York: Springer), in press (astro-ph/9907050)
 Fender, R. P., Garrington, S. T., McKay, D. J., Muxlow, T. W. B., Pooley, G. G., Spencer, R. E., Stirling, A. M., & Waltman, E. B. 1999, *MNRAS*, 304, 865
 Frater, R. H., Brooks, J. W., & Whiteoak, J. B. 1992, *J. EEEA*, 12, 103
 Hjellming, R. M., & Han, X. H. 1995, in *X-Ray Binaries*, ed. W. H. G. Lewin, J. van Paradijs, & E. P. J. van den Heuvel (Cambridge: Cambridge Univ. Press), 308
 Hjellming, R. M., & Johnston, K. J. 1981, *ApJ*, 246, L141
 Kennett, M., & Melrose, D. 1998, *Publ. Astron. Soc. Australia*, 15, 211
 Kesteven, M. J. 1997, ATCA Tech. Doc. 39.3/071, Field Rotation Due to an Antenna's Pointing Model
 Kotani, T., Kawai, N., Matsuoka, M., & Brinkmann, W. 1996, *PASJ*, 48, 619
 Kundt, W. 1998, in *The Many Faces of Neutron Stars*, ed. R. Buccheri, J. van Paradijs, & M. A. Alpar (NATO ASI Ser. C, 515; Dordrecht: Kluwer), 461
 Legg, M. P. C., & Westfold, K. C. 1968, *ApJ*, 154, 499
 Margon, B. 1984, *ARA&A*, 22, 507
 Ostrowski, M., Sikora, M., Madejski, G., & Begelman, M., eds. 1997, *Relativistic Jets in AGNs* (Kraków: Jagiellonian Univ., Obs. Astron.)
 Pacholczyk, A. G. 1973, *MNRAS*, 163, 29
 Paragi, Z., Vermeulen, R. C., Fejes, I., Schilizzi, R. T., Spencer, R. E., & Stirling, A. M. 1999, *A&A*, 348, 910
 Rayner, D. P., Norris, R. P., & Sault, R. J. 2000, *MNRAS*, submitted
 Reynolds, J. E. 1994, ATCA Tech. Doc. 39.3/040, A Revised Flux Scale for the AT Compact Array
 Sault, R. J., Killeen, N. E. B., & Kesteven, M. J. 1991, ATCA Tech. Doc. 39.3/015, AT Polarization Calibration
 Sault, R. J., Hamaker, J. P., & Bregman, J. D. 1996, *A&AS*, 117, 149
 Sault, R. J., & Macquart, J.-P. 1999, *ApJ*, 526, L85
 Sault, R. J., Teuben, P. J., & Wright, M. C. H. 1995, in *ASP Conf. Ser. 77, Astronomical Data Analysis Software and Systems IV*, ed. R. A. Shaw, H. E. Payne, & J. J. E. Hayes (San Francisco: ASP), 433
 Shields, G. A. 1999, *PASP*, 111, 661
 Vermeulen, R. C. 1989, Ph.D. thesis, Univ. Leiden
 ———. 1993, in *Astrophysical Jets*, ed. D. Burgarella, M. Livio, & C. P. O'Dea (Cambridge: Cambridge Univ. Press), 241
 Waltman, E. B., Fiedler, R. L., Johnston, K. J., Spencer, J. H., Florkowski, D. R., Josties, F. J., McCarthy, D. D., & Matsakis, D. N. 1991, *ApJS*, 77, 379
 Wardle, J. F. C., Homan, D. C., Ojha, R., & Roberts, D. H. 1998, *Nature*, 395, 457
 Wilson, A. S., & Weiler, K. W. 1997, *ApJ*, 475, 661