

Discrimination of Semi-Quantitative Models by Experiment Selection: Method and Application in Population Biology*

Ivayla Vatcheva¹, Olivier Bernard², Hidde de Jong³, Jean-Luc Gouzé², and Nicolaas J.I. Mars¹

¹ Department of Computer Science, University of Twente, P.O. Box 217
7500 AE Enschede, the Netherlands

² INRIA Sophia Antipolis, 2004 Route des Lucioles, BP 93
06902 Sophia Antipolis, France

³ INRIA Rhône-Alpes, 655, avenue de l'Europe
38330 Montbonnot Saint Martin, France

Email: {ivayla, mars}@cs.utwente.nl, {Olivier.Bernard, Jean-Luc.Gouze}@sophia.inria.fr, Hidde.de-Jong@inrialpes.fr

Abstract

Modeling an experimental system often results in a number of alternative models that are justified equally well by the experimental data. In order to discriminate between these models, additional experiments are needed. We present a method for the discrimination of models in the form of semi-quantitative differential equations. The method is a generalization of previous work in model discrimination. It is based on an entropy criterion for the selection of the most informative experiment which can handle cases where the models predict multiple qualitative behaviors. The applicability of the method is demonstrated on a real-life example, the discrimination of a set of competing models of the growth of phytoplankton in a bioreactor.

Introduction

Obtaining an adequate model of an experimental system is a laborious and error-prone task. In many cases one arrives at a number of rival models that are justified equally well by the experimental data. In order to discriminate between these models, additional experiments are needed. Since in real-life applications the number of experiments to perform may be quite large, and the costs of each of them considerable, it is important that the experiments be selected carefully. In fact, experiments need to be chosen such that the set of possible models is maximally reduced at minimal costs.

For experimental systems described by differential equations, several approaches for *model discrimination* have been proposed in the literature (e.g. (Espie and Macchietto 1989)). With few exceptions (e.g. (Struss 1994; Vatcheva *et al.* 2000)), these methods apply to completely specified quantitative models. That is, they cannot be used when precise numerical values for the parameters are not available and the precise form of functional relations is unknown.

This has motivated the development of a method for model discrimination that is able to handle incompletely specified models in the form of *semi-quantitative differential equations* (SQDEs). The method is based on an entropy

criterion for the selection of the most informative discriminatory experiment. This experiment is determined from the behavioral predictions obtained from the competing models through simulation under various experimental conditions.

In earlier work, we have developed a method for the discrimination of semi-quantitative models (Vatcheva *et al.* 2000). However, the previously proposed approach is restricted to the case that all models predict the same qualitative behavior, a situation rarely occurring in the case of more complex models. The method described in this paper is a generalization of the approach above in that it allows one to deal with situations in which multiple qualitative behaviors are predicted.

The applicability of the method is demonstrated on a real problem in population biology, the selection of experiments to discriminate between competing models of the growth of phytoplankton in a bioreactor. The choice of good discriminatory experiments is critical in this application, since the experiments may take several weeks to complete. Semi-quantitative models are appropriate, because the available data is incomplete and imprecise, as for most biological systems. We have compared the optimal experiment as determined by our method with an experiment that has been actually carried out. Furthermore, taking into account the results of the latter experiment, the best next experiment to perform has been suggested.

The paper is organized as follows. The next section deals with the basic concepts of semi-quantitative modeling and simulation. Then, an outline of the method for model discrimination is given, focusing on the criterion for selecting the most informative experiment. Next, the results of the application of the method to the modeling of phytoplankton growth in a bioreactor are presented. The final section discusses achievements and limitations of our work in the context of related work on model discrimination and gives some perspectives on further research.

Semi-quantitative modeling and simulation

Semi-quantitative differential equations (SQDEs) are abstractions of ordinary differential equations (ODEs) that allow incompletely or imprecisely specified dynamical sys-

* A shorter version of this paper appears in the proceedings of the 17th International Joint Conference on Artificial Intelligence, IJCAI-01, Seattle, 4-10 August, 2001.

Monod

$$\begin{aligned}\dot{x} &= \mu(s)x - dx \\ \dot{s} &= d(s_{in} - s) - \rho(s)x \\ \mu(s) &= \mu_{max} \frac{s}{s + k_s} \\ \rho(s) &= \frac{1}{y} \mu(s)\end{aligned}$$

$$\begin{aligned}d &\in [0.095, 0.105], \quad s_{in} \in [80, 120] \\ y &\in [0.15, 0.6], \quad \mu_{max} \in [1.2, 1.6], \quad k_s \in [0.01, 0.2]\end{aligned}$$

Figure 1: An example of an SQDE describing the growth of phytoplankton in a bioreactor, the *Monod* model. The physical meaning of the variables and parameters is given in the caption of Figure 3.

tems to be modeled (Berleant and Kuipers 1997). In an SQDE, bounding envelopes are defined for unknown functions, as well as numerical intervals to bound the values of parameters and initial conditions.

Figure 1 shows an example of a second-order SQDE describing the growth of the microalgae *Dunaliella tertiolecta* under nutrient limitation in a bioreactor (Monod 1942). The state variables are the biomass x and the concentration of the limiting nutrient s . The intervals bounding the model parameters μ_{max} , k_s , and y have been estimated from preliminary experiments.


For the simulation of SQDEs we employ the techniques Q2+Q3 (Berleant and Kuipers 1997), which refine the qualitative behavior tree produced by the QSIM algorithm (Kuipers 1994). The results of semi-quantitative simulation consist of one or more qualitative behaviors supplemented by ranges for the values of the variables at qualitatively significant time-points. The behaviors resulting from the simulation of the SQDE in Figure 1 are shown in Figure 2. In order to narrow down the interval predictions, we use the comparative analysis technique SQCA (Vatcheva and de Jong 1999). SQCA refines the simulation results by comparing a behavior predicted for one experiment with behaviors predicted for related experiments.

Method for model discrimination

The predictions obtained through semi-quantitative simulation can be exploited to maximally discriminate a set of competing models against minimal costs. The method for achieving this is based on a generalization of the entropy criterion for the most informative experiment developed in (Vatcheva *et al.* 2000).

Model discrimination and behavior predictions

Consider a set M of competing models of an experimental system. Each $m_i \in M$ has a probability $p(m_i)$ to be the correct model of the system. We assume that the set M is complete, that is $\sum_{m_i \in M} p(m_i) = 1$ (Fedorov 1972). The a



| | $s^* (\times 10^{-4})$ | x^* | s_{max} |
|-------|----------------------------|-----------|-----------------|
| b_1 | [6.31, 100] | [36, 72] | - |
| b_2 | [14.9, 100] | [36, 72] | [0.005, 0.0335] |
| b_3 | [6.31, 100] | [36, 59] | - |
| | $s_{min} (\times 10^{-5})$ | x_{max} | |
| b_1 | - | - | |
| b_2 | - | - | |
| b_3 | [5.85, 1000] | [36, 121] | |

Figure 2: Behaviors resulting from the simulation of the SQDE in Figure 1, for the initial conditions $x_0 \in [36, 39]$, $s_0 \in [0.005, 0.01]$. Behavior b_1 predicts that the system reaches its equilibrium (x^*, s^*) asymptotically. In b_2 , the nutrient concentration s reaches a maximum, before the system approaches its equilibrium. In b_3 , x reaches first a maximum, followed by a minimum of s and the equilibrium. The table summarizes the interval predictions for each of the three behaviors.

priori model probabilities are estimated from data obtained in preliminary experiments or assumed equal when no such data exist. Each time an experiment is executed, and new data becomes available, the model probabilities are being updated. If the data does not justify the predictions of some m_i , its *posteriori* probability becomes zero.

For the discrimination of the models in M , experiments from a set E of experiments need to be selected. The experiment that discriminates best between the models is estimated from the model predictions. For each experiment $e \in E$, the models in M are perturbed according to e , and then simulated to predict the behavior of the system under the experimental conditions. The prediction of m_i for some e is a set of behaviors B_i^e . The set of all qualitatively distinct behaviors resulting from the simulation of the models in M for e is denoted by B^e .

For discrimination purposes, only certain characteristics of the predicted behaviors $b \in B^e$ are taken into account.

This gives rise to a set of *behavioral features* F_b for b . The set of behavioral feature consists of minima, maxima and equilibria of the system variables. The behavioral features defined for b_3 in Figure 2, for instance, are the maximum of x and the minimum of s (x_{max} and s_{min}), and the steady state levels of these variables (s^* and x^*). Here we will assume that the value of a behavioral feature is an interval.

Intuitively, the experiment that can be expected to optimally discriminate between the models is the experiment for which the predicted values of the behavioral features overlap least. This intuition will be formalized below by defining the *most informative* experiment. A set of competing models can then be discriminated by repeatedly determining the most informative experiment, performing this experiment, and updating the model probabilities in the light of the outcomes.

Criterion for most informative experiment

A standard measure in information theory is the *information increment* of an experiment (Fedorov 1972). For every experiment $e \in E$ we define

$$\Delta H(b^e, \mathbf{y}^e) = - \sum_{m_i \in M} p(m_i) \ln p(m_i) + \sum_{m_i \in M} p(m_i | b^e, \mathbf{y}^e) \ln p(m_i | b^e, \mathbf{y}^e), \quad (1)$$

where $p(m_i)$ and $p(m_i | b^e, \mathbf{y}^e)$ are the *a priori* and *a posteriori* probabilities of m_i . b^e is the behavior of the system observed in e , and \mathbf{y}^e is the vector of midpoints of the observations for the behavioral features F_{b^e} . The observations are assumed to be intervals $Y_j^e = [y_j^e - \varepsilon_j/2, y_j^e + \varepsilon_j/2]$, with y_j^e the midpoint, and ε_j the estimated size of the confidence interval for the j th behavioral feature. For clarity of presentation, we will assume for the moment, that each behavior b is characterized by a single feature.

ΔH reaches its maximum when all posterior probabilities but one are zero. That is, when the observations obtained in e confirm the predictions of a single model. On the other hand, a minimal value is attained, when all posterior probabilities are equal.

Since the *a posteriori* probabilities of the models depend on the outcome of the experiment which is not yet known, ΔH cannot be computed directly. Instead, we can compute its expected value, or the *expected information increment* of e .

Assume, for the moment, that the system behavior in experiment e is known. The expected value of $\Delta H(b^e, \mathbf{y}^e)$ is defined as an integral of the function $\Delta H(b^e, y)$ weighted according to the possible values of y :

$$\Delta J(e, b^e) = \int_{y \in D} \Delta H(b^e, y) g^e(y|b^e) dy, \quad (2)$$

where D is the domain of the behavioral feature, and $g^e(y|b^e)$ the probability density function of its midpoint, provided b^e is the behavior of the system in e . In general, b^e would not be known. However, we know the set B^e of

the possible system behaviors. Recall, that B^e is the set of qualitatively distinct behaviors predicted by the competing models $m_i \in M$. From the assumption that M is complete, and the soundness of the simulation algorithms, it follows that B^e contains all possible behaviors the system can exhibit. Hence, the expected value of $\Delta J(e, b^e)$ can be computed as a weighted sum over the behaviors $b \in B^e$:

$$\Delta J(e) = \sum_{b \in B^e} \Delta J(e, b) p^e(b), \quad (3)$$

where $p^e(b)$ is the probability the system behavior observed in e to be b .

Combining (2) and (3), we obtain:

$$\Delta J(e) = \sum_{b \in B^e} \int_{y \in D} \Delta H(b, y) g^e(y|b) p^e(b) dy. \quad (4)$$

The probability of behavior b can be expressed as an weighted sum of the conditional behavior probabilities $p^e(b|m_i)$:

$$p^e(b) = \sum_{m_i \in M} p(m_i) p^e(b|m_i), \quad (5)$$

where $p^e(b|m_i)$ is the probability of behavior b in e , provided m_i is the correct model of the system. Similarly, we have

$$g^e(y|b) = \sum_{m_i \in M} p(m_i|b) g_i^e(y|b), \quad (6)$$

where $p(m_i|b)$ is the *a posteriori* probability of m_i if the observed behavior is b , and $g_i^e(y|b)$ is the model-specific probability density function of y , defined by

$$g_i^e(y|b) = \begin{cases} \frac{|[y - \frac{\varepsilon}{2}, y + \frac{\varepsilon}{2}] \cap V_i^e|}{\varepsilon |V_i^e|} & , [y - \frac{\varepsilon}{2}, y + \frac{\varepsilon}{2}] \cap V_i^e \neq \emptyset, \\ 0 & , [y - \frac{\varepsilon}{2}, y + \frac{\varepsilon}{2}] \cap V_i^e = \emptyset, \end{cases}$$

with V_i^e the interval prediction of m_i for the behavioral feature of b in experiment e , and $|\cdot|$ denoting interval length. The definition of $g_i^e(y|b)$ can be explained as follows. Assume Y is an interval, and $p(y^e \in Y)$ denotes the probability of y^e to be in Y . That is, the probability that the midpoint of the value of the behavioral feature belongs to Y . By definition, $p(y^e \in Y) = \int_{y \in Y} g_i^e(y|b) dy$. The definition of g_i^e expresses the idea that for intervals Y not overlapping with the model prediction V_i^e , the probability $p(y^e \in Y)$ is zero. Otherwise, $p(y^e \in Y)$ is weighted according to the size of the overlap between Y and V_i^e .

By substituting (5) and (6) in (4) and using the Bayes' rule:

$$p(m_i|b) = \frac{p(m_i) p^e(b|m_i)}{\sum_{m_k \in M} p(m_k) p^e(b|m_k)},$$

we obtain:

$$\Delta J(e) = \sum_{m_i \in M} p(m_i) \sum_{b \in B^e} p^e(b|m_i) \int_{y \in D} \Delta H(b, y) g_i^e(y|b) dy. \quad (7)$$

Finally, the substitution of the expression for ΔH in (7) and the use of the Bayes' rule

$$p(m_i|b, y) = \frac{p(m_i) p^e(b|m_i) g_i^e(y|b)}{\sum_{m_k \in M} p(m_k) p^e(b|m_k) g_k^e(y|b)},$$

leads to the following expression for the expected information increment of an experiment e :

$$\Delta J(e) = \sum_{m_i \in M} p(m_i) \sum_{b \in B_i^e} p^e(b|m_i) \int_{y \in D} g_i^e(y|b) \ln \frac{g_i^e(y|b) p(b|m_i)}{g^e(y|b)} dy. \quad (8)$$

The criterion ranks the experiments in E according to their expected informativeness. The optimal discriminatory experiment will be the *most informative experiment*, that is, the experiments for which $\Delta J(e)$ is maximal. Intuitively, experiments which give rise to predictions as different as possible will be favored. Figure 4(a)-(b) shows the predictions of the four models given in Figures 1 and 3 for two different experiments (see next section). In both cases, each of the models predicts two possible qualitative behaviors for the biomass x . The expected information increment for the first experiment, however, is higher than the expected information increment for the second ($\Delta J = 0.5927$ versus $\Delta J = 0.3290$), as the predicted intervals overlap less.

The expression for ΔJ can be simplified in a number of cases. For instance, if all models predict for a given experiment e the same qualitative behavior, (8) can be reduced to

$$\Delta J(e) = \sum_{m_i \in M} p(m_i) \int_{y \in D} g_i^e(y|b) \ln \frac{g_i^e(y|b)}{g^e(y|b)} dy,$$

which is the criterion previously derived by (Vatcheva *et al.* 2000).

On the other hand, if for a given e , each model predicts a different set of qualitative behaviors, we obtain:

$$\Delta J(e) = - \sum_{m_i \in M} p(m_i) \ln p(m_i),$$

which is the maximum value $\Delta J(e)$ can take.

The criterion (8) is easily generalizable to the case when each behavior b is characterized by more than one feature. In this case we have to substitute the probability density functions by joint probability density functions, and the integral by a multiple integral of the k behavioral features. For computational simplicity, we assume in this article that the behavioral features are independent. That is, joint probability density functions will be estimated by the product of the density functions of the involved behavioral features.

The algorithms for the simulation of SQDEs, outlined in the previous section have been proven sound. That is, all possible predictions are derived from a given SQDE model. If the results obtained in the experiment are correct, this guarantees that a model will never be falsely rejected. However, these algorithms do not exclude spurious predictions. As a consequence, an experimental result may corroborate a model while it should be ruled out. Spurious predictions, therefore, may prolong the discrimination process.

Computation of behavior probabilities

In order to compute $\Delta J(e)$, the conditional behavior probabilities $p(b_j|m_i)$ must be estimated. We have adopted the following approach. Let v_l be a parameter or initial condition in model m_i , and let $range(v_l, b_j)$ be the interval value of v_l for which behavior b_j is obtained. Define

$$r(b_j|m_i) = \frac{\prod_l |range(v_l, b_j)|}{\prod_l |\bigcup_{b_k \in B_i^e} range(v_l, b_k)|}.$$

$r(b_j|m_i)$ estimates the fraction of the interval volume of the model parameters which gives rise to b_j . The conditional probability of b_j is now given by normalizing the $r(b_j|m_i)$: $p(b_j|m_i) = r(b_j|m_i) / \sum_{b_k \in B_i^e} r(b_k|m_i)$.

Consider, for instance, the three behaviors given in Figure 2. b_1 , b_2 and b_3 have been obtained for different subintervals of the interval ranges for k_s and y :

| beh | k_s | y |
|-------|-----------------|---------------|
| b_1 | [0.01, 0.158] | [0.3, 0.6] |
| b_2 | [0.0236, 0.158] | [0.3, 0.6] |
| b_3 | [0.01, 0.158] | [0.15, 0.488] |

By using these values, the procedure outlined above gives $r(b_1) = 0.67$, $r(b_2) = 0.61$, $r(b_3) = 0.75$. Consequently, the behavior probabilities can be computed: $p(b_1) = 0.33$, $p(b_2) = 0.30$, $p(b_3) = 0.37$.

Alternative approaches for estimating the probabilities of qualitative behaviors have been suggested by (Leitch and Shen 1993; Berleant *et al.* 1992).

Application: phytoplankton growth

Phytoplankton play a crucial role in marine ecosystems. The lives of almost all animals that live in the sea depend on phytoplankton for food and life-sustaining oxygen, essential to the metabolism of all aerobic organisms. Furthermore, phytoplankton exert a global scale influence on our climate. The global carbon cycle, which regulates the temperature of our planet, is controlled by the actions of the phytoplankton. Large populations of these organisms can significantly lower atmospheric carbon dioxide levels and, in turn, lower average temperatures.

Understanding the regulation of phytoplankton growth is essential for predicting how life in the ocean may respond to climate changes and for learning more about how these organisms are contributing to, and affected by environmental changes. As these processes are difficult to study in the open

| Contois | Droop | Caperon-Meyer |
|--|---|--|
| $\dot{x} = \mu(s)x - dx$ | $\dot{x} = \mu(q)x - dx$ | $\dot{x} = \mu(q)x - dx$ |
| $\dot{s} = d(s_{in} - s) - \rho(s)x$ | $\dot{q} = \rho(s) - \mu(q)q$ | $\dot{q} = \rho(s) - \mu(q)q$ |
| $\mu(s) = \mu_{max} \frac{s}{s + k_x x}$ | $\dot{s} = d(s_{in} - s) - \rho(s)x$ | $\dot{s} = d(s_{in} - s) - \rho(s)x$ |
| $\rho(s) = \frac{1}{y} \mu(s)$ | $\mu(q) = \bar{\mu} \left(1 - \frac{k_q}{q}\right)$ | $\mu(q) = \mu_{max} \frac{q - k_q}{q - k_q + k_0}$ |
| | $\rho(s) = \rho_{max} \frac{s}{s + k_s}$ | $\rho(s) = \rho_{max} \frac{s}{s + k_s}$ |

$$y \in [0.15, 0.6], k_x \in [0.00014, 0.0167], \bar{\mu} \in [1.7, 2.3], k_q \in [1.6, 2.0], \mu_{max} \in [1.2, 1.6], \rho_{max} \in [9.25, 9.55], k_0 \in [2.0, 2.4]$$

Figure 3: Models for the growth of phytoplankton in a bioreactor. x is the amount of biomass per unit volume, s [$\mu\text{mol}/l$] the nutrient concentration, q [$\mu\text{mol}/l$] the internal quota. The Monod and Contois models assume constant growth yield y [$l/\mu\text{mol}$]. μ_{max} [day^{-1}] is the maximum growth rate of cells, and $\bar{\mu}$ [day^{-1}] a theoretical maximum growth rate obtained for infinite quota. k_s , k_x , and k_0 [$\mu\text{mol}/l$] are half-saturation constants, k_q [$\mu\text{mol}/l$] is the minimum cell quota, ρ_{max} [$\mu\text{mol}/l/\text{day}$] is the maximum uptake rate of nutrients. For all models $s_{in} \in [80, 120]$ [$\mu\text{mol}/l$] is the input nutrient concentration, and d [day^{-1}] the dilution rate to be controlled in the experiments. The initial conditions are $x_0 \in [36.0, 39.0]$, $s_0 \in [0.005, 0.01]$, $q_0 \in [2.4, 2.7]$ which are the equilibrium values reached for the initial dilution rate $d_0 = 0.4$.

sea, the growth conditions are recreated in the laboratory in a type of bioreactor known as chemostat. The technique basically consists of growing a population of microorganisms in a culture vessel. The nutrients and substrates required for cell growth are supplied continuously at a fixed rate to the culture vessel whose contents are continuously mixed. A medium, cells, and by-products are continually removed from the vessel, maintaining the culture in the chemostat chamber at a constant volume. In the context of phytoplankton, the chemostat is mainly used to study the growth of populations under nutrient limitation.

A variety of models can be used to describe the growth of phytoplankton in a chemostat. Which of these applies best in a given situation cannot be determined on *a priori* grounds. Therefore, experiments need to be performed to discriminate between the alternative models. Unfortunately, these experiments may take weeks to complete and are thus quite costly to perform.

We have applied the method of the previous section in the context of the microalgae *Dunaliella tertiolecta*, carried out by population biologists in a marine laboratory. Four alternative models to describe the system have been considered, which are shown in Figures 1 and 3. All models share the same basic idea: (a) at low nutrient concentration the rate of the nutrient consumption (uptake rate) ρ and the growth rate μ are limited by, and proportional to, the nutrient concentration, whereas (b) at high concentration the uptake and growth rates saturate and become constant, independent of nutrient concentration. However, the models make different assumptions about the nutrient consumption, the influence of the biomass on the growth rate of the population, and the relation between growth and uptake rates. The models are labeled after their originators: M (Monod 1942), C (Con-

tois 1959), D (Droop 1968), and CM (Caperon and Meyer 1972).

Because of coarse and noisy data, and evolution of the system in the time frame of the experiment, precise numerical estimations for the values of the parameters cannot be obtained. This motivates the use of semi-quantitative models. The interval values for the parameters required in the SQDEs have been estimated by the biologists, based on the outcome of preliminary experiments (see Figure 3).

In order to discriminate between the competing models, the value of the dilution rate d can be varied by the experimenter. Starting from an equilibrium, the dilution rate is changed and the transient behavior of the system towards a new equilibrium is observed. We have considered ten experiments, corresponding to equispaced values in the range $[0, 1.2]$: $d = 0.1, d = 0.2, \dots$. Taking into account 5% measurement uncertainty, the values of d become intervals.

The only variable that can be reliably measured in the course of the experiment is the biomass x . This determines the behavioral features that we have considered: the minimum and the maximum value of x (x_{min} and x_{max}), and the equilibrium value of x (x^*). In order to obtain the predicted values of the behavioral features required for the determination of the most informative experiment, the models have been simulated using the techniques outlined in the second section. For each experiment, all models predict multiple qualitative behaviors as a consequence of the large intervals for the parameter values. In total, four different behaviors for x are predicted. None of these behaviors is spurious, as we have been able to establish by comparing the predictions with the qualitative analysis of (Bernard and Gouzé 1995).

Starting from the assumption that the models are equiprobable in the beginning, we have calculated the ex-




Figure 4: (a) Interval predictions of the four competing models (Figures 1 and 3) for the behavioral features x^* (behavior b_1), and x^*, x_{min} (behavior b_2) in the predicted optimal discriminatory experiment ($d = 1.1$). (b) Interval predictions for x^* (behavior b_1), and x^*, x_{min} (behavior b_2) in the experiment that has been carried out ($d = 0.7$). The values for x^*, x_{min} measured in the experiment are also shown by a small rectangle. (c) Predictions for the features x^* (behavior b_1), and x^*, x_{max} (behavior b_2) in the next optimal experiment ($d = 0.1$). M, C, D, and CM stand for the Monod, Contois, Droop, and Caperon-Meyer models, respectively.

| | $d = 0.1$ | $d = 0.2$ | $d = 0.3$ | $d = 0.5$ | $d = 0.6$ |
|------------|-----------|-----------|-----------|-----------|-----------|
| ΔJ | 0.3204 | 0.3181 | 0.3519 | 0.3547 | 0.3323 |
| | $d = 0.7$ | $d = 0.8$ | $d = 0.9$ | $d = 1.0$ | $d = 1.1$ |
| ΔJ | 0.3290 | 0.3710 | 0.4668 | 0.5093 | 0.5927 |

Table 1: Values for the expected information increment ΔJ for each of the dilution rate experiments.

pected information increment (8) for each of the experiments (Table 1). The optimal discriminatory experiment is predicted to be $d = 1.1$. For this experiment, each of the four models predicts two behaviors, b_1 and b_2 , that differ with respect to the observable variable x . In b_1 the equilibrium of the system is reached asymptotically, whereas in b_2 , x reaches a minimum before the equilibrium is attained. Figure 4(a) shows the interval predictions of the behavioral features for all four models, and Table 2 lists the corresponding conditional behavior probabilities. Notice that in b_1 only one behavioral feature applies (x^*), whereas in b_2 predictions for x^* and x_{min} need to be taken into account.

The experiment $d = 1.1$ has not been performed, but data for the suboptimal experiment $d = 0.7$ was available from an earlier study. The predictions of the behavioral features for this experiment are shown in Figure 4(b) and the behavior probabilities are given in Table 2.

In the experiment $d = 0.7$, x was found to reach its equilibrium after passing through a minimum. This rules out b_1 . The measured values of the behavioral features, shown in Figure 4(b), are $x_{min} = [29.2, 32.2]$, and $x^* = [30.5, 33.5]$. Using these results, the *a posteriori* probabilities of the models have been computed via Bayes' rule: $p(M) = 0.34$,

| | $d = 1.1$ | | $d = 0.7$ | | $d = 0.1$ | |
|----|-----------|----------|-----------|----------|-----------|----------|
| | $p(b_1)$ | $p(b_2)$ | $p(b_1)$ | $p(b_2)$ | $p(b_1)$ | $p(b_2)$ |
| M | 0.47 | 0.53 | 0.47 | 0.53 | 0.50 | 0.50 |
| C | 0.47 | 0.53 | 0.47 | 0.53 | 0.51 | 0.49 |
| D | 0.57 | 0.43 | 0.57 | 0.43 | 0.54 | 0.46 |
| CM | 0.57 | 0.43 | 0.57 | 0.43 | 0.27 | 0.73 |

Table 2: Conditional probabilities of the behaviors b_1 and b_2 predicted by the four models for the experiments $d = 1.1$, $d = 0.7$, and $d = 0.1$ (see Figure 4). $p(b_1)$ and $p(b_2)$ have been estimated using the approach in the previous section.

$p(C) = 0.33$, $p(D) = 0.21$, and $p(CM) = 0.12$. In addition, the measurements have allowed the parameter values to be refined by means of the constraint propagation algorithm in Q2 (Berleant and Kuipers 1997).

The new model probabilities show that experiment $d = 0.7$ has not been very discriminating. Given the new model probabilities and parameter values, what would be the optimal experiment to perform next? The method advises that $d = 0.1$ be tried, as it has the highest expected information increment ($\Delta J = 0.7129$). For this experiment, each of the models predicts two possible behaviors for the biomass x : in b_1 , x increases asymptotically to its new equilibrium (giving rise to the behavioral feature x^*), and in b_2 , x first reaches a maximum before the equilibrium is attained (giving rise to the features x_{max} and x^*). The predicted values for the behavioral features of the two behaviors are shown in Figure 4(c). Table 2 gives the corresponding behavior probabilities. The experiment $d = 0.1$ has not been performed yet (we recall that each experiment takes weeks to complete). Notice, however, that $d = 0.1$ is likely to rule out at least

two of the four models due to the lack of overlap between $M - C$ and $D - CM$.

Discussion

We have proposed a method for the discrimination of semi-quantitative models of an experimental system. The method is based on an entropy criterion for the selection of the most informative experiment. The value of ΔJ for a particular experiment is calculated from the model predictions obtained through semi-quantitative simulation. The method generalizes upon a previous method (Vatcheva *et al.* 2000), in that it can handle cases where the models predict multiple qualitative behaviors. This occurs in the case of the phytoplankton growth models, which predict the biomass to asymptotically approach its equilibrium value or to pass through a maximum or a minimum first.

The applicability of the method has been demonstrated by having it predict the most informative experiment to discriminate between four models of the growth of *Dunaliella tertiolecta* in a bioreactor. This has been achieved in the presence of several complicating factors, in particular the nonlinearity of the models, the crude estimations of the parameter values, and the difficulty to observe the behavior of the system. The discrimination of bioreactor models has been attempted before (Espie and Macchietto 1989; Cooney and McDonald 1995), but unlike the method discussed in this paper, these approaches require precise numerical data to be available, a requirement that usually cannot be fulfilled in practice.

Within AI, methods for model discrimination have been developed in the field of model-based diagnosis (e.g., (Struss 1994; de Kleer 1990)). Basically, these methods determine which inputs need to be applied to a faulty device, and which measurements need to be made, in order to optimally discriminate between a number of diagnoses. In comparison with the approach in this paper, these methods have been adapted to qualitative models. By considering only qualitative distinctions, however, one may fail to discriminate between alternative behaviors. Although two models may predict the same qualitative behavior, their (semi-)quantitative predictions may be different, as clearly shown in Figure 4.

For the discrimination of the phytoplankton growth models, only one type of experiment was available, a change in the dilution rate. It should be emphasized, though, that the method is not restricted to parameter changes and may even involve structural changes of the models. A limitation of the method, however, is that the set of experiments needs to be specified in advance. In the case of the dilution rate experiments, for instance, ten possible values from a continuous range have been selected. There is obviously no guarantee that the optimal value is included in the list of prespecified experiments. A subject for further research would therefore be to handle continuous ranges of experiments, and more generally, to move away from the *selection* of experiments to the *design* of experiments. When the set of possible experiments is not given in advance, the criterion ΔJ can be defined as a function of the unknown parameter values. In this case, the determination of the optimal discriminatory ex-

periment will require the solution of a complex optimization problem.

References

- D. Berleant and B. Kuipers. Qualitative and quantitative simulation: Bridging the gap. *Artificial Intelligence*, 95:215–256, 1997.
- D. Berleant, A. Chandra, K. Bognaes, C.-G. Liaw, L. Sheng, and J. Ch'ng. Probabilities of qualitative behaviors for dependability analysis of a fault-tolerance model. In *Proceedings of the ACM/SIGAPP Symposium on Applied Computing*, pages 883–889, 1992.
- O. Bernard and J.-L. Gouzé. Transient behavior of biological loop models with application to the Droop model. *Mathematical Biosciences*, 127:19–43, 1995.
- J. Caperon and J. Meyer. Nitrogen-limited growth of marine phytoplankton. I. Changes in population characteristics with steady-state growth rate. *Deep-Sea Research*, 19:601–618, 1972.
- D.E. Contois. Kinetics of bacterial growth: relationship between population density and specific growth rate of continuous cultures. *Journal of General Microbiology*, 21:40–50, 1959.
- M.J. Cooney and K.A. McDonald. Optimal dynamic experiments for bioreactor model discrimination. *Applied Microbiology and Biotechnology*, 43:826–837, 1995.
- J. de Kleer. Using crude probability estimates to guide diagnosis. *Artificial Intelligence*, 45:381–391, 1990.
- M.R. Droop. Vitamin B12 and marine ecology. IV. The kinetics of uptake growth and inhibition in *Monochrysis lutheri*. *Journal of the Marine Biological Association*, 48(3):689–733, 1968.
- D. Espie and S. Macchietto. The optimal design of dynamic experiments. *AIChE Journal*, 35(2):223–229, 1989.
- V.V. Fedorov. *Theory of Optimal Experiments*. Academic Press, 1972.
- B. Kuipers. *Qualitative Reasoning: Modeling and Simulation with Incomplete Knowledge*. MIT Press, 1994.
- R. Leitch and Q. Shen. Prioritising behaviors in qualitative simulation. In *Proceedings of the 13th International Joint Conference on Artificial Intelligence, IJCAI'93*, pages 1523–1528, 1993.
- J. Monod. *Recherches sur la croissance des cultures bactériennes*. Herman and Cie, 1942.
- P. Struss. Testing for discrimination of diagnoses. In *Working Notes of the 5th International Workshop on Principles of Diagnosis, DX'94*, 1994.
- I. Vatcheva and H. de Jong. Semi-quantitative comparative analysis. In *Proceedings of the 16th International Joint Conference on Artificial Intelligence, IJCAI'99*, pages 1034–1040, 1999.
- I. Vatcheva, H. de Jong, and N.J.I. Mars. Selection of perturbation experiments for model discrimination. In *Proceedings of the 14th European Conference on Artificial Intelligence, ECAI-2000*, pages 191–195, 2000.