

# Distance to VY Canis Majoris with VERA

Yoon Kyung Choi (MPIfR)

In collaboration with:

Tomoya Hirota<sup>1</sup>, Mareki Honma<sup>1</sup>, Hideyuki Kobayashi<sup>1,2</sup>, and VERA members<sup>1,2,3</sup>

1 Mizusawa VERA Observatory, NAOJ


2 The University of Tokyo

3 Kagoshima University

# Outline

- Red Supergiants on HR diagram
- VY Canis Majoris
- Phase-referencing VLBI Observations with VERA
- Results & Discussion
  - Distance to VY CMa
  - Estimation of Stellar Position using SiO masers
  - Kinematics of the H<sub>2</sub>O masers in the Circumstellar Envelopes
- Summary

# Red Supergiants ①




Theoretical evolutionary model  
(Massey et al. 2005)

## Red Supergiants

- Evolved phase of  $9 M_{\odot} < M < 40 M_{\odot}$  stars on main sequence
- mass-loss rate  $10^{-4}$ - $10^{-5} M_{\odot} \text{ yr}^{-1}$
- lifetime  $10^{5-6}$  yr on RSG
- luminosity  $10^{4-5} L_{\odot}$
- effective temperature 3000 K

# Red Supergiants ②



(Massey et al. 2005)

## Problem

- There is a **discrepancy** between theoretically predicted and observed locations of RSGs on the HR diagram

luminosity depends on the (distance)<sup>2</sup>


→ accurate distance measurements are essential

- The distance to RSGs

→ too far to obtain reliable distance with trigonometric parallax measurements

The distance measurements of red supergiants are important to study properties of evolved massive stars.

# VY Canis Majoris ①




R.A. offset (arcsecond)  
HST images (Smith et al. 2001)

## Properties

- distance 1.5 kpc (Lada & Reid 1978)  
**with 30% accuracy !**
- luminosity  $5 \times 10^5 L_{\odot}$   
(Humphreys & Davidson 1994)
- mass-loss rate  $3 \times 10^{-4} M_{\odot} \text{ yr}^{-1}$   
(Danchi et al. 1994)
- effective temperature 2800 K  
(Monnier et al. 1999)

# VY Canis Majoris ②



(Massey et al. 2006)


distance 1.5 kpc  
Luminosity from the SED:  $2-5 \times 10^5 L_{\odot}$   
Temperature based on the spectral type: 2800 K

Aperture masking interferometry  $R_{*} \sim 8.3$  AU  
Temperature based on the spectral type: 2800 K

distance 1.5 kpc  
 $V=8.5$  (AAVSO)  
 $A_V$  &  $T_{\text{eff}}$  (MARCS Model)  
(4) TiO (red/NIR bands) :  $3650 \pm 25$  K  
(5) TiO (blue/yellow bands) :  $3450 \pm 25$  K  
(6) V-K:  $> 3475 \pm 35$  K  
(7) V-J:  $3705 \pm 90$  K  
 $A_V=3.20$

We need to measure a trigonometric parallax to obtain accurate luminosity.


# VY Canis Majoris ③



Bipolar outflow  
SiO ( $v=0$   $J=1-0$ ) emission  
VLA  
(Shinnaga et al. 2004)

## Mass-loss

- bipolar outflow (Shinnaga et al. 2004)
- asymmetric mass loss (Humphreys et al. 2007)



Asymmetric mass loss by HST observations  
(Humphreys et al. 2007)

# Aim of this study

- Measure the distance to VY CMa with a trigonometric parallax method.
- Reveal the structure and the 3-dimensional kinematics of the circumstellar envelopes around VY CMa using H<sub>2</sub>O and SiO masers.



# Observations

- Phase-referencing VLBI observations with VERA
- 10 epochs for 13 months since April 2006
- H<sub>2</sub>O masers (22 GHz) & SiO masers (43 GHz)
- Simultaneous dual-beam observations


Target source: VY CMA

Reference source:



J0725-2640

(S.A. 1.059 degrees)

Two receivers  
in each antenna





- Angular resolution (2270 km baseline)
  - 1.2 mas at 22 GHz
  - 0.6 mas at 43 GHz
- Velocity resolution  $\sim 0.21 \text{ km s}^{-1}$



uv coverage


# H<sub>2</sub>O masers of VY CMa

April, 2006





(Choi et al. 2008, PASJ in press)  
astro-ph arXiv:0808.0641

# Parallax Measurements



Measured positions = parallax + proper motion  
 $\text{H}_2\text{O}$  maser component at  $V_{\text{LSR}} \sim 0.55 \text{ km s}^{-1}$



Best fit for R.A. :


$$\pi = 0.88 \pm 0.08 \text{ mas} \rightarrow d = 1.14^{+0.11}_{-0.09} \text{ kpc}$$

Proper motion :  $-2.09 \pm 0.16 \text{ mas yr}^{-1}$  in R.A.

$1.02 \pm 0.61 \text{ mas yr}^{-1}$  in DEC

(Choi et al. 2008, PASJ in press)  
[astro-ph arXiv:0808.0641](https://arxiv.org/abs/0808.0641)

# Luminosity of VY CMa



The SED of VY CMa  
(Humphreys 2006)


- With our distance, we re-estimate the luminosity

$$L = 4 \pi d^2 F_{\text{bol}}$$

$$d = 1.14^{+0.11}_{-0.09} \text{ kpc}$$

$$L = (3.0 \pm 0.5) \times 10^5 L_{\odot}$$

# The location of VY CMa on the HR diagram




- Re-estimated luminosity with our distance of 1.14 kpc  
 $L \sim (3 \pm 0.5) \times 10^5 L_{\odot}$
- When we adopt the effective temperature of 3650 K (Massey et al. 2006), our result is consistent with the theoretical evolutionary track of initial mass of 25  $M_{\odot}$ .
- There is still uncertainty in the estimation of temperature.

(Choi et al. 2008, PASJ in press)  
astro-ph arXiv:0808.0641

# Inner motions of H<sub>2</sub>O masers

(Choi et al. in prep.)



$\alpha$  (J2000) 07h22m58.3315s  
 $\delta$  (J2000) -25d46'03.174"


Yoon Kyung Choi

- Subtract averaged absolute proper motions
- Average motion  
 $-3.24 \pm 0.16 \text{ mas yr}^{-1}$   
in right ascension  
 $2.06 \pm 0.60 \text{ mas yr}^{-1}$   
in declination

9<sup>th</sup> EVN Symposium, Bologna

# Superposition of masers


(Choi et al. in prep.)




- The circumstellar structure is revealed by phase-referencing VLBI observations with different frequencies of masers in detail.
- The SiO masers are tools to estimate the stellar position in the obscured dusty region by mass-loss with the highest resolution.

(0,0)  $\alpha$  (J2000) 07h22m58.3315s  
 $\delta$  (J2000) -25d46'03.174"

# Coordinate System




For H<sub>2</sub>O masers, we know  
① positions on 2-dimension, and  
② velocities on 3-dimension.



X-axis : right ascension  
Y-axis : declination  
Z-axis : radial direction





# 3-dim. Kinematics of the H<sub>2</sub>O Masers



- Our results show the bipolar outflow along to the line of sight.

# 3-dim Kinematics of the H<sub>2</sub>O Masers



- Our results show the bipolar outflow along to the line of sight.
- This is consistent with the result from Shinnaga et al. (2004).

# Summary

- We measured a distance to VY CMa with a trigonometric parallax.  
 $\pi = 0.866 \pm 0.075 \text{ mas} \rightarrow d = 1.14^{+0.11}_{-0.09} \text{ kpc}$
- We re-estimated the luminosity of VY CMa  
 $\rightarrow L = (3.0 \pm 0.5) \times 10^5 L_{\odot}$
- When we adopt the temperature of 3650 K, the location of VY CMa on HR diagram is consistent with the evolutionary track of initial mass of  $25 M_{\odot}$  star.
- The maps of the H<sub>2</sub>O and SiO masers are superposed, and we estimated a stellar position.
- 3-dimensional kinematics of the circumstellar envelopes of the H<sub>2</sub>O masers suggest a bipolar outflow along the line of sight.