

Distribution of Major Health Risks: Findings from the Global Burden of Disease Study

Anthony Rodgers^{1*}, Majid Ezzati², Stephen Vander Hoorn¹, Alan D. Lopez³, Ruey-Bin Lin¹, Christopher J. L. Murray², Comparative Risk Assessment Collaborating Group

1 Clinical Trials Research Unit, School of Population Health, University of Auckland, New Zealand, **2** Harvard School of Public Health, Boston, Massachusetts, United States of America, **3** School of Population Health, University of Queensland, Brisbane, Australia

Competing Interests: The authors have declared that no competing interests exist. ADL is a member of the editorial board of *PLoS Medicine*.

Author Contributions: AR, ME, ADL and CJLM designed the study. SVH and RBL analyzed the data. AR, ME, SVH, and ADL contributed to writing the paper.

Academic Editor: Thomas Novotny, University of California at San Francisco, United States of America

Citation: Rodgers A, Ezzati M, Vander Hoorn S, Lopez AD, Lin RB, et al. (2004) Distribution of major health risks: Findings from the Global Burden of Disease study. *PLoS Med* 1(1): e27.

Received: June 2, 2004
Accepted: August 13, 2004
Published: October 19, 2004

DOI:
10.1371/journal.pmed.0010027

Copyright: © 2004 Rodgers et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Abbreviations: BMI, body mass index; DALY, disability-adjusted life-year; PAF, population attributable fraction; SBP, systolic blood pressure; SD, standard deviation; WHO, World Health Organization

*To whom correspondence should be addressed. E-mail: a.rodgers@ctr.u.auckland.ac.nz

ABSTRACT

Background

Most analyses of risks to health focus on the total burden of their aggregate effects. The distribution of risk-factor-attributable disease burden, for example by age or exposure level, can inform the selection and targeting of specific interventions and programs, and increase cost-effectiveness.

Methods and Findings

For 26 selected risk factors, expert working groups conducted comprehensive reviews of data on risk-factor exposure and hazard for 14 epidemiological subregions of the world, by age and sex. Age-sex-subregion-population attributable fractions were estimated and applied to the mortality and burden of disease estimates from the World Health Organization Global Burden of Disease database. Where possible, exposure levels were assessed as continuous measures, or as multiple categories. The proportion of risk-factor-attributable burden in different population subgroups, defined by age, sex, and exposure level, was estimated. For major cardiovascular risk factors (blood pressure, cholesterol, tobacco use, fruit and vegetable intake, body mass index, and physical inactivity) 43%–61% of attributable disease burden occurred between the ages of 15 and 59 y, and 87% of alcohol-attributable burden occurred in this age group. Most of the disease burden for continuous risks occurred in those with only moderately raised levels, not among those with levels above commonly used cut-points, such as those with hypertension or obesity. Of all disease burden attributable to being underweight during childhood, 55% occurred among children 1–3 standard deviations below the reference population median, and the remainder occurred among severely malnourished children, who were three or more standard deviations below median.

Conclusions

Many major global risks are widely spread in a population, rather than restricted to a minority. Population-based strategies that seek to shift the whole distribution of risk factors often have the potential to produce substantial reductions in disease burden.

Introduction

Reliable and comparable analysis of risks to health is an important component of evidence-based policies and programs for disease prevention [1,2]. An important feature of risk assessment, with implications for specific interventions as well as broad prevention policies, is the distribution of disease burden among population subgroups. These subgroups may be defined by demographic factors, such as age and sex, or by socioeconomic status. Subgroups can also be defined by the level of exposure to a risk factor, if exposures are defined in multiple categories or continuously.

Understanding the distribution of risk-factor burden is particularly important for targeting interventions. For example, the large number of road traffic accident injuries and deaths among young adult males may be largely associated with binge alcohol consumption by this group. Interventions that focus on this population subgroup and their specific drinking behaviors may be more effective or cost-effective than, for example, raising alcohol taxes, which would have a more diffuse impact on alcohol consumption. On the other hand, the majority of effects from



risk factors such as blood pressure have been found to be among those at moderately elevated levels, motivating interventions beyond those intended for clinical hypertension [3,4,5].

The distributions of the health effects of risk-factor exposure by age and sex or by exposure level have been studied in specific cohorts and for specific risk factors [6,7,8]. Most notably Rose's seminal work stated that "a large number of people exposed to a small risk may generate many more cases than a small number exposed to high risk" [9]. Using the data from a global and regional assessment of multiple major risk factors, this paper reports the distribution by exposure levels, age, and sex of disease burden attributable to several major risk factors. The findings of this analysis confirm that Rose's observations have global relevance and also illustrate important new patterns on specific distributions of disease burden for multiple risks, in different age groups, and in populations at various stages of economic development.

Methods

The methods and data sources for individual risk factors and for estimating population attributable fractions (PAFs) and disease burden attributable to them have been fully described elsewhere [1,2] and are summarized below. The contribution of a risk factor to disease or mortality relative to some alternative exposure scenario (i.e., PAF, defined as the proportional reduction in population disease or mortality that would occur if exposure to the risk factor were reduced to an alternative exposure scenario [10]) is given by the generalized "potential impact fraction":

$$PIF = \frac{\int_{x=0}^m RR(x)P(x)dx - \int_{x=0}^m RR(x)P'(x)dx}{\int_{x=0}^m RR(x)P(x)dx} \quad (1)$$

$RR(x)$ is relative risk at exposure level x , $P(x)$ is the population distribution of exposure, $P'(x)$ is the alternative or counterfactual distribution of exposure, and m is the maximum exposure level.

The alternative (counterfactual) scenario used in this work is the exposure distribution that would result in the lowest population risk, referred to as the theoretical minimum-risk exposure distribution (Table 1) [1,2]. The theoretical minimum exposure distribution was zero in most cases since zero exposure reflected minimum risk (e.g., no smoking). For some risk factors, zero exposure was an inappropriate choice as the theoretical minimum, either because it is physiologically impossible (e.g., body mass index [BMI] and cholesterol) or because there are physical lower limits to exposure reduction (e.g., ambient particulate matter concentration and occupational noise). For these risk factors, the lowest levels observed in specific populations and epidemiological studies were used as the theoretical minimum. For factors with protective effects (i.e., fruit and vegetable intake and physical activity), a counterfactual exposure distribution was chosen based on levels in high-intake populations (e.g., fruit and vegetable intake in Greece) and the level to which the benefits may continue given current scientific evidence. Using theoretical

minimum exposure distribution as the counterfactual has the advantage of providing a vision of potential gains in population health by risk reduction from all levels of suboptimal exposure in a consistent way across risk factors.

Estimates were made for eight age groups (0–4, 5–14, 15–24, 25–44, 45–59, 60–69, 70–79, and 80+ y), both sexes, and 14 Global Burden of Disease subregions (Table 2). PAFs were estimated for mortality and incidence and were applied to regional cause-specific mortality and disease burden from the World Health Organization (WHO) Global Burden of Disease database (Table 1). Burden of disease, reported annually in the annexes of the *World Health Report*, was expressed in disability-adjusted life-years (DALYs) with methods and assumptions described elsewhere [11]. Aggregate results (both mortality and disease burden) for all ages, sexes, and exposure levels have been reported elsewhere [1,2]. Many risks act simultaneously to cause disease, and joint effects have also been estimated [12], though the separate effects are presented in this paper. The aim of the analyses reported here was to obtain estimates of the distribution of disease burden by age, sex, and exposure level. To make separate estimates of disease burden by exposure level, the relationship in equation 1 was re-estimated with the entire exposure distribution divided into "narrow bands," with each band corresponding to one level of exposure, and the estimation repeated for each such exposure band.

Results

The distribution of deaths and DALYs attributable to the risk factors by age and sex is shown in Table 3. Disease burden attributable to being underweight and to micronutrient deficiencies in children was equally distributed among males and females. The total all-age disease burden from iron and vitamin A deficiencies was greater in females because of effects on maternal mortality and morbidity conditions. The disease burden of other diet-related risks, physical inactivity, environmental risks, and unsafe sex (sex with an infected partner without any measures to prevent infection) occurred almost equally in males and females. However, approximately 80% of disease burden from addictive substances and 60%–90% from occupational risks occurred among men (bearing in mind that the assessment considered only formal employment). Women experienced an estimated two-thirds of the disease burden from childhood sexual abuse and all of the burden caused by a lack of contraception.

The estimated disease burden from childhood and maternal undernutrition; unsafe water, sanitation, and hygiene; and global climate change (much of whose estimated effects are mediated through nutritional and water variables) was almost exclusively among children under 5 y of age. For these risks, more than 85% of total attributable burden occurred in this age group, with the exception of iron deficiency, for which 30% of burden was borne by women of childbearing age. Only a small fraction of disease burden from the risk factors considered was among the 5–14 y olds. This was because some of the leading diseases of this age group (e.g., injuries and depression) have complex causes that could not easily be included in the current risk-based framework [12]. For other leading diseases of this group (e.g., diarrhea and lower respiratory infections), most epidemiological studies have

Table 1. Leading Global Risk Factors, Exposure Variables, Theoretical Minima, and Attributable Deaths and Disease Burden (measured in DALYs) in 2000

Category	Risk Factor	Exposure Variable	Theoretical Minimum	Deaths (Thousands)	DALYs (Percent of Total Global Burden of Disease)
Childhood and maternal undernutrition	Underweight	Children <1 SD weight-for-age compared to the international reference group in 1-SD increments; maternal BMI <20 kg/m ²	Same fraction of children <1 SD weight-for-age compared to the international reference group; all women of childbearing age with body mass index >20 kg/m ²	3,748	9.5%
	Iron deficiency anemia	Hemoglobin level (g/dl) estimated from prevalence of anemia	Hemoglobin distributions that halve anemia prevalence, estimated to occur if all iron deficiency were eliminated ^a	841	2.4%
	Vitamin A deficiency	Prevalence of vitamin A deficiency, estimated as low serum retinol concentrations (<0.70 μmol/L) among children aged 0–4 y and among pregnant women (aged 15–44 y) Less than the United States recommended dietary allowances for zinc	No vitamin A deficiency	778	1.8%
	Zinc deficiency	Less than the United States recommended dietary allowances for zinc	The entire population consuming sufficient dietary zinc to meet physiological needs, taking into account routine and illness-related losses and bioavailability	789	1.9%
Other nutrition-related risk factors and physical activity	High blood pressure	Level of systolic blood pressure	115 ± 6 mm Hg	7,141	4.4%
	High cholesterol	Level of total blood cholesterol	3.8 ± 0.6 mmol/l (147 ± 23 mg/dl)	4,415	2.8%
	High BMI	BMI (height over weight squared)	21 ± 1 kg/m ²	2,591	2.3%
Sexual and reproductive health	Low fruit and vegetable consumption	Fruit and vegetable intake per day	600 ± 50 g intake per day for adults	2,726	1.8%
	Physical inactivity	Three categories: inactive, insufficiently active (<2.5 h/wk of moderate-intensity activity, or less than 4,000 KJ/wk), and sufficiently active; activity in discretionary-time, work, and transport considered	All having at least 2.5 h/wk of moderate-intensity activity or equivalent (4,000 KJ/wk)	1,922	1.3%
Sexual and reproductive health	Unsafe sex	Sex with an infected partner without any measures to prevent infection (represented as parameters of an HIV model)	No unsafe sex	2,886	6.3%
	Non-use and use of ineffective methods of contraception	Prevalence of traditional methods or non-use of contraception	Use of modern contraceptives for all women who want to space or limit future pregnancies	149	0.6%

Table 1. Continued

Category	Risk Factor	Exposure Variable	Theoretical Minimum	Deaths (Thousands)	DALYs (Percent of Total Global Burden of Disease)
Addictive substances	Tobacco	Current levels of smoking impact ratio (indirect indicator of accumulated smoking risk based on excess lung cancer mortality); oral tobacco use prevalence	No tobacco use	4,907	4.1%
	Alcohol	Current alcohol consumption volumes and patterns	No alcohol use ^b	1,804	4.0%
	Illicit drugs	Use of amphetamine, cocaine, heroin, or other opioids and intravenous drugs	No illicit drug use	204	0.8%
Environmental risks	Unsafe water, sanitation, and hygiene	Six scenarios, ranging from regulated water and sanitation with hygiene to no improved water supply and no improved sanitation	Absence of transmission of diarrheal disease through water, sanitation, and hygiene	1,730	3.7%
	Indoor smoke from household use of solid fuels	Household use of solid fuels and ventilation	No household solid fuel use	1619	2.6%
Occupational risks	Urban air pollution	Estimated annual average particulate matter concentration for particles with aerodynamic diameters less than 2.5 μm ($\text{PM}_{2.5}$) or 10 μm (PM_{10})	7.5 $\mu\text{g}/\text{m}^3$ for $\text{PM}_{2.5}$; 15 $\mu\text{g}/\text{m}^3$ for PM_{10}	799	0.4%
	Lead	Current blood lead levels	0.016 $\mu\text{g}/\text{dl}$ blood lead levels ^c	234	0.9%
	Global climate change	Climate scenarios based on various carbon emissions and concentrations	1961–1990 atmospheric carbon concentrations	154	0.4%
Occupational risks	Risk factors for injuries	Current proportions of workers exposed to injury risk factors	Exposure corresponding to lowest rate of work-related fatalities observed: one per million per year for 16–17-y-olds employed as service workers in the United States	310	0.7%
	Carcinogens	Proportions of workers exposed to background, low, and high levels of workplace carcinogens	No work-related exposure above background	109	0.1%
	Airborne particulates	Proportions of workers with background, low, and high levels of exposure	No work-related exposure above background	356	0.4%
	Ergonomic stressors	High, moderate, and low exposure based on occupational categories	Physical workload at the level of managers and professionals (low)	0	0.1%
	Noise	High (>95 dBA) and moderate (85–90 dBA) exposure categories	Less than 85 dBA on average over eight working hours	0	0.3%

Table 1. Continued

Category	Risk Factor	Exposure Variable	Theoretical Minimum	Deaths (Thousands)	DALYs (Percent of Total Global Burden of Disease)
Other selected risk factors	Contaminated injections in health-care settings Child sexual abuse	Exposure to least one contaminated injection Prevalence of noncontact abuse, contact abuse, and intercourse	No contaminated injections No abuse	501 79	0.7% 0.6%

See Table 1 in Ezzati et al. [2] for disease outcomes and data sources.

^a The resulting hemoglobin levels vary across regions and age-sex groups (from 11.66 g/dl in children under five in a region of Southeast Asia to greater than 14.5 g/dl in adult males in developed countries) because the other risks for anemia (e.g., malaria) vary.

^b Theoretical minimum for alcohol is zero, the global theoretical minimum. Specific subgroups may have a non-zero theoretical minimum.

^c Theoretical minimum for lead is the blood lead levels expected at background exposure levels. Health effects were quantified for blood lead levels above 5 µg/dl where epidemiological studies have quantified hazards.

DOI: 10.1371/journal.pmed.0010027.t001

focused on children under 5 y of age and do not provide hazard estimates for older children.

The disease burden from other diet-related risks, tobacco, and occupational risks (except injuries and back pain) was almost equally distributed among adults above and below the age of 60 y. For example, 43% of disease burden due to blood pressure and 61% of burden due to tobacco occurred in the 15–59-y age group. More than 90% of disease burden attributable to lack of contraception, illicit drugs, occupational ergonomic stressors, risk factors for injury, and childhood sexual abuse occurred in adults below the age of 60 y. Similarly, 67%–87% of the disease burden attributable to alcohol, unsafe sex, and unsafe health-care injections arose from events occurring between 15 and 59 y of age. Most of the risks whose hazards are concentrated in the younger adults are causes of injuries, neuropsychiatric diseases, maternal conditions, and HIV/AIDS. Assessment by the level of economic and demographic development illustrated that, with the exception of alcohol, which has global presence, the majority of disease burden from risks for mortality and disease among young adults is concentrated in developing countries (see Figure 1 in [2]). Stratification of economic and demographic development was also a determinant of the age distribution patterns for risk factors for chronic diseases, which occurred in younger age groups in developing countries than in developed regions. For example, in high-mortality developing regions, 69% of the tobacco burden occurred in people aged 15–59 y, whereas this share was 63% for lower-mortality developing countries and 55% for developed countries.

The distributions of attributable risk-factor burden by exposure levels are shown in Table 4 for those risks quantified using categorical variables and in Figure 1 for those with continuous variables. For most of these risks a substantial proportion of attributable disease burden occurred among those with modest elevations of risk. For example, while 35% of the large disease burden from being underweight in childhood, the leading risk factor for global loss of healthy life, occurred in severely underweight children who would be subject to clinical interventions (i.e., more than three standard deviations [SDs] from referent group median), the rest occurred in children only 1–3 SDs below the median. Even though the relative risks for the latter group are much lower, the number of children exposed to risk at this level is so great that the total attributable disease burden amounted to much more than that occurring in the severe category. Similarly, 52% of the attributable burden from physical inactivity occurred among those engaged in some but less than the recommended level of 2.5 h per week of moderate-intensity activity. For unsafe water, sanitation, and hygiene, almost all of the attributable disease burden was distributed among three of the five at-risk exposure categories, with approximately equal levels. This reflects the fact that the exposure categories were defined as the presence or absence of technology-based water and sanitation interventions. During decades of water and sanitation projects, many countries have “clustered” in a limited number of technology groups. There is likely to be large heterogeneity of exposure within each exposure category, however, because of factors such as quantity of water consumed and hygiene behavior [13].

Figure 1 shows that at the aggregated level a substantial proportion of the attributable disease burden for high blood

Table 2. Global Burden of Disease 2000 Subregions

WHO Region	Mortality Stratum ^a	Countries	Population (Thousands)
AFR	D	Algeria, Angola, Benin, Burkina Faso, Cameroon, Cape Verde, Chad, Comoros, Equatorial Guinea, Gabon, Gambia, Ghana, Guinea, Guinea-Bissau, Liberia, Madagascar, Mali, Mauritania, Mauritius, Niger, Nigeria, Sao Tome and Principe, Senegal, Seychelles, Sierra Leone, Togo	294,078
	E	Botswana, Burundi, Central African Republic, Congo, Côte d'Ivoire, Democratic Republic of the Congo, Eritrea, Ethiopia, Kenya, Lesotho, Malawi, Mozambique, Namibia, Rwanda, South Africa, Swaziland, Uganda, United Republic of Tanzania, Zambia, Zimbabwe	345,515
AMR	A	Canada, Cuba, United States of America	325,183
	B	Antigua and Barbuda, Argentina, Bahamas, Barbados, Belize, Brazil, Chile, Colombia, Costa Rica, Dominica, Dominican Republic, El Salvador, Grenada, Guyana, Honduras, Jamaica, Mexico, Panama, Paraguay, Saint Kitts and Nevis, Saint Lucia, Saint Vincent and the Grenadines, Suriname, Trinidad and Tobago, Uruguay, Venezuela	430,932
EMR	D	Bolivia, Ecuador, Guatemala, Haiti, Nicaragua, Peru	71,230
	B	Bahrain, Cyprus, Iran (Islamic Republic of), Jordan, Kuwait, Lebanon, Libyan Arab Jamahiriya, Oman, Qatar, Saudi Arabia, Syrian Arab Republic, Tunisia, United Arab Emirates	139,059
EUR	D	Afghanistan, Djibouti, Egypt, Iraq, Morocco, Pakistan, Somalia, Sudan, Yemen	342,576
	A	Andorra, Austria, Belgium, Croatia, Czech Republic, Denmark, Finland, France, Germany, Greece, Iceland, Ireland, Israel, Italy, Luxembourg, Malta, Monaco, Netherlands, Norway, Portugal, San Marino, Slovenia, Spain, Sweden, Switzerland, United Kingdom	411,889
	B	Albania, Armenia, Azerbaijan, Bosnia and Herzegovina, Bulgaria, Georgia, Kyrgyzstan, Poland, Romania, Slovakia, Tajikistan, The Former Yugoslav Republic of Macedonia, Turkey, Turkmenistan, Uzbekistan, Yugoslavia	218,458
SEAR	C	Belarus, Estonia, Hungary, Kazakhstan, Latvia, Lithuania, Republic of Moldova, Russian Federation, Ukraine	243,184
	B	Indonesia, Sri Lanka, Thailand	293,819
WPR	D	Bangladesh, Bhutan, Democratic People's Republic of Korea, India, Maldives, Myanmar, Nepal	1,241,806
	A	Australia, Brunei Darussalam, Japan, New Zealand, Singapore	154,354
	B	Cambodia, China, Cook Islands, Fiji, Kiribati, Lao People's Democratic Republic, Malaysia, Marshall Islands, Micronesia (Federated States of), Mongolia, Nauru, Niue, Palau, Papua New Guinea, Philippines, Republic of Korea, Samoa, Solomon Islands, Tonga, Tuvalu, Vanuatu, Viet Nam	1,532,933

High-mortality developing regions: AFR-D, AFR-E, AMR-D, EMR-D, and SEAR-D. Lower-mortality developing regions: AMR-B, EMR-B, SEAR-B, and WPR-B. Developed regions: AMR-A, EUR-A, EUR-B, EUR-C, and WPR-A.

^a A, very low child mortality and very low adult mortality; B, low child mortality and low adult mortality; C, low child mortality and high adult mortality; D, high child mortality and high adult mortality; E, high child mortality and very high adult mortality.

DOI: 10.1371/journal.pmed.0010027.t002

pressure, cholesterol, and BMI and low fruit and vegetable intake occurred in the “mid-range” exposures. For example, the second and third quartiles (i.e., half of attributable disease burden) occurred at the following levels: systolic blood pressure (SBP) of 130–150 mm Hg, cholesterol of 5.0–6.1 mmol/l, BMI of 25–32 kg/m², and fruit and vegetable intake of 150–300 g/d (2–4 servings/d). This was similar to or greater than the amount of disease burden occurring among individuals with risk-factor levels above the commonly used, but arbitrary, thresholds for hypertension, hypercholesterolemia, and obesity indicated in Figure 1.

Despite the above finding on the important role of moderately elevated levels in total disease burden, the actual patterns of how disease burden is distributed among exposure levels varied across different regions and risk factors (Figure 2). For example, Figure 2 shows that a comparatively larger fraction of the disease burden attribut-

able to elevated blood pressure, cholesterol, and BMI occurred at lower levels in developing regions compared to developed regions, mainly because of lower age-specific exposure levels in those populations. Figure 2 also shows that the skewness of the distribution of disease burden was not substantially different across different age groups for BMI. This is because the comparatively larger relative risk per unit BMI at younger ages (which leads to more right-hand skew) is counterbalanced by the comparatively lower BMI at younger ages (which leads to left-hand skew). Therefore, the different distributions of BMI-attributable disease burden by region resulted not from the different age structures of populations across major world regions, but rather from the lower *age-specific* BMI levels in those countries [14]. This is in contrast to blood pressure, for which disease burden in younger age groups occurred at lower exposures because the

Table 3. Distribution of Risk-Factor-Attributable Deaths and Disease Burden (DALYs) by Age and Sex

Category	Risk Factor	Mortality (Percent)					Disease Burden (Percent)						
		0–4	5–14	15–59	≥60	Males	Females	0–4	5–14	15–59	≥60	Males	Females
Childhood and maternal undernutrition	Childhood and maternal underweight	100	0	0	0	51	49	100	0	0	0	51	49
	Iron deficiency anemia	72	1	22	4	45	55	62	6	30	2	45	55
	Vitamin A deficiency	85	1	14	0	43	57	86	1	12	0	44	56
	Zinc deficiency	100	0	0	0	51	49	100	0	0	0	51	49
Other nutrition-related risk factors and physical inactivity	High blood pressure	0	0	19	81	49	51	0	0	43	57	54	46
	High cholesterol	0	0	22	78	48	52	0	0	50	50	55	45
	Overweight and obesity (high BMI)	0	0	26	74	45	55	0	0	57	43	47	53
	Low fruit and vegetable consumption	0	0	23	77	53	47	0	0	49	51	57	43
	Physical inactivity	0	0	21	79	50	50	0	0	48	52	53	47
Addictive substances	Tobacco	0	0	30	70	79	21	0	0	61	39	82	18
	Alcohol	1	1	65	33	91	9	1	3	87	9	85	15
	Illicit drugs	0	0	100	0	80	20	0	2	98	0	77	23
Sexual and reproductive health	Unsafe sex	16	1	77	6	47	53	18	1	79	2	46	54
	Non-use and use of ineffective methods of contraception	0	0	100	0	0	100	0	0	100	0	0	100
Environmental risks	Unsafe water, sanitation, and hygiene	68	5	13	14	52	48	77	8	13	3	51	49
	Urban air pollution	3	0	16	81	51	49	12	0	40	49	56	44
	Indoor air pollution from household use of solid fuels	56	0	5	38	41	59	83	0	8	9	49	51
	Lead	0	0	41	57	66	34	75	0	16	8	55	45
	Global climate change	86	3	6	5	49	51	88	5	6	1	49	51
Occupational risks	Risk factors for injuries	0	0	85	14	94	6	0	0	95	5	93	7
	Carcinogens	0	0	28	72	85	15	0	0	51	49	83	17
	Airborne particulates	0	0	17	83	74	26	0	0	65	35	77	23
	Ergonomic stressors	0	0	0	0	0	0	0	0	95	5	59	41
	Noise	0	0	0	0	0	0	0	0	89	11	67	33
Other selected risk factors	Contaminated injections in health-care settings	10	2	53	35	63	37	16	3	67	13	61	39
	Child sexual abuse	0	0	81	22	48	52	0	0	96	4	36	64

See Table 1, and Figure 1 in [2], for definition of each risk factor, data sources and methods, and total magnitude of mortality and DALYs.
DOI: 10.1371/journal.pmed.0010027.t003

age patterns of exposure and relative risk do not entirely compensate.

Discussion

The findings reported here should be considered within the context of limited available data and are subject to uncertainty, which varies across risk factors and is generally most marked in developing countries. Full discussion of uncertainty in the basic data sources and parameters is provided elsewhere and includes the uncertainties in estimates of disease incidence, duration, severity, and disability weighting [1,2]. Uncertainty in this risk assessment is by far

dominated by absence or limitations of direct studies on exposure, hazard, and background disease burden, rather than parameter uncertainty, such as sampling error. This has motivated innovative assumptions and extrapolations even in the case of the best-studied risk factors in a limited number of countries [6]. While estimates of hazard size in individual studies were as much as possible adjusted for confounding effects, it is likely that residual confounding effects remain to some extent, hence leading to errors in estimation. Extrapolation of hazard from a limited number of studies to other populations is another source of potential error. While the robustness of relative measures of risk has been confirmed for more proximal factors in studies across populations




Figure 1. Distribution by Exposure Level of Attributable Disease Burden Due to Selected Continuous Risk Factors

Figure 1 shows the distribution of the estimated cardiovascular disease (CVD) burden of disease (in DALYs) attributable to four major continuous risk factors, by exposure levels. Half the attributable burden occurs to the left of the solid vertical line and half occurs to the right. The dashed vertical lines indicate commonly used thresholds—150 mm Hg for hypertension, 6.0 mmol/l for hypercholesterolemia, and 30 kg/m² for obesity. The blood pressure and cholesterol levels plotted are the estimated usual levels [22], which tend to have a smaller SD than levels based on one-off measurements commonly used in population surveys, because of normal day-to-day and week-to-week fluctuations. For example, the distribution of usual blood pressure is about half as wide as the distribution of one-off blood pressure measures, and so many fewer people would be classified as hypertensive (or hypotensive) if classifications were based on usual rather than one-off blood pressure. Thus, if a population mean SBP was 134 mm Hg, the SD of once-only measures might be 17 mm Hg (with about 18% of the population having one-off SBP over 150 mm Hg), and the SD of usual SBP based on many measures would be about 9 mm Hg (hence about 5% of the population would have usual SBP over 150 mm Hg).

DOI: 10.1371/journal.pmed.0010027.g001

[15,16,17], their extrapolation is an important source of uncertainty for more distal risks (e.g., childhood sexual abuse) or those whose effects are heterogeneous (e.g., alcohol and injuries versus alcohol and cancer). Direct exposure data for many risk factors are limited because of both difficulties in their measurement and underinvestment in risk-factor surveillance, especially in developing countries.

Of particular relevance to the current analysis is the fact that, due to data limitations, some risks were measured using categorical variables (e.g., indoor smoke from solid fuels, underweight, and physical inactivity) even though the health effects occur along a continuum. Other risk factors were represented using indirect or aggregate indicators (e.g., smoking impact ratio for accumulated hazards of smoking, and motor vehicle accident registries for alcohol-caused accidents) that do not allow quantification of risks along continua of exposure. Two important sources of uncertainty with implications for interventions are correlations among multiple risk factors and the skewness of risk-factor distribution. Because risks are often correlated (e.g., undernutrition, poor water and sanitation, and the use of solid fuels are more common among poor households in developing countries, and smokers are more likely to have poor diets), the contributions of high-exposure groups are likely to be underestimated. In addition to being a source of underestimation at higher exposure levels, risk-factor correlation implies that the same individuals and groups are at the high end of multiple risk factors. Positive (rightward) skewness of

exposure distribution, not modeled here, would also lead to an underestimation of events occurring in those who were hypertensive, hypercholesterolemic, or obese. The importance of skewness is emphasized by the observation that the recent increase in the proportion of people who are overweight and obese (e.g., in the United States) has involved not only a shift in the distribution, but also increasing skewness, which has extended the high-exposure tail. Coupled with risk-factor correlation, this should motivate more systematic data collection and reporting to determine mean exposure as well as the complete shape of distribution.

The temporal nature of risk-factor exposure also has implications for the current cross-sectional estimates. There is an expectation that the size and rank order of risk-factor burden will alter in coming decades as a result of changes in exposure levels and delays between exposure and hazard. For example, it is predicted that by 2020 the leading risks to health will be (1) unsafe sex, principally because of HIV/AIDS and driven by increasing exposure, and (2) tobacco, because of market expansion of tobacco products in developing countries and delayed temporal effects on major health outcomes such as lung cancer [1].

This analysis in multiple age and exposure categories, or along a continuum of exposures, showed that, globally, a considerable proportion of the disease burden attributable to major risk factors occurred among those with only moderately raised levels, not the extremes such as those that define hypertension, obesity, or severe malnutrition. Further for

Table 4. Distribution by Exposure Level of Attributable Burden Due to Selected Categorical Risk Factors

Category Type	Risk Factor					
	Physical Inactivity		Underweight		Water, Sanitation, and Hygiene	
	Category Description	Proportion of Total Attributable Burden	Category Description	Proportion of Total Attributable Burden	Category Description	Proportion of Total Attributable Burden
Referent category	At least 2.5 h/wk of moderate-intensity activity or equivalent (400 KJ/wk)	–	Same fraction of children below 1 SD weight-for-age as the international reference group	–	Absence of transmission of diarrheal disease through water, sanitation, and hygiene	–
Exposure categories	Some activity, but less than 2.5 h/wk of moderate-intensity activity	49%	1–2 SD below standard	20%	Regulated water supply and full sanitation coverage, with partial treatment for sewage	0%
	Little or no physical activity	51%	2–3 SD below standard	46%	Improved water supply, basic sanitation, improved access to drinking water, improved personal hygiene, and water disinfected at point of use	39%
			3+ SD below standard	35%	Improved water supply and basic sanitation	3%
					Basic sanitation but no improved water supply	28%
				No improved water supply and no basic sanitation	30%	

DOI: 10.1371/journal.pmed.0010027.t004

many chronic-disease risk factors, such as tobacco and high blood pressure, as well as risk factors for injuries and sexual and reproductive health, significant proportions of disease burden occurred from events in middle ages, especially in developing countries. The concentration of disease burden from such a large number of risk factors in those below 60

years of age illustrates the large, and at times neglected, disease burden from risks that affect young adults in developing nations, with important consequences for economic development [18].

The distribution of risks and their determinants in a population have major implications for strategies of pre-




Figure 2. Distribution of Attributable Cardiovascular Disease Burden Due to BMI, Blood Pressure, and Cholesterol by Exposure Level, Age, and Level of Development

Conventions as for Figure 1.

DOI: 10.1371/journal.pmed.0010027.g002

vention. As stated by Rose, risk typically increases across the spectrum of a risk factor [8]. Rose's work led to one of the most fundamental axioms in disease prevention across risk factors: "A large number of people exposed to a small risk may generate many more cases than a small number exposed to high risk." The analysis in this work showed that the risk factors for many of the major global diseases, such as lower respiratory infections, diarrhea, ischemic heart disease, and stroke exhibit such behavior, because they are caused by risks that occur along a continuum. Therefore, managing individual, high-risk crises, while appropriate for individuals, can only have a limited preventive effect at the population level and over long time periods because it relies on having adequate discriminative ability to predict future disease, and requires continued and expensive screening for new high-risk individuals. In contrast, population-based strategies that seek to shift the whole distribution of risk factors have the

potential to substantially reduce total disease burden, and possibly over long time periods, if the interventions alter the underlying risk behaviors or their socioeconomic causes. This is particularly relevant in the context of risk factors such as those related to under- or overnutrition that affect societies in all stages of development. For example, a policy to reduce salt content in manufactured foods would result in a leftward shift in the population distribution of blood pressures and a surprisingly large reduction in cardiovascular disease [5]. Another example would be population-level measures that affected energy intake (such as the availability and price of energy-dense foods) and/or expenditure (such as the level of motorization and mechanization)—these can be expected to determine the distribution of a population's BMI levels, and hence largely determine its level of type 2 diabetes [1].

There were distinct patterns for the distribution of disease burden across risk factors, and across regions at various

stages of development. At the extreme of these diverse patterns, for risk factors with acute exposure and acute outcomes, the underlying relationship is considerably more complex. For example while in many societies most alcohol-attributable injury (e.g., traffic accidents) involves people who on average drink moderate amounts [19], these people would be on the more extreme, high-risk spectrum in a different dimension: volume of drinking right before the injury. Therefore, if exposure to risk factors is clustered or the risk relationship does not follow a linear pattern, high-exposure groups may indeed play a disproportionately important role [20,21].

In summary, the analysis presented in this paper confirms and extends to a global level previous work indicating that the impact of many major risks is important across their exposure levels, not just among people with particularly high levels [8]. This analysis also illustrates that, beyond the general principle of population-wide shifts, there are important population-specific and risk-factor-specific patterns in how the burden of disease attributable to risk factors is distributed. Systematic assessment of multiple risks within any given population can provide the basis for selecting packages of interventions that include population-wide measures as well as highly targeted interventions provided to much smaller subsections of the population with constellations of major risks [1,5,18].

Acknowledgments

This work was sponsored by the National Institute of Aging grant PO1-AG17625. The sponsor had no influence on analysis and results. Anthony Rodgers holds a National Heart Foundation Senior Fellowship. We thank Clarissa Gould-Thorpe for manuscript preparation. We thank Mie Inoue, Doris Ma Fat, Susan Piccolo, Chalapati Rao, Kenji Shibuya, Niels Tomijima, and Marie-Claude von Rulach for assistance with global burden of disease databases and project management.

Comparative Risk Assessment Collaborating Group

Core, methodology, statistical analysis, editorial and peer review, and writing: Majid Ezzati (Harvard University), Anthony Rodgers (University of Auckland), Alan D. Lopez (University of Queensland), Stephen Vander Hoorn (University of Auckland), Christopher J. L. Murray (Harvard University).

Childhood and maternal underweight: Steven Fishman (Johns Hopkins University), Laura E. Caulfield (Johns Hopkins University), Mercedes de Onis (WHO), Monika Blössner (WHO), Adnan A. Hyder (Johns Hopkins University), Luke Mullany (Johns Hopkins University), Robert E. Black (Johns Hopkins University).

Iron deficiency anemia: Rebecca J. Stoltzfus (Cornell University), Luke Mullany (Johns Hopkins University), Robert E. Black (Johns Hopkins University).

Vitamin A deficiency: Amy J. Rice (Johns Hopkins University), Keith P. West, Jr. (Johns Hopkins University), Robert E. Black (Johns Hopkins University).

Zinc deficiency: Laura E. Caulfield (Johns Hopkins University), Robert E. Black (Johns Hopkins University).

High blood pressure: Carlene Lawes (University of Auckland), Stephen Vander Hoorn (University of Auckland), Malcolm Law (St Bartholomew's and Royal London School of Medicine), Paul Elliott (Imperial College School of Medicine), Stephen MacMahon (University of Sydney), Anthony Rodgers (University of Auckland).

High cholesterol: Carlene Lawes (University of Auckland), Stephen Vander Hoorn (University of Auckland), Malcolm Law (St Bartholomew's and Royal London School of Medicine), Stephen MacMahon (University of Sydney), Anthony Rodgers (University of Auckland).

Overweight and obesity (high BMI): W. Philip T. James (International Obesity Task Force), Rachel Jackson-Leach (International Obesity Task Force), Cliona Ni Mhurchu (University of Auckland), Eleni Kalamara (International Obesity Task Force), Maryam Shayeghi

(International Obesity Task Force), Neville J. Rigby (International Obesity Task Force), Chizuru Nishida (WHO), Anthony Rodgers (University of Auckland).

Low fruit and vegetable consumption: Karen Lock (London School of Hygiene and Tropical Medicine), Joceline Pomerleau (London School of Hygiene and Tropical Medicine), Louise Causer (London School of Hygiene and Tropical Medicine), Martin McKee (London School of Hygiene and Tropical Medicine).

Physical inactivity: Fiona C. Bull (Loughborough University), Tim Armstrong (WHO), Tracy Dixon (Australian Institute of Health and Welfare), Sandra Ham (United States Centers for Disease Control and Prevention), Andrea Neiman (United States Centers for Disease Control and Prevention), Michael Pratt (United States Centers for Disease Control and Prevention).

Tobacco: Majid Ezzati (Harvard University), Alan D. Lopez (University of Queensland).

Alcohol: Jürgen Rehm (Addiction Research Institute and University of Toronto), Robin Room (Stockholm University), Maristela Monteiro (WHO), Gerhard Gmel (Swiss Institute for the Prevention of Alcohol and Other Drug Problems), Kathryn Graham (University of Western Ontario), Nina Rehm (WHO), Christopher T. Sempos (University of Buffalo), Ulrich Frick (Psychiatric University Hospital Zurich), David Jernigan (Georgetown University).

Illicit drugs: Louisa Degenhardt (University of New South Wales), Wayne Hall (University of Queensland), Matthew Warner-Smith (University of New South Wales), Michael Lynskey (University of New South Wales).

Unsafe sex: Emma Slaymaker (London School of Hygiene and Tropical Medicine), Neff Walker (Joint United Nations Programme on HIV/AIDS), Basia Zaba (London School of Hygiene and Tropical Medicine), Martine Collumbien (London School of Hygiene and Tropical Medicine).

Non-use and use of ineffective methods of contraception: Martine Collumbien (London School of Hygiene and Tropical Medicine), Makeda Gerressu (London School of Hygiene and Tropical Medicine), John Cleland (London School of Hygiene and Tropical Medicine).

Unsafe water, sanitation, and hygiene: Annette Prüss-Ustun (WHO), David Kay (University of Wales), Lorna Fewtrell (University of Wales), Jamie Bartram (WHO).

Urban air pollution: Aaron Cohen (Health Effects Institute), Ross Anderson (St George's Hospital Medical School and University of London), Bart Ostro (California Environmental Protection Agency), Kiran Dev Pandey (World Bank), Michal Krzyzanowski (WHO), Nino Künzli (University of Southern California), Kersten Gutschmidt (WHO), Arden Pope (Brigham Young University), Isabelle Romieu (Instituto Nacional de Salud Publica de Mexico), Jonathan Samet (Johns Hopkins University), Kirk Smith (University of California at Berkeley).

Indoor smoke from household use of solid fuels: Kirk R. Smith (University of California at Berkeley), Sumi Mehta (Health Effects Institute), Mirjam Feuz (Federal Office of Public Health of Switzerland).

Lead: Annette Prüss-Ustun (WHO), Lorna Fewtrell (University of Wales), Philip Landrigan (Mount Sinai School of Medicine), José Luis Ayuso (Universidad Autonoma de Madrid).

Global climate change: Anthony McMichael (Australian National University), Diarmid Campbell-Lendrum (WHO), Sari Kovats (London School of Hygiene and Tropical Medicine), Sally Edwards (London School of Hygiene and Tropical Medicine), Paul Wilkinson (London School of Hygiene and Tropical Medicine), Frank Tanser (Medical Research Council of South Africa), David Le Sueur (deceased) (Medical Research Council of South Africa), Michael Schlesinger (University of Illinois at Urbana-Champaign), Natasha Andronova (University of Illinois at Urbana-Champaign), Robert Nicholls (University of Middlesex), Theresa Wilson (University of Middlesex), Simon Hales (University of Otago).

Occupational risk factors for injuries: Marisol Concha (Asociacion Chilena de Seguridad of Chile), Deborah Imel Nelson (WHO), Marilyn Fingerhut (WHO), Annette Prüss-Ustun (WHO), James Leigh (University of Sydney), Carlos Corvalan (WHO).

Occupational carcinogens: Tim Driscoll (University of Sydney), Deborah Imel Nelson (WHO), N. Kyle Steenland (United States National Institute for Occupational Safety and Health), James Leigh (University of Sydney), Marisol Concha (Asociacion Chilena de Seguridad), Annette Prüss-Ustun (WHO), Marilyn Fingerhut (WHO), Carlos Corvalan (WHO).

Occupational airborne particulates: Tim Driscoll (University of

Sydney), N. Kyle Steenland (United States National Institute for Occupational Safety and Health), Deborah Imel Nelson (WHO), James Leigh (University of Sydney), Marisol Concha (Asociacion Chilena de Seguridad), Marilyn Fingerhut (WHO).

Occupational ergonomic stressors: Annette Prüss-Ustun (WHO), Laura Punnett (University of Massachusetts at Lowell), SangWoo Tak (University of Massachusetts at Lowell), Deborah Imel Nelson (WHO), Marilyn Fingerhut (WHO), James Leigh (University of Sydney), Sharonne Phillips (Occupational Ergonomics).

Occupational noise: Deborah Imel Nelson (WHO), Marisol Concha (Asociacion Chilena de Seguridad), Marilyn Fingerhut (WHO).

Contaminated injections in health-care settings: Anja M. Hauri (Staatliches Untersuchungsamt Hessen), Gregory L. Armstrong (United States Centers for Disease Control and Prevention), Yvan J. F. Hutin (WHO).

Child sexual abuse: Gavin Andrews (University of New South Wales), Justine Corry (University of New South Wales), Cathy Issakidis (University of New South Wales), Tim Slade (University of New South Wales), Heather Swanson (University of New South Wales).

Poverty and risk: Tony Blakely (University of Otago), Simon Hales (University of Otago), Charlotte Kieft (University of Otago), Nick Wilson (University of Otago), Alistair Woodward (University of Auckland). ■

References

- World Health Organization(2002) The world health report 2002: Reducing risks, promoting healthy life. Geneva: World Health Organization. 250 p.
- Ezzati M, Lopez AD, Rodgers A, Vander Hoorn S, Murray CJ, Comparative Risk Assessment Collaborating Group. (2002) Selected major risk factors and global and regional burden of disease. *Lancet* 360: 1347–1360.
- Rodgers A, Lawes C, MacMahon S (2000) The global burden of cardiovascular disease conferred by raised blood pressure. Benefits of reversal of blood pressure-related cardiovascular risk in Eastern Asia. *J Hypertens* 18: S3–S5.
- Cook NR, Cohen J, Hebert P, Taylor JO, Hennekens CH (1995) Implications of small reductions in diastolic blood pressure for primary prevention. *Arch Intern Med* 155: 701–709.
- Murray CJL, Lauer JA, Hutubessy RCW, Niessen L, Tomijima N, et al. (2003) Effectiveness and costs of interventions to reduce systolic blood pressure and cholesterol: A global and regional analysis on reduction of cardiovascular-disease risk. *Lancet* 361: 717–725.
- Peto R, Lopez AD, Boreham J, Thun M, Heath CW (1992) Mortality from tobacco in developed countries: Indirect estimates from national vital statistics. *Lancet* 339: 1268–1278.
- Rodgers A, MacMahon S (1999) Blood pressure and the global burden of cardiovascular disease. *Clin Exp Hypertens* 21: 543–552.
- Rose G (1992) The strategy of preventive medicine. Oxford: Oxford University Press. 138 p.
- Rose G (1985) Sick individuals and sick populations. *Int J Epidemiol* 14: 32–38.
- Miettinen OS (1974) Proportion of disease caused or prevented by a given exposure, trait or intervention. *Am J Epidemiol* 99: 325–332.
- Murray CJL, Lopez AD, editors (1996) The global burden of disease. Cambridge (Massachusetts): Harvard University Press. 990 p.
- Ezzati M, Vander Hoorn S, Rodgers A, Lopez AD, Mathers CD, et al. (2003) Estimates of global and regional potential health gains from reducing multiple selected major risk factors. *Lancet* 362: 271–280.
- Curtis V, Cairncross S, Yonli R (2000) Domestic hygiene and diarrhoea—Pinpointing the problem. *Trop Med Int Health* 5: 22–32.
- James WPT, Jackson-Leach R, Ni Mhurchu C, Kalamara E, Shayeghi M, et al. (2004) Body mass index and the global burden of disease. In: Ezzati M, Lopez A, Rodgers A, Vander Hoorn S, Murray C, editors. Comparative quantification of health risks: Global and regional burden of disease due to selected major risk factors. Geneva: World Health Organization. pp. 497–596.
- Eastern Stroke and Coronary Heart Disease Collaborative Group(1998) Blood pressure, cholesterol and stroke in eastern Asia. *Lancet* 352: 1801–1807.
- Law MR, Wald NJ, Thompson SG (1994) By how much and how quickly does reduction in serum cholesterol concentration lower risk of ischaemic heart disease? *BMJ* 308: 367–373.
- Horton R (2000) Common sense and figures: The rhetoric of validity in medicine (Bradford Hill Memorial Lecture 1999). *Stat Med* 19: 3149–3164.
- Commission on Macroeconomics and Health(2001) Macroeconomics and health: Investing in health for economic development. World Health Organization. Available: <http://www.cid.harvard.edu/cidcmh/CMHReport.pdf>. Accessed 25 August 2004.
- Kreitman N (1986) Alcohol consumption and the preventive paradox. *Br J Addict* 81: 353–363.
- Lemmens P (2001) Relationship of alcohol consumption and alcohol problems at the population level. In: Heather N, Peters TJ, Stockwell T, editors. International handbook of alcohol dependence and problems. Chichester (United Kingdom): John Wiley and Sons.
- Skog OJ (1999) Prevention paradox revisited. *Addiction* 94: 751–757.
- MacMahon S, Peto R, Cutler J, Collins R, Sorlie P, et al. Blood pressure, stroke, and coronary heart disease. Part I, prolonged differences in blood pressure: Prospective observational studies corrected for the regression dilution bias. *Lancet* 335: 765–774.

Patient Summary

Background. Health policy makers must understand existing health risks and which diseases cause the most health problems. The Global Burden of Disease database, maintained by the World Health Organization, collects information from around the world on risk factors such as malnutrition, childbirth, tobacco use, and cholesterol levels, as well as on diseases such as depression, blindness, and diarrhea. This information can be used to target health interventions.

What Did the Researchers Do and Find? These researchers determined for 26 major risk factors the distribution of disease burden by age, sex, and exposure level. They found that many risk factors (such as high blood pressure and high cholesterol levels) occur across populations, and are not confined to one sex or age group. And for most risk factors, exposure to moderate risks (which is usually more common than exposure to severe risk) is responsible for causing most disease.

What Do the Results Mean, and Who Will Use Them? Measures that reduce exposure to risk factors across whole populations, if only by a modest extent, can achieve large reductions in disease burden. This information is important for people who set global health policies and priorities.

What Are the Problems with the Study? There are many challenges in estimating the impact of different major risks, each of which has distinct effects on a number of diseases. In addition, exposure to some risk factors today will cause disease only many years from now, so the picture will change over time. The major finding of the global distribution of many major risks is secure, but the exact extent of this remains uncertain due to the paucity of data in developing countries.



Minerva Access is the Institutional Repository of The University of Melbourne

Author/s:

Rodgers, A;Ezzati, M;Hoorn, SV;Lopez, AD;Lin, RB;Murray, CJL

Title:

Distribution of major health risks: Findings from the global burden of disease study

Date:

2004-10-01

Citation:

Rodgers, A., Ezzati, M., Hoorn, S. V., Lopez, A. D., Lin, R. B. & Murray, C. J. L. (2004).
Distribution of major health risks: Findings from the global burden of disease study. PLOS
MEDICINE, 1 (1), pp.44-55. <https://doi.org/10.1371/journal.pmed.0010027>.

Persistent Link:

<http://hdl.handle.net/11343/265441>

License:

[CC BY](#)