

WestminsterResearch

<http://www.westminster.ac.uk/westminsterresearch>

DNA damage in obesity: Initiator, promoter and predictor of cancer

Usman, M. and Volpi, E.

NOTICE: this is the authors' version of a work that was accepted for publication in Mutation Research/Reviews in Mutation Research. Changes resulting from the publishing process, such as peer review, editing, corrections, structural formatting, and other quality control mechanisms may not be reflected in this document. Changes may have been made to this work since it was submitted for publication. A definitive version was subsequently published in Mutation Research/Reviews in Mutation Research, doi: 10.1016/j.mrrev.2018.08.002, 2018.

The final definitive version in Mutation Research/Reviews in Mutation Research is available online at:

<https://dx.doi.org/10.1016/j.mrrev.2018.08.002>

© 2018. This manuscript version is made available under the CC-BY-NC-ND 4.0 license <https://creativecommons.org/licenses/by-nc-nd/4.0/>

The WestminsterResearch online digital archive at the University of Westminster aims to make the research output of the University available to a wider audience. Copyright and Moral Rights remain with the authors and/or copyright owners.

Whilst further distribution of specific materials from within this archive is forbidden, you may freely distribute the URL of WestminsterResearch: (<http://westminsterresearch.wmin.ac.uk/>).

In case of abuse or copyright appearing without permission e-mail repository@westminster.ac.uk

Accepted Manuscript

Title: DNA damage in obesity: Initiator, promoter and predictor of cancer

Authors: Moonisah Usman, Emanuela V. Volpi

PII: S1383-5742(18)30055-3
DOI: <https://doi.org/10.1016/j.mrrev.2018.08.002>
Reference: MUTREV 8249

To appear in: *Mutation Research*

Received date: 19-6-2018
Revised date: 29-7-2018
Accepted date: 15-8-2018



Please cite this article as: Usman M, Volpi EV, DNA damage in obesity: Initiator, promoter and predictor of cancer, *Mutation Research-Reviews in Mutation Research* (2018), <https://doi.org/10.1016/j.mrrev.2018.08.002>

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

DNA damage in obesity: initiator, promoter and predictor of cancer

Moonisah Usman and Emanuela V. Volpi

Department of Biomedical Sciences, Faculty of Science and Technology, University of Westminster,
115 New Cavendish Street, London W1W 6UW, U.K.

Correspondence should be addressed to: e.volpi@westminster.ac.uk

Abstract

Epidemiological evidence linking obesity with increased risk of cancer is steadily growing, although the causative aspects underpinning this association are only partially understood. Obesity leads to a physiological imbalance in the regulation of adipose tissue and its normal functioning, resulting in hyperglycaemia, dyslipidaemia and inflammation. These states promote the generation of oxidative stress, which is exacerbated in obesity by a decline in anti-oxidant defence systems. Oxidative stress can have a marked impact on DNA, producing mutagenic lesions that could prove carcinogenic.

Here we review the current evidence for genomic instability, sustained DNA damage and accelerated genome ageing in obesity. We explore the notion of genotoxicity, ensuing from systemic oxidative stress, as a key oncogenic factor in obesity. Finally, we advocate for early, pre-malignant assessment of genome integrity and stability to inform surveillance strategies and interventions.

Key words: DNA Damage, Obesity, Inflammation, Oxidative Stress, Cancer

1. Connecting obesity and cancer via oxidative stress and DNA damage

The association of obesity with cancer is naturally one of the most concerning aspects and the object of intense scrutiny. A number of prospective and case-control studies have recognised an epidemiological correlation between obesity and cancer in a sex and site-specific manner. Published evidence has been extensively reviewed [1, 2]. A recent paper by Kyrgiou and colleagues [3], in which over two-hundred meta-analyses of observational studies linking adiposity to cancer risk were systematically evaluated, has confirmed definitive evidential association between obesity and eleven types of cancer, including oesophageal adenocarcinoma, multiple myeloma, and cancers of the gastric cardia, colon, rectum, biliary tract system, pancreas, breast, endometrium, ovary and kidney. Obesity manifests with inflammation, hyperglycaemia and dyslipidaemia, maintained to be drivers of carcinogenesis as they may endorse the biological mechanisms associated with tumour development, sustainability and metastasis [4]. These pathological states in obesity may have a key role in the initiation of cancer as chronic inflammation and the metabolic disturbances in obesity have a common pathological consequence termed oxidative stress, a long-recognised threat for genome stability and integrity [5]. Cellular function, viability and fate are predominantly determined by the genomic stability and integrity of the cell. Conversely, cancer genomes are highly rearranged and genomic instability has been described as an enabling characteristic of cancer [6]. Oxidative stress has a physiological relevance as the excessive generation of reactive oxygen species (ROS) may lead to oxidative DNA damage and induce tumorigenic alterations [7], as well as drive genomic instability by mimicking replication stress [8].

ROS are produced via many fundamental cellular processes. Endogenous sources of ROS include mitochondrial respiration and enzymatic reactions within phagocytes and peroxisomes. ROS are

stimulated by growth factors and serve as necessary signalling molecules [9]. Concentrations of ROS are tightly regulated under a redox (reduction oxidation) system to prevent toxicity. Obesity is known to disrupt the redox system by elevating the production of ROS, and decreasing the availability of anti-oxidant defence enzymes [10]. A calorie-rich diet and an abundance of macronutrients in obesity, calls for continuity in the Krebs cycle that consequently leads to a leak of more electrons from the mitochondrial electron transport chain. These electrons promote a reduction of oxygen molecules – a process that will generate more ROS. However, the over-generation of ROS in obese individuals is brought about by more than just a continuum in the Krebs cycle.

2. The oxidative burden of inflammation

Chronic inflammation is a well-known etiological factor for DNA damage and promotor of neoplastic transformations in cells [11]. In a state of obesity, the release of pro-inflammatory molecules including CRP, TNF-alpha and IL-8 from adipose tissue is well documented [12–15]. Adipocytokines particularly TNF-alpha, IL-6 and IL-1 promote the recruitment of various cells including neutrophils, macrophages and dendritic cells, indicating the beginning of an inflammatory process [16]. These cells not only generate more pro-inflammatory cytokines but also encourage production of ROS/RNS via NADPH oxidases, particularly the NOX1, NOX2 and NOX4 isoforms [17]. Further production of ROS/RNS is generated as obesity develops with hypertrophy and hyperplasia of adipocytes, resulting in tissue hypoxia and cell necrosis [10].

Cell necrosis causes altered adipokine expression, including reduced levels of the anti-inflammatory hormone adiponectin [18]. Adiponectin is secreted by adipose tissue and may inhibit the actions of TNF-alpha and prevent recruitment of inflammatory cells [19]. Adiponectin deficiency may play a pathological role further than inflammation as it has been associated with obesity and cancers of the breast [20], GI tract [21,22], blood [23] and prostate [24].

Increased death of adipose cells has also been indicated as a path to release cell-free DNA into systemic circulation [25]. Nishimoto and collaborators [26] have recently identified that release of cell-free DNA caused by adipocytes' degeneration promotes macrophages accumulation in adipose tissue via Toll-like Receptor 9 (TLR9), originally known as a sensor of exogenous DNA fragments. Macrophage infiltration into adipose tissue is characteristic of obesity and may have further pathological consequences when they secrete high concentrations of nitrous oxide (NO) [27]. In high concentration, NO can cause deamination, oxidation and strand breakages in DNA [28]. Chronic inflammation and its ability to inflict DNA damage has been demonstrated in models of *H.pylori* associated gastrointestinal cancer [29] and ulcerative colitis associated colon cancer [30], as well as HCV mediated liver cancer [31]. Unresolved inflammation has also been reported to cause pre-malignant lesions in the mouth [32]. Izano and colleagues [33] investigated the risk of increased IL-6, CRP and TNF-alpha with colon cancer and other obesity related cancers in a follow-up study of 2490 participants. Their findings evoke the causative relationship between chronic inflammation and colon cancer in obese adults, but evidence for other obesity related cancers is non-conclusive. Overall, there is substantial evidence that chronic inflammation is a causative factor for DNA damage. This phenomenon should be further explored in obesity to elucidate the risks of carcinogenesis.

3. Dyslipidaemia and Lipid Peroxidation

Obesity coincides with an increased circulation of free fatty acids (FFA) and the deposition of excess fat in white adipose tissue (WAT), marked as dyslipidaemia [34]. Excessive plasma lipids are susceptible to oxidative modification which generates more ROS, and can also activate a protein kinase C pathway resulting in elevated production of nitroxide – a potent ROS [35]. The deposition of FFA in WAT also contributes to ROS generation by attracting leukocytes and causing inflammation [35].

Dyslipidemia - mainly the elevation in low density lipoproteins and fall in high-density lipoproteins, disrupts the redox system and leads to lipid peroxidation in obesity [36]. Lipid peroxidation is a mechanism for generating malondialdehyde (MDA) that can react with DNA to produce mutagenic MDA-DNA adducts such as MDA-deoxyguanosine [36]. MDA has the potential to cause inter-strand cross links with DNA [37] and point mutations [38]. The first reaction can be highly cytotoxic, whereas the latter could prove carcinogenic [39]. Elevated levels of MDA have been identified in both adult [40] and child [41] models of obesity. As MDA levels can also be a remarkable finding in breast and lung cancer patients [42], a potential causative association between lipid peroxidation, DNA damage and cancer risk in obesity can be argued.

4. Hyperglycaemia and ROS generation

In obese individuals, insulin signalling can be significantly impaired. Glucose transport, protein C kinase and other enzymatic activity can be defective, resulting in increased plasma glucose levels (hyperglycaemia) [43]. FFA circulation and uptake in the liver can further contribute towards hyperglycaemia when FFA is selected as an energy source over glucose [44]. Hyperglycaemia has been associated with increasing ROS generation four-fold in adipocytes and thus promoting inflammation [45]. These factors also contribute towards hyperinsulinemia and insulin resistance in obesity [46]. Hyperinsulinemia correlates with decreased levels of IGF binding protein 1 and IGF binding protein 2, thereby readily causing an increase in the availability of IGF-1 [46].

Excess free circulating IGF-1 has been associated with an increased risk of breast cancer [47] and colorectal cancer [48]. Although, IGF-1 was found to have a role in activating DNA repair pathways against oxidative DNA damage [49], elevated levels of IGF-1 in obesity may be a physiological response to oxidative stress [50] but this avenue is open to question.

Moreover, IGF-1 has been linked with down-regulation of sex hormone binding globulin (SHBG) [51], which could result in an increase in the availability of free oestrogen [52]. The association between obesity and increased risk of endometrial and breast cancer may owe to increased oestrogen

levels in obese individuals [53,54]. Traditionally, sustained exposure to oestrogen has been associated with upregulation of genes that control cell proliferation and cell cycle progression, marking an important role in tumorigenesis [55]. In addition, oestrogen may endogenously induce DNA modifications, create oestrogen-DNA adducts and generate oxidative DNA damage [56]. Oxygen radicals can also be generated as a result of the redox cycling of oestrogen metabolites [56]. Therefore, it can be inferred that hyperglycaemia in obese subjects may promote ROS generation via IGF-1, insulin and oestrogen, thus increasing the likelihood of DNA damage and cancer initiation.

5. Inadequate Anti-oxidant Defence

Increased BMI has been associated with a decline over time in essential regulators of the redox system. Ozata and collaborators [57] identified decreased activity of SOD and GPx in erythrocytes of 76 male obese subjects, in relation to age-matched healthy weight controls. Similarly, recent investigations have confirmed lower activities of SOD and GPx in obese women, although no significant difference in CAT activity was observed [58]. Interestingly, Erdevi and collaborators [59] reported the SOD enzyme to be increasingly active in a cohort of obese children. However, this finding was in parallel to identification of increased ROS production in the same cohort, suggesting that obesity may trigger an early physiological response to oxidative stress in children by increasing the activity of anti-oxidant enzymes to scavenge the ROS. Similarly, total oxidant status (TOS) and total anti-oxidant status (TAS) were both reported to be higher in a cohort of 37 obese children compared to health weight controls [60]. Other findings suggest that the implications of oxidative stress worsen with the severity of obesity, placing obese children at higher risk of oxidative stress. Albuali and colleagues [61] identified elevated activity of SOD and CAT in overweight children but found that this level of functioning did not exist in the obese cohort. Anti-oxidant status in obesity appears to be an interesting paradox whereby anti-oxidant defence mechanisms may be maximised in order to counteract oxidative stress [62], but whether this effect is substantial in controlling ROS and

preventing pathological consequences is unknown. Overall, there are substantial indications of a weakened anti-oxidant defence mechanism in both adult and childhood obesity [57,58,63].

A decrease in anti-oxidant activity has been linked with DNA damage and acute lymphoblastic leukaemia in children, although it is complicated to establish whether this association is causative [64]. A study conducted on mice deficient in SOD activity presented them to have a decreased lifespan and increased likelihood of liver malignancies, [65]. A latter review by Asaduzzaman and colleagues [66] explored the paradoxical roles of anti-oxidant enzymes in carcinogenesis. They highlighted that CAT, SOD and GPx activity is heterogeneous in a variety of cancers. Recent studies in oral [67] and breast cancer [68] have given a deeper insight into these roles and suggest that elevated activity of some of these enzymes may also be a cause for concern. Overall, it is plausible that inadequate anti-oxidant mechanisms may predispose to malignancy, possibly via increased DNA damage.

It is important to consider that obesity often coincides with a nutrient-poor diet and a lack of dietary intake of anti-oxidants [69]. Low levels of particular plasma micronutrients, including folate, vitamin B, vitamin C, iron and zinc, may imitate the effects of radiation on DNA, potentially resulting in DNA strand breaks or the production of oxidative lesions [70]. It has also recently emerged that obesity may cause vitamin D deficiency [71]. Vitamin D may play a significant role in managing the redox system, as it was found to upregulate the anti-oxidant enzyme G6PD in human prostate epithelial cells [72] and is associated with reducing the incidence of prostate cancer [73]. Thus, low levels of anti-oxidant enzymes in obesity may also be attributed to micro-nutritional deficiencies. Collectively these disturbances in the redox system and antioxidant defences may increase the potential for DNA damage and risk of cancer, although further research is required to confirm any causative association.

6. Markers of DNA damage in obesity

6.1 Micronuclei

Micronuclei (MNI) are well-recognised markers of genotoxic stress and genomic instability. By definition, a micronucleus is an extra-nuclear body within the cell cytoplasm containing chromosomal fragments or whole chromosomes, resulting either from *clastogenic* events, such as DNA double strand breaks (DSBs), or *aneugenic* events, such as segregation defects giving rise to abnormal numbers of chromosomes. There are indications that retention of micronuclei containing whole chromosomes can proceed into several cell generations [74]. Recently, the relevance of micronuclei as potential drivers of malignant transformation has come to the fore in relation to the phenomenon of chromothripsis, a newly described mutational process whereby micronuclei division cycles enable accumulation of mutagenic re-arrangements in a single or few chromosomes [75]. These localized chromosomal re-arrangements may be transferred to daughter nuclei in subsequent mitotic cycles and play a role in generating a pre-cancerous genome. Micronuclei may display a lack of nuclear membrane integrity when occurring in normal or cancer cells. Therefore, the nuclear envelope of a micronucleus is more likely to rupture, causing exposure of self-DNA to the cytosol. Possible immuno-stimulatory consequences of this event have recently been reported in a mouse model and human cancer cells [76].

Evidence for micronuclei frequency as a possible predictor of cancer is being explored. A series of observational and follow-up studies have consistently demonstrated raised incidence of micronuclei in 10 different types of cancer (examples in Table 1.). Most relevantly, prospective research approaches have associated an elevated frequency of micronuclei with an increased risk of developing cancer. Early research by Bonassi and colleagues followed up 6718 participants across 20 sites after their participation in the Human Micronucleus (HUMN) project and reported a statistically significant risk of stomach and urogenital cancer to be associated with a high baseline frequency of micronuclei in lymphocytes [77]. Another longitudinal study reported evidence to support a possible link between micronuclei and risk of cancer. This research followed 1650 initially disease free adults

for up to 14 years. Retrospective analysis indicates that participants that developed cancer had, on average, a three-fold increase in micronucleus frequency at baseline [78].

Ageing appears to be a major contributing factor to chromosomal instability and micronuclei formation. The influence of age on micronuclei frequency has been quite extensively researched. In an initial study on a large population of 791 participants, micronuclei frequency was found to increase with age and was higher in women than men [79]. Subsequent investigations have detected a biphasic relationship between age and induction of micronuclei in lymphocytes, with a positive correlation between micronuclei and age up to 50 years, but an inverted association beyond that age threshold [80,81]. This biphasic response may owe to the decreased proliferative capacity of cells in older individuals, that may exist as a mechanism to limit possible genomic instabilities [81].

Moreover, the frequency of micronuclei was reported as at least three times higher in the buccal mucosa of adults aged over 60, compared to cells from a cohort of 19-29 year olds [82]. More recent research draws out that micronuclei frequency in lymphocytes can be an indicator of frailty in an ageing cohort, however, whether micronuclei frequency in buccal epithelial cells is positively associated with age is yet to be confirmed [83].

There is increasing evidence to suggest a link between adiposity and micronuclei frequency (Table 2.). The first indication of a positive correlation between micronuclei frequency and BMI was provided by a research on polycystic ovarian syndrome (PCOS) [84]. Further evidence was provided by Andreassi *et al.*, [85], highlighting the correlations between BMI, DNA damage and insulin resistance in women with PCOS. Later, BMI was related with an almost four-fold increase in micronuclei frequency within the PBLs of 83 obese adults, when compared with normal-BMI controls [86]. A positive correlation between BMI and the number of apoptotic and necrotic cells in plasma was identified suggesting the elevated frequency of dying cells to be a result of them possessing a damaged and unstable genomic state. Furthermore, an increased association between BMI and micronuclei frequency was observed in a cohort of obese Italian children [87]. This was the

first study in which the occurrence of genome damage has been investigated in childhood obesity. Compelling evidence on positive associations between weight and length at birth, and three markers of chromosomal instability (MN frequency, nucleoplasmic bridges and nuclear buds) has been reported [88].

Some studies in human and animal models have highlighted discrepancies in terms of influence of adiposity on micronuclei frequency. Although MN frequency was positively associated with age in 300 Korean adults, no correlations were drawn with BMI [89]. An in-depth meta-analysis by Croco and collaborators [90] discovered body mass to be a determining factor for genomic stability, demonstrated by erythrocyte micronucleus frequencies across 47 mammalian species. However, more recently a, research on a diet-induced mouse model of obesity did not find a significant difference in the levels of micronuclei in erythrocytes or reticulocytes, albeit the obese mice sustained a greater frequency of *Pig-a* mutant erythrocytes, a newly described biomarker for somatic mutations [91].

The origins of micronuclei formation in obesity need to be extensively researched. Although, chronic inflammation is a possible pathological factor that may underpin the pathway to micronuclei formation, as identified in patients with cirrhotic liver as a result of hepatitis C infection [92].

Whether this concept can be applied to low-grade, systemic inflammation identified in obesity is an important, unanswered question.

6.2 8-OH-2-deoxy Guanosine (8-OHdG)

8-oxo-deoxy-guanosine (8-OHdG) is the product of free-radical induced oxidation of guanosine. 8-OHdG can be detected in serum, saliva and urine, making it an ideal DNA damage biomarker for large population based studies. The possible mutational consequences of this lesion, including mispairing with adenosine, inhibiting methylation of adjacent cytosines and deleterious GàT transversion mutations have been recently reviewed [93]. Ramon and colleagues [94] first identified that 8-OHdG may also directly interfere with the gene transcription process when found at promoter

regions, by demonstrating that 8-OHdG alters binding of the transcription factor Sp1. The role of Sp1 transcription factors in cell growth, development, and death have been extensively reviewed concluding Sp1 as key players in carcinogenesis [95]. In addition, a recent review highlighted that 8-Oxoguanine DNA Glycosylase 1 (OGG1) – a key repair enzyme, may also play a role in carcinogenesis by recruiting 8-OHdG as a ligand for gene regulation and activation of the Ras pathway [96].

The clinical significance of 8-OHdG has been extensively discussed by Cooke and colleagues, bringing to light that 8-OHdG is a well-established risk marker for age-related pathologies such as cancer [97]. The GT to AC substitution is a commonly described mutation that is derived from the oxidation of guanosine and is also prominent in gastrointestinal tumours [98]. It is not surprising that elevated serum or urinary 8-OHdG lesions are undisputedly associated with malignant tumours at multiple sites (Table 3.).

Interestingly, levels of 8-OHdG are also significantly higher in patients with pre-cancerous lesions of buccal mucosa [99] and dysplastic cervical cells [100]. Therefore, there may be a potential for 8-OHdG to be utilised as a clinical risk marker to detect early carcinogenic processes.

For long, the phenomenon of ageing has been described as a process that incorporates free-radical mediated attacks on nuclear DNA, leading to oxidative stress, telomere attrition and a halt in cell division and proliferation [101]. 8-OHdG lesions may have implications in cellular senescence and ageing caused by oxidative stress. This is because early research recognised 8-OHdG to emerge from senescing human fibroblasts in higher concentrations than their younger counterparts [102]. More recent studies have recognized increasing concentrations of 8-OHdG to be associated with age-related cataracts [103], Alzheimer's [104], Parkinson's [105], and vascular complications in type 2 diabetes [106]. This suggests that 8-OHdG is a specific biomarker for oxidative stress that is consistent with ageing.

Multiple studies have assessed the correlation of 8-OHdG with adiposity with emerging findings not devoid of discrepancies (Table 4.). de la Maza and colleagues were the first to report an association between increased body fat and oxidative DNA damage by assessing 8-OHdG in skeletal muscle [107]. These associations were based upon self-reported weight gain over the last 10 years by patients undergoing a hernia operation. It is intriguing that increased 8-OHdG in these weight gainers coincided with increased TNF-alpha – an undisputed marker of inflammation. Another study assessed 8-OHdG concentrations in lymphocytes and was unable to identify a correlation with BMI in adults [108]. In contrast, there are findings that indicate a correlation between leanness and increased oxidative stress. A one-point reduction in BMI was reported to coincide with a 2.7% increase in urinary 8-OHdG [109]. Similarly, Donmez-Altuntas and collaborators reported decreased levels of 8-OHdG in plasma of obese adults compared to adults of healthy weight, although other biomarkers of DNA damage including micronuclei frequency was elevated [86]. More recently, a study conducted in over 100 obese and healthy weight men concluded that there were no significant differences in urinary 8-OHdG [110]. In summary, the evidence for 8-OHdG as a marker of oxidative stress in adult obesity is inconclusive, this outcome likely due to discrepancies arising from differences in the degree of obesity in participants between studies.

Similarly, discordant are the results of investigations carried out across adolescent cohorts. The first evidence showing a positive association between BMI and oxidative DNA damage in obese adolescents, quantified 8-OHdG in serum via an ELISA method [111]. Across 103 adolescents, a BMI greater than the 95th percentile was associated with increased 8-OHdG. However, Protano and collaborators assessed 8-OHdG in urine across 159 children via liquid chromatography–tandem mass spectrometry and found an inverse association [112]. Yet more recent research has reported a higher concentration of 8-OHdG in urine from a much smaller cohort - 24 obese children [113]. However, the children in this cohort also presented with insulin resistance and metabolic syndrome (MS). The absence or presence of insulin resistance or MS may account for discrepancies in the measurement of

oxidative DNA damage, as MS has been suggested to disrupt anti-oxidant defences and encourage oxidative stress [114].

6.3 γ -H2AX foci

Induction of nuclear γ -H2AX foci or gamma foci signal the start of a crucial repair process that follows a potentially carcinogenic DNA lesion [115]. DNA double strand breaks (DSBs) are detected by the MRN repair complex consisting of NBS1, MRE11 and Rad50. This recognition process leads to activation and recruitment of ATM and a subsequent phosphorylation of γ -H2AX and other targets including BRCA1. As reviewed by Khanna & Jackson, DSBs can have lethal consequences for a cell as they can drive the cell into apoptosis or towards carcinogenesis [116]. Although the suitability of γ -H2AX as a DNA damage marker has been questioned as its presence has been observed in the absence of recognisable DNA damage [117], evidence against the use of γ -H2AX as a marker of DSBs remains limited. Recent research indicates the induction of γ -H2AX to coincide with unrepaired DSBs as well as blocked replication forks [118,119]. Indeed, the suitability of gamma foci continues to be explored as a diagnostic and prognostic marker for cancer (Table 5).

A higher frequency of γ -H2AX foci have been found expressed in breast cancer cells from the more aggressive triple negative breast cancer (TNBC) when compared to cells from non-TNBC [120].

Gamma foci have also been suggested as an appropriate diagnostic and prognostic marker for colon cancer [121] and more recently, epithelial ovarian cancer [122]. The detection of γ -H2AX in pre-cancerous lesions of the bladder *in vivo* [121] and pre-cancerous lesions of the lung and skin *in vitro* gives reason to explore the use of γ -H2AX as a possible predictor of cancer [123].

The DNA damage response to DSBs has been described as a major pathway to cellular senescence in mouse models [124,125]. Sedelnikova and collaborators discovered γ -H2AX foci at sites of persistent DSBs that led to cellular senescence in ageing mice and human cell lines, indicating that γ -H2AX foci may be a marker of un-repairable DNA damage and ageing in mammals [124]. They also

tested this association in cell lines from patients with Werner syndrome – a disorder characterised with premature ageing, and identified a comparable association, although recruitment of DNA proteins was slower to γ -H2AX foci. A more recent study also documented a linear correlation between γ -H2AX foci and ageing in human cell lines [126]. Schurman and colleagues further studied the association of γ -H2AX foci with age related disorders and found a positive correlation with cataracts and hypertension, but not with cancer [126]. This indicates the need for further research identifying whether there is a causative role for γ -H2AX foci in ageing and cancer.

Redon and colleagues reviewed the practical applications of H2AX foci in population studies, and identified their extensive use to draw associations between chronic inflammation and DNA damage [127]. It was emphasised that the long-term consequences of identifying elevated γ -H2AX foci remains an open avenue.

There is consistent evidence to suggest γ -H2AX foci are elevated with adiposity (Table 6).

Villaret and colleagues reported premature senescence of visceral adipose tissue cells in obese subjects, and found the endothelial cells within to be marked with a state of inflammation and up-regulated hypoxia genes in coexistence with increased γ -H2AX foci [128]. Interestingly, this phenomenon was not documented in sub-cutaneous adipose tissue to the same extent. In addition, for the first time, γ -H2AX foci were documented as 8x fold higher in cultured PBLs of obese children when compared to healthy weight controls in a study led by Scarpato and collaborators [87]. They also assessed chromosomal aberrations, via the micronucleus assay, in a large cohort of obese children, alongside markers of chronic inflammation and attributed an increased rate of DSBs to overweightness. The confounding, observed increase in TNF- α , CRP and IL-6 manifests a causative link between chronic low-grade inflammation and DNA damage in obese children. In 2016, the same investigators deepened their investigations into the repair efficiency of DSBs between obese and normal weight adolescents by measuring the response of PBLs to mitomycin C [129]. An almost 2 fold increase was found in mitomycin C induced DSBs in the obese cohort when compared to

healthy controls, measured by γ -H2AX foci. Intriguingly, the level of DNA repair in the obese cohort was repaired faster and more efficiently than in the healthy weight counterparts. This research indicates that there may be a potential for DNA repair mechanisms to be able to cope with the emerging accounts of DNA damage in obesity, but warrants further investigations.

Furthermore, the documented evidence for increased DSB repair initiation linked with childhood obesity has also been implied in obese Zucker rats - a model of genetically predisposed obesity [130]. Obese Zucker rats presented a greater expression of γ -H2AX foci in lung, pancreatic, oesophageal, kidney and gut tissues when compared to lean counterparts. It was also noted that older rats (9-13 weeks) had a more prominent increase in the expression of γ -H2AX foci than younger rats (3-4), further implying that DSB events may correlate with age. Nonetheless, it is important to comprehend that the induction of γ -H2AX in tissue is a well-regulated proper response to reversal of DNA Double strand breaks, although, the relationship between the number of foci present and number of DSBs repaired by γ -H2AX foci is questionable.

6.4 Comet Tails

Comet tails are sensitive biomarkers of deteriorating chromosomal integrity as they enable inference of chromosomal breakages. The comet assay (single cell gel electrophoresis) used to detect these tails is a well-established laboratory assay that was first pioneered by Ostling & Johanson in 1984 [131]. Single cells are embedded in agarose and treated to remove nuclear membranes and histone material prior to electrophoresis. The application of a current allows broken, uncoiled DNA loops to migrate faster towards the anode, creating the shape of a comet, with intact DNA remaining in the comet head whilst damaged, 'broken' DNA migrates in the comet tail. Subsequently, the DNA is treated with an intercalating dye and visualised with fluorescence microscopy. The length of the tail and percentage of DNA inside the tail is taken as a directly proportional measure of DNA damage [132]. The comet assay has been extensively applied to measure genotoxicity, and responsiveness to

chemotherapy, and is now evolving into a diagnostic marker of cancer [133]. Several cases of malignancy have been assessed via the comet assay and found to display significant DNA damage (Table 7.). Furthermore, Katarkar and colleagues identified two remarkable observations via a case-control study using 260 participants. The first is the applicability of the comet assay in PBLs as a biomarker for pre-malignant lesions of the buccal mucosa [134]. Secondly, they uncovered that DNA damage in PBLs correlates with DNA damage in cells extracted from the buccal mucosa, as measured by the presence of micronuclei. This evidence leads to the conclusion that the comet assay in PBLs may possibly act as a cancer susceptibility biomarker for non-haematological malignancies. This avenue requires further exploration and warrants studies to investigate the relevance of non-invasive assays; such as the buccal micronucleus assay, in susceptibility to systemic DNA damage and carcinogenesis.

The comet assay has also been applied to evaluate DNA damage in ageing [135,136]. Research has suggested the comet assay may be a functional biomarker for pre-mature ageing, as increased DNA damage was observed in cells from patients with Hutchinson-Gilford progeria syndrome (HGPS) when compared to normally ageing human fibroblasts [136]. These investigations intensify the link between DNA damage and ageing, when explored using the comet assay.

Body fat can be proposed as a pathological factor for chromosomal breaks following investigations with the comet assay in rat models of obesity and in humans (Table 8.) A correlation study first indicated a positive association between BMI in females and oxidative DNA damage via the comet assay [108]. Later, A positive correlation between increased chromosomal damage and BMI (>30) was demonstrated by the comet assay in pregnant women of a Pakistani population [137] and was subsequently confirmed in an Indian population too [138]. Remarkably, DNA damage was also identified via the comet assay in women with a normal BMI but elevated fat mass [139]. This 'normal-weight-obese' cohort also had higher levels of oxidative DNA damage, measured by 8-OHdG, than the obese cohort of women. These findings warrant for investigations of DNA damage

in obesity to consider measures of adiposity wider than BMI, as fat mass may have a stronger correlation with genomic instability. DNA damage has also been confirmed via the comet assay in liver cells of obese female Zucker rats [140]. This research identified age to be an important factor when interpreting results from the comet assay, since Zucker rats younger than 15 months showed no evidence of comet tails. Comet tails are indicative of DNA damage in adult and animal models of obesity. However, there is a lack of studies investigating the expression of comet tails in adolescent or childhood obesity.

More recently, a nutritional intervention study outlined a decrease in DNA single strand breaks with weight reduction in 50 participants, even though the obese/overweight participants did not have high levels of DNA damage at baseline [141]. Intriguingly, DNA damage was associated with increased BMI and breast cancer risk in a case-control study [142]. However, in the cancer-free control cohort, BMI was inversely associated with DNA damage [142]. It has been suggested that this discrepancy could be an outcome of increased metabolic rate in lean individuals and a possible generation of more ROS. In the breast cancer cohort, the positive association between BMI and DNA damage could be a consequence of defective DNA repair and marked chromosomal instabilities that are consistently found in cancer. Furthermore, this study analysed BMI data for only 45 cancer free patients, indicating that these findings may lack statistical power. Nonetheless, the role of the comet assay as a biomarker for genomic instability in increasing adiposity is propitious.

6.5 Telomere Length

Telomeres play a fundamental role in maintaining chromosomal integrity and can substantially underpin the genomic instabilities associated with carcinogenesis [143]. Telomeres are vital segments of non-coding DNA repeat sequences (TTAGGG) that are located at the ends of chromosomes and lose their DNA bases at every round of the cell cycle, consequently shortening in length. Telomere length can be assessed in multiple tissue types via a variety of techniques that include quantitative Polymerase Chain Reaction (qPCR), fluorescence probes, Southern blot and the

gold standard - Terminal Restriction Fragmentation (TRF). Telomere shortening to a critical length should ultimately result in replicative senescence, hence the reason for telomeres long being linked to the phenomenon of cellular ageing [144,145]

Telomere shortening has been discussed as a biomarker for detecting the onset of multiple age related disorders including cardiovascular disease and diabetes [146], Alzheimer's [147], Parkinson's [148] and arthritis [149]. Telomere length has also been assessed in disease models of accelerated ageing including Werner's syndrome [150] and Hutchinson-Gilford Progeria (HGPS) [151]. It is of interest that despite telomere shortening being a phenomenon common to both disorders, patients with Werner's syndrome have an increased susceptibility to acquiring cancer, whereas HGPS is associated with a resistance to oncogenic transformations [152]. To conclude, telomere shortening has become a well-known, undisputed biomarker for ageing, but its role as a biomarker for detecting early pathological changes in age-related disorders such as cancer, requires clarification.

Case-control studies in cancer patients have strengthened the associations between telomere attrition, genomic instability and possible increased risk of cancer (Table 9.). Interestingly, this phenomenon may be identified systemically in cancer patients, as Broberg and colleagues reported significantly shorter telomeres in buccal cells harvested from patients with bladder cancer, when compared to cancer free controls [153]. A recent comprehensive review of over 23,000 cases has brought to light that the significance of telomere length as a marker for cancer is more complex [154]. This research presented a heterogeneous relationship across different cancer types, including a statistically significant correlation between decreased telomere length and increased risk of GI tract, head and neck cancers, but not skin cancers. Contrastingly, it was also reported longer telomere length was more likely to increase the risk of skin cancer, indicating that telomeres may play a more diverse, tissue-specific role [154]. Similarly, other studies have noted a non-linear relationship for cancers of the breast [155], pancreas [156] and oesophagus [157]. In the case of lung cancer, both telomere shortening [158] and lengthening [159] have been described as risk factors. The findings

from these studies indicate that telomere length is an important marker of genomic instability where excessive shortening or lengthening may both be drivers or indicators of cancer development.

Telomere attrition may promote generation of cancer stem cells by causing a series of chromosomal fusions, anaphase bridge formations and breakage-fusion-breakage cycles that enable DNA damage to accumulate [143]. Whereas lengthened telomeres may enable the cell to enter into more division cycles, thus increasing the susceptibility of the genome to abnormalities, lethal mutations as well as immortality. Increased telomere dysfunction has also been identified alongside chromothripsis in patients with increased susceptibility to developing cancer (sufferers of L-Fraumeni syndrome) [160] suggesting an additional role for telomere attrition in increased catastrophic DNA re-arrangements. However, there is a greater need for more case-control studies to confirm these speculations and to test the applicability of telomere length as a biomarker for tissue-specific cancer risk.

Initial reports of telomere shortening in obesity were presented in a study conducted on 561 female twins and linked increased adiposity with a risk of up to 8.8 years of ageing [161]. Interestingly, a subsequent study published by Nordfjäll and colleagues confirmed these findings with a variety of obesity parameters including BMI, weight, and waist and hip circumference [162]. Although, the study reported a gender bias as these findings were only statistically significant in women. In 2009, an extensive study by Kim and collaborators also found an inverse correlation between weight gain and telomere length in 647 women [163]. Shorter telomere length has also been reported in cells of subcutaneous adipose tissue extracted from obese men and women [164]. A number of subsequent studies have built evidence to support the hypothesis that increased body fat in adults is an independent causative factor for accelerated ageing (Table 8). However, discrepancies in the evidence do exist, as some studies have found little or no association between multiple measures of adiposity and telomere dysfunction in women [165]. This highlights that telomere maintenance is a highly regulated and complex process affected by other physiological factors such as menstrual status, as well as environmental factors.

A study that established a link between obesity and telomere shortening in adults, was unable to verify this phenomenon in obese children [166]. However, subsequent studies notoriously reported findings of telomere shortening in obese children. Al-Attas and collaborators presented an inverse correlation between obesity and telomere length in boys, but not girls [167]. A subsequent, large case-control study (n=793) identified a 25% reduction in telomere length in obese children when compared to non-obese healthy controls [168]. Interestingly, Zhu and colleagues did not find an association between adiposity and telomere length in 667 adolescents aged between 14 and 18 years [169]. An extensive meta-analysis by Park and colleagues raised an interesting association between obesity in childhood and increased risk of morbidities and mortality in adulthood, independent of weight loss [170]. The associations between increased adiposity in childhood and telomere shortening are compelling, and further evidence is required to conclude upon the degree and consequences of telomere dysfunction in such obesity.

It is of interest that increased markers of inflammation have been exhibited alongside telomere attrition in obese cancer patients [171]. A high BMI in individuals with obesity-associated polymorphisms correlated with increased levels of CRP and decreased telomere length [172].

Telomere shortening was also seen in conjunction with increased CRP and adiposity in a more recent investigation [173]. This research showed BMI, waist circumference, and body fat percentage to be inversely correlated with telomere length in adults. These findings strengthen the hypothesis for chronic, low-grade inflammation as a possible etiological route to DNA damage in obese subjects.

7. Genetic Determinants of Obesity and DNA damage

Genome wide association studies (GWAS) have indicated a number of single nucleotide polymorphisms (SNPs) to be associated with obesity. Of these, particular polymorphisms (such as rs9939609) in the fat mass and obesity-associated (FTO) gene, have been confirmed to increase the

risk of non-syndromic obesity [174]. Such genetic polymorphisms are likely to alter the activity of metabolic hormones, leading to reduced feelings of satiety and an increased caloric intake [175]. It has been investigated whether such genetic determinants can be responsible for the increasing evidence of DNA damage in obesity. A study evaluated the frequency of polymorphisms in genes coding for ghrelin (rs26802), ghrelin receptor (rs572169), leptin (rs7799039), leptin receptor (rs1137101) and FTO (rs9939609) in 300 morbidly obese Caucasian women alongside 300 controls [176]. These individuals were also investigated for oxidative DNA damage via the comet assay. Obesity was associated with a significantly increased frequency of the AA-FTO genotype allele A, decreased frequency of the TT-FTO genotype, and elevated DNA strand breaks. However, the degree of DNA damage was independent of genotype, indicating that these particular polymorphisms may not play an etiological role in the development of DNA damage in obese subjects, and suggests the damage may be caused by other pathological states such as low grade inflammation and oxidative stress. However, variants of the FTO have also been associated with an increased risk of breast cancer in African American [177] but not Caucasian women [178]. This suggests SNP associations and their consequences in obesity display strong ethnic variations.

A population-based study discovered variants in the DNA repair genes *MGMT* (rs12917), *MSH2* (rs3732182 and rs4583514), *MSH3* (rs1650663), *XPG* (rs17655), and *XRCC1* (rs25487) that code for BER, NER and MMR pathways to elevate the risk of postmenopausal breast cancer in obese women [179]. Simultaneously, the research found obese women to be at increased risk of oxidative stress if they possess a polymorphism in the *MPO* gene. Similar findings were outlined in a meta-analysis that concluded polymorphisms of *GPX1* and *GPX7* genes that are associated with higher body fat and insulin resistance, contribute to oxidative stress in obesity, by lowering anti-oxidant defence [180]. Therefore, there is a possibility that certain genetic variants that predispose to obesity also promote pathogenesis of oxidative stress and DNA damage. If this pathological process is coupled with a lack of sufficient DNA repair, then it may increase the risk of certain cancers.

Since genetic determinants of TL in adults do not correspond with TL in children, and genetic determinants of TL remain unconfirmed [181], there is a likelihood that telomere length shortening in children may be associated with obesity related ROS rather than genetic make-up. However, there is a greater need for further studies to identify the relationship between telomere length in obesity and the long-term risk of pathological conditions.

8. Weight loss interventions and DNA Damage

Decreasing adiposity to a certain degree may improve oxidative stress and DNA health in obesity.

Bariatric surgery is an established method of weight-loss which has shown positive implications on obesity associated complications, including cardiovascular disease [182], obstructive sleep apnoea [183], non-alcoholic fatty liver disease [184] and a rapid improvement in type 2 diabetes [185].

Bariatric surgery has also been suggested to reduce the risk of cancer by restoring genome stability and integrity [186–188]. Laparoscopic gastric band application has shown a reversal in oxidative DNA damage, as reduced levels of urinary 8-OHdG were identified 6 months after this surgical procedure in a group of morbidly obese patients [189]. The same surgical procedure has demonstrated a reduction in DNA double strand breaks with decreased levels of γ -H2AX positive cells up to 7 years after surgery [190] and restoration of telomere length in obese men [191].

However, a study conducted on 107 obese subjects with and without metabolic syndrome reported no extension in telomere length after a 12 month follow up of bariatric surgery [192]. Although more recently, Bankoglu and colleagues reported a reduction in DNA breaks assessed by the comet assay in 56 blood samples 12 months after surgery [193]. Therefore, bariatric surgery appears to be propitious for improving DNA integrity and stability in obese adults. Whether such is the case in severe adolescent obesity could be explored.

Dietary and lifestyle modifications have also been associated with a reduction of DNA damage in obesity. Following a two month intervention programme, governed by improvements in diet and physical activity levels, increased telomere length in 74 obese adolescents [194]. Similarly, dietary

modifications and consequent weight loss also reduced DNA damage in obese women with PCOS [195]. From these studies it can be deduced that DNA damage in obesity may be a reversible phenomenon. Although, it is also possible that gene mutations that have been induced and accumulated during the phase of obesity may persist beyond weight-loss.

9. Conclusion

Oxidative stress in obesity poses a significant threat to DNA stability and integrity as indicated by the growing number of investigations reporting a positive correlation between markers of oxidative DNA damage and increased adiposity. These associations raise concerns that the obese state may manifest with accelerated genome ageing and promote carcinogenesis. Monitoring of acquired DNA damage may become an important part of clinical investigations in the management of obesity, alongside markers of inflammation and micro nutritional deficiencies in order to detect pre-pathological changes. Potential benefits of monitoring 'genome health' in obesity include early detection of pre-cancerous alterations to establish urgency of personalised intervention measures and assess their progress. These avenues should be explored as DNA damage in obesity appears to be a reversible phenomenon.

Declaration of Interest: none

Acknowledgments

This work was supported by the University of Westminster. The Authors are grateful for funding received by the University through the FST StartUp Scheme (E.V.) and FST PhD Scholarship (M.U.).

References

- [1] Renehan AG, Tyson M, Egger M, Heller RF, Zwahlen M. Body-mass index and incidence of cancer: a systematic review and meta-analysis of prospective observational studies. *Lancet*. 2008;371(9612):569–78.
- [2] Arnold M, Leitzmann M, Freisling H, Bray F, Romieu I, Renehan A, et al. Obesity and cancer: An update of the global impact. *Cancer Epidemiol*. 2016 Apr;41:8–15.
- [3] Kyrgiou M, Kalliala I, Markozannes G, Gunter MJ, Paraskevaidis E, Gabra H, et al. Adiposity and cancer at major anatomical sites: umbrella review of the literature. *BMJ [Internet]*. 2017 Feb 28;356. Available from: <http://www.bmj.com/content/356/bmj.j477.abstract>
- [4] Mazzeo L. Why does obesity promote cancer? *Epidemiology, biology, and open questions. Ecancermedicallscience [Internet]*. 2015 Jul 23;9:554. Available from: <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC4531127/>
- [5] Fernández-Sánchez A, Madrigal-Santillán E, Bautista M, Esquivel-Soto J, Morales-González Á, Esquivel-Chirino C, et al. Inflammation, oxidative stress, and obesity. *Int J Mol Sci*. 2011;12(5):3117–32.
- [6] Hanahan D, Weinberg RA. Hallmarks of Cancer: The Next Generation. *Cell [Internet]*. 2016 Jun 8;144(5):646–74. Available from: <http://dx.doi.org/10.1016/j.cell.2011.02.013>
- [7] Wiseman H, Halliwell B. Damage to DNA by reactive oxygen and nitrogen species: role in inflammatory disease and progression to cancer. *Biochem J*. 1996;313(Pt 1):17.
- [8] Xu B, Wang W, Guo H, Sun Z, Wei Z, Zhang X, et al. Oxidative stress preferentially induces a subtype of micronuclei and mediates the genomic instability caused by p53 dysfunction. *Mutat Res Mol Mech Mutagen*. 2014;770:1–8.
- [9] Finkel T. Signal transduction by reactive oxygen species. *J Cell Biol*. 2011 Jul;194(1):7 LP-15.
- [10] Codoñer-Franch P, Valls-Bellés V, Arilla-Codoñer A, Alonso-Iglesias E. Oxidant mechanisms in childhood obesity: the link between inflammation and oxidative stress. *Transl Res [Internet]*. 2011 Dec [cited 2015 Feb 26];158(6):369–84. Available from: <http://www.sciencedirect.com/science/article/pii/S1931524411002647>
- [11] Ohshima H, Tatemichi M, Sawa T. Chemical basis of inflammation-induced carcinogenesis. *Arch Biochem Biophys [Internet]*. 2003 Sep 1;417(1):3–11. Available from: <http://www.sciencedirect.com/science/article/pii/S0003986103002832>
- [12] Chielle EO, de Souza WM, da Silva TP, Moresco RN, Moretto MB. Adipocytokines, inflammatory and oxidative stress markers of clinical relevance altered in young overweight/obese subjects. *Clin Biochem*. 2016;49(7):548–53.
- [13] Ellulu MS, Khaza'ai H, Rahmat A, Patimah I, Abed Y. Obesity can predict and promote systemic inflammation in healthy adults. *Int J Cardiol*. 2016;215:318–24.
- [14] Park HS, Park JY, Yu R. Relationship of obesity and visceral adiposity with serum concentrations of CRP, TNF- α and IL-6. *Diabetes Res Clin Pract*. 2005;69(1):29–35.
- [15] Schipper HS, Nuboer R, Prop S, Van Den Ham HJ, De Boer FK, Kesmir C, et al. Systemic inflammation in childhood obesity: circulating inflammatory mediators and activated CD14⁺⁺ monocytes. *Diabetologia*. 2012;55(10):2800–10.
- [16] Lee B-C, Lee J. Cellular and molecular players in adipose tissue inflammation in the development of obesity-induced insulin resistance. *Biochim Biophys Acta [Internet]*. 2014 Mar [cited 2015 Jan 19];1842(3):446–62. Available from: <http://www.sciencedirect.com/science/article/pii/S0925443913001798>
- [17] Mittal M, Siddiqui MR, Tran K, Reddy SP, Malik AB. Reactive oxygen species in inflammation and tissue injury. *Antioxid Redox Signal*. 2014;20(7):1126–67.
- [18] Kern PA, Di Gregorio GB, Lu T, Rassouli N, Ranganathan G. Adiponectin Expression From Human Adipose Tissue. *Diabetes [Internet]*. 2003 Jul 1;52(7):1779–85. Available from: <http://diabetes.diabetesjournals.org/content/52/7/1779.abstract>
- [19] Ouchi N, Kihara S, Arita Y, Maeda K, Kuriyama H, Okamoto Y, et al. Novel modulator for endothelial adhesion molecules adipocyte-derived plasma protein adiponectin. *Circulation*. 1999;100(25):2473–6.
- [20] Mantzoros C, Petridou E, Dessypris N, Chavelas C, Dalamaga M, Alexe DM, et al. Adiponectin and breast cancer risk. *J Clin Endocrinol Metab*. 2004;89(3):1102–7.

- [21] Ishikawa M, Kitayama J, Kazama S, Hiramatsu T, Hatano K, Nagawa H. Plasma adiponectin and gastric cancer. *Clin Cancer Res*. 2005;11(2):466–72.
- [22] Wei EK, Giovannucci E, Fuchs CS, Willett WC, Mantzoros CS. Low plasma adiponectin levels and risk of colorectal cancer in men: a prospective study. *J Natl Cancer Inst*. 2005;97(22):1688–94.
- [23] Petridou E, Mantzoros CS, Dessypris N, Dikaloti SK, Trichopoulos D. Adiponectin in relation to childhood myeloblastic leukaemia. *Br J Cancer*. 2006;94(1):156–60.
- [24] Goktas S, Yilmaz MI, Caglar K, Sonmez A, Kilic S, Bedir S. Prostate cancer and adiponectin. *Urology*. 2005;65(6):1168–72.
- [25] Haghiaç M, Vora NL, Basu S, Johnson KL, Presley L, Bianchi DW, et al. Increased Death of Adipose Cells, a Path to Release Cell- Free DNA Into Systemic Circulation of Obese Women. *Obesity*. 2012;20(11):2213–9.
- [26] Nishimoto S, Fukuda D, Higashikuni Y, Tanaka K, Hirata Y, Murata C, et al. Obesity-induced DNA released from adipocytes stimulates chronic adipose tissue inflammation and insulin resistance. *Sci Adv*. 2016;2(3):e1501332.
- [27] Weisberg SP, McCann D, Desai M, Rosenbaum M, Leibel RL, Ferrante AW. Obesity is associated with macrophage accumulation in adipose tissue. *J Clin Invest* [Internet]. 2003 Dec 15;112(12):1796–808. Available from: <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC296995/>
- [28] Tamir S, Burney S, Tannenbaum SR. DNA damage by nitric oxide. *Chem Res Toxicol* [Internet]. 1996;9(5):821–7. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/8828916>
- [29] Baik S-C, Youn H-S, Chung M-H, Lee W-K, Cho M-J, Ko G-H, et al. Increased Oxidative DNA Damage in Helicobacter pylori-infected Human Gastric Mucosa. *Cancer Res* [Internet]. 1996 Mar 1;56(6):1279–82. Available from: <http://cancerres.aacrjournals.org/content/56/6/1279.abstract>
- [30] Bernstein CN, Blanchard JF, Kliewer E, Wajda A. Cancer risk in patients with inflammatory bowel disease. *Cancer* [Internet]. 2001 Feb 15;91(4):854–62. Available from: [http://dx.doi.org/10.1002/1097-0142\(20010215\)91:4%3C854::AID-CNCR1073%3E3.0.CO](http://dx.doi.org/10.1002/1097-0142(20010215)91:4%3C854::AID-CNCR1073%3E3.0.CO)
- [31] Shawki SM, Meshaal SS, El Dash AS, Zayed NA, Hanna MOF. Increased DNA damage in hepatitis C virus-related hepatocellular carcinoma. *DNA Cell Biol* [Internet]. 2014 Dec [cited 2015 Feb 25];33(12):884–90. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/25211328>
- [32] Tezal M, Grossi SG, Genco RJ. Is Periodontitis Associated With Oral Neoplasms? *J Periodontol* [Internet]. 2005 Mar 1;76(3):406–10. Available from: <http://dx.doi.org/10.1902/jop.2005.76.3.406>
- [33] Izano M, Wei EK, Tai C, Swede H, Gregorich S, Harris TB, et al. Chronic inflammation and risk of colorectal and other obesity- related cancers: The health, aging and body composition study. *Int J Cancer*. 2016;138(5):1118–28.
- [34] Boden G. Obesity and free fatty acids. *Endocrinol Metab Clin North Am*. 2008;37(3):635–46.
- [35] Vincent HK, Taylor AG. Biomarkers and potential mechanisms of obesity-induced oxidant stress in humans. *Int J Obes*. 2006;30(3):400–18.
- [36] Olusi SO. Obesity is an independent risk factor for plasma lipid peroxidation and depletion of erythrocyte cytoprotective enzymes in humans. *Int J Obes Relat Metab Disord*. 2002;26(9).
- [37] Niedernhofer LJ, Daniels JS, Rouzer CA, Greene RE, Marnett LJ. Malondialdehyde, a product of lipid peroxidation, is mutagenic in human cells. *J Biol Chem*. 2003;278(33):31426–33.
- [38] VanderVeen LA, Hashim MF, Shyr Y, Marnett LJ. Induction of frameshift and base pair substitution mutations by the major DNA adduct of the endogenous carcinogen malondialdehyde. *Proc Natl Acad Sci*. 2003;100(24):14247–52.
- [39] Del Rio D, Stewart AJ, Pellegrini N. A review of recent studies on malondialdehyde as toxic molecule and biological marker of oxidative stress. *Nutr Metab Cardiovasc Dis*. 2005;15(4):316–28.
- [40] Chen S, Sun L, Gao H, Ren L, Liu N, Song G. Visfatin and oxidative stress influence endothelial progenitor cells in obese populations. *Endocr Res*. 2015;40(2):83–7.
- [41] Ustundag B, Gungor S, Aygün AD, Turgut M, Yilmaz E. Oxidative status and serum leptin levels in obese prepubertal children. *Cell Biochem Funct* [Internet]. 2007 Sep 1;25(5):479–83. Available from: <http://dx.doi.org/10.1002/cbf.1334>
- [42] Gönenç A, Özkan Y, Torun M, Şimşek B. Plasma malondialdehyde (MDA) levels in breast and lung cancer patients. *J Clin Pharm Ther*. 2001;26(2):141–4.
- [43] Friedman JE, Dohm GL, Leggett-Frazier N, Elton CW, Tapscott EB, Pories WP, et al. Restoration of insulin responsiveness in skeletal muscle of morbidly obese patients after weight loss. Effect on muscle glucose transport and glucose transporter GLUT4. *J Clin Invest*. 1992;89(2):701.

- [44] Shah P, Vella A, Basu A, Basu R, Adkins A, Schwenk WF, et al. Elevated Free Fatty Acids Impair Glucose Metabolism in Women Decreased Stimulation of Muscle Glucose Uptake and Suppression of Splanchnic Glucose Production During Combined Hyperinsulinemia and Hyperglycemia. *Diabetes*. 2003;52(1):38–42.
- [45] Lin Y, Berg AH, Iyengar P, Lam TKT, Giacca A, Combs TP, et al. The hyperglycemia-induced inflammatory response in adipocytes the role of reactive oxygen species. *J Biol Chem*. 2005;280(6):4617–26.
- [46] Renehan AG, Frystyk J, Flyvbjerg A. Obesity and cancer risk: the role of the insulin-IGF axis. *Trends Endocrinol Metab*. 2006;17(8):328–36.
- [47] Li BDL, Khosravi MJ, Berkel HJ, Diamandi A, Dayton MA, Smith M, et al. Free insulin-like growth factor- I and breast cancer risk. *Int J cancer*. 2001;91(5):736–9.
- [48] Renehan AG, O’Connell J, O’Halloran D, Shanahan F, Potten CS, O’Dwyer ST, et al. Acromegaly and colorectal cancer: a comprehensive review of epidemiology, biological mechanisms, and clinical implications. *Horm Metab Res*. 2003;35(11/12):712–25.
- [49] Yang S, Chintapalli J, Sodagum L, Baskin S, Malhotra A, Reiss K, et al. Activated IGF-1R inhibits hyperglycemia-induced DNA damage and promotes DNA repair by homologous recombination. *Am J Physiol Physiol*. 2005;289(5):F1144–52.
- [50] Frystyk J, Vestbo E, Skjaerbaek C, Mogensen CE, Ørskov H. Free insulin-like growth factors in human obesity. *Metabolism*. 1995;44:37–44.
- [51] Kaaks R, Lukanova A, Sommersberg B. Plasma androgens, IGF-1, body size, and prostate cancer risk: a synthetic review. *Prostate Cancer Prostatic Dis*. 2000;3(3):157–72.
- [52] Selby C. Sex hormone binding globulin: origin, function and clinical significance. *Ann Clin Biochem An Int J Biochem Med*. 1990;27(6):532–41.
- [53] Clemons M, Goss P. Estrogen and the risk of breast cancer. *N Engl J Med*. 2001;344(4):276–85.
- [54] Siiteri PK, Nisker JA, Hammond GL. Hormonal basis of risk factors for breast and endometrial cancer. *Hormones and cancer*. Raven Press, NY, USA; 1980. p. 499–505.
- [55] Caldon CE, Sutherland RL, Musgrove EA. Cell cycle proteins in epithelial cell differentiation: Implications for breast cancer. *Cell Cycle*. 2010 May;9(10):1918–28.
- [56] Roy D, Liehr JG. Estrogen, DNA damage and mutations. *Mutat Res Mol Mech Mutagen*. 1999;424(1):107–15.
- [57] Ozata M, Mergen M, Oktenli C, Aydin A, Sanisoglu SY, Bolu E, et al. Increased oxidative stress and hypozincemia in male obesity. *Clin Biochem*. 2002;35(8):627–31.
- [58] Amirkhizi F, Siassi F, Minaie S, Djalali M, Rahimi A, Chamari M. Is obesity associated with increased plasma lipid peroxidation and oxidative stress in women? *ARYA Atheroscler*. 2010;2(4).
- [59] Erdeve O, Siklar Z, Kocaturk PA, Dallar Y, Kavas GO. Antioxidant superoxide dismutase activity in obese children. *Biol Trace Elem Res*. 2004;98(3):219–27.
- [60] Kilic E, Özer ÖF, Ereğ AT, Erman H, Torun E, Ayhan SK, et al. Oxidative Stress Status in Childhood Obesity: A Potential Risk Predictor. *Med Sci Monit*. 2016 Oct;22:3673–9.
- [61] Albuali WH. Evaluation of oxidant-antioxidant status in overweight and morbidly obese Saudi children. *World J Clin Pediatr [Internet]*. 2014 Feb 8;3(1):6–13. Available from: <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC4145643/>
- [62] Sfar S, Boussoffara R, Sfar MT, Kerkeni A. Antioxidant enzymes activities in obese Tunisian children. *Nutr J*. 2013 Jan;12:18.
- [63] Zhu YOUEN, Zhang SHUEI, Wang JIUE, Xiao WEII. Overweight and Obesity — Induced Oxidative Stress in Children Obesity is one of the most common nutritional. 2006;359:353–9.
- [64] Sentürker S, Karahalil B, Inal M, Yilmaz H, Gedikoglu G, Dizdaroglu M. Oxidative DNA base damage and antioxidant enzyme levels in childhood acute lymphoblastic leukemia. *FEBS Lett*. 1997;416(3):286–90.
- [65] Elchuri S, Oberley TD, Qi W, Eisenstein RS, Roberts LJ, Van Remmen H, et al. CuZnSOD deficiency leads to persistent and widespread oxidative damage and hepatocarcinogenesis later in life. *Oncogene*. 2005;24(3):367–80.
- [66] Asaduzzaman Khan M, Tania M, Zhang D, Chen H. Antioxidant enzymes and cancer. *Chinese J Cancer Res [Internet]*. 2010;22(2):87–92. Available from: <http://dx.doi.org/10.1007/s11670-010-0087-7>
- [67] Srivastava KC, Austin RD, Shrivastava D. Evaluation of oxidant-antioxidant status in tissue samples in

- oral cancer: A case control study. *Dent Res J (Isfahan)* [Internet]. 2016 Apr;13(2):181–7. Available from: <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC4810917/>
- [68] Hecht F, Pessoa CF, Gentile LB, Rosenthal D, Carvalho DP, Fortunato RS. The role of oxidative stress on breast cancer development and therapy. *Tumor Biol* [Internet]. 2016;37(4):4281–91. Available from: <http://dx.doi.org/10.1007/s13277-016-4873-9>
- [69] Via M. The Malnutrition of Obesity: Micronutrient Deficiencies That Promote Diabetes. *ISRN Endocrinol* [Internet]. 2012 Mar 15;2012:103472. Available from: <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC3313629/>
- [70] Ames BN. DNA damage from micronutrient deficiencies is likely to be a major cause of cancer. *Mutat Res*. 2001 Apr;475(1–2):7–20.
- [71] Vimaleswaran KS, Berry DJ, Lu C, Tikkanen E, Pilz S, Hiraki LT, et al. Causal relationship between obesity and vitamin D status: bi-directional Mendelian randomization analysis of multiple cohorts. *PLoS Med*. 2013;10(2):e1001383.
- [72] Bao B, Ting H, Hsu J, Lee Y. Protective role of 1 α , 25- dihydroxyvitamin D3 against oxidative stress in nonmalignant human prostate epithelial cells. *Int J cancer*. 2008;122(12):2699–706.
- [73] Hanchette CL, Schwartz GG. Geographic patterns of prostate cancer mortality. Evidence for a protective effect of ultraviolet radiation. *Cancer*. 1992;70(12):2861–9.
- [74] Crasta K, Ganem NJ, Dagher R, Lantermann AB, Ivanova E V, Pan Y, et al. DNA breaks and chromosome pulverization from errors in mitosis. *Nature* [Internet]. 2012 Feb 2;482(7383):53–8. Available from: <http://dx.doi.org/10.1038/nature10802>
- [75] Hatch EM, Fischer AH, Deerinck TJ, Hetzer MW. Catastrophic Nuclear Envelope Collapse in Cancer Cell Micronuclei. *Cell* [Internet]. 2013;154(1):47–60. Available from: <http://dx.doi.org/10.1016/j.cell.2013.06.007>
- [76] Mackenzie KJ, Carroll P, Martin C-A, Murina O, Fluteau A, Simpson DJ, et al. cGAS surveillance of micronuclei links genome instability to innate immunity. *Nature*. 2017 Aug;548(7668):461–5.
- [77] Bonassi S, Znaor A, Ceppi M, Lando C, Chang WP, Holland N, et al. An increased micronucleus frequency in peripheral blood lymphocytes predicts the risk of cancer in humans. *Carcinog* [Internet]. 2006 Sep 14;28(3):625–31. Available from: <http://carcin.oxfordjournals.org/content/28/3/625.abstract>
- [78] Murgia E, Ballardini M, Bonassi S, Rossi AM, Barale R. Validation of micronuclei frequency in peripheral blood lymphocytes as early cancer risk biomarker in a nested case-control study. *Mutat Res - Fundam Mol Mech Mutagen*. 2008;639(1–2):27–34.
- [79] Bolognesi C, Lando C, Forni A, Landini E, Scarpato R, Migliore L, et al. Chromosomal damage and ageing: effect on micronuclei frequency in peripheral blood lymphocytes. *Age Ageing* [Internet]. 1999 Jul 1;28(4):393–7. Available from: <http://ageing.oxfordjournals.org/content/28/4/393.abstract>
- [80] Milosevic-Djordjevic O, Grujicic D, Novakovic T, Arsenijevic S, Marinkovic D. Micronuclei and Ageing in a Sample of Yugoslavian Population. *Russ J Genet* [Internet]. 2002;38(2):201–4. Available from: <http://dx.doi.org/10.1023/A:1014342312959>
- [81] Orta T, Günebakan S. The effect of aging on micronuclei frequency and proliferation in human peripheral blood lymphocytes. *Indian J Hum Genet* [Internet]. 2012;18(1):95–100. Available from: <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC3385189/>
- [82] Ferraz GA, Costa Neto A de O, Cerqueira E de MM, Meireles JRC. Effects of age on the frequency of micronuclei and degenerative nuclear abnormalities. *Rev Bras Geriatr e Gerontol* [Internet]. 2016;19(4):627–34. Available from: http://www.scielo.br/scielo.php?script=sci_arttext&pid=S1809-98232016000400627&lng=en&tlng=en
- [83] Sanchez-Flores M, Marcos-Perez D, Lorenzo-Lopez L, Maseda A, Millan-Calenti JC, Bonassi S, et al. Frailty Syndrome and Genomic Instability in Older Adults: Suitability of the Cytome Micronucleus Assay As a Diagnostic Tool. *J Gerontol A Biol Sci Med Sci*. 2018 Jun;73(7):864–72.
- [84] Yesilada E, Sahin I, Ozcan H, Yildirim IH, Yologlu S, Taskapan C. Increased micronucleus frequencies in peripheral blood lymphocytes in women with polycystic ovary syndrome. *Eur J Endocrinol* [Internet]. 2006;154(4):563–8. Available from: <http://www.eje-online.org/content/154/4/563> <http://www.eje-online.org/content/154/4/563.long> <http://www.eje-online.org/content/154/4/563.full.pdf>
- [85] Andreassi MG, Barale R, Iozzo P, Picano E. The association of micronucleus frequency with obesity, diabetes and cardiovascular disease. *Mutagenesis* [Internet]. 2011 Jan [cited 2015 Feb 6];26(1):77–83. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/21164186>

- [86] Donmez-Altuntas H, Sahin F, Bayram F, Bitgen N, Mert M, Guclu K, et al. Evaluation of chromosomal damage, cytostasis, cytotoxicity, oxidative DNA damage and their association with body-mass index in obese subjects. *Mutat Res Toxicol Environ Mutagen* [Internet]. 2014 Sep 1;771:30–6. Available from: <http://www.sciencedirect.com/science/article/pii/S138357181400179X>
- [87] Scarpato R, Verola C, Fabiani B, Bianchi V, Saggese G, Federico G. Nuclear damage in peripheral lymphocytes of obese and overweight Italian children as evaluated by the γ -H2AX focus assay and micronucleus test. *FASEB J* [Internet]. 2011 Feb 1;25(2):685–93. Available from: <http://www.fasebj.org/content/25/2/685.abstract>
- [88] Dass Singh M, Thomas P, Hor M, Almond T, Owens J, Hague W, et al. Infant birth outcomes are associated with DNA damage biomarkers as measured by the cytokinesis block micronucleus cytome assay: the DADHI study. *Mutagenesis* [Internet]. 2017 May 1;32(3):355–70. Available from: <http://dx.doi.org/10.1093/mutage/gex001>
- [89] Cho NY, Kim KW, Kim KK. Genomic health status assessed by a cytokinesis-block micronucleus cytome assay in a healthy middle-aged Korean population. *Mutat Res*. 2017 Feb;814:7–13.
- [90] Croco E, Marchionni S, Lorenzini A. Genetic instability and aging under the scrutiny of comparative biology: a meta-analysis of spontaneous micronuclei frequency. *Mech Ageing Dev* [Internet]. 2016 Jun;156:34–41. Available from: <http://www.sciencedirect.com/science/article/pii/S004763741630046X>
- [91] Wickliffe JK, Dertinger SD, Torous DK, Avlasevich SL, Simon- Friedt BR, Wilson MJ. Diet-induced obesity increases the frequency of Pig- a mutant erythrocytes in male C57BL/6J mice. *Environ Mol Mutagen*. 2016;57(9):668–77.
- [92] Almeida T, Leitão RMC, Carrilho FJ, Sonohara S. Micronuclei formation in liver fibrosis samples from patients infected by hepatitis C virus. *Genet Mol Biol*. 2010;33(3):418–21.
- [93] Suzuki T, Kamiya H. Mutations induced by 8-hydroxyguanine (8-oxo-7, 8-dihydroguanine), a representative oxidized base, in mammalian cells. *Genes Environ*. 2017;39(1):2.
- [94] Ramon O, Sauvaigo S, Gasparutto D, Faure P, Favier A, Cadet J. Effects of 8-oxo-7, 8-dihydro-2'-deoxyguanosine on the binding of the transcription factor Sp1 to its cognate target DNA sequence (GC box). *Free Radic Res*. 1999;31(3):217–29.
- [95] Black AR, Black JD, Azizkhan-Clifford J. Sp1 and krüppel-like factor family of transcription factors in cell growth regulation and cancer. *J Cell Physiol* [Internet]. 2001 Aug 1;188(2):143–60. Available from: <http://dx.doi.org/10.1002/jcp.1111>
- [96] Ba X, Boldogh I. 8-Oxoguanine DNA glycosylase 1: Beyond repair of the oxidatively modified base lesions. *Redox Biol*. 2018 Apr;14:669–78.
- [97] Cooke MS, Olinski R, Evans MD. Does measurement of oxidative damage to DNA have clinical significance? *Clin Chim Acta* [Internet]. 2006 Mar;365(1–2):30–49. Available from: <http://www.sciencedirect.com/science/article/pii/S0009898105005577>
- [98] Burns MB, Temiz NA, Harris RS. Evidence for APOBEC3B mutagenesis in multiple human cancers. *Nat Genet* [Internet]. 2013 Sep 14;45(9):977–83. Available from: <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC3902892/>
- [99] Kaur J, Politis C, Jacobs R. Salivary 8-hydroxy-2'-deoxyguanosine, malondialdehyde, vitamin C, and vitamin E in oral pre-cancer and cancer: diagnostic value and free radical mechanism of action. *Clin Oral Investig*. 2016 Mar;20(2):315–9.
- [100] Romano G, Sgambato A, Mancini R, Capelli G, Giovagnoli MR, Flamini G, et al. 8-Hydroxy-2'-deoxyguanosine in cervical cells: correlation with grade of dysplasia and human papillomavirus infection. *Carcinogenesis* [Internet]. 2000 Jun 1;21(6):1143–7. Available from: <http://dx.doi.org/10.1093/carcin/21.5.143>
- [101] Oberdoerffer P, Sinclair D a. The role of nuclear architecture in genomic instability and ageing. *Nat Rev Mol Cell Biol*. 2007;8(9):692–702.
- [102] Chen Q, Fischer A, Reagan JD, Yan LJ, Ames BN. Oxidative DNA damage and senescence of human diploid fibroblast cells. *Proc Natl Acad Sci U S A* [Internet]. 1995;92(10):4337–41. Available from: <http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=41939&tool=pmcentrez&rendertype=abstract>
- [103] Xu B, Kang L, Zhang G, Wu J, Zhu R, Yang M, et al. The Changes of 8-OHdG, hOGG1, APE1 and Pol β in Lenses of Patients with Age-Related Cataract. *Curr Eye Res*. 2015;40(4):378–85.
- [104] Rubio-Perez JM, Albaladejo MD, Zafrilla P, Vidal-Guevara ML, Morillas-Ruiz JM. Effects of an

- antioxidant beverage on biomarkers of oxidative stress in Alzheimer's patients. *Eur J Nutr* [Internet]. 2015;1–12. Available from: <http://dx.doi.org/10.1007/s00394-015-1024-9>
- [105] Kikuchi A, Takeda A, Onodera H, Kimpara T, Hisanaga K, Sato N, et al. Systemic Increase of Oxidative Nucleic Acid Damage in Parkinson's Disease and Multiple System Atrophy. *Neurobiol Dis* [Internet]. 2002 Mar;9(2):244–8. Available from: <http://www.sciencedirect.com/science/article/pii/S0969996102904663>
- [106] Nishikawa T, Sasahara T, Kiritoshi S, Sonoda K, Senokuchi T, Matsuo T, et al. Evaluation of urinary 8-hydroxydeoxy-guanosine as a novel biomarker of macrovascular complications in type 2 diabetes. *Diabetes Care*. 2003;26(5):1507–12.
- [107] de la Maza M-P, Olivares D, Hirsch S, Sierralta W, Gattas V, Barrera G, et al. Weight increase and overweight are associated with DNA oxidative damage in skeletal muscle. *Clin Nutr*. 2006 Dec;25(6):968–76.
- [108] Hofer T, Karlsson HL, Möller L. DNA oxidative damage and strand breaks in young healthy individuals: A gender difference and the role of life style factors. *Free Radic Res* [Internet]. 2006 Jan 1;40(7):707–14. Available from: <https://doi.org/10.1080/10715760500525807>
- [109] Mizoue T, Tokunaga S, Kasai H, Kawai K, Sato M, Kubo T. Body mass index and oxidative DNA damage: A longitudinal study. *Cancer Sci* [Internet]. 2007 Aug 1;98(8):1254–8. Available from: <http://dx.doi.org/10.1111/j.1349-7006.2007.00511.x>
- [110] Cejvanovic V, Asferg C, Kjær LK, Andersen UB, Linneberg A, Frystyk J, et al. Markers of oxidative stress in obese men with and without hypertension. *Scand J Clin Lab Invest* [Internet]. 2016 Nov 16;76(8):620–5. Available from: <https://doi.org/10.1080/00365513.2016.1230776>
- [111] El Wakkad A, Giza AO, Elwakkad A, Hassan NE, Sibaii H, el Zayat S. Relationship Between Obesity and 8—hydr0xy—2—de0xy Guanosine as an Oxidative Marker in Obese. *J Med Sci*. 2011;11(5):231–5.
- [112] Protano C, Andreoli R, Mutti A, Petti S, Vitali M. Biomarkers of oxidative stress to nucleic acids: Background levels and effects of body mass index and life-style factors in an urban paediatric population. *Sci Total Environ* [Internet]. 2014;500–501:44–51. Available from: <http://www.sciencedirect.com/science/article/pii/S0048969714012741>
- [113] Ramachandra N, Rodriguez L, Devaraj S. Urinary Biomarkers of Oxidative Stress and Insulin Resistance in Childhood Obesity. *FASEB J* [Internet]. 2015 Apr 1;29(1 Supplement). Available from: http://www.fasebj.org/content/29/1_Supplement/927.2.abstract
- [114] Demirbag R, Yilmaz R, Gur M, Celik H, Guzel S, Selek S, et al. DNA damage in metabolic syndrome and its association with antioxidative and oxidative measurements. *Int J Clin Pract*. 2006 Oct;60(10):1187–93.
- [115] Kobayashi J. Molecular mechanism of the recruitment of NBS1/hMRE11/hRAD50 complex to DNA double-strand breaks: NBS1 binds to gamma-H2AX through FHA/BRCT domain. *J Radiat Res* [Internet]. 2004;45(4):473–8. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/15635255>
- [116] Khanna KK, Jackson SP. DNA double-strand breaks: signaling, repair and the cancer connection. *Nat Genet* [Internet]. 2001;27(3):247–54. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/11242102>
- [117] Tu WZ, Li B, Huang B, Wang Y, Liu XD, Guan H, et al. ??h2AX foci formation in the absence of DNA damage: Mitotic H2AX phosphorylation is mediated by the DNA-PKcs/CHK2 pathway. *FEBS Lett* [Internet]. 2013;587(21):3437–43. Available from: <http://dx.doi.org/10.1016/j.febslet.2013.08.028>
- [118] Nikolova T, Dvorak M, Jung F, Adam I, Kramer E, Gerhold-Ay A, et al. The gammaH2AX assay for genotoxic and nongenotoxic agents: comparison of H2AX phosphorylation with cell death response. *Toxicol Sci*. 2014 Jul;140(1):103–17.
- [119] Nikolova T, Marini F, Kaina B. Genotoxicity testing: Comparison of the gammaH2AX focus assay with the alkaline and neutral comet assays. *Mutat Res*. 2017 Oct;822:10–8.
- [120] Zhou L, Li K, Luo Y, Tian L, Wang M, Li C, et al. Novel prognostic markers for patients with triple-negative breast cancer. *Hum Pathol*. 2013;44(10):2180–7.
- [121] Bartkova J, Horejsí Z, Koed K, Krämer A, Tort F, Zieger K, et al. DNA damage response as a candidate anti-cancer barrier in early human tumorigenesis. *Nature*. 2005;434(7035):864–70.
- [122] Mei L, Hu Q, Peng J, Ruan J, Zou J, Huang Q, et al. Phospho-histone H2AX is a diagnostic and prognostic marker for epithelial ovarian cancer. *Int J Clin Exp Pathol* [Internet]. 2015 May 1;8(5):5597–602. Available from: <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC4503141/>
- [123] Gorgoulis VG, Vassiliou L-VF, Karakaidos P, Zacharatos P, Kotsinas A, Liloglou T, et al. Activation

- of the DNA damage checkpoint and genomic instability in human precancerous lesions. *Nature* [Internet]. 2005 Apr 14;434(7035):907–13. Available from: <http://dx.doi.org/10.1038/nature03485>
- [124] Sedelnikova OA, Horikawa I, Zimonjic DB, Popescu NC, Bonner WM, Barrett JC. Senescing human cells and ageing mice accumulate DNA lesions with unreparable double-strand breaks. *Nat Cell Biol* [Internet]. 2004 Feb;6(2):168–70. Available from: <http://dx.doi.org/10.1038/ncb1095>
- [125] Wang C, Jurk D, Maddick M, Nelson G, Martin-Ruiz C, Von Zglinicki T. DNA damage response and cellular senescence in tissues of aging mice. *Aging Cell* [Internet]. 2009 Jun 1;8(3):311–23. Available from: <http://dx.doi.org/10.1111/j.1474-9726.2009.00481.x>
- [126] Schurman SH, Dunn CA, Greaves R, Yu B, Ferrucci L, Croteau DL, et al. Age-Related Disease Association of Endogenous γ -H2AX Foci in Mononuclear Cells Derived from Leukapheresis. *PLoS One*. 2012;7(9):3–10.
- [127] Redon CE, Nakamura AJ, Martin OA, Parekh PR, Weyemi US, Bonner WM. Recent developments in the use of γ -H2AX as a quantitative DNA double-strand break biomarker. *Aging (Albany NY)* [Internet]. 2011 Feb 11;3(2):168–74. Available from: <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC3082012/>
- [128] Villaret A, Galitzky J, Decaunes P, Estève D, Marques M-A, Sengenès C, et al. Adipose Tissue Endothelial Cells From Obese Human Subjects: Differences Among Depots in Angiogenic, Metabolic, and Inflammatory Gene Expression and Cellular Senescence. *Diabetes* [Internet]. 2010 Oct 27;59(11):2755–63. Available from: <http://diabetes.diabetesjournals.org/content/59/11/2755.abstract>
- [129] Azzarà A, Pirillo C, Giovannini C, Federico G, Scarpato R. Different repair kinetic of DSBs induced by mitomycin C in peripheral lymphocytes of obese and normal weight adolescents. *Mutat Res Mol Mech Mutagen* [Internet]. 2016 Jul;789:9–14. Available from: <http://www.sciencedirect.com/science/article/pii/S0027510716300562>
- [130] Azzarà A, Chiamonte A, Filomeni E, Pinto B, Mazzoni S, Piaggi S, et al. Increased level of DNA damage in some organs of obese Zucker rats by γ -H2AX analysis. *Environ Mol Mutagen*. 2017 Aug;58(7):477–84.
- [131] Ostling O, Johanson KJ. Microelectrophoretic study of radiation-induced DNA damages in individual mammalian cells. *Biochem Biophys Res Commun*. 1984;123(1):291–8.
- [132] Olive PL, Banath JP. The comet assay: a method to measure DNA damage in individual cells. *Nat Protoc* [Internet]. 2006 Jun;1(1):23–9. Available from: <http://dx.doi.org/10.1038/nprot.2006.5>
- [133] Apostolou P, Toloudi M, Kourtidou E, Mimikakou G, Vlachou I, Chatziioannou M, et al. Use of the comet assay technique for quick and reliable prediction of in vitro response to chemotherapeutics in breast and colon cancer. *J Biol Res* [Internet]. 2014 Dec 1;21(1):14. Available from: <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC4389674/>
- [134] Katarkar A, Mukherjee S, Khan MH, Ray JG, Chaudhuri K. Comparative evaluation of genotoxicity by micronucleus assay in the buccal mucosa over comet assay in peripheral blood in oral precancer and cancer patients. *Mutagen* [Internet]. 2014 Sep 1;29(5):325–34. Available from: <http://mutage.oxfordjournals.org/content/29/5/325.abstract>
- [135] López-Diazguerrero NE, Luna-López A, Gutiérrez-Ruiz MC, Zentella A, Königsberg M. Susceptibility of DNA to oxidative stressors in young and aging mice. *Life Sci* [Internet]. 2005 Oct 14;77(22):2840–54. Available from: <http://www.sciencedirect.com/science/article/pii/S0024320505005266>
- [136] Piperakis SM, Kontogianni K, Karanastasi G, Iakovidou-Kritsi Z, Piperakis MM. The use of comet assay in measuring DNA damage and repair efficiency in child, adult, and old age populations. *Cell Biol Toxicol* [Internet]. 2009;25(1):65–71. Available from: <http://dx.doi.org/10.1007/s10565-007-9046-6>
- [137] Bukhari SA, Rajoka MI, Nagra SA, Rehman ZU. Plasma homocysteine and DNA damage profiles in normal and obese subjects in the Pakistani population. *Mol Biol Rep* [Internet]. 2010;37(1):289–95. Available from: <http://dx.doi.org/10.1007/s11033-009-9686-0>
- [138] Gandhi G, Kaur G. Assessment of DNA Damage in Obese Individuals. *Res J Biol*. 2012;02(02):37–44.
- [139] Tomasello B, Malfa G, Galvano F, Renis M. DNA damage in normal-weight obese syndrome measured by Comet assay. *Med J Nutrition Metab* [Internet]. 2011;4(2):99–104. Available from: <http://dx.doi.org/10.1007/s12349-010-0035-6>
- [140] Tenorio NM, Ribeiro DA, Alvarenga TA, Fracalossi ACC, Carlin V, Hirotsu C, et al. The influence of sleep deprivation and obesity on DNA damage in female Zucker rats. *Clinics* [Internet]. 2013 Mar

- 30;68(3):385–9. Available from: <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC3611896/>
- [141] Ibero-Baraibar I, Azqueta A, Lopez de Cerain A, Martinez JA, Zulet MA. Assessment of DNA damage using comet assay in middle-aged overweight/obese subjects after following a hypocaloric diet supplemented with cocoa extract. *Mutagen* [Internet]. 2015 Jan 1;30(1):139–46. Available from: <http://mutage.oxfordjournals.org/content/30/1/139.abstract>
- [142] Smith TR, Miller MS, Lohman KK, Case LD, Hu JH. DNA damage and breast cancer risk. *Carcinogenesis*. 2003;24(5):883–9.
- [143] De Lange T. 10 Telomere Dynamics and Genome Instability in Human Cancer. *Cold Spring Harb Monogr Arch*. 1995;29:265–93.
- [144] Olovnikov AM. A theory of marginotomy. The incomplete copying of template margin in enzymic synthesis of polynucleotides and biological significance of the phenomenon. *J Theor Biol*. 1973 Sep;41(1):181–90.
- [145] Allsopp RC, Vaziri H, Patterson C, Goldstein S, Younglai E V, Futcher AB, et al. Telomere length predicts replicative capacity of human fibroblasts. *Proc Natl Acad Sci U S A* [Internet]. 1992 Nov 1;89(21):10114–8. Available from: <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC50288/>
- [146] Xi H, Li C, Ren F, Zhang H, Zhang L. Telomere, aging and age-related diseases. *Aging Clin Exp Res* [Internet]. 2013;25(2):139–46. Available from: <https://doi.org/10.1007/s40520-013-0021-1>
- [147] Hochstrasser T, Marksteiner J, Humpel C. Telomere length is age-dependent and reduced in monocytes of Alzheimer patients. *Exp Gerontol* [Internet]. 2012 Feb 20;47(2):160–3. Available from: <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC3278593/>
- [148] Wang H, Chen H, Gao X, McGrath M, Deer D, De Vivo I, et al. Telomere length and risk of Parkinson’s disease. *Mov Disord* [Internet]. 2008 Jan 30;23(2):302–5. Available from: <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC2387100/>
- [149] Steer SE, Williams FMK, Kato B, Gardner JP, Norman PJ, Hall MA, et al. Reduced telomere length in rheumatoid arthritis is independent of disease activity and duration. *Ann Rheum Dis* [Internet]. 2007 Apr 17;66(4):476–80. Available from: <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC1856061/>
- [150] Ishikawa N, Nakamura K-I, Izumiyama-Shimomura N, Aida J, Ishii A, Goto M, et al. Accelerated in vivo epidermal telomere loss in Werner syndrome. *Aging (Albany NY)* [Internet]. 2011 Apr 25;3(4):417–29. Available from: <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC3117457/>
- [151] Decker ML, Chavez E, Vulto I, Lansdorp PM. Telomere length in Hutchinson-Gilford Progeria Syndrome. *Mech Ageing Dev* [Internet]. 2009;130(6):377–83. Available from: <http://www.sciencedirect.com/science/article/pii/S0047637409000396>
- [152] Fernandez P, Scaffidi P, Markert E, Lee J-H, Rane S, Misteli T. Transformation Resistance in a Premature Aging Disorder Identifies a Tumor-Protective Function of BRD4. *Cell Rep* [Internet]. 2014;9(1):248–60. Available from: <http://www.sciencedirect.com/science/article/pii/S221112471400761X>
- [153] Broberg K, Björk J, Paulsson K, Höglund M, Albin M. Constitutional short telomeres are strong genetic susceptibility markers for bladder cancer. *Carcinogenesis* [Internet]. 2005 Jul 1 [cited 2016 May 29];26(7):1263–71. Available from: <http://carcin.oxfordjournals.org/cgi/content/long/26/7/1263>
- [154] Zhu X, Han W, Xue W, Zou Y, Xie C, Du J, et al. The association between telomere length and cancer risk in population studies. *Sci Rep* [Internet]. 2016;6:22243. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/26915412>
- [155] Qu S, Wen W, Shu X-O, Chow W-H, Xiang Y-B, Wu J, et al. Association of Leukocyte Telomere Length With Breast Cancer Risk: Nested Case-Control Findings From the Shanghai Women’s Health Study. *Am J Epidemiol* [Internet]. 2013 Apr 1;177(7):617–24. Available from: <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC3657533/>
- [156] Skinner HG, Gangnon RE, Litzelman K, Johnson RA, Chari ST, Petersen GM, et al. Telomere length and pancreatic cancer: a case-control study. *Cancer Epidemiol Biomarkers Prev* [Internet]. 2012 Nov 23;21(11):2095–100. Available from: <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC3493789/>
- [157] Risques RA, Vaughan TL, Li X, Odze RD, Blount PL, Ayub K, et al. Leukocyte telomere length predicts cancer risk in Barrett’s esophagus. *Cancer Epidemiol Biomarkers Prev*. 2007;16(12):2649–55.
- [158] Jang JS, Choi YY, Lee WK, Choi JE, Cha SI, Kim YJ, et al. Telomere length and the risk of lung cancer. *Cancer Sci* [Internet]. 2008 Jul 1;99(7):1385–9. Available from: <http://dx.doi.org/10.1111/j.1349-7006.2008.00831.x>
- [159] Lan Q, Cawthon R, Gao Y, Hu W, Hosgood III HD, Barone-Adesi F, et al. Longer Telomere Length

- in Peripheral White Blood Cells Is Associated with Risk of Lung Cancer and the rs2736100 (CLPTM1L-TERT) Polymorphism in a Prospective Cohort Study among Women in China. *PLoS One* [Internet]. 2013 Mar 26;8(3):e59230. Available from: <https://doi.org/10.1371/journal.pone.0059230>
- [160] Ernst A, Jones DTW, Maass KK, Rode A, Deeg KI, Jebaraj BMC, et al. Telomere dysfunction and chromothripsis. *Int J Cancer* [Internet]. 2016 Jun 15;138(12):2905–14. Available from: <http://dx.doi.org/10.1002/ijc.30033>
- [161] Valdes AM, Andrew T, Gardner JP, Kimura M, Oelsner E, Cherkas LF, et al. Obesity, cigarette smoking, and telomere length in women. *Lancet* (London, England). 2005 Aug;366(9486):662–4.
- [162] Nordfjäll K, Eliasson M, Stegmayr B, Melander O, Nilsson P, Roos G. Telomere Length Is Associated With Obesity Parameters but With a Gender Difference. *Obesity* [Internet]. 2008 Dec 1;16(12):2682–9. Available from: <http://dx.doi.org/10.1038/oby.2008.413>
- [163] Kim S, Parks CG, DeRoo LA, Chen H, Taylor JA, Cawthon RM, et al. Obesity and Weight Gain in Adulthood and Telomere Length. *Cancer Epidemiol Biomarkers Prev* [Internet]. 2009 Mar;18(3):816. Available from: <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC2805851/>
- [164] Moreno-Navarrete JM, Ortega F, Sabater M, Ricart W, Fernández-Real JM, Fernandez-Real JM. Telomere length of subcutaneous adipose tissue cells is shorter in obese and formerly obese subjects. *Int J Obes*. 2010;34(8):1345–8.
- [165] Diaz VA, Mainous a G, Player MS, Everett CJ. Telomere length and adiposity in a racially diverse sample. *Int J Obes* [Internet]. 2010;34(2):261–5. Available from: <http://dx.doi.org/10.1038/ijo.2009.198>
- [166] Zannolli R, Mohn A, Buoni S, Pietrobelli A, Messina M, Chiarelli F, et al. Telomere length and obesity. *Acta Paediatr*. 2008;97(7):952–4.
- [167] Al-Attas OS, Al-Daghri N, Bamakhramah A, Shaun Sabico S, McTernan P, Huang T-K. Telomere length in relation to insulin resistance, inflammation and obesity among Arab youth. *Acta Paediatrica* [Internet]. 2010 Jun 1;99(6):896–9. Available from: <http://dx.doi.org/10.1111/j.1651-2227.2010.01720.x>
- [168] Buxton JL, Walters RG, Visvikis-Siest S, Meyre D, Froguel P, Blakemore AIF. Childhood obesity is associated with shorter leukocyte telomere length. *J Clin Endocrinol Metab*. 2011;96(5):1500–5.
- [169] Zhu H, Wang X, Gutin B, Davis CL, Keeton D, Thomas J, et al. Leukocyte Telomere Length in Healthy White and Black Adolescents: Relations to Race, Sex, Adiposity, Adipokines and Physical Activity. *J Pediatr* [Internet]. 2011 Feb 19;158(2):215–20. Available from: <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC3010324/>
- [170] Park MH, Falconer C, Viner RM, Kinra S. The impact of childhood obesity on morbidity and mortality in adulthood: a systematic review. *Obes Rev*. 2012;13(11):985–1000.
- [171] Hardikar S, Song X, Risques RA, Montine TJ, Duggan C, Blount PL, et al. Obesity and inflammation markers in relation to leukocyte telomere length in a cross-sectional study of persons with Barrett's esophagus. *BMC Obes* [Internet]. 2015 Sep 10;2:32. Available from: <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC4566310/>
- [172] Rode L, Nordestgaard BG, Weischer M, Bojesen SE. Increased Body Mass Index, Elevated C-reactive Protein, and Short Telomere Length. *J Clin Endocrinol Metab* [Internet]. 2014 Apr 24;99(9):E1671–5. Available from: <http://dx.doi.org/10.1210/jc.2014-1161>
- [173] Rehkopf DH, Needham BL, Lin J, Blackburn EH, Zota AR, Wojcicki JM, et al. Leukocyte Telomere Length in Relation to 17 Biomarkers of Cardiovascular Disease Risk: A Cross-Sectional Study of US Adults. *PLOS Med* [Internet]. 2016 Nov 29;13(11):e1002188. Available from: <https://doi.org/10.1371/journal.pmed.1002188>
- [174] Frayling TM, Timpson NJ, Weedon MN, Zeggini E, Freathy RM, Lindgren CM, et al. A common variant in the FTO gene is associated with body mass index and predisposes to childhood and adult obesity. *Science*. 2007 May;316(5826):889–94.
- [175] Rhee KE, Phelan S, McCaffery J. Early Determinants of Obesity: Genetic, Epigenetic, and In Utero Influences. *Int J Pediatr* [Internet]. 2012 May 31;2012:463850. Available from: <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC3371343/>
- [176] Luperini BCO, Almeida DC, Porto MP, Marcondes JPC, Prado RP, Rasera I, et al. Gene polymorphisms and increased DNA damage in morbidly obese women. *Mutat Res*. 2015 Jun;776:111–7.
- [177] Long J, Zhang B, Signorello LB, Cai Q, Deming-Halverson S, Shrubsole MJ, et al. Evaluating

- genome-wide association study-identified breast cancer risk variants in African-American women. *PLoS One*. 2013;8(4):e58350.
- [178] Kusinska R, Górniak P, Pastorczak A, Fendler W, Potemski P, Mlynarski W, et al. Influence of genomic variation in FTO at 16q12.2, MC4R at 18q22 and NRXN3 at 14q31 genes on breast cancer risk. *Mol Biol Rep* [Internet]. 2012 Mar 19;39(3):2915–9. Available from: <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC3271204/>
- [179] McCullough LE, Eng SM, Bradshaw PT, Cleveland RJ, Steck SE, Terry MB, et al. Genetic polymorphisms in DNA repair and oxidative stress pathways may modify the association between body size and postmenopausal breast cancer. *Ann Epidemiol*. 2015 Apr;25(4):263–9.
- [180] Rupérez AI, Gil A, Aguilera CM. Genetics of Oxidative Stress in Obesity. *Int J Mol Sci* [Internet]. 2014 Feb 20;15(2):3118–44. Available from: <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC3958901/>
- [181] Stathopoulou MG, Petrelis AM, Buxton JL, Froguel P, Blakemore AIF, Visvikis-Siest S. Genetic determinants of leucocyte telomere length in children: a neglected and challenging field. *Paediatr Perinat Epidemiol*. 2015 Mar;29(2):146–50.
- [182] Kwok CS, Pradhan A, Khan MA, Anderson SG, Keavney BD, Myint PK, et al. Bariatric surgery and its impact on cardiovascular disease and mortality: a systematic review and meta-analysis. *Int J Cardiol*. 2014 Apr;173(1):20–8.
- [183] Ashrafian H, Toma T, Rowland SP, Harling L, Tan A, Efthimiou E, et al. Bariatric Surgery or Non-Surgical Weight Loss for Obstructive Sleep Apnoea? A Systematic Review and Comparison of Meta-analyses. *Obes Surg*. 2015 Jul;25(7):1239–50.
- [184] Lassailly G, Caiazzo R, Buob D, Pigeyre M, Verkindt H, Labreuche J, et al. Bariatric Surgery Reduces Features of Nonalcoholic Steatohepatitis in Morbidly Obese Patients. *Gastroenterology*. 2015 Aug;149(2):376–9.
- [185] Rubino F, Schauer PR, Kaplan LM, Cummings DE. Metabolic surgery to treat type 2 diabetes: clinical outcomes and mechanisms of action. *Annu Rev Med*. 2010;61:393–411.
- [186] Casagrande DS, Rosa DD, Umpierre D, Sarmiento RA, Rodrigues CG, Schaan BD. Incidence of cancer following bariatric surgery: systematic review and meta-analysis. *Obes Surg*. 2014 Sep;24(9):1499–509.
- [187] Himbert C, Thompson H, Ulrich CM. Effects of Intentional Weight Loss on Markers of Oxidative Stress, DNA Repair and Telomere Length - a Systematic Review. *Obes Facts* [Internet]. 2017;10(6):648–65. Available from: <https://www.karger.com/DOI/10.1159/000479972>
- [188] Fejfer K, Buczeko P, Niczyporuk M, Ładny JR, Hady HR, Knaś M, et al. Oxidative Modification of Biomolecules in the Nonstimulated and Stimulated Saliva of Patients with Morbid Obesity Treated with Bariatric Surgery. *Biomed Res Int*. 2017;2017.
- [189] Kocael A, Erman H, Zengin K, Kocael PCA, Korkmaz GG, Gelisgen R, et al. The effects on oxidative DNA damage of laparoscopic gastric band applications in morbidly obese patients. *Can J Surg*. 2014;57(3):183.
- [190] Mitterberger MC, Mattesich M, Zwerschke W. Bariatric surgery and diet-induced long-term caloric restriction protect subcutaneous adipose-derived stromal/progenitor cells and prolong their life span in formerly obese humans. *Exp Gerontol*. 2014;56:106–13.
- [191] O'Callaghan NJ, Clifton PM, Noakes M, Fenech M. Weight loss in obese men is associated with increased telomere length and decreased abasic sites in rectal mucosa. *Rejuvenation Res*. 2009;12(3):169–76.
- [192] Formichi C, Cantara S, Ciuoli C, Neri O, Chiofalo F, Selmi F, et al. Weight loss associated with bariatric surgery does not restore short telomere length of severe obese patients after 1 year. *Obes Surg*. 2014;24(12):2089–93.
- [193] Bankoglu EE, Seyfried F, Arnold C, Soliman A, Jurowich C, Germer CT, et al. Reduction of DNA damage in peripheral lymphocytes of obese patients after bariatric surgery-mediated weight loss. *Mutagenesis*. 2017 Dec;
- [194] García-Calzón S, Moleres A, Marcos A, Campoy C, Moreno LA, Azcona-Sanjulián MC, et al. Telomere Length as a Biomarker for Adiposity Changes after a Multidisciplinary Intervention in Overweight/Obese Adolescents: The EVASYON Study. *PLoS One* [Internet]. 2014 Feb 24;9(2):e89828. Available from: <https://doi.org/10.1371/journal.pone.0089828>
- [195] Soares NP, Santos ACS dos, Costa EC, Azevedo GD, Damasceno DC, Fayh APT, et al. Diet-Induced Weight Loss Reduces DNA Damage and Cardiometabolic Risk Factors in Overweight/Obese Women

- with Polycystic Ovary Syndrome. *Ann Nutr Metab.* 2016;68(3):220–7.
- [196] Pardini B, Viberti C, Naccarati A, Allione A, Oderda M, Critelli R, et al. Increased micronucleus frequency in peripheral blood lymphocytes predicts the risk of bladder cancer. *Br J Cancer* [Internet]. 2017;116(2):202–10. Available from: <http://dx.doi.org/10.1038/bjc.2016.411>
- [197] Varga D, Hoegel J, Maier C, Jainta S, Hoehne M, Patino-Garcia B, et al. On the difference of micronucleus frequencies in peripheral blood lymphocytes between breast cancer patients and controls. *Mutagenesis.* 2006 Sep;21(5):313–20.
- [198] Santos RA, Teixeira AC, Mayorano MB, Carrara HHA, Andrade JM, Takahashi CS. Basal levels of DNA damage detected by micronuclei and comet assays in untreated breast cancer patients and healthy women. *Clin Exp Med.* 2010;10(2):87–92.
- [199] Bolognesi C, Bruzzi P, Gismondi V, Volpi S, Viassolo V, Pedemonte S, et al. Clinical Application of Micronucleus Test: A Case-Control Study on the Prediction of Breast Cancer Risk/Susceptibility. Scarfi MR, editor. *PLoS One* [Internet]. 2014 Nov 21;9(11):e112354. Available from: <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC4240584/>
- [200] Maffei F, Zolezzi Moraga JM, Angelini S, Zenesini C, Musti M, Festi D, et al. Micronucleus frequency in human peripheral blood lymphocytes as a biomarker for the early detection of colorectal cancer risk. *Mutagen* [Internet]. 2014 May 1;29(3):221–5. Available from: <http://mutage.oxfordjournals.org/content/29/3/221.abstract>
- [201] Aires GMA, Meireles JRC, Oliveira PC, Oliveira JL, Araújo EL, Pires BC, et al. Micronuclei as biomarkers for evaluating the risk of malignant transformation in the uterine cervix. *Genet Mol Res.* 2011;10(3):1558–64.
- [202] Kiraz A, Açmaz G, Uysal G, Unal D, Dönmez-Altuntas H. Micronucleus testing as a cancer detector: endometrial hyperplasia to carcinoma. *Arch Gynecol Obstet.* 2016;293(5):1065–71.
- [203] El-Zein RA, Schabath MB, Etzel CJ, Lopez MS, Franklin JD, Spitz MR. Cytokinesis-Blocked Micronucleus Assay as a Novel Biomarker for Lung Cancer Risk. *Cancer Res* [Internet]. 2006 Jun 15;66(12):6449–56. Available from: <http://cancerres.aacrjournals.org/content/66/12/6449.abstract>
- [204] Chang P, Li Y, Li D. Micronuclei levels in peripheral blood lymphocytes as a potential biomarker for pancreatic cancer risk. *Carcinog* [Internet]. 2011 Feb 1;32(2):210–5. Available from: <http://carcin.oxfordjournals.org/content/32/2/210.abstract>
- [205] Bonassi S, Znaor A, Ceppi M, Lando C, Chang WP, Holland N, et al. An increased micronucleus frequency in peripheral blood lymphocytes predicts the risk of cancer in humans. *Carcinogenesis.* 2007 Mar;28(3):625–31.
- [206] Moran LJ, Noakes M, Clifton PM, Norman RJ, Fenech MF. Genome instability is increased in lymphocytes of women with polycystic ovary syndrome and is correlated with insulin resistance. *Mutat Res Mol Mech Mutagen.* 2008;639(1):55–63.
- [207] Murugaiyan SB, Ramasamy R, Nakkeeran M, Rangdhol V, Srinivasan AR, Niranjana G. Urinary 8-hydroxydeoxyguanosine as a marker of oxidative stress induced genetic toxicity in oral cancer patients. *Indian J Dent Res.* 2015;26(3):226.
- [208] Kuo H-W, Chou S-Y, Hu T-W, Wu F-Y, Chen D-J. Urinary 8-hydroxy-2'-deoxyguanosine (8-OHdG) and genetic polymorphisms in breast cancer patients. *Mutat Res Toxicol Environ Mutagen.* 2007;631(1):62–8.
- [209] Himmetoglu S, Dincer Y, Ersoy YE, Bayraktar B, Celik V, Akcay T. DNA oxidation and antioxidant status in breast cancer. *J Investig Med.* 2009;57(6):720–3.
- [210] Chang D, Wang FAN, Zhao Y-S, Pan H-Z. Evaluation of oxidative stress in colorectal cancer patients. *Biomed Environ Sci.* 2008;21(4):286–9.
- [211] Guo C, Li X, Wang R, Yu J, Ye M, Mao L, et al. Association between Oxidative DNA Damage and Risk of Colorectal Cancer: Sensitive Determination of Urinary 8-Hydroxy-2'-deoxyguanosine by UPLC-MS/MS Analysis. *Sci Rep* [Internet]. 2016 Sep 2;6:32581. Available from: <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC5009303/>
- [212] Diakowska D, Lewandowski A, Kopeć W, Diakowski W, Chrzanowska T. Oxidative DNA damage and total antioxidant status in serum of patients with esophageal squamous cell carcinoma. *Hepatogastroenterology.* 2007;54(78):1701–4.
- [213] PYLVÁS M, PUISTOLA U, LAATIO L, KAUPPILA S, KARIHTALA P. Elevated Serum 8-OHdG Is Associated with Poor Prognosis in Epithelial Ovarian Cancer. *Anticancer Res* [Internet]. 2011 Apr 1;31(4):1411–5. Available from: <http://ar.iarjournals.org/content/31/4/1411.abstract>

- [214] MIYAKE H, HARA I, KAMIDONO S, ETO H. Oxidative Dna Damage in Patients With Prostate Cancer and Its Response To Treatment. *J Urol* [Internet]. 2004;171(4):1533–6. Available from: <http://linkinghub.elsevier.com/retrieve/pii/S0022534705623361>
- [215] Zhao L, Chang DW, Gong Y, Eng C, Wu X. Measurement of DNA damage in peripheral blood by the γ -H2AX assay as predictor of colorectal cancer risk. *DNA Repair (Amst)* [Internet]. 2015;53:24–30. Available from: <http://www.sciencedirect.com/science/article/pii/S1568786416304189>
- [216] Lee YI-C, Yin TC, Chen YI-T, Chai C-Y, Wang JY, Liu M-C, et al. High expression of phospho-H2AX predicts a poor prognosis in colorectal cancer. *Anticancer Res*. 2015;35(4):2447–53.
- [217] Kim JH, Kim SS, Byun SW, Chang YJ, Kim JS, Kim JK, et al. Double strand break of DNA in gastric adenoma and adenocarcinoma. *Korean J Gastroenterol*. 2010;55(1):19–25.
- [218] Wasco MJ, Pu RT, Yu L, Su L, Ma L. Expression of gamma-H2AX in melanocytic lesions. *Hum Pathol*. 2008 Nov;39(11):1614–20.
- [219] Hu J, Hu S, Hou X, Zhu X, Cao J, Jiang L, et al. Abnormal Expression of DNA Double-Strand Breaks Related Genes, ATM and GammaH2AX, in Thyroid Carcinoma. *Int J Endocrinol* [Internet]. 2015 Mar 16;2015:136810. Available from: <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC4378699/>
- [220] Kattimuthu P, Ramachandra RK, Chand P, Kadambari D. DNA Damage in Patients of Carcinoma Breast a Clinical Study by using Comet Assay. *J Anat Soc India* [Internet]. 2012;61(2):149–54. Available from: <http://www.sciencedirect.com/science/article/pii/S0003277812800220>
- [221] Schabath MB, Spitz MR, Grossman HB, Zhang K, Dinney CP, Zheng P-J, et al. Genetic Instability in Bladder Cancer Assessed by the Comet Assay. *J Natl Cancer Inst* [Internet]. 2003 Apr 2;95(7):540–7. Available from: <http://jnci.oxfordjournals.org/content/95/7/540.abstract>
- [222] Rawat G, Urs A, Chakravarti A, Kumar P. Evaluation of DNA Damage in Peripheral Blood Leukocytes in Oral Potentially Malignant and Malignant Disorders by Comet Assay. *Clin Cancer Investig J* [Internet]. 2018 Mar 1;7(2):50–5. Available from: <http://www.cci-j-online.org/article.asp?issn=2278-0513>
- [223] Buchynska LG, Brieieva O V, Iurchenko NP, Protsenko V V, Nespryadko S V. DNA damage in tumor cells and peripheral blood lymphocytes of endometrial cancer patients assessed by the comet assay. *Exp Oncol*. 2017 Dec;39(4):299–303.
- [224] Shen J, Terry MB, Gurchich I, Liao Y, Senie RT, Santella RM. Short Telomere Length and Breast Cancer Risk: A Study in Sister Sets. *Cancer Res* [Internet]. 2007 Jun 1;67(11):5538 LP-5544. Available from: <http://cancerres.aacrjournals.org/content/67/11/5538.abstract>
- [225] Liu X, Bao G, Huo T, Wang Z, He X, Dong G. Constitutive telomere length and gastric cancer risk: Case-control analysis in Chinese Han population. *Cancer Sci* [Internet]. 2009 Jul 1;100(7):1300–5. Available from: <http://dx.doi.org/10.1111/j.1349-7006.2009.01169.x>
- [226] Mirabello L, Garcia-Closas M, Cawthon R, Lissowska J, Brinton LA, Peplowska B, et al. Leukocyte telomere length in a population-based case-control study of ovarian cancer: a pilot study. *Cancer Causes Control* [Internet]. 2010;21(1):77–82. Available from: <https://doi.org/10.1007/s10552-009-9436-6>
- [227] Meeker AK, Hicks JL, Platz EA, March GE, Bennett CJ, Delannoy MJ, et al. Telomere shortening is an early somatic DNA alteration in human prostate tumorigenesis. *Cancer Res*. 2002;62(22):6405–9.
- [228] Han J, Qureshi AA, Prescott J, Guo Q, Ye L, Hunter DJ, et al. A Prospective Study of Telomere Length and the Risk of Skin Cancer. *J Invest Dermatol* [Internet]. 2009;129(2):415–21. Available from: <http://www.sciencedirect.com/science/article/pii/S0022202X15342202>
- [229] Cassidy A, De Vivo I, Liu Y, Han J, Prescott J, Hunter DJ, et al. Associations between diet, lifestyle factors, and telomere length in women. *Am J Clin Nutr* [Internet]. 2010 May 10;91(5):1273–80. Available from: <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC2854902/>
- [230] LEE M, MARTIN H, FIRPO MA, DEMERATH EW. Inverse Association Between Adiposity and Telomere Length: The Fels Longitudinal Study. *Am J Hum Biol* [Internet]. 2011;23(1):100–6. Available from: <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC3245638/>
- [231] Njajou OT, Cawthon RM, Blackburn EH, Harris TB, Li R, Sanders JL, et al. Shorter telomeres are associated with obesity and weight gain in the elderly. *Int J Obes* [Internet]. 2012 Sep;36(9):1176–9. Available from: <http://dx.doi.org/10.1038/ijo.2011.196>
- [232] Cui Y, Gao Y-T, Cai Q, Qu S, Cai H, Li H-L, et al. Associations of leukocyte telomere length with body anthropometric indices and weight change in Chinese women. *Obesity (Silver Spring)* [Internet]. 2013 Dec 29;21(12):2582–8. Available from: <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC3676725/>

Table 1. Associations between micronuclei (MNi) frequency and site-specific cancers.

Cancer Site	Study Population	Cell type USED for Micronuclei (MNi) test	Key Findings	Authors, Date
Bladder	158 patients/ 158 controls (age matched)	PBLs	Increase in MN frequency associated with increased risk of cancer	Pardini <i>et al.</i> , 2017 [196]
Breast	91 patients/ 96 controls	PBLs	Higher frequency of MNi in breast cancer patients	Varga <i>et al.</i> , 2006 [197]
	45 patients/ 85 controls (age matched)	PBLs	Higher frequency of MNi in breast cancer patients	Santos <i>et al.</i> , 2010. [198]
	220 patients/ 295 controls	PBLs	No significant difference of MNi frequency between patients and controls	Bolognesi <i>et al.</i> , 2014 [199]
	Follow up of 1650 adults	PBLs	Higher frequency of MNi at baseline in cancer patients.	Murgia <i>et al.</i> , 2008 [78]
Colorectal	25 cancer patients/ 26 polyp patients/ 31 controls	PBLs	2.1x higher frequency of MNi in cancer patients than controls. Polyp patients had 1.5x higher frequency of MNi than controls.	Maffei <i>et al.</i> , 2014 [200]
Encephalon	Follow up of 1650 adults	PBLs	Higher frequency of MNi at baseline in cancer patients.	Murgia <i>et al.</i> , 2008 [78]
Endometrial Cancer	59 patients/ 59 controls	Cervical squamous epithelial cells	20% higher frequency of MNi in patients.	Aires <i>et al.</i> , 2011 [201]
	20 endometrial hyperplasia patients/ 20 cancer patients/ 20 controls	PBLs	2.9x higher frequency of MNi in women with endometrial cancer.	Kiraz <i>et al.</i> , 2016 [202]
Lung	216 small cell lung cancer patients/ 173 Non-small cell lung cancer	PBLs	Higher MNi frequency in lung cancer patients.	El-Zein <i>et al.</i> , 2006 [203]

	patients/ 204 controls			
Lymphoma	Follow up of 1650 adults	PBLs	Higher frequency of MNi at baseline in cancer patients.	Murgia et al., 2008 [78]
Oral	Follow up of 1650 adults	PBLs	Higher frequency of MNi at baseline in cancer patients.	Murgia et al., 2008 [78]
Pancreatic	346 patients/ 449 controls	PBLs	1.6x higher MNi frequency in pancreatic cancer patients.	Chang, Li and Li, 2011 [204]
Stomach	6718 adult participants across 20 sites	PBLs	Higher frequencies of MNi were associated with increased risk of stomach cancer.	Bonassi <i>et al.</i> , 2007 [205]
Urogenital	6718 adult participants across 20 sites	PBLs	Higher frequencies of MNi were associated with increased risk of urogenital cancer.	Bonassi <i>et al.</i> , 2007 [205]

Table 2. Associations between micronuclei (MNi) frequency and adiposity

Date	Study type	Study population	Cell type for MNi test	Key findings	Authors
2006	Case control	19 PCOS patients/19 controls	PBLs	Higher frequency of MNi in PCOS patients. Positive correlation between BMI and MNi frequency.	Yesilada <i>et al.</i> , [84]
2008	Case control	14 PCOS patients/16 controls (Age & overweight matched)	PBLs	Higher frequency of MNi in overweight PCOS patients. Insulin resistance was positively associated with aneuploidy of X chromosomes.	Moran <i>et al.</i> , [206]
2010	Case control	36 overweight PCOS patients/ 29 controls	PBLs	Higher frequency of MNi in overweight PCOS patients.	Hamurcu <i>et al.</i> , [86]
2011	Case-control	60 obese/ 20 overweight/ 38 children controls	PBLs	Higher frequency of MNi and γ -H2AX foci in overweight and obese children.	Scarpato <i>et al.</i> , [87]

2014	Case-control	83 obese/ 21 overweight/ 21 healthy weight adult controls	PBLs	BMI not WHR is positively correlated with MNi frequency in all subjects.	Donmez-Altuntas <i>et al.</i> , [86]
2016	Meta-analysis	8 studies covering 47 mammalian species	Erythrocytes	Body mass is positively correlated with MNi frequency.	Croco <i>et al.</i> , [90]
2016	Case-control	18 Diet-induced obesity (DIO) mice/ 18 non-DIO control mice	Erythrocytes and reticulocytes	No difference in MNi frequency between DIO and non-DIO mice. Erythrocytes showed an increase in mutant cells (<i>Pig-a</i>).	Wickliffe <i>et al.</i> , [91]
2017	Longitudinal study	87 pregnant women and their fetal offspring	PBLs	Positive correlation of birth weight and length with MNi frequency, nucleoplasmic bridges and nuclear buds.	Singh <i>et al</i> [88]
2017	Correlation	300 Korean men and women aged 30-59 years	PBLs	Positive correlation between age and MNi but not BMI.	Cho <i>et al</i> [89]

Table 3. Associations between 8-OHdG concentration and site-specific cancers.

Cancer Site	Study Population	Tissue type for 8-OHdG test	Key Findings	Authors, Date
Buccal mucosa	30 patients/ 30 controls	Urine	8-OHdG higher in cancer patients.	Murugaiyan <i>et al.</i> , 2015 [207]
Breast	49 patients/ 49 controls	Serum	8-OHdG higher in cancer patients.	Himmetoglu <i>et al.</i> , 2009 [209]
	60 patients/ 60 controls	Urine	8-OHdG higher in cancer patients.	Kuo <i>et al.</i> , 2007 [208]
Colo-rectum	36 patients/ 40 controls	Serum	8-OHdG higher in cancer patients.	Chang <i>et al.</i> , 2008 [210]
	84 patients/ 142 controls	Urine	8-OHdG higher in cancer patients.	Guo <i>et al.</i> , 2016 [211]

Oesophagus	18 patients/ 12 controls	Serum	8-OHdG higher in cancer patients.	Diakowska <i>et al.</i> , 2007 [212]
Ovaries	84 patients	Serum and tumour tissue	High 8-OHdG levels (>140pg/ml) are associated with poorer prognosis.	Pylvas <i>et al.</i> , 2011 [213]
Prostate	82 patients/ 82 controls	Urine	8-OHdG higher in cancer patients.	Miyake <i>et al.</i> , 2004 [214]

Table 4. Associations between 8-OHdG concentration and adiposity

Date	Study type	Study population	Tissue type for 8-ohdG test	Key findings	Authors
2006	Case control	27 non-obese hernia patients	Skeletal Muscle	Weight gain was associated with increased 8-OHdG.	<i>De la Maza et al.</i> , [107]
2006	Correlation	99 young adults	PBLs	No correlation between BMI and 8-OHdG.	Hofer, Karlsson and Möller, [108]
2007	Longitudinal	174 healthy adults	Urine	A decrease in BMI by 1.0 was associated with a 2.7% increase in 8-OHdG.	Mizoue <i>et al.</i> , [109]
2011	Correlation	103 adolescents	Serum	BMI was positively correlated with 8-OHdG.	El Wakkad <i>et al.</i> , [111]
2014	Case control	83 obese/ 21 overweight/ 21 control adults	Plasma	8-OHdG concentration was lower in the obese group.	Donmez-Altuntas <i>et al.</i> , [86]
2014	Correlation	159 healthy children	Urine	BMI was inversely correlated 8-OHdG.	Protano <i>et al.</i> , [112]
2015	Case control	24 obese/28 control children	Urine	Higher 8-OHdG in obese children.	Ramachandra <i>et al.</i> [113]
2016	Case control	63 obese hypertensive/40 obese-normotensive/ 27 control men	Urine	No significant difference in 8-OHdG.	Cejvanovic <i>et al.</i> , [110]

Table 5. Associations between γ -H2AX frequency and site-specific cancers.

Cancer Site	Study Population	Tissue/Cell type for γ -H2AX test	Key Findings	Authors, Date
Breast	31 patients and 89 control	Breast	γ -H2AX higher in cancer patients.	Zhou <i>et al.</i> , 2013 [120]
Bladder	94 patients and 8 controls	Bladder	γ -H2AX higher in cancer patients.	Bartkova <i>et al.</i> , 2005 [121]
GI tract	320 normal and 320 controls	Lymphocytes	γ -H2AX higher in colorectal cancer patients.	Zhao <i>et al.</i> , 2017 [215]
	92 cancer patients	Colorectal	γ -H2AX associated with poor cancer prognosis.	Lee <i>et al.</i> , 2015 [216]
	121 cancer patients	Gastric epithelial cells	γ -H2AX higher in cancerous epithelial tissue.	Kim <i>et al.</i> , 2010 [217]
Ovaries	87 patients and 20 controls	Epithelial ovarian	γ -H2AX higher in cancer patients.	Mei <i>et al.</i> , 2015 [122]
Skin	9 benign and 108 cancer tissues	Melanocytes	γ -H2AX higher in cancer tissue.	Wasco <i>et al.</i> , 2008 [218]
Thyroid	30 benign and 153 cancer tissues	Thyroid	γ -H2AX higher in cancer tissue.	Hu <i>et al.</i> , 2015 [219]

Table 6. Associations between γ -H2AX frequency and adiposity

Date	Study type	Study population	Tissue/Cell type for γ -H2AX test	Key findings	Authors
2010	Correlation	29 males/1 female undergoing bariatric surgery	Adipocytes and endothelial cells	Endothelial cells from visceral adipose tissue had greater γ -H2AX positive foci than Subcutaneous adipose tissue.	Villaret <i>et al.</i> , [128]
2011	Case control	61 obese/ 20 overweight/ 38 control children	PBLs	Higher γ -H2AX positive foci in overweight and obese cohorts.	Scarpato <i>et al.</i> , [87]
2016	Case control	23 obese/ 15 control adolescents	PBLs	Higher γ -H2AX positive foci in obese cohort.	Azzarà <i>et al</i> [129]
2017	Case control	Obese and healthy weight 3-4 weeks old	Pancreas Lung	Higher γ -H2AX-positive foci in obese 9 to 13-week-old obese	Azzarà <i>et al</i> [130]

		rats and 9-13 weeks old rats.	Esophagus Kidney Gut	rats. No differences in the 3-4 week cohorts.	
--	--	-------------------------------	----------------------------	-----------------------------------------------	--

Table 7. Associations between comet tails and and site-specific cancers

Cancer Site	Study Population	Cell type for comet assay	Key Findings	Authors, Date
Breast	70 patients and 70 controls	PBLs	Comet tails were significantly higher in cancer patients.	Smith <i>et al.</i> , 2003 [142]
	35 patients, 15 first degree relatives and 15 controls	PBLs	Comet tails were significantly higher in cancer patients and first degree relatives.	Kattimuthu <i>et al.</i> , 2012 [220]
Bladder	114 patients and 145 controls	PBLs	Comet tails were significantly higher in cancer patients.	Schabath <i>et al.</i> , 2003 [221]
Buccal Mucosa	54 patients and 52 controls	PBLs	Comet tails were significantly higher in cancer patients.	Katarkar <i>et al.</i> , 2014 [134]
	55 patients and 55 controls	PBLs	Comet tails were significantly higher in cancer patients.	Rawat <i>et al.</i> , 2018 [222]
Uterus	106 patients and 30 controls	PBLs	Comet tails were significantly higher in cancer patients.	Buchynska <i>et al.</i> , 2017 [223]

Table 8. Associations between comet tails and adiposity

Date	Study type	Study population	Cell type for comet assay	Key findings	Authors
2003	Case control	70 breast cancer patients/ 70 controls	PBLs	BMI was positively correlated with comet tails in cancer patients but inversely correlated in controls.	Smith <i>et al.</i> , [142]
2006	Correlation	99 females (aged 19-31)	PBLs	BMI was positively correlated with comet tails.	Hofer <i>et al.</i> , [108]
2010	Case control	20 pregnant women/ 10 controls	Leukocytes	BMI was positively correlated with comet tails in pregnant women.	Bukhari <i>et al.</i> , [137]
2011	Case control	20 obese/ 20 normal weight	PBLs	Higher frequency of comet tails in obese and 'normal-	Tomasello <i>et al.</i> , [139]

		+ fat mass > 30%/ 20 control females		weight obese' cohorts.	
2012	Case Control	36 obese/15 control adults	Leukocytes	Higher frequency of comet tails in obese cohort.	Gandhi and Kaur [138]
2013	Case Control	Lean and obese Zucker rats at 3, 6, and 15 months old assigned to a sleep-deprived group or home- cage control (n=5 in each group)	PBLs	Higher frequency of comet tails in obese rats at 15 months old, irrespective of sleep status.	Tenorio et al., [140]
2015	Randomized, double-blind placebo controlled trial	50 adults	PBLs	Calorie restriction was associated with weight-loss and a decrease in comet tails.	Ibero-Baraibar <i>et al.</i> , [141]

Table 9. Associations between telomere length (TL) and site-specific cancers.

Cancer Site	Study Population	Cell type and technique for telomere length test	Key Findings	Authors, Date
Bladder	63 patients/ 158 controls	Buccal cells via qPCR	Shorter TL in cancer patients.	Broberg <i>et al.</i> , 2005 [153]
Breast	287 patients/ 350 sister controls	PBLs via qPCR	No significant difference in TL between groups.	Shen <i>et al.</i> , 2007 [224]
	601 patients/ 695 controls	PBLs via qPCR	Shorter TL in cancer patients.	Qu <i>et al.</i> , 2013 [155]
GI tract	396 patients/ 378 controls	PBLs via qPCR	Shorter TL in cancer patients.	Liu <i>et al.</i> , 2009 [225]
Lung	243 patients/ 243 controls	PBLs via qPCR	Shorter TL in cancer patients.	Jang <i>et al.</i> , 2008 [158]

	215 patients/ 215 controls	PBLs via qPCR	Shorter TL in cancer patients.	Lan <i>et al.</i> , 2013 [159]
Ovarian	99 patients/ 100 controls (age matched)	Leukocytes via qPCR	Shorter TL in cancer patients.	Mirabello <i>et al.</i> , 2010 [226]
Pancreas	499 patients/ 963 controls	PBLs via qPCR	Shorter telomeres and extremely long telomeres both related to increased pancreatic cancer risk.	Skinner <i>et al.</i> , 2012 [156]
Prostate	6 patients with prostate adenocarcinomas	Telomere Fluorescence in Situ Hybridization of prostate cells	Cells with high-grade prostatic intraepithelial neoplasia had shorter telomeres than normal cells in same tissue.	Meeker <i>et al.</i> , 2002 [227]
Skin	218 melanoma/ 285 squamous-cell carcinoma/ 300 basal-cell carcinoma (BCC) cases/ 870 controls	PBLs via qPCR	Short TL was associated with a decreased number of moles and decreased risk of melanoma and BCC.	Han <i>et al.</i> , 2009 [228]

Table 10. Associations between telomere length (TL) and adiposity

Date	Study type	Study population	Cell type & technique for telomere length test	Key findings	Authors
2005	Case control	119 obese/ 85 healthy weight female twins	PBLs via TRF	TRF length was lowest in the obese cohort and correlated with BMI and serum leptin.	Valdes <i>et al.</i> , [161]
2008	Correlation	989 adults	Leucocytes via qPCR	Inverse correlation between BMI and TL in women.	Nordfjäll <i>et al.</i> , [162]
2008	Case control	53 children/ 23 adults	PBLs via TRF	No difference in TL in children. Obese adults had shorter TL.	Zannolli <i>et al.</i> , [166]
2009	Correlation follow-up	647 female adults	Leucocytes via qPCR	High BMI and hip circumference inversely correlated with TL. Obese	Kim <i>et al.</i> , [163]

				females had shortest TL.	
2010	Case control	51 obese/ 21 non-obese adults	Subcutaneous adipose tissue via Southern blotting	BMI inversely correlated with TL. Formerly obese patients had shorter TL than never-obese.	Moreno-Navarrete <i>et al.</i> , [164]
2010	Correlation	317 adults (aged 40-64 years)	Leukocytes via qPCR	No significant correlations between BMI or visceral adipose tissue and TL.	Diaz <i>et al.</i> , [165]
2010	Correlation	2284 females	Leukocytes via qPCR	Waist circumference was inversely correlated with TL.	Cassidy <i>et al.</i> , [229]
2011	Correlation	309 non-Hispanic white participants aged 8 to 80 years	Leukocytes via qPCR	BMI, waist circumference, hip circumference, total body fat, and visceral adipose tissue volume inversely correlated with TL.	Lee <i>et al.</i> , [230]
2011	Case control	793 children (aged 2-17 years)	Leukocytes via qPCR	23.9% shorter TL in obese children.	Buxton <i>et al.</i> , [168]
2012	Correlation at baseline and 7yr follow up	2721 elderly subjects	Leukocytes via qPCR	BF% and subcutaneous fat inversely correlated with TL. No correlation between BMI and TL. 7 year follow up showed inverse correlation with BMI and BF%.	Njajou <i>et al.</i> , [231]
2013	Correlation	2,912 females (aged 40-70 years)	Leukocytes via qPCR	TL inversely correlated with BMI, waist circumference, waist-to-height ratio, weight, and hip circumference but not waist to hip ratio.	Cui <i>et al.</i> , [232]
2016	Correlation	7527 adults (aged 20-84)	Leukocytes via qPCR	Telomere length inversely correlated with BMI, waist circumference, BF% and C-reactive protein.	Rehkopf <i>et al.</i> , [173]