

# DNA Damage Repair in the Context of Plant Chromatin<sup>1</sup>

Mattia Donà and Ortrun Mittelsten Scheid\*

Gregor Mendel Institute of Molecular Plant Biology, Austrian Academy of Sciences, Vienna Biocenter, 1030 Vienna, Austria

ORCID IDs: 0000-0002-0812-7319 (M.D.); 0000-0002-7757-4809 (O.M.S.).

The integrity of DNA molecules is constantly challenged. All organisms have developed mechanisms to detect and repair multiple types of DNA lesions. The basic principles of DNA damage repair (DDR) in prokaryotes and unicellular and multicellular eukaryotes are similar, but the association of DNA with nucleosomes in eukaryotic chromatin requires mechanisms that allow access of repair enzymes to the lesions. This is achieved by chromatin-remodeling factors, and their necessity for efficient DDR has recently been demonstrated for several organisms and repair pathways. Plants share many features of chromatin organization and DNA repair with fungi and animals, but they differ in other, important details, which are both interesting and relevant for our understanding of genome stability and genetic diversity. In this Update, we compare the knowledge of the role of chromatin and chromatin-modifying factors during DDR in plants with equivalent systems in yeast and humans. We emphasize plant-specific elements and discuss possible implications.

A DNA molecule in a eukaryotic chromosome has a diameter of 2 nm but a length in the range of centimeters. Being  $10^7$  times longer than it is wide would make it highly sensitive to breakage if it were not organized in compact and dynamic chromatin, with nucleosomes as basic units followed by multiple higher order levels of organization. Within this packaging, replication and transcription constitute an endogenous mechanical strain, while exposure of cells to physical or chemical hazard creates a wide range of DNA damage induced by external factors, including photoproducts, pyrimidine dimers, interstrand cross-linking, and single and double-strand breaks (DSBs). All organisms possess mechanisms that repair DNA molecules. DNA damage repair (DDR) systems for the various types of damage are overlapping as well as complementary, and their prevalence depends on the organism, the stage of the cell cycle, the site and the amount of damage, and the availability of intact templates. A coarse categorization distinguishes nucleotide excision repair, base excision repair, nonhomologous end joining (NHEJ), and homologous recombination (HR). Many schematic models for these DDR pathways describe the action and interaction of several repair components, but they are usually misleading in one aspect: DNA strands are illustrated as straight lines, neglecting the emerging role of chromatin for the location and the fate of DNA lesions. More realistically, we should envisage DNA lesions in the chromatin context

like the leakage or burst of an in-ground pipe. Recognizing the defect, localizing it, excavating the broken parts, cleaning the ends, reconnecting them, and restoring the original state are mirrored in the subsequent steps of DDR: signaling, labeling, accessing, resecting, religating, and reassembling. Chromatin is expected to especially affect the access, resection, and restoration of the original arrangement, and several factors that can dissolve higher order structures, slide, evict, or exchange nucleosomes, or restore chromatin have been implicated in DDR (for review, see Lukas et al., 2011; Czaja et al., 2012; Dinant et al., 2012; Euskirchen et al., 2012; Lans et al., 2012; Soria et al., 2012; Altmeyer and Lukas, 2013; Dion and Gasser, 2013; Gospodinov and Herceg, 2013a, 2013b; Ohsawa et al., 2013; Papamichos-Chronakis and Peterson, 2013; Peterson and Almouzni, 2013; Price and D'Andrea, 2013; Stanley et al., 2013; House et al., 2014; Jeggo and Downs, 2014; Swygert and Peterson, 2014; Polo, 2015). The repair process in plants has been the focus of several reviews (Roth et al., 2012; Donà et al., 2013; Knoll et al., 2014a; Missirian et al., 2014). In this Update, we summarize recent literature describing the connection of repair and chromatin in plants, adding to earlier reviews (Balestrazzi et al., 2011; Zhu et al., 2011; Roy, 2014). Although plants are surprisingly much less investigated in this context when compared with other multicellular organisms, they are potentially rewarding for further research, as some repair- and chromatin-related genes have extended gene families, and several complete loss-of-function mutations that are lethal in animals are viable in plants. Due to length restriction, we will not cover replication-associated DNA repair, the role of chromatin for long-range interactions, or the recently emerging role of small RNA in DDR; instead, we focus on defects during interphase, local chromatin configuration, and DNA-associated proteins.


<sup>1</sup> This work was supported by the Austrian Academy of Sciences, the Austrian Science Fund (grant nos. FWF I1107, FWF I1477, and FWF W1238), and the Vienna Science and Technology Fund (grant no. LS13-057).

\* Address correspondence to [ortrun.mittelsten\\_scheid@gmi.oeaw.ac.at](mailto:ortrun.mittelsten_scheid@gmi.oeaw.ac.at).

[www.plantphysiol.org/cgi/doi/10.1104/pp.15.00538](http://www.plantphysiol.org/cgi/doi/10.1104/pp.15.00538)

Chromatin consists mainly of nucleosomes, spherical octamers formed by two molecules of each of the four histone types H2A, H2B, H3, and H4 with 1.7 turns of DNA wrapped around their surface and sealed by H1 linker histones. Firm interaction between histones and DNA together with regular nucleosome spacing results in stable and periodic structures, but

histone octamers can also become less tightly bound, change their position along the DNA, or be completely dismantled. The strength of the association can be modulated in three ways: (1) by energy-consuming remodeling processes that shift or remove nucleosomes; (2) by exchanging core histones with histone variants; and (3) by introducing posttranslational modifications



**Figure 1.** Schematic representation of chromatin changes connected with DNA damage. DDR is associated with chromatin at three different levels: nucleosomes can be shifted, evicted, or exchanged; their composition can be varied; and their subunits can be covalently modified. Arabidopsis proteins or protein complexes participating in these processes are listed, those with a known repair-related role in green, those with a reported chromatin connection in yellow; factors in gray have only sequence similarity to yeast or human genes necessary for DDR.

(PTMs) in the histone subunits (Fig. 1). In the following three paragraphs, we review the role of these chromatin changes during DNA repair in plants.

## CHROMATIN-REMODELING COMPLEXES CONNECTED WITH DDR

Chromatin remodeling complexes (remodelers) use ATP hydrolysis to slide, evict, or unwrap nucleosomes and can alter histone composition (for review, see Clapier and Cairns, 2009). They are grouped into different classes with unique features and specificity (Fig. 2) and are able to recognize a plethora of chromatin marks, either through diverse protein domains and/or through interaction with a large set of different subunits (Hopfner et al., 2012). This complexity reflects the multiple functions of remodelers in DNA-related processes, and it is not surprising that mutants lacking functional chromatin remodelers often exhibit increased sensitivity to DNA damage. However, only a few complexes have been linked unambiguously and directly to DDR. DNA repair requires nucleosome remodeling from the early steps through the final repair (Seeber et al., 2013; Tsabar and Haber, 2013). Recruitment of remodelers to the damage site changes the nucleosome occupancy of the surrounding area, ensuring accessibility for the repair factors. Moreover, as shown in the yeast *Saccharomyces cerevisiae*, chromatin remodeling is essential for modulating DNA strand resection, resolving intermediate structures during HR, and orchestrating the mobility of chromatin to enhance HR efficiency (for review, see Gospodinov and Herceg, 2013a; Seeber et al., 2013). Although unique features distinguish yeast and mammalian chromatin-remodeling complexes, the mechanisms underlying nucleosome alteration during DNA repair are highly conserved throughout evolution (Jasin and Rothstein, 2013; Seeber et al., 2013; Tsabar and Haber, 2013). Most of the repair-related chromatin-remodeling complexes described in yeast and mammals have orthologs in plants, and early characterization of several mutants indicated similar roles in repair (Shaked et al., 2006). However, the individual components were most often identified through their role in transcriptional regulation rather than in repair, as their loss of function caused striking deviations from normal development. Compared with the knowledge of yeast and mammalian complexes, there is relatively little known about the role of chromatin remodeling in DNA repair in plants. Unique structural and functional features of plant chromatin-remodeling complexes justify and highlight the need for more detailed studies (Knizewski et al., 2008). Therefore, we extend a recent comparison of the repair functions of chromatin remodelers in yeast and mammals (Seeber et al., 2013) to include current insights into their role in plant DDR (Fig. 2). We list the full gene names only for the most important components in our context; for the expansion of all abbreviated gene and protein complex names, see Supplemental Text S1.

The SWR1 (Swi2/Snf2-RELATED1) and INO80 (INOSITOL-REQUIRING MUTANT80) complexes are members of the family of SWR1-like chromatin remodelers. They share a conserved role in genome-wide regulation of transcription via installation and removal of the H2A.Z variant at transcriptional start sites. H2A.Z deposition affects genome stability, as SWR-C activity is coordinated with replicative polymerases to reduce mutation rates (Van et al., 2015). Both complexes also impact DDR by remodeling the nucleosomal neighborhood surrounding DSBs in yeast and mammals (for review, see Gerhold and Gasser, 2014). SWR1-C is also required during the early steps of mammalian HR to prevent the propagation of  $\gamma$ H2A.X, the phosphorylated histone variant that labels the site of damage, and at later steps for loading RAD51 onto single-stranded DNA to promote successful repair (Courilleau et al., 2012). Conversely, loss of SWR1-C in yeast does not affect resection, which suggests a role in NHEJ rather than HR. Both complexes are also involved in larger scale mobility: the yeast SWR1-C and INO80-C affect the relocation of unrepaired DSBs to anchor sites at the nuclear envelope in a cell cycle-dependent manner (Horigome et al., 2014). Unrepaired DSBs in mammalian cells are also relocated to the nuclear envelope to prevent erroneous recombination and to activate alternative repair pathways that can overcome persistent damage (Lemaître and Soutoglou, 2015). The specificity of INO80-C and SWR1-C function is provided by different subunits that mediate the complexes' interaction with DNA and proteins (Yen et al., 2013).

Also in plants, SWR1-C mediates the incorporation of the H2A.Z histone variant, besides being the most well-studied remodeling complex in the repair context. In *Arabidopsis* (*Arabidopsis thaliana*), the complex has a major role in transcriptional regulation, as SWR1-C mutants exhibit a characteristic early-flowering phenotype due to the dysregulation of several developmental genes (Noh and Amasino, 2003; Choi et al., 2005; Deal et al., 2005; Lázaro et al., 2008; March-Díaz and Reyes, 2009). A role of *Arabidopsis* SWR1-C in DNA repair has only recently been documented. Loss of one of several SWR1-C subunits, namely ACTIN-RELATED PROTEIN6 (ARP6), PHOTOPERIOD-INDEPENDENT EARLY FLOWERING1, and SWR1 COMPONENT6, results in increased DNA damage sensitivity. This phenotype is aggravated by the simultaneous impairment of NHEJ, while SWR1-C and HR mutants are epistatic, suggesting a role of SWR1-C in HR specifically. The involvement in somatic HR is supported by reduced recombination frequencies between incomplete but overlapping parts of a marker gene in the AtSWR1-C mutant background. Furthermore, reduced fertility of the mutants also indicates defects in meiosis (Rosa et al., 2013). The proximate cause, however, is difficult to separate from the role of ARP6 for temporal and spatial regulation of meiotic genes (Qin et al., 2014).

The INO80 complex in *Arabidopsis* is also involved in developmental regulation as well as in DNA repair

Grouping		Organism							
Family	Subfamily & composition	<i>Saccharomyces cerevisiae</i>	<i>Homo sapiens</i>	<i>Arabidopsis thaliana</i>					
Ino80	Complex	INO80	INO80	INO80					
	ATPase	Ino80	hINO80	INO80 (AT3G57300)					
	Orthologous subunits	Rvb1, Rvb2	TIP49a	TIP49a, TIP49b	RIN1 (AT5G22330), AT3G49830				
		Arp4, Arp5, Arp8, Act1	BAF53a, ARP5, ARP8		ARP4 (AT1G18450), ARP5 (AT3G12380), ARP9 (AT5G43500)				
		Taf14			TAF14 (AT2G18000)				
		les2		hIES2	AT2G47350				
		les6		hIES6	AT4G38495				
	Unique	les3-5, Nhp10	Amida, NFRKB, FLJ90652, FLJ20309						
	Swr1 like	Complex	SWR1	SRCAP	TRRAP/Tip60	SWR1	AT2G17930, AT4G36080/HAM1,2		
		ATPase	Swr1	SRCAP	p400	PIE1 (AT3G12810)			
		Orthologous subunits	Rvb1, Rvb2	TIP49a	TIP49b	TIP49a, TIP49b	RIN1 (AT5G22330), AT3G49830	RIN1 (AT5G22330), AT3G49830	
			Arp6, Act1	ARP6	BAF53a, Actin		ARP6 (AT3G33520)		
			Arp4	BAF53a			ARP4 (AT1G18450)	ARP4 (AT1G18450)	
			Yaf9	GAS41	GAS41		GAS41	GAS41	
			Swc4/Eaf2	DMAP1	DMAP1		AT2G47210	AT2G47210	
			Swc2/Vps72	YL-1	YL-1		SWC2 (AT2G36740)	SWC2 (AT2G36740)	
						BRD8/TRCP120			
			H2A, Z, H2B	H2A, Z, H2B			H2A, Z (AT1G52740, AT2G38810, AT3G54560)		
Swc6/Vps71		Zfn-HIT1			SWC6 (AT5G37055)				
				Tip60		HAM1 (AT5G64610), HAM2 (AT5G09740)			
			MRG15		MRG1 (AT4G37280)				
			ING3		ING1 (AT3G24010)				
Unique	Swc3, 5, 7		MRGx, FLJ11730, MRGBP, EPC1, EPC-like	TAF14 (AT2G18000)					
Etl1	Complex	FUN30	SMARCAD1		ETL1				
	ATPase	Fun30	SMARCAD1		ETL1/CHR19 (AT2G02090)				
	Orthologous subunits	not identified	not identified						
Mi-2	Complex	no homolog	NurD		PKL				
	ATPase		CHD3/MI-2α CHD3/MI-2β		PKL (AT2G25170)				
	Orthologous subunits			HDAC1		PKR2 (AT4G31900)			
				HDAC2		HDA1 (AT4G38130)			
				RbAp46 or 48		HDA6 (AT5G63110)			
				p66α, p66β		FVE/MSI4 (AT2G19520)			
	Chd1	Complex	CHD1	CHD1		CHR5			
	ATPase	Chd1	CHD1		CHR5 (AT2G13370)				
	Orthologous subunits	monomeric	monomeric		unknown				
A1c1	Complex	no homolog	ALC1		CHR10				
	ATPase		ALC1		CHR10 (AT2G44980)				
	Orthologous subunits		not identified						
Snf2 like	Complex	SWI/SNF	RSC	BAF	PBAF	BRM	CHR12		
	ATPase	Swi2/Snf2	Sth1	BRG1/SMARCA4 or hBRM/SMARCA2	BRG1	BRM (AT2G46020)	ATCHR12 (AT3G06010), AT5G19310		
	Orthologous subunits				BAF250a/ARID1A or BAF250b/ARID1B				
				Rsc1, Rsc2, Rsc4	BAF180, BAF200/ARID2	BAF180, BAF200/ARID2			
			Swi3	Rsc8	BAF155/SMARCC1 and/or BAF170/SMARCC2	BAF155/SMARCC1 and/or BAF170/SMARCC2	ATSWI3A (AT2G47620), ATSWI3B (AT2G33610), ATSWI3C (AT1G21700), CHB3 (AT4G34430)	ATSWI3A (AT2G47620), ATSWI3B (AT2G33610), ATSWI3C (AT1G21700), CHB3 (AT4G34430)	
			Swp73	Rsc6	BAF60a/SMARCD1	BAF60a/SMARCD1	BAF60/CHC1 (AT5G14170), AT3G01890	BAF60/CHC1 (AT5G14170), AT3G01890	
			Arp7, Arp9	Arp7, Arp9	BAF53a	BAF53a	ARP4 (AT1G18450)	ARP4 (AT1G18450)	
			Snf5	Sth1	BAF47/hSNF5/SMARCB1	BAF47/hSNF5/SMARCB1	BSH (AT3G17590)	BSH (AT3G17590)	
					BAF57/SMARCE1	BAF57/SMARCE1			
	Unique	Swi1/Adr6, Swp82, Taf14, Snf6, Snf11	Rsc3, Rsc5, Rsc7, Rsc9, Rsc10, Rsc30, Htt1, Lbd7, Rtt102	BAF45a or BAF45d					
	Iswi	Complex	ISW1a	ISW1b	ISW2	ACF	CHRAC	NURF	ISW2
		ATPase	Isw1	Isw1	Isw2	hSNF2H		hSNF2L	CHR11 (AT3G06400), CHR17 (AT5G18620)
Orthologous subunits				Itc1	WCRF180/hACF1	WCRF180/hACF1	BPTF		
				Dpb4		hCHRAC17	RbAp46	MSI4 (AT2G19520)	
Unique		loc3	loc2, loc4	Dis		hCHRAC15	RbAp48	MSI4 (AT2G19520)	
Rad54 like	Complex	Rad54		Rad54				RAD54	
	ATPase	Rad54		Rad54				ATRAD54 (AT3G19210)	

**Figure 2.** Overview of DNA damage-related chromatin-remodeling complexes. The data for the budding yeast and human complexes from Seeber et al. (2013) were complemented by orthologous complexes, subunits, or genes in Arabidopsis. Acronyms are followed by Arabidopsis Genome Initiative gene identity codes in parentheses; genes without acronyms have only their Arabidopsis Genome Initiative code. Green background, experimental evidence for a role in plant DDR; yellow background, connection with chromatin; gray background, only sequence similarity to yeast or human genes necessary for DDR.

Downloaded from https://academic.oup.com/phys/article/168/4/1206/6113912 by guest on 20 August 2022

(Kandasamy et al., 2009; Zhang et al., 2015). Loss of INO80, the ATP-dependent helicase of the complex, results in impairment of HR under standard growth conditions, but, in contrast to the SWR1 mutants, HR is increased after induced DNA damage (Fritsch et al., 2004; Zhang et al., 2015). Transfer DNA integration, which relies upon illegitimate repair pathways such as NHEJ, is not affected (Fritsch et al., 2004). Loss of INO80 increases sensitivity to genotoxic treatments (Zhang et al., 2015), and also Arabidopsis mutants lacking the INO80 subunit ARP5, which is necessary for H2A.Z removal in yeast (Yen et al., 2013), are strongly affected by DNA-damaging agents (Kandasamy et al., 2009). Other plant INO80 subunits have not been challenged for their role in DDR.

Yeast and mammalian components of a third SWR1-like remodeler subfamily with a role in strand resection at DSBs, FUN30 and SMARCAD1, respectively, share the ETL1 complex like the Arabidopsis equivalent. Its ATPase CHR19 was identified as an interactor of SUVR2 that is involved in transcriptional silencing (Han et al., 2014), but a link to DDR has yet to be described.

Snf2-like complexes are the second large group of remodelers; however, their connection to DNA repair is less well established. The human NurD complex is not tethered to, but rather removed from, sites of lesions (Goodarzi et al., 2011), a response that may allow access for other complexes. Several subunits have equivalents in Arabidopsis, with prominent chromatin-interacting (PKL, PKR2, and FVE/MULTICOPY SUPPRESSOR OF IRA1 4 [MSI4]) and chromatin-modifying (HDA1 and HDA6) components. While central roles in developmental regulation (determination of cell identity, flowering time regulation, and several pleiotropic effects) are well documented (Ogas et al., 1999; Tian and Chen, 2001; Ausín et al., 2004; Kim et al., 2004; Aichinger et al., 2009; Hu et al., 2014), no repair-related function has been described. Interaction of the mammalian CHD1 and ALC1 complexes is positively correlated with DNA damage (for review, see Seeber et al., 2013), and their ATPase subunits are encoded as *CHR5* and *CHR10* in Arabidopsis. Again, only a developmentally relevant role is known (Shen et al., 2015). The same holds true for orthologs of the Snf2 complex subunits. They form two complexes regulating plant development, differing in their ATPase subunits BRM and CHR12. While the mammalian equivalents have a clear role in genome stability, especially in the context of cancer genesis (Wilson and Roberts, 2011), their involvement in DNA repair in plants remains to be investigated; the same holds true for all other subunits of the complexes. Among these subunits, BAF60 has a potential role in repair, as it is involved in the formation of DNA loops at the *FLC* locus controlling flowering time (Jégu et al., 2014). Such structural arrangements could also be important during repair processes, especially during intrachromosomal recombination. The only SNF2 family member investigated outside Arabidopsis is the ATPase ALT1 in rice (*Oryza sativa*), which regulates

alkaline tolerance through the modulation of genes involved in reactive oxygen species metabolism and DNA repair. In spite of this rather indirect involvement in DDR, it is interesting that the same nucleosome remodeler acts in both ROS and DNA damage response, strengthening the link between these two stresses (Guo et al., 2014).

The ISWI family of remodelers has a prominent role in several pathways of DNA repair in mammals (for review, see Aydin et al., 2014). The human ATPase SMARCA5/SNF2H is targeted by PARP1 upon activation of DDR. CHR11 and CHR17 are the Arabidopsis orthologs, for which a function in plant development and chromatin remodeling is documented (Li et al., 2012b, 2014), although a direct link with DNA repair needs to be investigated. MSI4 of Arabidopsis is related to subunits of the CHRAC complex that is involved in DDR in *Drosophila melanogaster* and mammals (Lan et al., 2010; Mathew et al., 2014) and is a potential target of the repair-related kinases ATM/ATR (see below).

RAD54, another Snf2-related chromatin remodeler with a prominent role in HR, is so well conserved that the yeast and Arabidopsis genes can reciprocally complement mutants with regard to their DNA damage sensitivity (Klutstein et al., 2008), and overexpression of the yeast gene enhances gene targeting in Arabidopsis (Shaked et al., 2005). More recently, RAD54 was specified to participate mainly in the synthesis-dependent strand-annealing mechanism of DSB repair by HR (Roth et al., 2012).

Chromatin organization also affects DNA repair beyond the nucleosome level. Genes encoding STRUCTURAL MAINTENANCE OF CHROMOSOMES (SMC) complexes in Arabidopsis, namely *AtRAD18*, *AtRAD21.1*, and *AtRAD21.3*, are transcriptionally activated in response to DNA damage to increase the pool of cohesins in the nucleus. When mutated, the number of DSBs increases prior to and after DNA damage, and repair is severely delayed. Additional loss of KU80, a component of NHEJ, exacerbates this repair deficiency, suggesting a role of SMCs during early phases of DSB repair, likely for initiating de novo cohesion between sister chromatids to facilitate HR (Kozak et al., 2009; da Costa-Nunes et al., 2014). AtMMS21, interacting with SMC5, participates in DNA damage reduction in the root stem cell niche to preserve genome integrity (Xu et al., 2013).

RECQ helicases take part in the resolution of Holliday junctions that arise during HR or branch migration during replication. They have multiple functions, reflected in their representation by seven genes in Arabidopsis. AtRECQ4A counteracts recombination during the DNA damage response, while AtRECQ4B is involved in HR, unique among RECQ helicases (Knoll and Puchta, 2011). The AtRTEL1 helicase disrupts the D-loop structures arising from HR recombination intermediates or stalled replication forks to prevent inappropriate recombination between chromosome ends (Recker et al., 2014).

MORC2 ATPases contribute to the DNA damage response in human cells. Their nucleosome-remodeling activity at damage sites is triggered by PAK1-dependent phosphorylation, resulting in a subsequent decondensation of chromatin that provides accessibility to the lesion (Li et al., 2012a). Although MORC orthologs are present in Arabidopsis, their only described role is to establish heterochromatin regions (Lorković, 2012; Moissiard et al., 2012).

While not necessarily a remodeler, the REPLICATION PROTEIN A (RPA) complex is associated with DDR in several groups of organisms. Arabidopsis and rice have multiple genes for the three subunit types, and mutants of the RPA1 types show different DNA sensitivity phenotypes, indicating specialization and redundancy at the same time (Chang et al., 2009; Aklilu et al., 2014). *RPA2* mutants are affected in crossover formation in rice (Li et al., 2013). The fact that RPA2 can be phosphorylated by kinases involved in both cell cycle and DNA damage signaling, as well as its epigenetic control function in gene silencing (Elmayan et al., 2005; Kapoor et al., 2005), make it a very interesting central component of the regulation and integration of several cellular processes. The same could be true for two other replication-associated proteins, the catalytic subunit of DNA polymerase- $\alpha$  and DNA replication factor C1, which upon mutation result in DNA damage sensitivity (Liu et al., 2010a, 2010b).

### HISTONES AND HISTONE CHAPERONES CONNECTED WITH DDR

The basic nucleosome composition of  $4 \times 2$  subunits and the sequences of the most abundant canonical histones are highly conserved across all organisms, but several more diverse histone variants (and histone modifications; see next paragraph) can modulate the nucleosome interaction with DNA and other proteins. Prior to their integration into chromatin, histones associate with chaperone complexes that prevent uncontrolled assembly or association and deliver them to the DNA. The mobility of nucleosomes along DNA, and transport of their building blocks to and from DNA, are also connected with the repair of DNA lesions (for review, see Ransom et al., 2010; Biterge and Schneider, 2014).

Among all histones, the H2A family is most diverse and expanded in plants (for review, see Kawashima et al., 2015), and several variants are demonstrated or assumed to be connected to DDR. H2A.X is a variant of the H2A subunit conserved between yeast, mammals, and plants. It is discussed below due to its phosphorylation during DNA damage signaling. Another H2A variant, H2A.Z, was already mentioned in connection with the SWR1 complex, but its involvement in loading repair proteins was so far shown only for mammalian cells (Xu et al., 2012b). It is not yet clear whether the role of SWR1 in DSB repair via HR is coupled to its role in H2A.Z incorporation (Rosa et al., 2013). MacroH2A variants, associated with repressive

chromatin in mammals, are incorporated at DSBs in a PARP1-dependent manner (Xu et al., 2012a). MacroH2A is not present in plants, but it shares a conserved SPKK motif with another, plant-specific, variant, H2A.W, that is associated with heterochromatin in Arabidopsis (Yelagandula et al., 2014). Whether H2A.W is connected with repair is not yet known, but modification by PARP1 is also repair relevant in Arabidopsis (Jia et al., 2013). Histone H3 has two major variants in plants, H3.1 and H3.3, which are associated with different transcriptional activities. Although H3 chaperones have been shown to have a role in regulating DDR efficiency (see below), it is difficult to determine whether this role is mediated by the chaperone or by H3 directly. The special features of the H3 variant at centromeric repeats, Cse4 in budding yeast, CENP-A in mammals, and CenH3 in plants, likely necessitate an adapted repair process for lesions in these chromosomal regions. Again, the complex interaction of diverse centromeric proteins and their role in DNA repair and recombination (for review, see Osman and Whitby, 2013) make it difficult to dissect the involvement of individual components in all systems, including plants.

Besides the histone subunits within the nucleosomes, the presence, absence, and modification of linker histones are additional factors expected to influence DNA repair processes. A specific variant of H1 was connected with DSB repair via HR in chicken cells (Hashimoto et al., 2007). Arabidopsis has three H1 genes, and the role of H1 in chromatin dynamics is well documented (for review, see Over and Michaels, 2014); however, its role during DDR remains to be specified.

Although histones are by far the most abundant chromatin proteins, a group of other mostly chromatin-associated proteins should not be neglected: the HIGH MOBILITY GROUP (HMG) proteins, which are characterized by a common motif, the HMG box. Their multiplicity and variation along with partially contradictory experimental evidence impede a precise description of their involvement in DNA repair in mammalian cells (for review, see Stros, 2010). They also represent an interesting and diverse gene family in plants and, among many other functions, participate in recombination (Grasser et al., 2007; Antosch et al., 2012). Their mechanistic contribution to DDR deserves further investigation.

Besides the proteins described above, which are more or less tightly associated with DNA at all times, there are several complexes serving as chaperones that transport these proteins to and from the DNA. CHROMATIN ASSEMBLY FACTOR1 (CAF-1) is an H3/H4 chaperone with a major role in nucleosome assembly during replication. An active role for nucleosome dissociation prior to repair has not been documented, but the expression of its components is induced by genotoxic stress. The complex might serve as an acceptor for disassembled subunits, and it is clearly necessary for reassembly after repair in a manner similar to that during replication (for review, see Ransom et al., 2010). The CAF-1 complex is conserved in plants, and its impairment causes multiple

phenotypes, including DDR defects. Mutants in the *FAS1* gene, encoding the largest CAF-1 subunit, display increased rates of HR (Kirik et al., 2006) and transfer DNA integration (Endo et al., 2006), elevated expression levels of ATM-dependent repair genes (Hisanaga et al., 2013), shorter telomeres, and a reduced number of ribosomal DNA copies (Muchová et al., 2014). Some of these phenotypes were also found in *fas2* mutants, devoid of the middle-sized CAF-1 subunit. MSI-1, the small subunit, is a member of several other complexes as well; therefore, the severe mutant phenotypes it displays are difficult to study. The interaction of CAF-1 with RecQ helicases in human cells (for review, see Ransom et al., 2010) could also be relevant in plants, as several RecQ family members in Arabidopsis and rice have been connected to genotoxic stress resistance and DDR repair pathway choice (Knoll and Puchta, 2011; Kwon et al., 2012, 2013; Knoll et al., 2014b; Recker et al., 2014; Schröpfer et al., 2014). HISTONE REGULATOR A (HIRA), another H3/H4 chaperone, acts mainly independent of replication and was recently shown to deposit nucleosome subunits at sites of DNA repair after UV-C light-induced damage in mammalian cells (Adam et al., 2013). All known HIRA subunits are conserved in plants (Nie et al., 2014), and the complex is necessary for regular nucleosome loading in Arabidopsis (Duc et al., 2015). Its link with plant DNA repair needs to be established. Another histone chaperone termed ANTI-SILENCING FUNCTION1 (ASF1), which works upstream of both major H3/H4 chaperones, is also conserved in plants and represented by two partially redundant variants in Arabidopsis that are involved in replication and cell cycle control (Zhu et al., 2011). ASF1 promotes nucleosome reassembly after UV light irradiation (Lario et al., 2013) and interacts with the histone acetyltransferases HAM1/2, which are also connected with efficient repair (Campi et al., 2012).

The H2A/H2B chaperone complex NUCLEOSOME ASSEMBLY PROTEIN (NAP) has also been identified in Arabidopsis and is important for HR (Gao et al., 2012; Zhou et al., 2015), as are the NAP-related proteins NRP1 and NRP2 (Zhu et al., 2006). The hyporecombinogenic phenotype of NAP mutants, in contrast to increased HR in CAF-1 mutants, points to important mechanistic differences between H3/H4 and H2A/H2B delivery during DNA repair processes. FACILITATES CHROMATIN TRANSCRIPTION is a histone chaperone that also mainly associates with H2A/H2B dimers. From the two human subunits, only SPT16 (but not the HMG box-containing SSRP1) is required for the efficient repair of UV light damage (Dinant et al., 2013; Oliveira et al., 2014). Both subunits have orthologs in Arabidopsis, and these are important for regular development (Lolas et al., 2010) and epigenetic regulation (Ikeda et al., 2011), but a link to DNA damage is not described.

#### CHROMATIN MODIFIERS CONNECTED WITH DDR

DNA can become covalently modified, most prominently in eukaryotes by the methylation of cytosine

residues. Additionally, multiple amino acid residues of all the histone subunits can undergo PTMs, leading to steric or electrostatic changes with impacts on nucleosome/DNA association or on the recruitment of other proteins. The presence or absence of Ser and Thr phosphorylation, Lys acetylation, Lys and Arg methylation, Lys ubiquitylation, biotinylation and sumoylation, as well as poly-ADP-ribosylation at Arg and Glu residues offer a plethora of combinations. PTMs usually define larger chromatin regions associated with different functions, in first approximation between transcriptionally active and open euchromatin and less active and more condensed heterochromatin. Dynamic regulation of histone PTMs is tuned by specific writer and eraser enzymes with antagonistic functions, like histone acetyltransferases and histone deacetylases (HDACs) or histone methyltransferases and histone demethylases (Berr et al., 2011). Selected PTMs have been described in connection with DNA damage (for review, see Méndez-Acuña et al., 2010).

Phosphorylation of Ser-139 in the C-terminal portion of the H2A.X histone variant is strongly correlated to DDR, as it is one of the earliest events occurring after DSB induction. Phosphatidylinositol-3-kinases, such as ATM, ATR, and DNA-PKc, control the formation and propagation of phosphorylated  $\gamma$ H2A.X next to the DSB site, which is essential for DNA damage signaling and repair factor recruitment. Swr1 and Ino80 chromatin-remodeling complexes (discussed above) interact with  $\gamma$ H2A.X through the shared subunit Arp4 (Downs et al., 2004). Similar to yeast and animals, phosphorylation of H2A.X in Arabidopsis occurs early after DSB formation and is a trigger for both HR and NHEJ pathways (Charbonnel et al., 2011). It also interacts with the transcriptional activator E2F, likely in a cell cycle-dependent manner (Lang et al., 2012). H2A and H2A.X in mammalian cells are also targets for monoubiquitylation in the context of DDR. This modification is made at DSBs by PRC1, a ubiquitin ligase and component of the POLYCOMB REPRESSIVE COMPLEX. H2A ubiquitylation contributes to transcriptional repression in the proximity of the lesion and prompts the activation of the RNF8-RNF168-ubiquitin pathway, a key event in the DNA repair cascade (Vissers et al., 2012). Whether PRC1 has a similar role in plants remains to be investigated.

In mammals, ATM is further involved in fine-tuning of the chromatin state at DSBs, as it becomes activated by Tip60, an acetyltransferase that, in turn, is recruited by larger domains of H3K9me3 that are transiently formed around DSBs. The latter involves a complex containing the methyltransferase SUV39h1 and a self-reinforcing loop, due to the recruitment of the heterochromatin-binding proteins HP1 and kap-1, to mark larger domains (Ayrapetov et al., 2014). How these transient sites are distinguished from other regions with more permanent heterochromatin is unclear. The Arabidopsis TIP60 orthologs, HAM1 and HAM2, regulate developmental gene expression through the acetylation of H4K5 (Xiao et al., 2013), but

their function is also connected to genotoxic stress. The *ham1* and *ham2* mutants accumulate persistent cyclobutane pyrimidine dimers after UV-B light treatment, resulting in increased UV-B light sensitivity (Campi et al., 2012). Recently, MSI4, a component of a histone deacetylase complex, was identified as a target of ATM/ATR kinases in response to DNA damage (Roitinger et al., 2015).

The histone deacetylase SIRTUIN6 (SIRT6) plays a central role in DNA repair in human cells. SIRT6 has a dual function during early events after DSB formation: enzymatically, it reduces the level of H3K56Ac, and as a scaffold protein, it recruits SNF2H, an example of how PTMs can interact with chromatin remodelers and control access to DSBs (Toiber et al., 2013). Histone acetylation in the context of DDR in plants is both varied and complex. For example, x-ray irradiation of *Arabidopsis* causes H3 hyperacetylation and H4 hypoacetylation, partially dependent on ATM (Drury et al., 2012). In contrast,  $\gamma$ -irradiation of wheat (*Triticum aestivum*) seedlings results in H3 hypoacetylation and H4 hyperacetylation (Raut and Sainis, 2012). Both studies demonstrate the need for further, detailed studies with higher resolution of modification specificity, location, kinetics, dependence on and interaction with other DDR pathways, as well as taking into account evolutionary conservation or species-specific adaptation. Histone methylation or demethylation, which has prominent roles in developmental regulation, has not been directly connected with DNA repair, as the suggested role for the Polycomb group protein CURLY LEAF in somatic recombination may be exerted by the up-regulation of DNA repair genes (Chen et al., 2014).

Methylation of the DNA itself can also have a strong effect on the ability to repair lesions. It has been known for a long time that *Arabidopsis* mutants that are directly or indirectly affected in DNA methylation display a DNA damage-sensitive phenotype (Gong et al., 2002; Elmayan et al., 2005; Shaked et al., 2006; Yao et al., 2012). However, the relation is complex, as it depends on both the type of damage and the type of repair pathway. Repair of UV-B light-induced damage in *repressor of silencing1* (*ros1*), mutated in a gene for a glycosylase that specifically removes methylated cytosines from DNA, is impaired only in the dark, as the negative effect is compensated in light-grown *ros1* mutants by the up-regulation of photorepair-specific genes (Qüesta et al., 2013). Interestingly, the demethylation process results in DNA lesions, as it requires base excision repair to restore the DNA sequence (Lee et al., 2014). The *decreased in DNA methylation1* (*ddm1*) mutant, without an SNF2 chromatin-remodeling factor that is crucial for establishing a correct methylation pattern, is generally affected in the repair of UV light-induced damage (Qüesta et al., 2013). Whether or how these effects are linked to the methylation per se is not yet clear, as the repair function of LSH1, the mammalian protein closest to DDM1, is independent from its role in methylation but rather connected to the

formation of  $\gamma$ H2A.X foci (Burrage et al., 2012). One possibility is that DDM1 and LSH could modulate the accessibility of the lesion, as discussed previously for the other chromatin-remodeling factors.

The many different substrates of CK2, a Ser/Thr kinase, include several chromatin components (Stemmer et al., 2002; Krohn et al., 2003). CK2 is involved in DDR in *Arabidopsis*, as a dominant-negative mutant causes increased sensitivity to DNA-damaging agents and decreased HR efficiency (Moreno-Romero et al., 2012). The mutant plants have additional phenotypes similar to mutants of the histone chaperones NRP1 and NRP2, where chromatin decondensation leads to the overexpression of silenced loci upon genotoxic treatment. Therefore, it is likely that phosphorylation by CK2 controls genetic and epigenetic stability in some still to be investigated way. Although not yet shown to have chromatin modification activity, BRUSHY, a protein that is responsible for genetic stability and epigenetic maintenance in specific parts of the genome, shares the CK2 mutant phenotype of increased sensitivity to DNA damage combined with a release of transcriptional silencing (Takeda et al., 2004; Ohno et al., 2011).

#### THE ROLE OF CHROMATIN DURING PROGRAMMED DSB REPAIR DURING MEIOSIS

In addition to random DNA lesions resulting from adverse conditions, sexual propagation and gamete formation are preceded by numerous programmed DSBs during meiosis. These occur to promote HR between homologous chromosomes. The induction of DSBs is a finely tuned process controlled by different proteins, and chromatin organization is an important parameter for the location and fate of meiotic DSBs. After the synthesis of sister chromatids in the S phase prior to the first meiotic division, chromosomes are arranged in a particular structure consisting of chromatin loops that are anchored in a proteinaceous matrix termed the chromosome axis. SPO11, highly conserved among eukaryotes, catalyzes DSBs on DNA regions exposed by this loop configuration. In yeast, low-density nucleosome regions, such as promoters, are targeted more frequently by SPO11. In mammals, DSBs occur especially in regions marked by histones carrying Set1-dependent H3K4me3, as the chromatin reader PRDM9 recognizes this modification and recruits SPO11, thereby determining the localization of DSBs and recombinational hot spots (for review, see Borde and de Massy, 2013). In *Arabidopsis*, very few of the approximately 200 initial DSBs created by SPO11 per cell are finally converted into crossover events. Their location seems to be determined by a mix of genetic (sequence-determined) factors and chromatin parameters (for review, see Choi and Henderson, 2015). Genome-wide analysis of crossover events in *Arabidopsis* revealed that they are concentrated in regions enriched for CTT-repeat DNA motifs and epigenetic marks, including low nucleosome density,

high H2A.Z and H3K4me3, and low DNA methylation (Choi et al., 2013; Wijnker et al., 2013). The role of H2A.Z is plausible due to its role in the HR repair pathway (Choi et al., 2013). Unexpectedly, however, higher transcription in heterochromatic regions in hypomethylated mutants did not result in an overall higher recombination rate but instead shifted the position of crossovers and the interference between them (Colomé-Tatché et al., 2012; Melamed-Bessudo and Levy, 2012; Mirouze et al., 2012; Yelina et al., 2012). In barley (*Hordeum vulgare*), the borders between euchromatin and heterochromatin appear as preferential recombination sites (Higgins et al., 2012). A recent study in maize (*Zea mays*) revealed that DNA sequence diversity, distance from telomeres, DNA methylation, GC content, and repeat content can predict the location of crossover events with high confidence (Rodgers-Melnick et al., 2015). Therefore, it is likely that recombinational hot spots are determined by many different parameters in a complex interplay between synergistic and competing factors.

## CONCLUSION AND FUTURE CHALLENGES

The configuration of chromatin at the site of a DNA lesion and the factors that shape this configuration are important for the efficiency, and likely also for the outcome, of the DDR process (Aymard et al., 2014; Burgess et al., 2014). Reciprocally, chromatin organization might influence the frequency and type of damage along the genome. Besides the role of chromatin in a narrow sense, the involvement of other epigenetic regulators is emerging, such as the participation of coding, noncoding, and small RNAs (Wei et al., 2012; Chen et al., 2013; Keskin et al., 2014). Therefore, genetic and epigenetic (in)stability should be perceived as tightly interconnected in all organisms, including plants. It will remain difficult to distinguish whether apparently heritable epigenetic changes occurring after DNA damage (Mueller-Xing et al., 2014) are indeed independent from the accompanying genetic modifications, but the idea of stress-induced long-term plant adaption without genome sequence modification is widely discussed (for review, see Becker and Weigel, 2012; Pecinka and Mittelsten Scheid, 2012). It should be kept in mind that, especially for plants outside of well-controlled laboratory conditions, in natural habitats DNA-damaging conditions like UV light irradiation are usually connected with high light intensity, high temperature, or oxidizing conditions. These additional stress factors can directly or indirectly modify chromatin configurations and make the interplay between genetic and epigenetic effects even more complex (Turner and Caspari, 2014). Besides the role that chromatin features play in the repair of randomly occurring DNA damage, the emerging potential of genome-editing procedures (Cantos et al., 2014; Baltes and Voytas, 2015) might be enhanced, if the organization and accessibility of the target DNA in its chromatin context are considered.

The combined application of tools to create targeted lesions together with sequence-specific chromatin remodelers or modifiers might further improve the precision and efficiency of gene-replacement attempts.

## Supplemental Data

The following supplemental materials are available.

**Supplemental Text S1.** Expansions of all abbreviated gene and protein complex names and other abbreviations.

## ACKNOWLEDGMENTS

Due to the nature of an Update, we had to omit many other relevant publications, and we apologize to authors who might not find their work directly cited here. We thank J. Matthew Watson for careful editing.

Received April 10, 2015; accepted June 17, 2015; published June 18, 2015.

## LITERATURE CITED

- Adam S, Polo SE, Almouzni G (2013) Transcription recovery after DNA damage requires chromatin priming by the H3.3 histone chaperone HIRA. *Cell* **155**: 94–106
- Aichinger E, Villar CBR, Farrona S, Reyes JC, Hennig L, Köhler C (2009) CHD3 proteins and polycomb group proteins antagonistically determine cell identity in Arabidopsis. *PLoS Genet* **5**: e1000605
- Aklilu BB, Soderquist RS, Culligan KM (2014) Genetic analysis of the Replication Protein A large subunit family in Arabidopsis reveals unique and overlapping roles in DNA repair, meiosis and DNA replication. *Nucleic Acids Res* **42**: 3104–3118
- Altmeyer M, Lukas J (2013) To spread or not to spread: chromatin modifications in response to DNA damage. *Curr Opin Genet Dev* **23**: 156–165
- Antosch M, Mortensen SA, Grasser KD (2012) Plant proteins containing high mobility group box DNA-binding domains modulate different nuclear processes. *Plant Physiol* **159**: 875–883
- Ausín I, Alonso-Blanco C, Jarillo JA, Ruiz-García L, Martínez-Zapater JM (2004) Regulation of flowering time by FVE, a retinoblastoma-associated protein. *Nat Genet* **36**: 162–166
- Aydin OZ, Vermeulen W, Lans H (2014) ISWI chromatin remodeling complexes in the DNA damage response. *Cell Cycle* **13**: 3016–3025
- Aymard F, Bugler B, Schmidt CK, Guillou E, Caron P, Briois S, Iacovoni JS, Daburon V, Miller KM, Jackson SP, et al (2014) Transcriptionally active chromatin recruits homologous recombination at DNA double-strand breaks. *Nat Struct Mol Biol* **21**: 366–374
- Ayrapetov MK, Gursoy-Yuzugullu O, Xu C, Xu Y, Price BD (2014) DNA double-strand breaks promote methylation of histone H3 on lysine 9 and transient formation of repressive chromatin. *Proc Natl Acad Sci USA* **111**: 9169–9174
- Balestrazzi A, Confalonieri M, Macovei A, Donà M, Carbonera D (2011) Genotoxic stress and DNA repair in plants: emerging functions and tools for improving crop productivity. *Plant Cell Rep* **30**: 287–295
- Baltes NJ, Voytas DF (2015) Enabling plant synthetic biology through genome engineering. *Trends Biotechnol* **33**: 120–131
- Becker C, Weigel D (2012) Epigenetic variation: origin and transgenerational inheritance. *Curr Opin Plant Biol* **15**: 562–567
- Berr A, Shafiq S, Shen WH (2011) Histone modifications in transcriptional activation during plant development. *Biochim Biophys Acta* **1809**: 567–576
- Biterge B, Schneider R (2014) Histone variants: key players of chromatin. *Cell Tissue Res* **356**: 457–466
- Borde V, de Massy B (2013) Programmed induction of DNA double strand breaks during meiosis: setting up communication between DNA and the chromosome structure. *Curr Opin Genet Dev* **23**: 147–155
- Burgess RC, Burman B, Kruhlak MJ, Misteli T (2014) Activation of DNA damage response signaling by condensed chromatin. *Cell Rep* **9**: 1703–1717

- Burrage J, Termanis A, Geissner A, Myant K, Gordon K, Stancheva I (2012) The SNF2 family ATPase LSH promotes phosphorylation of H2AX and efficient repair of DNA double-strand breaks in mammalian cells. *J Cell Sci* **125**: 5524–5534
- Campi M, D'Andrea L, Emiliani J, Casati P (2012) Participation of chromatin-remodeling proteins in the repair of ultraviolet-B-damaged DNA. *Plant Physiol* **158**: 981–995
- Cantos C, Francisco P, Trijatmiko KR, Slamet-Loedin I, Chadha-Mohanty PK (2014) Identification of “safe harbor” loci in indica rice genome by harnessing the property of zinc-finger nucleases to induce DNA damage and repair. *Front Plant Sci* **5**: 302
- Chang Y, Gong L, Yuan W, Li X, Chen G, Li X, Zhang Q, Wu C (2009) Replication protein A (RPA1a) is required for meiotic and somatic DNA repair but is dispensable for DNA replication and homologous recombination in rice. *Plant Physiol* **151**: 2162–2173
- Charbonnel C, Allain E, Gallego ME, White CI (2011) Kinetic analysis of DNA double-strand break repair pathways in Arabidopsis. *DNA Repair (Amst)* **10**: 611–619
- Chen H, Kobayashi K, Miyao A, Hirochika H, Yamaoka N, Nishiguchi M (2013) Both OsRecQ1 and OsRDR1 are required for the production of small RNA in response to DNA-damage in rice. *PLoS ONE* **8**: e55252
- Chen N, Zhou WB, Wang YX, Dong AW, Yu Y (2014) Polycomb-group histone methyltransferase CLF is required for proper somatic recombination in Arabidopsis. *J Integr Plant Biol* **56**: 550–558
- Choi K, Henderson IR (2015) Meiotic recombination hotspots: a comparative view. *Plant J* **83**: 52–61
- Choi K, Kim S, Kim SY, Kim M, Hyun Y, Lee H, Choe S, Kim SG, Michaels S, Lee I (2005) SUPPRESSOR OF FRIGIDA3 encodes a nuclear ACTIN-RELATED PROTEIN6 required for floral repression in Arabidopsis. *Plant Cell* **17**: 2647–2660
- Choi K, Zhao X, Kelly KA, Venn O, Higgins JD, Yelina NE, Hardcastle TJ, Ziolkowski PA, Copenhaver GP, Franklin FCH, et al (2013) Arabidopsis meiotic crossover hot spots overlap with H2A.Z nucleosomes at gene promoters. *Nat Genet* **45**: 1327–1336
- Clapier CR, Cairns BR (2009) The biology of chromatin remodeling complexes. *Annu Rev Biochem* **78**: 273–304
- Colomé-Tatché M, Cortijo S, Wardenaar R, Morgado L, Lahouze B, Sarazin A, Etcheverry M, Martin A, Feng S, Duvernois-Berthet E, et al (2012) Features of the Arabidopsis recombination landscape resulting from the combined loss of sequence variation and DNA methylation. *Proc Natl Acad Sci USA* **109**: 16240–16245
- Courilleau C, Chailleux C, Jauneau A, Grimal F, Briois S, Boutet-Robinet E, Boudsocq F, Trouche D, Canitrot Y (2012) The chromatin remodeler p400 ATPase facilitates Rad51-mediated repair of DNA double-strand breaks. *J Cell Biol* **199**: 1067–1081
- Czaja W, Mao P, Smerdon MJ (2012) The emerging roles of ATP-dependent chromatin remodeling enzymes in nucleotide excision repair. *Int J Mol Sci* **13**: 11954–11973
- da Costa-Nunes JA, Capitão C, Kozak J, Costa-Nunes P, Ducasa GM, Pontes O, Angelis KJ (2014) The AtRAD21.1 and AtRAD21.3 Arabidopsis cohesins play a synergistic role in somatic DNA double strand break damage repair. *BMC Plant Biol* **14**: 353
- Deal RB, Kandasamy MK, McKinney EC, Meagher RB (2005) The nuclear actin-related protein ARP6 is a pleiotropic developmental regulator required for the maintenance of *FLOWERING LOCUS C* expression and repression of flowering in Arabidopsis. *Plant Cell* **17**: 2633–2646
- Dinant C, Ampatzidis-Michailidis G, Lans H, Tresini M, Lagarou A, Grosbart M, Theil AF, van Cappellen WA, Kimura H, Bartek J, et al (2013) Enhanced chromatin dynamics by FACT promotes transcriptional restart after UV-induced DNA damage. *Mol Cell* **51**: 469–479
- Dinant C, Bartek J, Bekker-Jensen S (2012) Histone displacement during nucleotide excision repair. *Int J Mol Sci* **13**: 13322–13337
- Dion V, Gasser SM (2013) Chromatin movement in the maintenance of genome stability. *Cell* **152**: 1355–1364
- Donà M, Macovei A, Faè M, Carbonera D, Balestrazzi A (2013) Plant hormone signaling and modulation of DNA repair under stressful conditions. *Plant Cell Rep* **32**: 1043–1052
- Downs JA, Allard S, Jobin-Robitaille O, Javaheri A, Auger A, Bouchard N, Kron SJ, Jackson SP, Côté J (2004) Binding of chromatin-modifying activities to phosphorylated histone H2A at DNA damage sites. *Mol Cell* **16**: 979–990
- Drury GE, Dowle AA, Ashford DA, Waterworth WM, Thomas J, West CE (2012) Dynamics of plant histone modifications in response to DNA damage. *Biochem J* **445**: 393–401
- Duc C, Benoit M, Le Goff S, Simon L, Poulet A, Cotterell S, Tatout C, Probst AV (2015) The histone chaperone complex HIR maintains nucleosome occupancy and counterbalances impaired histone deposition in CAF-1 complex mutants. *Plant J* **81**: 707–722
- Elmayan T, Proux F, Vaucheret H (2005) Arabidopsis RPA2: a genetic link among transcriptional gene silencing, DNA repair, and DNA replication. *Curr Biol* **15**: 1919–1925
- Endo M, Ishikawa Y, Osakabe K, Nakayama S, Kaya H, Araki T, Shibahara K, Abe K, Ichikawa H, Valentine L, et al (2006) Increased frequency of homologous recombination and T-DNA integration in Arabidopsis CAF-1 mutants. *EMBO J* **25**: 5579–5590
- Euskirchen G, Auerbach RK, Snyder M (2012) SWI/SNF chromatin-remodeling factors: multiscale analyses and diverse functions. *J Biol Chem* **287**: 30897–30905
- Fritsch O, Benvenuto G, Bowler C, Molinier J, Hohn B (2004) The INO80 protein controls homologous recombination in Arabidopsis thaliana. *Mol Cell* **16**: 479–485
- Gao J, Zhu Y, Zhou W, Molinier J, Dong A, Shen WH (2012) NAP1 family histone chaperones are required for somatic homologous recombination in Arabidopsis. *Plant Cell* **24**: 1437–1447
- Gerhold CB, Gasser SM (2014) INO80 and SWR complexes: relating structure to function in chromatin remodeling. *Trends Cell Biol* **24**: 619–631
- Gong Z, Morales-Ruiz T, Ariza RR, Roldán-Arjona T, David L, Zhu JK (2002) ROS1, a repressor of transcriptional gene silencing in Arabidopsis, encodes a DNA glycosylase/lyase. *Cell* **111**: 803–814
- Goodarzi AA, Kurka T, Jeggo PA (2011) KAP-1 phosphorylation regulates CHD3 nucleosome remodeling during the DNA double-strand break response. *Nat Struct Mol Biol* **18**: 831–839
- Gospodinov A, Herceg Z (2013a) Chromatin structure in double strand break repair. *DNA Repair (Amst)* **12**: 800–810
- Gospodinov A, Herceg Z (2013b) Shaping chromatin for repair. *Mutat Res* **752**: 45–60
- Grasser KD, Launholt D, Grasser M (2007) High mobility group proteins of the plant HMGB family: dynamic chromatin modulators. *Biochim Biophys Acta* **1769**: 346–357
- Guo M, Wang R, Wang J, Hua K, Wang Y, Liu X, Yao S (2014) ALT1, a Snf2 family chromatin remodeling ATPase, negatively regulates alkaline tolerance through enhanced defense against oxidative stress in rice. *PLoS ONE* **9**: e112515
- Han YF, Dou K, Ma ZY, Zhang SW, Huang HW, Li L, Cai T, Chen S, Zhu JK, He XJ (2014) SUV2 is involved in transcriptional gene silencing by associating with SNF2-related chromatin-remodeling proteins in Arabidopsis. *Cell Res* **24**: 1445–1465
- Hashimoto H, Sonoda E, Takami Y, Kimura H, Nakayama T, Tachibana M, Takeda S, Shinkai Y (2007) Histone H1 variant, H1R is involved in DNA damage response. *DNA Repair (Amst)* **6**: 1584–1595
- Higgins JD, Perry RM, Barakate A, Ramsay L, Waugh R, Halpin C, Armstrong SJ, Franklin FCH (2012) Spatiotemporal asymmetry of the meiotic program underlies the predominantly distal distribution of meiotic crossovers in barley. *Plant Cell* **24**: 4096–4109
- Hisanaga T, Ferjani A, Horiguchi G, Ishikawa N, Fujikura U, Kubo M, Demura T, Fukuda H, Ishida T, Sugimoto K, et al (2013) The ATM-dependent DNA damage response acts as an upstream trigger for compensation in the *fas1* mutation during Arabidopsis leaf development. *Plant Physiol* **162**: 831–841
- Hopfner KP, Gerhold CB, Lakomek K, Wollmann P (2012) Swi2/Snf2 remodelers: hybrid views on hybrid molecular machines. *Curr Opin Struct Biol* **22**: 225–233
- Horigome C, Oma Y, Konishi T, Schmid R, Marcomini I, Hauer MH, Dion V, Harata M, Gasser SM (2014) SWR1 and INO80 chromatin remodelers contribute to DNA double-strand break perinuclear anchorage site choice. *Mol Cell* **55**: 626–639
- House NCM, Koch MR, Freudenreich CH (2014) Chromatin modifications and DNA repair: beyond double-strand breaks. *Front Genet* **5**: 296
- Hu Y, Lai Y, Zhu D (2014) Transcription regulation by CHD proteins to control plant development. *Front Plant Sci* **5**: 223
- Ikeda Y, Kinoshita Y, Susaki D, Ikeda Y, Iwano M, Takayama S, Higashiyama T, Kakutani T, Kinoshita T (2011) HMG domain containing SSRP1 is required for DNA demethylation and genomic imprinting in Arabidopsis. *Dev Cell* **21**: 589–596
- Jasin M, Rothstein R (2013) Repair of strand breaks by homologous recombination. *Cold Spring Harb Perspect Biol* **5**: a012740

- Jeggo PA, Downs JA (2014) Roles of chromatin remodellers in DNA double strand break repair. *Exp Cell Res* **329**: 69–77
- Jégu T, Latrasse D, Delarue M, Hirt H, Domenichini S, Ariel F, Crespi M, Bergounioux C, Raynaud C, Benhamed M (2014) The BAF60 subunit of the SWI/SNF chromatin-remodeling complex directly controls the formation of a gene loop at *FLOWERING LOCUS C* in *Arabidopsis*. *Plant Cell* **26**: 538–551
- Jia Q, den Dulk-Ras A, Shen H, Hooykaas PJJ, de Pater S (2013) Poly (ADP-ribose) polymerases are involved in microhomology mediated back-up non-homologous end joining in *Arabidopsis thaliana*. *Plant Mol Biol* **82**: 339–351
- Kandasamy MK, McKinney EC, Deal RB, Smith AP, Meagher RB (2009) *Arabidopsis* actin-related protein ARP5 in multicellular development and DNA repair. *Dev Biol* **335**: 22–32
- Kapoor A, Agarwal M, Andreucci A, Zheng X, Gong Z, Hasegawa PM, Bressan RA, Zhu JK (2005) Mutations in a conserved replication protein suppress transcriptional gene silencing in a DNA-methylation-independent manner in *Arabidopsis*. *Curr Biol* **15**: 1912–1918
- Kawashima T, Lorković Z, Nishihama R, Ishizaki K, Axelsson E, Yelagandula R, Kohchi T, Berger F (2015) Diversification of histone H2A variants during plant evolution. *Trends Plant Sci pii*: S1360–S1385
- Keskin H, Shen Y, Huang F, Patel M, Yang T, Ashley K, Mazin AV, Storic F (2014) Transcript-RNA-templated DNA recombination and repair. *Nature* **515**: 436–439
- Kim HJ, Hyun Y, Park JY, Park MJ, Park MK, Kim MD, Kim HJ, Lee MH, Moon J, Lee I, et al (2004) A genetic link between cold responses and flowering time through FVE in *Arabidopsis thaliana*. *Nat Genet* **36**: 167–171
- Kirik A, Pecinka A, Wendeler E, Reiss B (2006) The chromatin assembly factor subunit FASCIATA1 is involved in homologous recombination in plants. *Plant Cell* **18**: 2431–2442
- Klutstein M, Shaked H, Sherman A, Avivi-Ragolsky N, Shema E, Zenvirth D, Levy AA, Simchen G (2008) Functional conservation of the yeast and *Arabidopsis* RAD54-like genes. *Genetics* **178**: 2389–2397
- Knizewski L, Ginalski K, Jerzmanowski A (2008) Snf2 proteins in plants: gene silencing and beyond. *Trends Plant Sci* **13**: 557–565
- Knoll A, Fauser F, Puchta H (2014a) DNA recombination in somatic plant cells: mechanisms and evolutionary consequences. *Chromosome Res* **22**: 191–201
- Knoll A, Puchta H (2011) The role of DNA helicases and their interaction partners in genome stability and meiotic recombination in plants. *J Exp Bot* **62**: 1565–1579
- Knoll A, Schröpfer S, Puchta H (2014b) The RTR complex as caretaker of genome stability and its unique meiotic function in plants. *Front Plant Sci* **5**: 33
- Kozak J, West CE, White C, da Costa-Nunes JA, Angelis KJ (2009) Rapid repair of DNA double strand breaks in *Arabidopsis thaliana* is dependent on proteins involved in chromosome structure maintenance. *DNA Repair (Amst)* **8**: 413–419
- Krohn NM, Stemmer C, Fojan P, Grimm R, Grasser KD (2003) Protein kinase CK2 phosphorylates the high mobility group domain protein SSRP1, inducing the recognition of UV-damaged DNA. *J Biol Chem* **278**: 12710–12715
- Kwon YI, Abe K, Endo M, Osakabe K, Ohtsuki N, Nishizawa-Yokoi A, Tagiri A, Saika H, Toki S (2013) DNA replication arrest leads to enhanced homologous recombination and cell death in meristems of rice *OsRecQ4* mutants. *BMC Plant Biol* **13**: 62
- Kwon YI, Abe K, Osakabe K, Endo M, Nishizawa-Yokoi A, Saika H, Shimada H, Toki S (2012) Overexpression of *OsRecQ4* and/or *OsExo1* enhances DSB-induced homologous recombination in rice. *Plant Cell Physiol* **53**: 2142–2152
- Lan L, Ui A, Nakajima S, Hatakeyama K, Hoshi M, Watanabe R, Janicki SM, Ogiwara H, Kohno T, Kanno S, et al (2010) The ACF1 complex is required for DNA double-strand break repair in human cells. *Mol Cell* **40**: 976–987
- Lang J, Smetana O, Sanchez-Calderon L, Lincker F, Genestier J, Schmit AC, Houlné G, Chabouté ME (2012) Plant γH2AX foci are required for proper DNA DSB repair responses and colocalize with E2F factors. *New Phytol* **194**: 353–363
- Lans H, Marteiijn JA, Vermeulen W (2012) ATP-dependent chromatin remodeling in the DNA-damage response. *Epigenetics Chromatin* **5**: 4
- Lario LD, Ramirez-Parra E, Gutierrez C, Spampinato CP, Casati P (2013) ANTI-SILENCING FUNCTION1 proteins are involved in ultraviolet-induced DNA damage repair and are cell cycle regulated by E2F transcription factors in *Arabidopsis*. *Plant Physiol* **162**: 1164–1177
- Lázaro A, Gómez-Zambrano A, López-González L, Piñeiro M, Jarillo JA (2008) Mutations in the *Arabidopsis* SWC6 gene, encoding a component of the SWR1 chromatin remodeling complex, accelerate flowering time and alter leaf and flower development. *J Exp Bot* **59**: 653–666
- Lee J, Jang H, Shin H, Choi WL, Mok YG, Huh JH (2014) AP endonucleases process 5-methylcytosine excision intermediates during active DNA demethylation in *Arabidopsis*. *Nucleic Acids Res* **42**: 11408–11418
- Lemaître C, Soutoglou E (2015) DSB (im)mobility and DNA repair compartmentalization in mammalian cells. *J Mol Biol* **427**: 652–658
- Li DQ, Nair SS, Ohshiro K, Kumar A, Nair VS, Pakala SB, Reddy SDN, Gajula RP, Eswaran J, Aravind L, et al (2012a) MORC2 signaling integrates phosphorylation-dependent, ATPase-coupled chromatin remodeling during the DNA damage response. *Cell Rep* **2**: 1657–1669
- Li G, Liu S, Wang J, He J, Huang H, Zhang Y, Xu L (2014) ISWI proteins participate in the genome-wide nucleosome distribution in *Arabidopsis*. *Plant J* **78**: 706–714
- Li G, Zhang J, Li J, Yang Z, Huang H, Xu L (2012b) Imitation Switch chromatin remodeling factors and their interacting RINGLET proteins act together in controlling the plant vegetative phase in *Arabidopsis*. *Plant J* **72**: 261–270
- Li X, Chang Y, Xin X, Zhu C, Li X, Higgins JD, Wu C (2013) Replication protein A2c coupled with replication protein A1c regulates crossover formation during meiosis in rice. *Plant Cell* **25**: 3885–3899
- Liu J, Ren X, Yin H, Wang Y, Xia R, Wang Y, Gong Z (2010a) Mutation in the catalytic subunit of DNA polymerase alpha influences transcriptional gene silencing and homologous recombination in *Arabidopsis*. *Plant J* **61**: 36–45
- Liu Q, Wang J, Mikki D, Xia R, Yu W, He J, Zheng Z, Zhu JK, Gong Z (2010b) DNA replication factor C1 mediates genomic stability and transcriptional gene silencing in *Arabidopsis*. *Plant Cell* **22**: 2336–2352
- Lolas IB, Himanen K, Grönlund JT, Lynggaard C, Houben A, Melzer M, Van Lijsebettens M, Grasser KD (2010) The transcript elongation factor FACT affects *Arabidopsis* vegetative and reproductive development and genetically interacts with HUB1/2. *Plant J* **61**: 686–697
- Lorković ZJ (2012) MORC proteins and epigenetic regulation. *Plant Signal Behav* **7**: 1561–1565
- Lukas J, Lukas C, Bartek J (2011) More than just a focus: the chromatin response to DNA damage and its role in genome integrity maintenance. *Nat Cell Biol* **13**: 1161–1169
- March-Díaz R, Reyes JC (2009) The beauty of being a variant: H2A.Z and the SWR1 complex in plants. *Mol Plant* **2**: 565–577
- Mathew V, Pauleau AL, Steffen N, Bergner A, Becker PB, Erhardt S (2014) The histone-fold protein CHRAC14 influences chromatin composition in response to DNA damage. *Cell Rep* **7**: 321–330
- Melamed-Bessudo C, Levy AA (2012) Deficiency in DNA methylation increases meiotic crossover rates in euchromatic but not in heterochromatic regions in *Arabidopsis*. *Proc Natl Acad Sci USA* **109**: E981–E988
- Méndez-Acuña L, Di Tomaso MV, Palitti F, Martínez-López W (2010) Histone post-translational modifications in DNA damage response. *Cytogenet Genome Res* **128**: 28–36
- Mirouze M, Lieberman-Lazarovich M, Aversano R, Bucher E, Nicolet J, Reinders J, Paszkowski J (2012) Loss of DNA methylation affects the recombination landscape in *Arabidopsis*. *Proc Natl Acad Sci USA* **109**: 5880–5885
- Missirian V, Conklin PA, Culligan KM, Huefner ND, Britt AB (2014) High atomic weight, high-energy radiation (HZE) induces transcriptional responses shared with conventional stresses in addition to a core “DSB” response specific to clastogenic treatments. *Front Plant Sci* **5**: 364
- Moissiard G, Cokus SJ, Cary J, Feng S, Billi AC, Stroud H, Husmann D, Zhan Y, Lajoie BR, McCord RP, et al (2012) MORC family ATPases required for heterochromatin condensation and gene silencing. *Science* **336**: 1448–1451
- Moreno-Romero J, Armengot L, Marqués-Bueno MM, Britt A, Martínez MC (2012) CK2-defective *Arabidopsis* plants exhibit enhanced double-strand break repair rates and reduced survival after exposure to ionizing radiation. *Plant J* **71**: 627–638
- Muchová V, Amiard S, Mozgová I, Dvořáčková M, Gallego ME, White C, Fajkus J (2015) Homology-dependent repair is involved in 45S rDNA loss in plant CAF-1 mutants. *Plant J* **81**: 198–209

- Mueller-Xing R, Xing Q, Goodrich J (2014) Footprints of the sun: memory of UV and light stress in plants. *Front Plant Sci* 5: 474
- Nie X, Wang H, Li J, Holec S, Berger F (2014) The HIRA complex that deposits the histone H3.3 is conserved in Arabidopsis and facilitates transcriptional dynamics. *Biol Open* 3: 794–802
- Noh YS, Amasino RM (2003) *PIE1*, an ISWI family gene, is required for FLC activation and floral repression in Arabidopsis. *Plant Cell* 15: 1671–1682
- Ogas J, Kaufmann S, Henderson J, Somerville C (1999) PICKLE is a CHD3 chromatin-remodeling factor that regulates the transition from embryonic to vegetative development in Arabidopsis. *Proc Natl Acad Sci USA* 96: 13839–13844
- Ohno Y, Narangajavana J, Yamamoto A, Hattori T, Kagaya Y, Paszkowski J, Grussem W, Hennig L, Takeda S (2011) Ectopic gene expression and organogenesis in Arabidopsis mutants missing BRU1 required for genome maintenance. *Genetics* 189: 83–95
- Ohsawa R, Seol JH, Tyler JK (2013) At the intersection of non-coding transcription, DNA repair, chromatin structure, and cellular senescence. *Front Genet* 4: 136
- Oliveira DV, Kato A, Nakamura K, Ikura T, Okada M, Kobayashi J, Yanagihara H, Saito Y, Tsuchi H, Komatsu K (2014) Histone chaperone FACT regulates homologous recombination by chromatin remodeling through interaction with RNF20. *J Cell Sci* 127: 763–772
- Osman F, Whitby MC (2013) Emerging roles for centromere-associated proteins in DNA repair and genetic recombination. *Biochem Soc Trans* 41: 1726–1730
- Over RS, Michaels SD (2014) Open and closed: the roles of linker histones in plants and animals. *Mol Plant* 7: 481–491
- Papamichos-Chronakis M, Peterson CL (2013) Chromatin and the genome integrity network. *Nat Rev Genet* 14: 62–75
- Pecinka A, Mittelsten Scheid O (2012) Stress-induced chromatin changes: a critical view on their heritability. *Plant Cell Physiol* 53: 801–808
- Peterson CL, Almouzni G (2013) Nucleosome dynamics as modular systems that integrate DNA damage and repair. *Cold Spring Harb Perspect Biol* 5: 5
- Polo SE (2015) Reshaping chromatin after DNA damage: the choreography of histone proteins. *J Mol Biol* 427: 626–636
- Price BD, D'Andrea AD (2013) Chromatin remodeling at DNA double-strand breaks. *Cell* 152: 1344–1354
- Qin Y, Zhao L, Skaggs MI, Andruzza S, Tsukamoto T, Panoli A, Wallace KN, Smith S, Siddiqi I, Yang Z, et al (2014) ACTIN-RELATED PROTEIN6 regulates female meiosis by modulating meiotic gene expression in Arabidopsis. *Plant Cell* 26: 1612–1628
- Qüesta JI, Fina JP, Casati P (2013) DDM1 and ROS1 have a role in UV-B induced and oxidative DNA damage in *A. thaliana*. *Front Plant Sci* 4: 420
- Ransom M, Dennehey BK, Tyler JK (2010) Chaperoning histones during DNA replication and repair. *Cell* 140: 183–195
- Raut VV, Sainis JK (2012) <sup>60</sup>Co- $\gamma$  radiation induces differential acetylation and phosphorylation of histones H3 and H4 in wheat. *Plant Biol (Stuttg)* 14: 110–117
- Recker J, Knoll A, Puchta H (2014) The Arabidopsis thaliana homolog of the helicase RTEL1 plays multiple roles in preserving genome stability. *Plant Cell* 26: 4889–4902
- Rodgers-Melnick E, Bradbury PJ, Elshire RJ, Glaubitz JC, Acharya CB, Mitchell SE, Li C, Li Y, Buckler ES (2015) Recombination in diverse maize is stable, predictable, and associated with genetic load. *Proc Natl Acad Sci USA* 112: 3823–3828
- Roitinger E, Hofer M, Köcher T, Pichler P, Novatchkova M, Yang J, Schlögelhofer P, Mechtler K (2015) Quantitative phosphoproteomics of the ataxia telangiectasia-mutated (ATM) and ataxia telangiectasia-mutated and rad3-related (ATR) dependent DNA damage response in Arabidopsis thaliana. *Mol Cell Proteomics* 14: 556–571
- Rosa M, Von Harder M, Cigliano RA, Schlögelhofer P, Mittelsten Scheid O (2013) The Arabidopsis SWR1 chromatin-remodeling complex is important for DNA repair, somatic recombination, and meiosis. *Plant Cell* 25: 1990–2001
- Roth N, Klimesch J, Dukowicz-Schulze S, Pacher M, Mannuss A, Puchta H (2012) The requirement for recombination factors differs considerably between different pathways of homologous double-strand break repair in somatic plant cells. *Plant J* 72: 781–790
- Roy S (2014) Maintenance of genome stability in plants: repairing DNA double strand breaks and chromatin structure stability. *Front Plant Sci* 5: 487
- Schröpfer S, Kobbe D, Hartung F, Knoll A, Puchta H (2014) Defining the roles of the N-terminal region and the helicase activity of RECO4A in DNA repair and homologous recombination in Arabidopsis. *Nucleic Acids Res* 42: 1684–1697
- Seeber A, Hauer M, Gasser SM (2013) Nucleosome remodelers in double-strand break repair. *Curr Opin Genet Dev* 23: 174–184
- Shaked H, Avivi-Ragolsky N, Levy AA (2006) Involvement of the Arabidopsis SWI2/SNF2 chromatin remodeling gene family in DNA damage response and recombination. *Genetics* 173: 985–994
- Shaked H, Melamed-Bessudo C, Levy AA (2005) High-frequency gene targeting in Arabidopsis plants expressing the yeast RAD54 gene. *Proc Natl Acad Sci USA* 102: 12265–12269
- Shen Y, Devic M, Lepiniec L, Zhou D (2015) Chromodomain, helicase and DNA-binding CHD1 protein, CHR5, are involved in establishing active chromatin state of seed maturation genes. *Plant Biotechnol J* 13: 811–820
- Soria G, Polo SE, Almouzni G (2012) Prime, repair, restore: the active role of chromatin in the DNA damage response. *Mol Cell* 46: 722–734
- Stanley FKT, Moore S, Goodarzi AA (2013) CHD chromatin remodelling enzymes and the DNA damage response. *Mutat Res* 750: 31–44
- Stemmer C, Schwander A, Bauw G, Fojan P, Grasser KD (2002) Protein kinase CK2 differentially phosphorylates maize chromosomal high mobility group B (HMGB) proteins modulating their stability and DNA interactions. *J Biol Chem* 277: 1092–1098
- Stros M (2010) HMGB proteins: interactions with DNA and chromatin. *Biochim Biophys Acta* 1799: 101–113
- Swygert SG, Peterson CL (2014) Chromatin dynamics: interplay between remodeling enzymes and histone modifications. *Biochim Biophys Acta* 1839: 728–736
- Takeda S, Tadele Z, Hofmann I, Probst AV, Angelis KJ, Kaya H, Araki T, Mengiste T, Mittelsten Scheid O, Shibahara K, et al (2004) BRU1, a novel link between responses to DNA damage and epigenetic gene silencing in Arabidopsis. *Genes Dev* 18: 782–793
- Tian L, Chen ZJ (2001) Blocking histone deacetylation in Arabidopsis induces pleiotropic effects on plant gene regulation and development. *Proc Natl Acad Sci USA* 98: 200–205
- Toiber D, Erdel F, Bouazoune K, Silberman DM, Zhong L, Mulligan P, Sebastian C, Cosentino C, Martinez-Pastor B, Giacosa S, et al (2013) SIRT6 recruits SNF2H to DNA break sites, preventing genomic instability through chromatin remodeling. *Mol Cell* 51: 454–468
- Tsabar M, Haber JE (2013) Chromatin modifications and chromatin remodeling during DNA repair in budding yeast. *Curr Opin Genet Dev* 23: 166–173
- Turner T, Caspari T (2014) When heat casts a spell on the DNA damage checkpoints. *Open Biol* 4: 140008
- Van C, Williams JS, Kunkel TA, Peterson CL (2015) Deposition of histone H2A.Z by the SWR-C remodeling enzyme prevents genome instability. *DNA Repair (Amst)* 15: 9–14
- Visser JHA, van Luizen M, Citterio E (2012) The emerging role of Polycomb repressors in the response to DNA damage. *J Cell Sci* 125: 3939–3948
- Wei W, Ba Z, Gao M, Wu Y, Ma Y, Amiard S, White CI, Rendtlew Danielsen JM, Yang YG, Qi Y (2012) A role for small RNAs in DNA double-strand break repair. *Cell* 149: 101–112
- Wijnker E, Velikkakam James G, Ding J, Becker F, Klasen JR, Rawat V, Rowan BA, de Jong DF, de Snoo CB, Zapata L, et al (2013) The genomic landscape of meiotic crossovers and gene conversions in Arabidopsis thaliana. *eLife* 2: e01426
- Wilson BG, Roberts CWM (2011) SWI/SNF nucleosome remodellers and cancer. *Nat Rev Cancer* 11: 481–492
- Xiao J, Zhang H, Xing L, Xu S, Liu H, Chong K, Xu Y (2013) Requirement of histone acetyltransferases HAM1 and HAM2 for epigenetic modification of FLC in regulating flowering in Arabidopsis. *J Plant Physiol* 170: 444–451
- Xu C, Xu Y, Gursoy-Yuzugullu O, Price BD (2012a) The histone variant macroH2A1.1 is recruited to DSBs through a mechanism involving PARP1. *FEBS Lett* 586: 3920–3925
- Xu P, Yuan D, Liu M, Li C, Liu Y, Zhang S, Yao N, Yang C (2013) AtMMS21, an SMC5/6 complex subunit, is involved in stem cell niche maintenance and DNA damage responses in Arabidopsis roots. *Plant Physiol* 161: 1755–1768
- Xu Y, Arapetov MK, Xu C, Gursoy-Yuzugullu O, Hu Y, Price BD (2012b) Histone H2A.Z controls a critical chromatin remodeling step required for DNA double-strand break repair. *Mol Cell* 48: 723–733

- Yao Y, Bilichak A, Golubov A, Kovalchuk I** (2012) ddm1 plants are sensitive to methyl methane sulfonate and NaCl stresses and are deficient in DNA repair. *Plant Cell Rep* **31**: 1549–1561
- Yelagandula R, Stroud H, Holec S, Zhou K, Feng S, Zhong X, Muthurajan UM, Nie X, Kawashima T, Groth M, et al** (2014) The histone variant H2A.W defines heterochromatin and promotes chromatin condensation in *Arabidopsis*. *Cell* **158**: 98–109
- Yelina NE, Choi K, Chelysheva L, Macaulay M, de Snoo B, Wijnker E, Miller N, Drouaud J, Grelon M, Copenhaver GP, et al** (2012) Epigenetic remodeling of meiotic crossover frequency in *Arabidopsis thaliana* DNA methyltransferase mutants. *PLoS Genet* **8**: e1002844
- Yen K, Vinayachandran V, Pugh BF** (2013) SWR-C and INO80 chromatin remodelers recognize nucleosome-free regions near +1 nucleosomes. *Cell* **154**: 1246–1256
- Zhang C, Cao L, Rong L, An Z, Zhou W, Ma J, Shen WH, Zhu Y, Dong A** (2015) The chromatin-remodeling factor AtINO80 plays crucial roles in genome stability maintenance and in plant development. *Plant J* **82**: 655–668
- Zhou W, Zhu Y, Dong A, Shen WH** (2015) Histone H2A/H2B chaperones: from molecules to chromatin-based functions in plant growth and development. *Plant J* **83**: 78–95
- Zhu Y, Dong A, Meyer D, Pichon O, Renou JP, Cao K, Shen WH** (2006) *Arabidopsis* *NRP1* and *NRP2* encode histone chaperones and are required for maintaining postembryonic root growth. *Plant Cell* **18**: 2879–2892
- Zhu Y, Dong A, Shen WH** (2013) Histone variants and chromatin assembly in plant abiotic stress responses. *Biochim Biophys Acta* **1819**: 343–348
- Zhu Y, Weng M, Yang Y, Zhang C, Li Z, Shen WH, Dong A** (2011) *Arabidopsis* homologues of the histone chaperone ASF1 are crucial for chromatin replication and cell proliferation in plant development. *Plant J* **66**: 443–455