

Original Research

Cite this article: Carmassi C, Dell'Oste V, Barberi FM, Pedrinelli V, Cordone A, Cappelli A, Cremone IM, Rossi R, Bertelloni CA, and Dell'Osso L (2021). Do somatic symptoms relate to PTSD and gender after earthquake exposure? A cross-sectional study on young adult survivors in Italy. *CNS Spectrums* 26(3), 268–274.

<https://doi.org/10.1017/S1092852920000097>

Received: 18 November 2019

Accepted: 09 January 2020

Key words:


Post-traumatic stress disorder (PTSD); natural disaster; migraine; gastrointestinal diseases; painful stimuli hypo- or hypersensitivity; DSM-5

Author for correspondence:

*Valerio Dell'Oste, MD

Email: valerio.delloste@gmail.com

Do somatic symptoms relate to PTSD and gender after earthquake exposure? A cross-sectional study on young adult survivors in Italy

Claudia Carmassi¹, Valerio Dell'Oste^{1*} , Filippo M. Barberi¹, Virginia Pedrinelli¹, Annalisa Cordone¹, Andrea Cappelli¹, Ivan M. Cremone¹, Rodolfo Rossi², Carlo A. Bertelloni¹ and Liliana Dell'Osso¹

¹Department of Clinical and Experimental Medicine, University of Pisa, Pisa, Italy, and ²Department of Systems Medicine, University of Rome Tor Vergata, Rome, Italy

Abstract

Objective. Increasing evidence confirms a strict relationship between mental disorders and physical health. Particularly, stressful life events and post-traumatic stress disorder (PTSD) have been closely correlated with various physical disorders and somatic symptoms, such as chronic pain, gastrointestinal disorders, and headaches. The aim of this study was to investigate the emergence of somatic symptoms in a sample of young adult survivors 21 months after exposure to the L'Aquila 2009 earthquake, with particular attention to PTSD and gender impact.

Methods. Four hundred and fifty high-school senior students (253 male and 197 female) exposed to the 2009 L'Aquila earthquake, 21 months earlier, were enrolled and evaluated by the Trauma and Loss Spectrum Self-Report (TALS-SR), for symptomatological PTSD, and the Mood Spectrum Self-Report-Lifetime Version (MOODS-SR) “*rhythmicity and vegetative functions*” domain, for somatic symptoms.

Results. Significantly higher rates of endorsement of the MOODS-SR somatic symptoms emerged in survivors with PTSD compared to those without. Females reported higher rates of endorsement of at least one MOODS-SR somatic symptom compared to males; however, a Decision Tree model and a two-way analysis of variance model confirmed a significant effect of PTSD only. A multivariate logistical regression showed a significant association between the presence of at least one MOOD-SR somatic symptom and re-experiencing and maladaptive coping TALS-SR domains.

Conclusion. This study corroborates a relevant impact of symptomatological PTSD, across both the genders, on somatic symptoms occurring in young adults after months from exposure to a massive earthquake.

Introduction

Post-traumatic stress disorder (PTSD) typically arises following exposure, both direct and indirect, to a traumatic event, and is characterized by the onset and persistence of a series of clinical symptoms that can often be profoundly incapacitating and tendentially chronic.¹ There is evidence that PTSD is a chronic, drug-resistant disorder, associated with high suicidal risk (about 25% of patients have attempted suicide), substance abuse, and maladaptive behaviors, that can also be insidious beginning even after years of silence.² The lifetime prevalence of PTSD can vary considerably depending on the populations considered and the year in which the study was carried out for the adopted diagnostic criteria. Nowadays, the prevalence in the general population corresponds to about 8%, that can arise up to even 40% in aftermath of a natural disaster, depending on its severity and impact on exposed populations, with twice or much of the rates in women compared to men.^{3,4}

In recent years, several studies have examined and confirmed the strict relationship between mental disorders and physical health. Suffering from mental disorder, in fact, may determine biochemical and hormonal alterations in the body that can lead to various physical illnesses. In particular, stressful life events and PTSD have been closely correlated with various physical disorders, especially somatic symptoms such as chronic pain, gastrointestinal (GI) disorders, and headache.^{5,6} High rates of somatic symptoms have been reported among war-exposed civilians,^{7,8} as well as in special populations, who regularly exposed to traumatic events because of their employment states, such as military veterans and emergency services personnel.^{6,9,10} Increasing data highlighted the importance to detect full and partial PTSD in general populations exposed to natural disasters in order to provide effective support and treatment, but despite the emerging evidence on a growing use of painkillers in

© The Author(s) 2020. Published by Cambridge University Press. This is an Open Access article, distributed under the terms of the Creative Commons Attribution licence (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted re-use, distribution, and reproduction in any medium, provided the original work is properly cited.

such populations even months after exposure,^{11–13} scant data explored the onset of somatic severe symptoms. The studies available investigated special populations, such as children and adolescents, exposed to natural disasters such as earthquake. In particular, Zhang et al¹⁴ investigated somatic symptoms in 2299 children and adolescent survivors of the Lushan 2013 earthquake with probable PTSD, demonstrating a strong association between PTSD and somatic symptoms.

The exact mechanism underlying the link between PTSD and somatic symptoms have not yet been identified though they are believed to be the result of both biological and psychological changes occurring after the exposure to a traumatic event.^{15,16} PTSD is characterized by a failure to physiologically adapt to stressors and reminders of the stressors, such that the long-term activation of the stress pathways like the sympathetic nervous system and hypothalamic-pituitary-adrenal (HPA) axis with consequent influence on immune activity that places individuals at higher risk for developing or exacerbating physical medical conditions.^{17–19}

A link between chronic pain and PTSD or trauma exposure has in fact been hypothesized to be underlined by a chronic altered cortisol or inflammatory response.^{18,20} In a variety of samples, most of the literature regarding the relationship between PTSD and chronic pain indicates that individuals with PTSD report greater frequency or severity of overall pain than those without PTSD.^{21–23} More specifically, these conditions include mainly fibromyalgia, back pain and headaches, and arthritis pain.^{2,24–28}

A positive relationship between PTSD and GI problems, including irritable bowel syndrome (IBS), vomiting, ulcer, and constipation/diarrhea has also been demonstrated.^{2,27–31} Further studies analyzed the relationship between migraine and PTSD, suggesting PTSD to be more common in patients with migraine than in the general population.³²

There is evidence that PTSD usually occurs almost twice as much in women compared to men and this has been also confirmed among general population exposed to earthquake.^{33–34} However, little or no data are currently available on the possible gender differences in somatic symptoms emerging in patients with PTSD in the aftermath of trauma exposure. A first study from McCall-Hosenfeld et al³⁵ in a sample of 597 urban primary care patients with chronic pain showed women to report significantly more somatic symptoms than men when exposed to three interpersonal trauma types (sexual trauma, intimate partner violence, and childhood trauma history), although somatic symptoms were increased among all interpersonal trauma survivors. More recently, Morina et al³⁶ showed women reported significantly higher levels of somatic symptoms than men in sample of 142 civilian war survivors, whereas levels of PTSD symptoms were similar in both the genders. Conversely, a systematic review including 26 studies showed that individuals who reported exposure to trauma, especially men, reported an increased prevalence of functional somatic syndromes. The magnitude of the association of these latter with PTSD was significantly stronger than that with trauma exposure alone; however, the gender differences lost the statistical significance by performing more conservative analyses in the same sample.³⁷

In light of these data, the aim of the present study was to investigate the possible impact of gender and PTSD on the onset of somatic symptoms, complained in the aftermath of the event, among young adult survivors of the 2009 L'Aquila earthquake in Italy.

Methods

Participants

The study sample consisted in 450 high-school senior students, 253 (56.2%) male and 197 (47.8%) female (mean age \pm SD: 17.64 \pm 0.78 years), who had been exposed to the 2009 L'Aquila earthquake, enrolled 21 months after the event. Sociodemographic characteristics of the study sample were detailed elsewhere.³⁸ PTSD was diagnosed, according to DSM-5 criteria, in 162 (36.0%) subjects, specifically in 61 men and 101 women (24.1% vs 51.3%, $P < .001$).

The Ethics Committee of the University of L'Aquila and the school councils approved all recruitment and assessment procedures. In accordance with the Declaration of Helsinki, subjects provided written informed consent after receiving a complete description of the study.

Instruments and assessments

Subjects were assessed by the modified versions of the Mood Spectrum Self-Report-Lifetime Version (MOODS-SR) and the Trauma and Loss Spectrum Self-Report (TALS-SR),^{39,40} to evaluate symptoms occurred in the aftermath of the earthquake exposure.

The MOODS-SR is a 140-item questionnaire exploring mood spectrum symptoms, coded dichotomously, as present or absent, for one or more periods of at least three to five days.³⁹ According to the aims of the present study, we adopted a modified version of the instrument assessing symptoms developed in the aftermath of earthquake. Items are organized into three manic and three depressive domains, exploring "mood," "energy," and "cognition," besides a *rhythmicity and vegetative functions* domain. This latter explores alterations in the circadian rhythms and vegetative functions, including changes in energy, physical wellbeing, mental and physical efficiency related to the weather and season, and changes in appetite, sleep, and sexual activities across 29 items. Five of the MOODS-SR items belonging to the rhythmicity and vegetative functions domain, assessing changes in somatic symptoms, were examined: item n = 159a (...you repeatedly had distressing physical symptoms, for instance: frequent headaches?), item n = 159b (...you repeatedly had distressing physical symptoms, for instance: your mouth felt dry?), item n = 159c (...you repeatedly had distressing physical symptoms, for instance: you were constipated?), item n = 159d (...you repeatedly had distressing physical symptoms, for instance: you had nausea or other stomach or bowel problems?), item n = 160 (...you were more sensitive or less sensitive than usual to heat, cold or pain?). As we adopted a modified version of the MOODS-SR, as described before, we assessed these symptoms when occurring in the aftermath of earthquake exposure.

The TALS-SR includes 116 items exploring the lifetime experience of a range of loss and/or traumatic events and lifetime symptoms, behaviors and personal characteristics that might represent manifestations and/or risk factors for the development of a stress response syndrome.⁴⁰ The instrument is organized into nine domains, of which four exploring PTSD symptomatology, namely, re-experiencing, avoidance and numbing, maladaptive coping, and arousal. A modified TALS-SR version, which specifically evaluated PTSD symptomatology related to the earthquake exposure, was used. According previous studies,^{38,41} DSM-5 PTSD diagnosis was assessed utilizing the following matching between symptom criteria and TALS-SR items: criterion B (B1 = 80; B2 = 77; B3 = 79; B4 = 78; and B5 = 81); criterion C (C1 = 86; C2 = 87 and/or 88 and/or 89); criterion D (D1 = 90; D2 = 95; D3 = 85; D4 = 96; D5 = 91; D6 = 93; and D7 = 92); and

criterion E (E1 = 108; E2 = 99 and/or 100 and/or 102 and/or 103 and/or 104; E3 = 106; E4 = 107; E5 = 105; and E6 = 109). Due to the sample characteristics, criterion A was considered satisfied.

Statistical analysis

Chi-square tests (or Fisher when appropriate) were computed in order to compare endorsement rates of each MOODS-SR somatic symptom item in male vs female and PTSD vs NO-PTSD survivors. To study all the possible effects of gender and PTSD diagnosis (and their possible interaction) on the presence of at least one MOODS-SR somatic symptom item a Decision Tree procedure based on the Chi-squared automated interaction detection growing method was utilized.

Finally, the effects of gender and PTSD on the total MOODS-SR somatic symptoms score (derived by the sum of the scores of item 159a, 159b, 159c, 159d, and 160), and of their possible interactions were analyzed by a model of a two-way analysis of variance (ANOVA).

In order to study the post-traumatic stress symptomatology that is more strongly related to the endorsement of at least one MOODS-SR somatic symptom, and the possible impact of gender, we conducted a multivariate logistic regression that reported the presence of at least one MOODS-SR somatic symptom as a dependent variable and gender and the four TALS-SR domains exploring post-traumatic stress symptoms as independent variables.

All statistical analyses were carried out using the Statistical Package for Social Science (SPSS Inc., Chicago, IL), version 25.0.

Results

Full data on the MOODS-SR were available for 450 survivors, the same that also fulfilled the TALS-SR. MOOD-SR somatic symptoms scores were significantly higher in survivors with PTSD than

in those without, except for the MOOD-SR item n = 159b (...you repeatedly had distressing physical symptoms, for instance: your mouth felt dry?). PTSD survivors also showed higher ratio of endorsing at least one MOODS-SR somatic symptom than those without (Table 1). No gender differences emerged in MOODS-SR somatic symptoms, albeit females reported higher rates in each item as well as a statistically significant higher rate of endorsement of at least one MOODS-SR somatic symptom compared to males (Table 2).

A Decision Tree procedure was utilized to locate the strongest interactions with at least one MOODS-SR somatic symptom (considered as dependent or target variable). The tree-based classification model, able to predict the values of the dependent variable based on the values of the independent ones (gender and PTSD in this case), had an acceptable misclassification risk of the model (value = 0.14). Subjects with PTSD showed a significantly higher ratio of endorsing at least one MOODS-SR somatic symptom compared to those without (25.3% vs 7.6%, $\chi^2 = 26.89, P < .001$). According to this model, a significant impact of the gender on the presence of at least one MOODS-SR somatic symptom did not emerged. See Figure 1 for details.

The two-way ANOVA model confirmed only a significant main effect on the total MOODS-SR somatic symptoms score of PTSD [$F(1446) = 13.40, P < .001$] with a mean score significantly higher in PTSD group compared to No PTSD group (0.40 ± 0.58 vs 0.14 ± 0.54). No significant main effect of gender [$F(1446) = 0.48, P = .489$], nor of gender \times PTSD interaction [$F(1446) = 0.43, P = .514$], emerged.

The multivariate logistical regression showed a significant association between the presence of at least one MOOD-SR somatic symptom and re-experiencing (odds ratio = 1.37, $P = .001$) and maladaptive coping (odds ratio = 1.26, $P = .029$) TALS-SR symptomatologic domains (Table 3).

Table 1. Comparison of MOODS-SR Somatic Symptom Items Endorsement Rates among L'Aquila Earthquake Young Adult Survivors With (n = 162) and Without (n = 288) PTSD.

MOODS-SR Somatic Symptoms Item	Total N (%)	No PTSD (n [%])	PTSD (n [%])	P
n = 159a (...you repeatedly had distressing physical symptoms, for instance: frequent headaches?)	21 (4.7%)	8 (2.8%)	13 (8.0%)	.021
n = 159b (...you repeatedly had distressing physical symptoms, for instance: your mouth felt dry?)	12 (2.7%)	5 (1.7%)	7 (4.3%)	.129
n = 159c (...you repeatedly had distressing physical symptoms, for instance: you were constipated?)	17 (3.8%)	6 (2.1%)	11 (6.8%)	.024
n = 159d (...you repeatedly had distressing physical symptoms, for instance: you had nausea or other stomach or bowel problems?)	22 (4.9%)	9 (3.1%)	13 (8.0%)	.037
n = 160 (...you were more sensitive or less sensitive than usual to heat, cold, or pain?)	32 (7.1%)	12 (4.2%)	20 (12.3%)	.002
At least one MOODS-SR somatic symptom	63 (14.0%)	22 (7.6%)	41 (25.3%)	<.001

Table 2. Gender Differences in MOODS-SR Somatic Symptoms Items Endorsement Rates among L'Aquila Earthquake Young Adult Survivors (Male = 253 and Female = 197).

MOODS-SR Somatic Symptoms Item	Total N (%)	Male (n [%])	Female (n [%])	P
n = 159a (...you repeatedly had distressing physical symptoms, for instance: frequent headaches?)	21 (4.7%)	9 (3.6%)	12 (6.1%)	.299
n = 159b (...you repeatedly had distressing physical symptoms, for instance: your mouth felt dry?)	12 (2.7%)	6 (2.4%)	6 (3.0%)	.884
n = 159c (...you repeatedly had distressing physical symptoms, for instance: you were constipated?)	17 (3.8%)	6 (2.4%)	11 (5.6%)	.128
n = 159d (...you repeatedly had distressing physical symptoms, for instance: you had nausea or other stomach or bowel problems?)	22 (4.9%)	10 (4.0%)	12 (6.1%)	.410
n = 160 (...you were more sensitive or less sensitive than usual to heat, cold, or pain?)	32 (7.1%)	14 (5.5%)	18 (9.1%)	.197
At least one MOODS-SR somatic symptom	63 (14.0%)	26 (10.3%)	37 (18.8%)	.015




Figure 1. Decision Tree showing post-traumatic stress disorder (PTSD) interaction with Mood Spectrum Self-Report-Lifetime Version (MOODS-SR) somatic symptoms as target variable.

Table 3. Multivariate Logistic Regression Reporting the Presence of at least One MOODS-SR Somatic Symptom as Dependent Variable and Gender and the Four TALS-SR Domains Exploring Post-Traumatic Stress Symptoms as Independent Variables.

TALS-SR Somatic Symptoms Item	B (SE)	OR	IC _{95%}	P
k	-3.82 (0.54)	-	-	<.001
Gender	0.19 (0.34)	1.21	0.62; 2.33	.578
Reexperiencing	0.32 (0.09)	1.37	1.15; 1.64	.001
Avoidance and numbing	0.05 (0.08)	1.05	0.91; 1.22	.521
Maladaptive coping	0.23 (0.10)	1.26	1.02; 1.54	.029
Arousal	0.00 (0.13)	1.00	0.78; 1.28	.982

Hosmer and Lemeshow test: $\chi^2 = 8.60, P = .377$; Cox and Snell $R^2 = 0.10$; Nagelkerke $R^2 = 0.18$ overall percentage of correct classification: 85.3%.
 Abbreviations: OR, odds ratio; TALS-SR, Trauma and Loss Spectrum Self-Report; SE, standard error.

Discussion

To the best of our knowledge, the present study is the first to investigate the impact of PTSD and gender on somatic symptoms among civilian young adult survivors to a massive earthquake. According to our data, up to 14% of survivors to the L’Aquila 2009 earthquake complained of somatic symptoms after 21 months from exposure. Complaints included headaches, GI problems, and thermal/painful stimuli hypo- or hypersensitivity, with PTSD survivors showing significantly higher prevalence of such symptoms compared with subjects without PTSD.

In this regard, our results are in line with prior evidence reporting a relationship between PTSD and somatic symptoms, such as headache and stomach pain, in children and adolescents exposed to the Lushan 2013 earthquake.⁴² In particular, 4.7% of the total sample reported frequent headaches, with statistically significant higher rates, more than two times, in patients with PTSD compared with those without. Growing evidence suggests a relationship between PTSD symptoms and headaches. Smitherman et al,⁴³ in a cross-sectional study on 1051 young adults, compared patients with and without migraine, showing PTSD to be a predictor of migraine. Among veterans, PTSD has been repeatedly associated

with increased consumption of medications for headaches, more in men than in women.⁴⁴ Furthermore, subjects with migraine and comorbid PTSD were found to report a greater disability and more lost workdays due to physical or mental health or substance abuse, and more difficulties with developing and maintaining a social life than subjects with migraine but without PTSD.^{45,46} On the other hand, it has also been pointed out that patients with migraine may suffer from lower resilience and subsequent maladaptive stress responses, which might predispose them to PTSD.⁴⁷ Current hypotheses on neurobiological mechanism underlying the association between PTSD and migraines include serotonergic, autonomic nervous system, and HPA axis dysfunction.⁴⁸

The results of the present study also showed 3.8% of total sample to report constipation and 4.9% nausea or other bowel problems, with almost twice as much rates in PTSD subjects compared with those without. The GI distress related to PTSD has been hypothesized to be linked to the hyperarousal response that is usually associated with enhanced cortical arousal, which can elicit sensations in the gut.^{49,50} About 36% of patients seeking treatment for IBS met the criteria for PTSD.⁵¹ Recently, one study correlated neuropeptide Y (NPY) levels with bowel function in patients with PTSD; in particular, low levels of NPY were reported in the cerebrospinal fluid and plasma of male combat veterans with PTSD, and this is negatively related with sympathetic nervous system (SNS) hyperreactivity, PTSD symptoms, and time to recovery.⁵² It should be noted that NPY regulation has not yet been evaluated in women with PTSD. NPY levels are reported to be low in bowel tissue of IBS subjects with diarrhea (IBS-D) vs IBS subjects with constipation. Greater density of ghrelin-containing cells of the gastric oxyntic mucosa has been reported in IBS-D. PTSD-related SNS hyperreactivity may interact with this mechanism to increase ghrelin release, which activates receptors in the lumbosacral spinal cord and basolateral amygdala increasing colonic motility and amygdala hyperreactivity, respectively. Loss of function in the adrenergic α_2 -autoreceptors gene and increased corticotropin-releasing hormone, as observed in PTSD, are also thought to contribute to IBS-D.^{52,53}

Our data also showed that 7.1% of the total sample reported thermal/painful stimuli hypo- or hypersensitivity, which represents a dysregulation leading to chronic pain. Also in this case,

statistically significant higher rates emerged in PTSD subjects than in those without. As previously reported in an Italian study concerning pain in survivors of L'Aquila 2009 earthquake, a high prevalence of somatic conditions such as headaches or lower back pain was detected in the five weeks after the event, associated to increased use of painkillers,¹¹ albeit in such a close observational frame time from the traumatic event, acute symptoms related to wounds or direct damage from the earthquake had affected the results. However, in our findings, the strong relationship between post-traumatic stress and somatic symptoms was well evident after 21 months from the devastating event and related to changing in the perception of somatic health in the aftermath of the event. Further, subjects with PTSD showed more severe chronic pain and this latter correlated with post-traumatic stress symptoms.^{54,55} Pain and PTSD are associated with similar neurobiological abnormalities,^{56,57} as hyperarousal,⁵⁸ dysregulations in stress responses,⁵⁹ and in pain modulation.⁶⁰ PTSD subjects show an intense and widespread chronic pain and a peculiar sensory profile of hyposensitivity to pain, related to increased dissociation, accompanied by hyperreactivity to suprathreshold noxious stimuli, linked to increased anxiety sensitivity.⁶¹ Chronic pain, identified as a persistent condition with an emotional impact, becoming in itself a self-perpetuating stressing event.^{62,63} Depending on the population studied and the disorder initially assessed, pain can contribute to the onset of PTSD and also maintain it. The same thing is true for PTSD, which can act as both a triggering factor and a maintenance factor of chronic pain.⁶⁴

It is also important to recall that in southern Mediterranean populations, somatic symptoms have been reported to be the prevalent manifestation of distress related to trauma.⁶⁵ The role of somatic complaints after trauma exposure on decreased quality of life seems consistent with the data suggesting that somatization may be a preferred mode of suffering expression in certain populations, thus playing the role of idioms of distress.^{66–68} This suggests that assessing somatic complaints after trauma exposure, in addition to PTSD symptoms, may help better evaluate the real impact of trauma in exposed populations.

To the best of our knowledge, our results first report on the gender differences in somatic complaints among earthquake survivors in the general population. In particular, no significant impact of gender seemed to emerge on the risk of developing somatic symptoms in the aftermath of a non-war-related massive traumatic event. Despite the fact that we found statistically significant higher rates of at least one MOODS-SR somatic symptom in females rather than males, we hypothesized that this was primarily related to the higher PTSD rates reported by females than in males, with almost twice as much the levels. This appears to be in contrast with a previous research on civilian war victims, where women showed higher levels of somatization symptoms than men, whereas the levels of PTSD symptoms were similar across the two genders.⁶⁹ However, a systematic review, examining the association of reported psychological trauma and PTSD with functional somatic syndromes, showed no significant gender effect on the prevalence of functional somatic syndromes in individuals who reported exposure to trauma, even a slightly but not significant higher rate in males.³⁷ In order to better clarify this issue, we performed a further analysis to examine the contribution of continuous PTSD symptoms on somatic symptoms above and beyond sex. Results showed re-experiencing and maladaptive coping TALS-SR domains to be the only related to the presence of at least one MOODS-SR somatic symptom. For what concern the re-experiencing symptoms, we argue that these

symptoms are usually related to prolonged rumination on the trauma and its consequences. Szabo *et al.*⁷⁰ showed a moderate, positive relationship between rumination and PTSD symptoms in trauma-exposed adults, with a stronger association between rumination and intrusive re-experiencing than avoidance or hyperarousal. Rumination is, in fact, defined as a form of perseverative cognition that focuses on negative content, generally past and present, and results in emotional distress.⁷¹ Emerging literature suggests the possible role of rumination in fully mediating the relationship between neuroticism and somatic complaints.⁷² Lohaus *et al.*,⁷³ in a sample of children and adolescents, showed ruminative response style to be related to increased frequency of experiencing somatic and psychological symptoms, particularly among girls. Our results may account for a primary role of post-traumatic stress reactions on the relationship with somatic symptoms beyond gender. For what concern our results on the relationship between TALS-SR maladaptive coping and somatic symptoms, we may suggest this relationship may have a dual label. If on the one hand, maladaptive symptoms and self-destructive behaviors are fully acknowledged as core PTSD DSM-5 symptoms (within diagnostic criterion E),^{1,74} our results may account for a relationship between an increased severity of the PTSD and the somatic complaints. On the other hand, in the maladaptive coping domain, the TALS-SR items include “having stopped taking care of themselves” and “the use of drugs or over the counter medications to relief symptoms,” and this may be consequent to the presence of somatic symptoms *per-se*. Previous studies on trauma-exposed adolescent showed high rates of drugs, alcohol, or medication (such as painkillers) to calm themselves or to relieve emotional or physical pain.^{12,13,75,76}

Some important limitations of the present study should be kept in mind when interpreting our results. The first limitation is the lack of information about psychiatric comorbidities in the sample, including prior psychiatric history; this could, in fact, have an impact on the development of PTSD or somatic symptoms. Second is the use of self-report instruments that could be considered less accurate than the rating of the clinician. Third is represented by the lack of information on the presence of somatic illness and on quality of life in the sample. Fourth is the fact that we did not use a specific scale to assessing chronic pain.

Conclusion

Despite these limitations, our results highlight the significant relationship between somatic symptoms developed in young adult population in the aftermath of a massive trauma exposure, such as an earthquake, suggesting a potential higher vulnerability in those subjects presenting PTSD without a clear prevalence of gender. Evaluating the burden of somatic symptoms seems essential for a more accurate assessment and clinical management of survivors to mass traumas, such as earthquakes or other natural catastrophic events, with post-traumatic stress symptoms, also in order to prevent inappropriate painkillers prescriptions. In light of our findings, in fact, treatment should focus on PTSD symptoms and on re-experiencing and rumination on the trauma and its consequence in order to provide adequate psychopharmacological and psychological treatment, to prevent somatizations and, potentially, the often-revealed abuse of painkillers and over the counter medications in such populations. However, scant trials are still available on effective treatments suggesting the need for further research in this specific field.

Funding. This research received no specific grant from any funding agency, commercial or not-for-profit sectors.

Disclosures. The authors report no conflicts of interest.

References

- American Psychiatric Association. *Diagnostic and Statistical Manual of Mental Disorders: DSM-5*. Washington, DC: American Psychiatric Publishing Incorporated; 2013.
- Sareen J, Cox BJ, Stein MB, et al. Physical and mental comorbidity, disability, and suicidal behavior associated with posttraumatic stress disorder in a large community sample. *Psychosom Med*. 2007;**69**(3):242–248.
- Kessler RC, Sonnega A, Bromet E, et al. Posttraumatic stress disorder in the National Comorbidity Survey. *Arch Gen Psychiatry*. 1995;**52**(12):1048–1060.
- Carmassi C, Dell’Osso L, Manni C, et al. Frequency of trauma exposure and post-traumatic stress disorder in Italy: analysis from the World Mental Health Survey Initiative. *J Psychiatr Res*. 2014;**59**:77–84.
- Barkmann C, Braehler E, Schulte-Markwort M, et al. Chronic somatic complaints in adolescents: prevalence, predictive validity of the parent reports, and associations with social class, health status and psychosocial distress. *Soc Psychiatry Psychiatr Epidemiol*. 2011;**46**(10):1003–1011.
- Pacella ML, Hruska B, Delahanty DL. The physical health consequences of PTSD and PTSD symptoms: a meta-analytic review. *J Anxiety Disord*. 2013;**27**(1):33–46.
- Morina N, Ford JD, Risch AK, et al. Somatic distress among Kosovar civilian war survivors: relationship to trauma exposure and the mediating role of experiential avoidance. *Soc Psychiatry Psychiatr Epidemiol*. 2010;**45**(12):1167–1177.
- Cheung A, Makhshvili N, Javakhishvili J, et al. Patterns of somatic distress among internally displaced persons in Ukraine: analysis of a cross-sectional survey. *Soc Psychiatry Psychiatr Epidemiol*. 2019;**54**(10):1265–1274.
- Hoge CW, Terhakopian A, Castro CA, et al. Association of posttraumatic stress disorder with somatic symptoms, health care visits, and absenteeism among Iraq war veterans. *Am J Psychiatry*. 2007;**164**(1):150–153.
- Milligan-Saville JS, Paterson HM, Harkness EL, et al. The amplification of common somatic symptoms by posttraumatic stress disorder in firefighters. *J Trauma Stress*. 2017;**30**(2):142–148.
- Angeletti C, Guetti C, Papola R, et al. Pain after earthquake. *Scand J Trauma Resusc Emerg Med*. 2012;**20**:43.
- Dell’Osso L, Carmassi C, Stratta P, et al. Gender differences in the relationship between maladaptive behaviors and post-traumatic stress disorder. A Study on 900 L’Aquila 2009 Earthquake Survivors. *Front Psychiatry*. 2013;**3**:111.
- Carmassi C, Rossi A, Pedrinelli V, et al. PTSD in the aftermath of a natural disaster: what we learned from the Pisa-L’Aquila Collaboration Project. *Journal of Psychopathology*. 2020;**26**(2):99–106.
- Zhang J, Zhu S, Du C, et al. Posttraumatic stress disorder and somatic symptoms among child and adolescent survivors following the Lushan earthquake in China: a six-month longitudinal study. *J Psychosom Res*. 2015;**79**(2):100–106.
- Schnurr PP, Green BL. Understanding relationships among trauma, post-traumatic stress disorder, and health outcomes. *Adv Mind Body Med*. 2004;**20**(1):18–29.
- Miao XR, Chen QB, Wei K, et al. Posttraumatic stress disorder: from diagnosis to prevention. *Mil Med Res*. 2018;**5**(1):32.
- Schnurr PP, Jankowski MK. Physical health and post-traumatic stress disorder: review and synthesis. *Semin Clin Neuropsychiatry*. 1999;**4**(4):295–304.
- Boscarino JA. Posttraumatic stress disorder and physical illness: results from clinical and epidemiologic studies. *Ann N Y Acad Sci*. 2004;**1032**:141–153.
- Dougall AL, Baum A. Psychoneuroimmunology and trauma. In Schnurr PP, Green BL eds. *Trauma and Health: Physical Health Consequences of Exposure to Extreme Stress*. Washington, DC: American Psychological Association; 2004:129–155.
- Kendall-Tackett K. Psychological trauma and physical health: a psychoneuroimmunology approach to etiology of negative health effects and possible intervention. *Psychol Trauma Theor Res Pract Policy*. 2009;**1**(1):35–48.
- Ouimette P, Cronkite R, Henson BR, et al. Posttraumatic stress disorder and health status among female and male medical patients. *J Trauma Stress*. 2004;**17**(1):1–9.
- Woods SJ, Hall RJ, Campbell JC, et al. Physical health and posttraumatic stress disorder symptoms in women experiencing intimate partner violence. *J Midwifery Womens Health*. 2008;**53**(6):538–546.
- Smith BN, Shepherd JC, Schuster JL, et al. Posttraumatic stress symptomatology as a mediator of the association between military sexual trauma and post-deployment physical health in women. *J Trauma Dissociation*. 2011;**12**(3):275–289.
- Vedantham K, Brunet A, Boyer R, et al. Posttraumatic stress disorder, trauma exposure, and the current health of Canadian bus drivers. *Can J Psychiatry*. 2001;**46**(2):149–155.
- Dell’Osso L, Carmassi C, Massimetti G, et al. Impact of traumatic loss on post-traumatic spectrum symptoms in high school students after the L’Aquila 2009 earthquake in Italy. *J Affect Disord*. 2011;**134**(1–3):59–64.
- Dell’Osso L, Bazzichi L, Consoli G et al. Manic spectrum symptoms are correlated to the severity of pain and the health-related quality of life in patients with fibromyalgia. *Clin Exp Rheumatol*. 2009;**27**(5 Suppl 56):S57–61.
- Lauterbach D, Vora R, Rakow M. The relationship between posttraumatic stress disorder and self-reported health problems. *Psychosom Med*. 2005;**67**(6):939–947.
- Kelly U. Intimate partner violence, physical health, posttraumatic stress disorder, depression, and quality of life in latinas. *West J Emerg Med*. 2010;**11**(3):247–251.
- Shalev A, Bleich A, Ursano RJ. Posttraumatic stress disorder: somatic comorbidity and effort tolerance. *Psychosomatics*. 1990;**31**(2):197–203.
- Andreski P, Chilcoat H, Breslau N. Post-traumatic stress disorder and somatization symptoms: a prospective study. *Psychiatry Res*. 1998;**79**(2):131–138.
- Schnurr PP, Spiro A, Paris AH. Physician-diagnosed medical disorders in relation to PTSD symptoms in older male military veterans. *Health Psychol*. 2000;**19**(1):91–97.
- Saunders K, Merikangas K, Low NC, et al. Impact of comorbidity on headache-related disability. *Neurology*. 2008;**70**(7):538–547.
- Dell’Osso L, Carmassi C, Massimetti G, et al. Full and partial PTSD among young adult survivors 10 months after the L’Aquila 2009 earthquake: gender differences. *J Affect Disord*. 2011;**131**(1–3):79–83.
- Dell’Osso L, Carmassi C, Rucci P, et al. Lifetime subthreshold mania is related to suicidality in posttraumatic stress disorder. *CNS Spectrums*. 2009;**14**(5):262–266.
- McCall-Hosenfeld J, Winter M, Heeren T, et al. The association of interpersonal trauma with somatic symptom severity in a primary care population with chronic pain: exploring the role of gender and the mental health sequelae of trauma. *J Psychosom Res*. 2014;**77**(3):196–204.
- Morina N, Schnyder U, Klaghofer R, et al. Trauma exposure and the mediating role of posttraumatic stress on somatic symptoms in civilian war victims. *BMC Psychiatry*. 2018;**18**(1):92.
- Afari N, Ahumada SM, Wright LJ, et al. Psychological trauma and functional somatic syndromes: a systematic review and meta-analysis. *Psychosom Med*. 2014;**76**(1):2–11.
- Carmassi C, Akiskal HS, Yong SS, et al. Post-traumatic stress disorder in DSM-5: estimates of prevalence and criteria comparison versus DSM-IV-TR in a non-clinical sample of earthquake survivors. *J Affect Disord*. 2013;**151**(3):843–848.
- Dell’Osso L, Armani A, Rucci P, et al. Measuring mood spectrum: comparison of interview (SCI-MOODS) and self-report (MOODS-SR) instruments. *Compr Psychiatry*. 2002;**43**(1):69–73.
- Dell’Osso L, Carmassi C, Rucci P, et al. A multidimensional spectrum approach to post-traumatic stress disorder: comparison between the Structured Clinical Interview for Trauma and Loss Spectrum (SCI-TALS) and

- the Self-Report instrument (TALS-SR). *Compr Psychiatry*. 2009;**50**(5):485–490.
41. Dell'osso L, Stratta P, Conversano C, Massimetti E, Akiskal KK, Akiskal HS, Rossi A, Carmassi C. Lifetime mania is related to post-traumatic stress symptoms in high school students exposed to the 2009 L'Aquila earthquake. *Compr Psychiatry*. 2014 Feb;**55**(2):357–62. doi:10.1016/j.comppsy.2013.08.017.
 42. Zhang Y, Zhang J, Zhu S, *et al.* Prevalence and predictors of somatic symptoms among child and adolescents with probable posttraumatic stress disorder: a cross-sectional study conducted in 21 primary and secondary schools after an earthquake. *PLoS One*. 2015;**10**(9):e0137101.
 43. Smitherman TA, Kolivas ED. Trauma exposure versus posttraumatic stress disorder: relative associations with migraine. *Headache*. 2013;**53**(5):775–786.
 44. Seng EK, Driscoll MA, Brandt CA *et al.* Prescription headache medication in OEF/OIF veterans: results from the Women Veterans Cohort Study. *Headache*. 2013;**53**(8):1312–1322.
 45. Peterlin BL, Tietjen GE, Brandes JL, *et al.* Posttraumatic stress disorder in migraine. *Headache*. 2009;**49**(4):541–551.
 46. Rao AS, Scher AI, Vieira RV *et al.* The impact of post-traumatic stress disorder on the burden of migraine: results from the National Comorbidity Survey-Replication. *Headache*. 2015;**55**(10):1323–1341.
 47. Peterlin BL, Nijjar SS, Tietjen GE. Post-traumatic stress disorder and migraine: epidemiology, sex differences, and potential mechanisms. *Headache*. 2011;**51**(6):860–868.
 48. Juang KD, Yang CY. Psychiatric comorbidity of chronic daily headache: focus on traumatic experiences in childhood, post-traumatic stress disorder and suicidality. *Curr Pain Headache Rep*. 2014;**18**(4):405.
 49. Levy F, Krebs PR. Cortical-subcortical re-entrant circuits and recurrent behaviour. *Aust N Z J Psychiatry*. 2006;**40**(9):752–758.
 50. Mayer EA, Bradesi S, Chang L, *et al.* Functional GI disorders: from animal models to drug development. *Gut*. 2008;**57**(3):384–404.
 51. Irwin C, Falsetti SA, Lydiard RB, *et al.* Comorbidity of posttraumatic stress disorder and irritable bowel syndrome. *J Clin Psychiatry*. 1996;**57**(12):576–578.
 52. Rasmussen AM. The gut peptide neuropeptide Y and post-traumatic stress disorder. *Curr Opin Endocrinol Diabetes Obes*. 2017;**24**(1):3–8.
 53. Dell'Osso L, Carmassi C, Mucci F, *et al.* Depression, Serotonin and Tryptophan. *Curr Pharm Des*. 2016;**22**(8):949–54.
 54. Defrin R, Ginzburg K, Solomon Z, *et al.* Quantitative testing of pain perception in subjects with PTSD—implications for the mechanism of the coexistence between PTSD and chronic pain. *Pain*. 2008;**138**(2):450–459.
 55. Bartel A, Jordan J, Correll D, *et al.* Somatic burden and perceived cognitive problems in trauma-exposed adults with posttraumatic stress symptoms or pain. *J Clin Psychol*. 2020;**76**(1):146–160.
 56. Scioli-Salter ER, Forman DE, Otis JD *et al.* The shared neuroanatomy and neurobiology of comorbid chronic pain and PTSD: therapeutic implications. *Clin J Pain*. 2015;**31**(4):363–374.
 57. Dell'Osso L, Da Pozzo E, Carmassi C, *et al.* Lifetime manic-hypomanic symptoms in post-traumatic stress disorder: relationship with the 18 kDa mitochondrial translocator protein density. *Psychiatry Res*. 2010;**177**(1–2):139–43.
 58. Daniels JK, McFarlane AC, Bluhm RL, *et al.* Switching between executive and default mode networks in posttraumatic stress disorder: alterations in functional connectivity. *J Psychiatry Neurosci*. 2010;**35**(4):258–266.
 59. McLean SA, Clauw DJ, Abelson JL, *et al.* The development of persistent pain and psychological morbidity after motor vehicle collision: integrating the potential role of stress response systems into a biopsychosocial model. *Psychosom Med*. 2005;**67**(5):783–790.
 60. Kosek E, Ekholm J, Hansson P. Sensory dysfunction in fibromyalgia patients with implications for pathogenic mechanisms. *Pain*. 1996;**68**(2–3):375–383.
 61. Defrin R, Schreiber S, Ginzburg K. Paradoxical pain perception in posttraumatic stress disorder: the unique role of anxiety and dissociation. *J Pain*. 2015;**16**(10):961–970.
 62. Chapman TRP, Song CW. Pain and stress in a systems perspective: reciprocal neural, endocrine and immune interactions. *J Pain*. 2008;**9**(2):122–145.
 63. Carmassi C, Dell'Oste V, Ceresoli D, *et al.* Frequent attenders in general medical practice in Italy: a preliminary report on clinical variables related to low functioning. *Neuropsychiatr Dis Treat*. 2018;**15**:115–125.
 64. Brennstuhl MJ, Tarquinio C, Montel S. Chronic pain and PTSD: evolving views on their comorbidity. *Perspect Psychiatr Care*. 2015;**51**(4):295–304.
 65. Hiar S, Thomas CL, Hinton DE, *et al.* Somatic symptoms mediate the relationship between trauma during the Arab spring and quality of life among Tunisians. *J Nerv Ment Dis*. 2016;**204**(2):153–155.
 66. Kirmayer LJ. Confusion of the senses: implications of ethnocultural variations in somatoform and dissociative disorders for PTSD. In: Marsella AJ, Friedman MJ, Gerrity ET, Scurfield RM, eds. *Ethnocultural Aspects of Posttraumatic Stress Disorder*. Washington, DC: American Psychological Association; 1996:131–163.
 67. Terheggen MA, Stroebe MS, Kleber RJ. Western conceptualizations and Eastern experience: a cross-cultural study of traumatic stress reactions among Tibetan refugees in India. *J Trauma Stress*. 2001;**14**:391–403.
 68. Hinton DE, Lewis-Fernández R. The cross-cultural validity of posttraumatic stress disorder: implications for DSM-5. *Depress Anxiety*. 2011;**28**:783–801.
 69. Morina N, Schnyder U, Klaghofer R, *et al.* Trauma exposure and the mediating role of posttraumatic stress on somatic symptoms in civilian war victims. *BMC Psychiatry*. 2018;**18**(1):92.
 70. Szabo YZ, Warnecke AJ, Newton TL, *et al.* Rumination and posttraumatic stress symptoms in trauma-exposed adults: a systematic review and meta-analysis. *Anxiety Stress Coping*. 2017;**30**(4):396–414.
 71. Sansone RA, Sansone LA. Rumination: relationships with physical health. *Innov Clin Neurosci*. 2012;**9**(2):29–34.
 72. Denovan A, Dagnall N, Lofthouse G. Neuroticism and somatic complaints: concomitant effects of rumination and worry. *Behav Cogn Psychother*. 2019;**47**(4):431–445.
 73. Lohaus A, Vierhaus M, Frevert A, *et al.* Rumination and symptom reports in children and adolescents: results of a cross-sectional and experimental study. *Psychol Health*. 2013;**28**(9):1032–1045.
 74. Carmassi C, Barberi FM, Cordone A, *et al.* Trauma, PTSD and post-traumatic stress spectrum: 15 years' experience on a multidimensional approach to trauma related psychopathology. *Journal of Psychopathology*. 2020;**26**:411. doi:10.36148/2284-0249-376.
 75. Glodich A, Allen JG. Adolescents exposed to violence and abuse: a review of the group therapy literature with an emphasis on preventing trauma reenactment. *J Child Adolesc Group Ther*. 1998;**8**:135–154.
 76. Steven SJ, Murphy BS, McKnight K. Traumatic stress and gender differences in relationship to substance abuse, mental health, physical health and HIV risk-taking in a sample of adolescents enrolled in drug treatment. *Child Maltreat*. 2003;**8**:46–57.