

Do tissues other than the kidney produce 1,25-dihydroxyvitamin D₃ *in vivo*? A reexamination

(vitamin D/metabolism/renal failure)

TERRY D. SHULTZ, JOHN FOX, HUNTER HEATH III, AND RAJIV KUMAR*

Endocrine Research Unit, Department of Medicine, Mayo Clinic and Mayo Medical School, Rochester, Minnesota 55905

Communicated by R. H. Wasserman, November 29, 1982

ABSTRACT Recent experiments have shown that 1,25-dihydroxyvitamin D₃-like material is produced in cultured nonrenal cells and may be present in the sera of anephric patients. We reexamined the question of whether 1,25-dihydroxyvitamin D₃ can be synthesized extrarenally in the rat *in vivo*. To intact, sham-operated, ureter-ligated, or acutely nephrectomized vitamin D-deficient rats raised on a diet normal in calcium and phosphorus, we gave a physiologic dose of high-specific-activity 25-hydroxy-[³H]vitamin D₃ (3.6–3.8 μCi; ≈ 25 pmol per rat). Twenty-four hours later we examined their tissues and plasma for the presence of radiolabeled 1,25-dihydroxyvitamin D₃. Large amounts of radioactivity that behaved chromatographically as identical with authentic 1,25-dihydroxyvitamin D₃ were present in the plasma, bone, and intestine of the intact, sham-operated, or ureter-ligated rats. However, no radioactivity eluting in a manner similar to 1,25-dihydroxyvitamin D₃ was found in plasma, bone, or intestine of acutely nephrectomized rats. We conclude that, in the acutely nephrectomized living rat, 1,25-dihydroxyvitamin D₃ is not present in plasma, bone, or intestine in quantities detectable by the sensitive techniques we have used. No conversion of 25-hydroxyvitamin D₃ to 1,25-dihydroxyvitamin D₃ was observed during a 24-hr period after nephrectomy of vitamin D-deprived rats. This fact casts doubt upon the significance of the *in vitro* production of 1,25-dihydroxyvitamin D₃ by nonrenal cells as an *in vivo* phenomenon.

Vitamin D₃ is metabolized to its most biologically active form, 1,25-dihydroxyvitamin D₃ [1,25(OH)₂D₃], via an intermediate, 25-hydroxyvitamin D₃ [25(OH)D₃] (1). Fraser and Kodicek (2), and subsequently Gray *et al.* (3), found that radiolabeled 1,25(OH)₂D₃ was absent from the tissues and plasma of acutely nephrectomized rats that had received low-specific-activity radiolabeled vitamin D₃ or 25(OH)D₃ immediately after nephrectomy. In human anephric subjects given radiolabeled vitamin D₃, Mawer *et al.* (4) failed to detect radioactive 1,25(OH)₂D₃. Gray *et al.* (5) administered 25(OH)[³H]D₃ to 10 anephric subjects; in none of the patients studied was radiolabeled 1,25(OH)₂D₃ detected. In these human studies, radiolabeled 1,25(OH)₂D₃ was clearly detectable in normal subjects (4, 5). Lambert *et al.* (6) found recently that a substance was present in the plasma of anephric subjects that displaced 1,25(OH)₂[³H]D₃ from its receptor; additionally, samples processed in a similar manner displaced ⁴⁵Ca from labeled fetal bone in a bioassay system for 1,25(OH)₂D₃. The authors concluded that 1,25(OH)₂D₃ was present in the plasma of anephric subjects. However, other investigators, using radioreceptor, radioimmune, and biological assays for 1,25(OH)₂D₃, have been unable to detect 1,25(OH)₂D₃ in anephric persons (7–12).

In certain situations, however, 1,25(OH)₂D₃ may be synthesized extrarenally. For example, a report by Barbour *et al.* (13)

showed that an anephric subject with sarcoidosis had detectable plasma 1,25(OH)₂D₃-like material at a time when he was hypercalcemic (13); hypercalcemia abated after prednisone treatment that presumably diminished the size of the sarcoid granulomas in the patient. It is likely that sarcoid granulomatous tissue produced the 1,25(OH)₂D₃. Pregnant nephrectomized rats may synthesize 1,25(OH)₂D₃ *in vivo* (14, 15). The site of production in pregnant animals could be the placenta because 25(OH)D₃ is converted to 1,25(OH)₂D₃ by placental tissue of rat and human origin *in vitro* (16). Whether the placenta is the sole site of 1,25(OH)₂D₃ production in the nephrectomized pregnant animal is not known because fetal kidneys could also serve as the source of it. Chicken calvarial cells and human bone cells metabolize 25(OH)[³H]D₃ to a material that behaves chromatographically like 1,25(OH)₂[³H]D₃ (17, 18). The nature of the material was not verified in the original publications, although a recent report suggests that the material is 1,25(OH)₂D₃ on the basis of mass spectroscopic analysis (†). Two other groups of investigators (19, ‡) have reported that macrophages and cultured intestinal cells can make 1,25(OH)₂D₃-like material *in vitro*.

Papers alleging such production of 1,25(OH)₂D₃ *in vitro* generally lack rigorous proof of the material's structure, and those studies failing to detect 1,25(OH)₂D₃ in anephric individuals suffer from potential problems such as use of low-specific-activity ³H-labeled 25(OH)D₃ or vitamin D₃ and use of pharmacologic amounts of precursor. Because this issue is important to the understanding of calcium metabolism, we have reexamined the ability of anephric rats to synthesize 1,25(OH)₂D₃ *in vivo* by using high-specific-activity labeled 25(OH)D₃ in physiologic amounts as the precursor to 1,25(OH)₂D₃.

MATERIALS AND METHODS

General. The methods used for HPLC, determination of ultraviolet absorbance, and measurement of radioactivity have been reported (20). Counting efficiency for ³H was ≈ 50%.

25(OH)[26,27-³H]D₃ (148 or 158 Ci/mmol; 1 Ci = 3.7 × 10¹⁰ Bq) and 1,25(OH)₂[26,27-³H]D₃ (160 Ci/mmol) were purchased from Amersham and New England Nuclear, respectively. All radiochemicals were purified on a 1 × 60 cm Sephadex LH-20 column with a chloroform/*n*-hexane, 65:35 (vol/vol), solvent system. In both cases, the 25(OH)[³H]D₃ and 1,25(OH)₂[³H]D₃ were the only peaks eluting from these col-

Abbreviations: 25(OH)D₃, 25-hydroxyvitamin D₃; 24R,25(OH)₂D₃, (24R)-24,25-dihydroxyvitamin D₃; 1,25(OH)₂D₃, 1,25-dihydroxyvitamin D₃; 25S,26(OH)₂D₃, (25S)-25,26-dihydroxyvitamin D₃.

* To whom reprint requests should be addressed.

† Howard, G. A., Turner, R. T., Puzas, J. E., Knapp, D. K. & Nichols, F., Fifth Workshop on Vitamin D, Williamsburg, VA, Feb. 18, 1982, p. 3 (abstracts).

‡ Roswell, R. H. & Young, M. J., Fourth Annual Scientific Meeting of the American Society of Bone Mineral Research, San Francisco, June 15, 1982, p. 58 (abstracts).

The publication costs of this article were defrayed in part by page charge payment. This article must therefore be hereby marked "advertisement" in accordance with 18 U. S. C. §1734 solely to indicate this fact.

umns. Radiochemical purity of each product was established by co-HPLC with authentic $25(\text{OH})\text{D}_3$ and $1,25(\text{OH})_2\text{D}_3$; the radiolabeled material was >99% pure. Nonradiolabeled $25(\text{OH})\text{D}_3$ was obtained from Upjohn; $(24R)$ - $24,25$ -dihydroxyvitamin D_3 [$24R,25(\text{OH})_2\text{D}_3$], $(25S)$ - $25,26$ -dihydroxyvitamin D_3 [$25S,26(\text{OH})_2\text{D}_3$], and $1,25(\text{OH})_2\text{D}_3$ were obtained from Hoffmann-LaRoche.

Animals. Male albino weanling rats (50–60 g) were purchased from Holtzman (Madison, WI). We maintained them in individual hanging wire cages in a room devoid of UV light and fed them ad lib a vitamin D-deficient diet containing 0.47% calcium and 0.3% phosphorus (21). After 7 weeks on this diet, seven bilaterally nephrectomized rats and six intact control rats were given $25(\text{OH})[^3\text{H}]\text{D}_3$ as described below. In a second experiment, another group of similar rats was maintained on an identical diet for 7 weeks. Six rats were bilaterally nephrectomized, seven rats underwent bilateral ureteral ligation, and seven rats were subjected to a sham operation. These rats received $25(\text{OH})[^3\text{H}]\text{D}_3$ as described below. Additionally, plasma was obtained from 10 vitamin D-deplete rats raised for 7 weeks on the diet described above and used for measurement of $25(\text{OH})\text{D}_3$ and $1,25(\text{OH})_2\text{D}_3$ levels by described methods (8, 22, 23). Another group of eight rats received 10 international units (250 ng) of vitamin D_3 per day orally for 7 weeks. They served as a positive control group for the measurement of plasma $25(\text{OH})\text{D}_3$ and $1,25(\text{OH})_2\text{D}_3$. Plasma creatinine and blood urea nitrogen values were measured by described techniques (24, 25).

Metabolism of $25(\text{OH})[26,27\text{-}^3\text{H}]\text{D}_3$ in Vivo. In the first experiment, six rats with intact kidneys and seven bilaterally nephrectomized rats were lightly anesthetized with ether and given (intrajugular injection) 25 pmol of $25(\text{OH})[^3\text{H}]\text{D}_3$ (3.6 μCi) dissolved in 68 μl of 95% ethanol. In the second experiment, six bilaterally nephrectomized, seven sham-operated, and seven ureter-ligated rats received (intrajugular injection) 25 pmol of $25(\text{OH})[^3\text{H}]\text{D}_3$ (3.8 μCi) dissolved in 88 μl of 95% ethanol. We took 0.3 ml of blood prior to administration of the dose in both experiments for measurement of serum calcium.

Twenty-four hours after the injection of $25(\text{OH})[^3\text{H}]\text{D}_3$, rats were exsanguinated and plasma was separated from the blood cells. We immediately removed the entire small intestine and the fore- and hindlimbs. We dissected the fore- and hindlimb bones free from adhering muscle and ligaments and weighed them. The bones were then cooled in liquid nitrogen and pulverized. We removed the intestinal contents by flushing with ice-cold saline and then scraped off the mucosa with a glass microscope slide. Radioactivity was measured in homogenates of intestine and in plasma prior to lipid extraction. Plasma, 50% aqueous suspensions of pulverized bone, and intestinal mucosa were extracted with methanol/chloroform, 2:1 (vol/vol), as described (26). The aqueous layer was reextracted with an equal volume of chloroform. We then evaporated the chloroform fractions to dryness under reduced pressure and resuspended them in the appropriate column solvent.

In the first set of experiments, the residues from plasma, intestine, or bone were suspended in 1 ml of chloroform/*n*-hexane, 65:35 (vol/vol), and chromatographed on Sephadex LH-20 columns (1 \times 60 cm) developed and eluted with 350 ml of the same solvent. Five-milliliter fractions of the eluate were collected, an aliquot of each fraction was transferred to a 5-ml scintillation vial and dried, and radioactivity was determined (20). After chromatography, the Sephadex LH-20 in the column was washed with methanol. The methanol wash was collected and dried in order to determine the presence of more polar radioactive material.

In the second set of experiments, the residues from the plasma, bone, and intestine were suspended in an appropriate volume

of *n*-hexane/isopropanol, 9:1 (vol/vol), and applied directly to a HPLC system as described below.

HPLC of Metabolites. We pooled and dried (under nitrogen) fractions containing radioactivity that eluted from Sephadex LH-20 columns in a manner identical to $1,25(\text{OH})_2\text{D}_3$ or $25S,26(\text{OH})_2\text{D}_3$. The radioactive material was redissolved in 0.1 ml of *n*-hexane/isopropanol, 9:1 (vol/vol). We added 90 ng of authentic $1,25(\text{OH})_2\text{D}_3$ to the radioactive samples prior to chromatography on a 0.4×30 cm $\mu\text{Porasil}$ column (Waters Associates) in an *n*-hexane/isopropanol, 9:1 (vol/vol), solvent system developed at a flow rate of 2 ml/min and a pressure of 800 psi. In order to determine the elution volume of $25S,26(\text{OH})_2\text{D}_3$, 100 ng was chromatographed in the same system. Absorbance of the effluent was monitored continuously at 265 nm.

In the second experiment, 2 μg of each authentic $25(\text{OH})\text{D}_3$, $24R,25(\text{OH})_2\text{D}_3$, $25S,26(\text{OH})_2\text{D}_3$, and $1,25(\text{OH})_2\text{D}_3$ was added to the plasma, intestine, and bone lipid extracts. Chromatography was performed on a 0.4×30 cm $\mu\text{Porasil}$ column with a *n*-hexane/isopropanol, 9:1 (vol/vol), solvent system run at a flow rate of 2 ml/min. Fractions were collected at 0.5-min intervals. Effluent from the $1,25(\text{OH})_2\text{D}_3$ region was further chromatographed on a reverse-phase 0.4×30 cm C_{18} $\mu\text{Bondapak}$ column with a methanol/water, 8:2 (vol/vol), solvent system




FIG. 1. Representative chromatograms of the lipid extracts of plasma, bone, and intestinal mucosa taken 24 hr after intact and nephrectomized rats had received 25 pmol of $25(\text{OH})[^3\text{H}]\text{D}_3$. Chromatography was performed on a Sephadex LH-20 column (1 \times 60 cm) developed in and eluted with 350 ml of chloroform/*n*-hexane, 65:35 (vol/vol). After 350 ml, the column was washed with methanol. Arrows show the elution position of authentic $25(\text{OH})[^3\text{H}]\text{D}_3$ and $1,25(\text{OH})_2[^3\text{H}]\text{D}_3$. (Inserts) Expanded scales for radioactivity in actual dpm.

run at a flow rate of 2 ml/min. Fractions were collected at 0.5-min intervals.

RESULTS

25(OH)₂D₃ levels were undetectable in the vitamin D₃-deficient group and the mean ± SEM was 9.2 ± 0.6 ng/ml in the group given D₃. 1,25(OH)₂D₃ levels were <5 pg/ml in the vitamin D-deficient group and 213.4 ± 16.5 pg/ml in the vitamin D₃-treated group. Mean (± SEM) serum calcium levels were 4.8 ± 0.2 mg/dl in the vitamin D-deficient sham-operated, ureter-ligated, and nephrectomized animals and 9.9 ± 0.1 mg/dl in the control vitamin D₃-treated group (*P* < 0.001). Plasma creatinine and blood urea nitrogen levels were significantly higher (*P* < 0.001) in the nephrectomized and ureter-ligated rats than in the sham-operated controls (nephrectomized, 4.83 ± 0.04 and 151.6 ± 3.4 mg/dl; ureter-ligated, 4.37 ± 0.08 and 133.4 ± 6.4 mg/dl; sham-operated, 0.47 ± 0.02 and 10.09 mg/dl).

The distributions of radioactivity in the plasma, bone, and intestine of vitamin D-deficient intact and nephrectomized rats were comparable after an intravenous injection of 25 pmol of 25(OH)[³H]D₃. Twenty-four hours after injection, plasma contained the largest percentage of the radioactivity administered per gram of tissue followed by intestine and then bone. From 98% to 104% of the radioactivity in the plasma and 72% to 90% of that in the intestine was chloroform extractable.

In plasma, bone, and intestinal tissues from intact rats a polar metabolite(s) appeared at the elution position of 1,25(OH)₂D₃ or 25S,26(OH)₂D₃ on Sephadex LH-20 chromatography (Fig. 1; Table 1). Plasma, intestine, and bone from intact rats also contained large amounts of radioactive material eluting in a manner identical to 1,25(OH)₂D₃ on HPLC (Table 2). On Sephadex LH-20 chromatography, the plasma lipid extracts from nephrectomized animals also contained a small radioactive peak eluting in the position of 1,25(OH)₂D₃ or 25S,26(OH)₂D₃ (Fig. 1; Table 1). However, when this peak was applied to the straight phase

Table 2. HPLC of Sephadex LH-20 elution peaks representing 1,25(OH)₂D₃ or 25S,26(OH)₂D₃

Tissue	dpm applied to column	dpm eluted as		Total recovered
		1,25(OH) ₂ D ₃	25S,26(OH) ₂ D ₃	
Plasma:				
Intact (n = 5)	5,923 ± 590	4,812 ± 492 (81.4 ± 2.5)	0	5,382 ± 490 (91.4 ± 3.0)
Nx (n = 5)	838 ± 64	0	577 ± 36 (69.6 ± 3.8)	789 ± 61 (94.7 ± 4.0)
Bone:				
Intact (n = 6)	2,852 ± 148	2,278 ± 156 (79.6 ± 2.6)	0	2,553 ± 164 (89.4 ± 2.6)
Nx (n = 7)	0	0	0	0
Intestine:				
Intact (n = 6)	1,961 ± 143	1,671 ± 99 (85.6 ± 1.4)	0	1,793 ± 101 (90.6 ± 1.8)
Nx (n = 7)	0	0	0	0

See legend to Table 1.

HPLC system, it migrated not with authentic 1,25(OH)₂D₃ but in a manner similar to authentic 25S,26(OH)₂D₃ (Table 2). Bone and intestine from nephrectomized vitamin D-deficient rats contained no detectable radioactivity in the elution position of 1,25(OH)₂D₃ or 25S,26(OH)₂D₃ (Fig. 1; Tables 1 and 2). 25(OH)D₃ was the major metabolite appearing in plasma, bone, and intestine of nephrectomized rats 24 hr after injection of radiolabeled 25(OH)D₃, accounting for 88–101% of the radioactivity.

Table 1. Radioactive vitamin D metabolites in plasma, bone, and intestine after injection of 25(OH)[26,27-³H]D₃ to intact and nephrectomized (Nx) rats

Tissue	dpm applied to column	dpm eluted					Total recovered
		Less polar than 25(OH)D ₃	25(OH)D ₃	Less polar than 1,25(OH) ₂ D ₃	1,25(OH) ₂ D ₃ or 25S,26(OH) ₂ D ₃	Column strip	
Plasma:							
Intact (n = 5)	512,964 ± 81,720	9,742 ± 4,569 (1.6 ± 0.6)	281,626 ± 33,089 (57.0 ± 3.5)	2,236 ± 1,308 (0.4 ± 0.2)	112,600 ± 8,297 (24.6 ± 4.5)	18,580 ± 6,372 (3.8 ± 1.3)	424,784 ± 43,432 (87.2 ± 7.2)
Nx (n = 5)	756,247 ± 43,115	16,511 ± 2,473 (2.2 ± 0.3)	666,645 ± 75,440 (87.7 ± 7.5)	727 ± 300 (0.1 ± 0.04)	2,489 ± 192 (0.33 ± 0.01)	5,598 ± 701 (0.8 ± 0.1)	692,095 ± 78,093 (91.0 ± 7.6)
Bone:							
Intact (n = 6)	30,201 ± 3,684	645 ± 118 (2.1 ± 0.2)	18,874 ± 2,085 (63.8 ± 4.5)	137 ± 72 (0.4 ± 0.2)	12,286 ± 2,013 (40.0 ± 3.0)	1,344 ± 165 (4.5 ± 0.3)	33,286 ± 3,928 (110.7 ± 4.7)
Nx (n = 7)	34,464 ± 3,746	1,232 ± 198 (3.5 ± 0.2)	34,927 ± 3,997 (101.3 ± 3.1)	0	0	1,246 ± 176 (3.7 ± 0.5)	37,272 ± 4,241 (108.1 ± 3.0)
Intestine:							
Intact (n = 6)	155,196 ± 21,112	3,555 ± 583 (2.3 ± 0.2)	14,064 ± 1,273 (9.8 ± 1.4)	4,564 ± 562 (3.0 ± 0.2)	119,320 ± 15,255 (77.5 ± 3.1)	11,618 ± 1,686 (7.6 ± 1.0)	153,146 ± 17,581 (100.2 ± 2.4)
Nx (n = 7)	37,929 ± 4,038	2,700 ± 434 (6.9 ± 0.4)	33,831 ± 4,292 (89.5 ± 7.0)	469 ± 146 (1.3 ± 0.5)	0	6,102 ± 1,113 (15.4 ± 1.4)	45,140 ± 5,801 (112.9 ± 7.2)

Metabolites are referred to by their elution positions (Sephadex LH-20 column) relative to authentic 25(OH)D₃ and 1,25(OH)₂D₃. n, Number of animals per experiment; Nx, nephrectomized. Results are shown as mean ± SEM. The numbers in parentheses represent percentage of total counts applied.

Table 3. HPLC of lipid extracts from plasma, bone, and intestine after injection of 25(OH)[³H]D₃ to sham-operated, ureter-ligated, and nephrectomized rats

Tissue	Distribution of ³ H among metabolites, %			
	25-(OH)D ₃	25S,26-(OH) ₂ D ₃	1,25-(OH) ₂ D ₃	Total recovered
Plasma:				
Sham-operated (n = 7)	60.5 ± 1.4	0.8 ± 0	29.0 ± 1.2	90.8 ± 1.3
Ureter-ligated (n = 6)	73.9 ± 1.3	0.7 ± 0.02	15.3 ± 0.6	90.3 ± 1.4
Nx (n = 6)	92.8 ± 3.9	0.6 ± 0.04	0	93.7 ± 3.8
Bone:				
Sham-operated (n = 6)	53.7 ± 1.8	0	38.1 ± 1.9	91.9 ± 1.3
Ureter-ligated (n = 7)	68.4 ± 0.8	0	26.5 ± 1.7	95.2 ± 1.4
Nx (n = 6)	93.5 ± 2.5	0	0	93.7 ± 2.5
Intestine:				
Sham-operated (n = 7)	9.2 ± 1.1	0	76.8 ± 4.0	86.2 ± 4.2
Ureter-ligated (n = 7)	17.9 ± 0.7	0	50.9 ± 3.6	69.2 ± 3.0
Nx (n = 6)	71.4 ± 2.3	0	0	72.5 ± 2.1

See legend to Table 1.

The results of the second experiment are shown in Table 3. Radioactivity coeluting with 1,25(OH)₂D₃ was present in the plasma, bone, and intestine of sham-operated and ureter-ligated rats. None was observed in the plasma, bone, or intestine of nephrectomized rats. Radioactivity coeluting with authentic 1,25(OH)₂D₃ from the straight phase HPLC of sham-operated or ureter-ligated rats also coeluted with 1,25(OH)₂D₃ on the reverse-phase system.

DISCUSSION

Our data are consistent with prior reports that 1,25(OH)₂D₃ is not detected in plasma and intestine of acutely nephrectomized rats (2, 3). In addition, our results show that 1,25(OH)₂D₃ is not present in the bone of nephrectomized rats. The animals used in our experiments had been maintained on a vitamin D-deficient diet for 7 weeks, resulting in serum calcium levels ≈50% of normal and no detectable 25(OH)D₃ and 1,25(OH)₂D₃ in plasma. Under these circumstances, the 25(OH)D₃ 1α-hydroxylase should be maximally stimulated (1). Furthermore, because we used high-specific-activity 25(OH)[³H]D₃ as the radiolabeled precursor, we were able to give physiologic amounts of 25(OH)D₃ (≈25 pmol) but large amounts of radioactivity (3.6–3.8 μCi) to each rat. The sensitivity of our technique would have allowed us to detect as little as 100–200 dpm of radioactive 1,25(OH)₂D₃ in various tissues. This is about 0.1% of the amount of radioactivity actually present in each intact rat plasma or intestinal sample (≈100,000 dpm; see Fig. 1 and Table 1) and about 1% of that present in a bone sample (≈12,000 dpm). Thus, if the nephrectomized rats had circulating plasma or intestinal tissue levels of 1,25(OH)₂D₃ that were even 0.1% of those present in intact rats, we would have been able to detect them. Similarly, had bone from nephrectomized rats produced 1,25(OH)₂D₃ in amounts 1% of those of intact rats we would have detected this metabolite.

That uremia *per se* is not responsible for the lack of 1,25(OH)₂D₃ production is shown by the ureter-ligated group of rats, in which 1,25(OH)₂D₃ production was observed (Table

3) despite blood urea nitrogen and plasma creatinine values similar to those in nephrectomized animals. Our data confirm and strengthen the observations of Fraser and Kodicek (2) and Gray *et al.* (3) in rats. Our observations are also supported by the fact that anephric man does not produce radiolabeled 1,25(OH)₂D₃ from an appropriate precursor (4, 5). The absence of circulating 1,25(OH)₂D₃ in anephric humans as measured by radioreceptor, radioimmunoassay, and biological assays also supports this contention (7–12).

Lambert *et al.* (6) recently detected material in the plasma of anephric patients that resembles 1,25(OH)₂D₃ in a radioreceptor assay and a bioassay. It is conceivable that some interfering substance was present in the sera from anephric persons that caused displacement of 1,25(OH)₂[³H]D₃ from the cytosolic intestinal receptor and mobilization of ⁴⁵Ca from fetal long bones. Certainly, vitamin D-deficient plasma contains materials that displace 1,25(OH)₂[³H]D₃ from its receptor—hence the need for extensive purification before assay (7–12). It should also be kept in mind that both laboratories involved in the Lambert *et al.* study (6) previously reported plasma 1,25(OH)₂D₃ levels to be undetectable in anephric subjects (10, 12). If, as one must assume, larger amounts of plasma were extracted than before, then one cannot dismiss the possibility that larger amounts of interfering material were also extracted. The authors did not present results from appropriate control samples—i.e., vitamin D-deficient plasma samples of a volume equal to that extracted in the anephric patients. It is also puzzling that, with the stated sensitivity of the bioassay method, ≈1–2 pg/ml, 1,25(OH)₂D₃ levels were not detectable earlier (12) in 2- to 5-ml plasma samples from anephric persons [the average value of 1,25(OH)₂D₃ reported later in the non-vitamin-D-supplemented anephrics was 3.2–7.9 pg/ml] (6). The report by Lambert *et al.* (6) should be interpreted cautiously because of the points raised above and because of contrary evidence from groups using diverse techniques (2–5, 7–12, 27). Despite these criticisms, we must emphasize that our experiments were performed in acutely nephrectomized rats and the possibility of induction of the 25(OH)D₃ 1α-hydroxylase in the chronic uremic state cannot be dismissed.

Several laboratories have reported that 1,25(OH)₂D₃ is made in tissues other than the kidney (14–19, †). In the pregnant animal this may well be the placenta or the fetoplacental unit (14–16). The *in vitro* data are supported by the *in vivo* observations that nephrectomized pregnant rats can synthesize a material resembling 1,25(OH)₂D₃. Chicken and human bone cells reportedly synthesize 1,25(OH)₂D₃ *in vitro*. The existence of 1,25(OH)₂D₃ in these cultures has also been confirmed recently by mass spectrometry (†). The cultures were maintained in serum-free media wherein vitamin D-binding protein is absent, a circumstance that does not occur *in vivo*. This might influence the ability of these cells to synthesize 1,25(OH)₂D₃. Reports showing that intestinal cells synthesize 1,25(OH)₂D₃ are preliminary; the identity of the putative 1,25(OH)₂D₃ has not been unambiguously established (‡). A report that macrophages synthesize 1,25(OH)₂D₃ has now been withdrawn (19, §). Our data show that 1,25(OH)₂D₃ is not detectable in the plasma, bone, or intestine of anephric rats. Although it is possible that certain cells can synthesize 1,25(OH)₂D₃ *in vitro*, it is unlikely that this phenomenon is of significance *in vivo*.

We conclude that the acutely nephrectomized rat does not form 1,25(OH)₂D₃ in amounts detectable by highly sensitive methods. If any 1,25(OH)₂D₃ is formed outside the kidney of

§ Gray, T. K., Maddux, F. W., Mentz, W. M. & Williams, M. E., Fourth Annual Scientific Meeting of the American Society of Bone Mineral Research, San Francisco, June 14, 1982, p. S52 (abstracts).

acutely nephrectomized rats, it must be in amounts less than 0.1% of normal. Our data may not be applicable to the anephric human because our experiments were performed in the acutely nephrectomized state which may not mimic the chronic anephric state. Our experiments, however, do question the role of *in vitro* production of $1,25(\text{OH})_2\text{D}_3$ by nonrenal cells.

This work was supported in part by National Institutes of Health Grants AM-25409 (R.K.), AM-19607 (H.H.), AM-27440 (H.H. and J.F.), and AM-07147 (T.S.).

1. DeLuca, H. F. (1979) *Nutr. Rev.* **37**, 161-193.
2. Fraser, D. R. & Kodicek, E. (1970) *Nature (London)* **228**, 764-766.
3. Gray, R., Boyle, I. & DeLuca, H. F. (1971) *Science* **172**, 1232-1234.
4. Mawer, E. B., Blackhouse, J., Lumb, G. A. & Stanbury, S. W. (1971) *Nature (London) New Biol.* **232**, 188-189.
5. Gray, R. W., Weber, H. P., Dominguez, J. H. & Lemann, J., Jr. (1974) *J. Clin. Endocrinol. Metab.* **39**, 1045-1056.
6. Lambert, P. W., Stern, P. H., Avioli, R. C., Brackett, N. C., Turner, R. T., Greene, A., Fu, I. Y. & Bell, N. H. (1982) *J. Clin. Invest.* **69**, 722-725.
7. Brumbaugh, P. F., Haussler, D. H., Bressler, R. & Haussler, M. R. (1974) *Science* **183**, 1089-1091.
8. Eisman, J. A., Hamstra, J., Kream, B. E. & DeLuca, H. F. (1976) *Science* **193**, 1021-1023.
9. Bouillon, R., De Moor, P., Baggiolini, E. G. & Uskokovic, M. R. (1980) *Clin. Chem.* **26**, 562-567.
10. Lambert, P. W., DeOreo, P. B., Hollis, B. W., Fu, I. Y., Ginsberg, D. J. & Roos, B. A. (1981) *J. Lab. Clin. Med.* **98**, 536-548.
11. Gray, T. K., McAdoo, T., Pool, D., Lester, G. E., Williams, M. E. & Jones, G. (1981) *Clin. Chem.* **27**, 458-463.
12. Stern, P. H., Phillips, T. E. & Mavreas, T. (1980) *Anal. Biochem.* **102**, 22-30.
13. Barbour, G. L., Coburn, J. W., Slatopolsky, E., Norman, A. W. & Horst, R. L. (1981) *N. Engl. J. Med.* **305**, 440-443.
14. Weisman, Y., Vargas, A., Duckett, G., Reiter, E. & Root, A. W. (1978) *Endocrinology* **103**, 1992-1996.
15. Gray, T. K., Lester, G. E. & Lorenc, R. S. (1979) *Science* **204**, 1311-1313.
16. Tanaka, Y., Halloran, B., Schnoes, H. K. & DeLuca, H. F. (1979) *Proc. Natl. Acad. Sci. USA* **76**, 5033-5035.
17. Turner, R. T., Puzas, J. E., Forte, M. D., Lester, G. E., Gray, T. K., Howard, G. A. & Baylink, D. J. (1980) *Proc. Natl. Acad. Sci. USA* **77**, 5720-5724.
18. Howard, G. A., Turner, R. T., Sherrard, D. J. & Baylink, D. J. (1981) *J. Biol. Chem.* **256**, 7738-7740.
19. Gray, T. K., Maddux, F. W., Mentz, W. M. & Williams, M. E. (1982) *Clin. Res.* **30**, 524 (abstr.).
20. Kumar, R., Nagubandi, S., Mattox, V. R. & Londowski, J. M. (1980) *J. Clin. Invest.* **65**, 277-284.
21. Suda, T., DeLuca, H. F. & Tanaka, Y. (1970) *J. Nutr.* **100**, 1049-1052.
22. Eisman, J. A., Shepard, R. M. & DeLuca, H. F. (1977) *Anal. Biochem.* **80**, 298-305.
23. Kumar, R., Cohen, W. R., Silva, P. & Epstein, F. H. (1979) *J. Clin. Invest.* **63**, 342-344.
24. DiGiorgio, J. (1974) in *Clinical Chemistry Principles and Techniques*, eds. Henry, R. J., Cannon, D. C. & Winkelman, J. W. (Harper & Row, New York), pp. 541-553.
25. Faulkner, W. R. & King, J. W. (1976) in *Fundamentals of Clinical Chemistry*, ed. Tietz, N. W. (Saunders, Philadelphia), pp. 991-994.
26. Bligh, E. G. & Dyer, W. J. (1959) *Can. J. Biochem. Physiol.* **37**, 911-917.
27. Stern, P. H., Hamstra, A. J., DeLuca, H. F. & Bell, N. H. (1978) *J. Clin. Endocrinol. Metab.* **46**, 891-896.