

Does the medical literature remain inadequately described despite having reporting guidelines for 21 years? – A systematic review of reviews: an update

Yanling Jin,^{1,*} Nitika Sanger,^{2,*} Ieta Shams,^{3,*} Candice Luo,^{4,*} Hamnah Shahid,^{5,*} Guowei Li,^{1,*} Meha Bhatt,¹ Laura Zielinski,⁶ Bianca Bantoto,⁷ Mei Wang,¹ Luciana PF Abbade,⁸ Ikunna Nwosu,⁴ Alvin Leenus,¹ Lawrence Mbuagbaw,¹ Muhammad Maaz,¹ Yaping Chang,¹ Guangwen Sun,¹ Mitchell AH Levine,^{1,9} Jonathan D Adachi^{1,9} Lehana Thabane,^{1,9} Zainab Samaan^{1,10}

¹Department of Health Research Methods, Evidence and Impact, McMaster University, Hamilton, ON, Canada; ²Department of Medical Science, Medical Sciences Graduate Program, McMaster University, Hamilton, ON, Canada; ³Department of Psychology, Neuroscience and Behaviour, McMaster University, Hamilton, ON, Canada; ⁴Faculty of Health Sciences, Bachelors of Health Sciences, McMaster University, Hamilton, ON, Canada; ⁵Department of Arts and Science, McMaster University, Hamilton, ON, Canada; ⁶Department of Neuroscience, McMaster Integrative Neuroscience Discovery and Study, McMaster University, Hamilton, ON, Canada; ⁷Department of Science, Honours Integrated Sciences Program, McMaster University, Hamilton, ON, Canada; ⁸Department of Dermatology and Radiotherapy, Botucatu Medical School, Universidade Estadual Paulista, UNESP, São Paulo, Brazil; ⁹St. Joseph's Healthcare Hamilton, Hamilton, ON, Canada; ¹⁰Department of Psychiatry and Behavioural Neurosciences, McMaster University, Hamilton, ON, Canada

*These authors contributed equally to this work

Correspondence: Zainab Samaan
Department of Psychiatry and Behavioural Neurosciences, McMaster University, 1280 Main Street West, L8S 4L8 Hamilton, ON, Canada
Tel +1 905 522 1155x35448
Fax +1 905 381 5629
Email samaanz@mcmaster.ca

Purpose: Reporting guidelines (eg, Consolidated Standards of Reporting Trials [CONSORT] statement) are intended to improve reporting standards and enhance the transparency and reproducibility of research findings. Despite accessibility of such guidelines, researchers are not required to adhere to them. Our goal was to determine the current status of reporting quality in the medical literature and examine whether adherence of reporting guidelines has improved since the inception of reporting guidelines.

Materials and methods: Eight reporting guidelines, such as CONSORT, Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA), STrengthening the Reporting of OBservational studies in Epidemiology (STROBE), Quality of Reporting of Meta-analysis (QUOROM), STAndards for Reporting of Diagnostic accuracy (STARD), Animal Research: Reporting In Vivo Experiments (ARRIVE), Consolidated Health Economic Evaluation Reporting Standards (CHEERS), and Meta-analysis of Observational Studies in Epidemiology (MOOSE) were examined. Our inclusion criteria included reviews published between January 1996 to September 2016 which investigated the adherence to reporting guidelines in the literature that addressed clinical trials, systematic reviews, observational studies, meta-analysis, diagnostic accuracy, economic evaluations, and preclinical animal studies that were in English. All reviews were found on Web of Science, Excerpta Medical Database (EMBASE), MEDLINE, and Cumulative Index to Nursing and Allied Health Literature (CINAHL).

Results: Among the general searching of 26,819 studies by using the designed searching method, 124 studies were included post screening. We found that 87.9% of the included studies reported suboptimal adherence to reporting guidelines. Factors associated with poor adherence included non-pharmacological interventions, year of publication, and trials concluding with significant results. Improved adherence was associated with better study designs such as allocation concealment, random sequence, large sample sizes, adequately powered studies, multiple authorships, and being published in journals endorsing guidelines.

Conclusion: We conclude that the level of adherence to reporting guidelines remains suboptimal. Endorsement of reporting guidelines by journals is important and recommended.

Keywords: guidelines, adherence, review, CONSORT

Introduction

Medical science is an evolving and dynamic field of research that impacts health care, disease outcomes, and health care systems in general. The evidence generated from millions of medical publications is meant to inform these dynamic changes

and therefore has to be presented in a clear, consistent, and transparent fashion. There are more than 26 million citations for biomedical literature in the PubMed¹ database alone. To understand and evaluate the evidence presented in these citations, a harmonized method of reporting the research findings is needed to ensure clarity, consistency, and the uptake and dissemination of knowledge.² Tremendous efforts have been made to provide guidelines for different types of research designs to assist in the process of transparent and clear reporting, eg, Enhancing the QUALity and Transparency Of health Research (EQUATOR) Network website.³ However, despite the wide availability of such guidelines since the inception of the Consolidated Standards of Reporting Trials (CONSORT⁴) statement in 1996, the uptake remains suboptimal in the face of the exponential volume of medical literature leaving the readers confused. For example, some studies show positive harmful results from eating red meat on the risk of having colorectal cancer,⁵ while others are showing inconsistent effect marked by substantial methodological differences, type of red meat investigated, and the population selection limitations.⁶ Therefore, the reader is unable to decide whether red meat has an effect on bowel cancer risk. Poor reporting without using well-designed guidelines in primary studies may lead to a bias in the treatment effects found in systematic reviews. In addition, poorly conducted systematic reviews may not be able to detect the bias effect that the studies included. In a previous study, we conducted a scoping review and examined the level of adherence to six reporting guidelines and found the level of adherence to be suboptimal in 86% of the included studies.⁷

The aim of this review was to conduct a systematic review of reviews to update the state of adherence to guidelines since 2012 and to identify factors associated with improved adherence. Our hypothesis was that the reporting standards have improved since our last examination in 2012 given that a longer period has passed after guideline statements were first introduced for researchers and more journals started to endorse the guidelines. Our search was looking at reviews published between January 1, 1996, and September 30, 2016.

Materials and methods

This systematic review was performed and reported according to the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) guidelines.⁸ A protocol for a series of three reviews including the current systematic review has been peer reviewed and published elsewhere.⁹

Study inclusion and exclusion criteria

Systematic reviews which investigated the adherence to commonly used reporting guidelines in medical literature that addressed clinical trials, systematic reviews, observational studies, meta-analysis, diagnostic accuracy, economic evaluations, and preclinical animal studies that have been reported in English were selected. Eight guidelines included in this review were as follows: CONSORT,⁴ PRISMA,⁸ STrengthening the Reporting of OBservational studies in Epidemiology (STROBE),¹⁰ Quality of Reporting of Meta-analysis (QUOROM),¹¹ STAndards for Reporting of Diagnostic accuracy (STARD),¹² Animal Research: Reporting In Vivo Experiments (ARRIVE),¹³ Consolidated Health Economic Evaluation Reporting Standards (CHEERS),¹⁴ and Meta-analysis of Observational Studies in Epidemiology (MOOSE).¹⁵

The exclusion criteria included studies that 1) were not systematic reviews; 2) did not explore adherence to the aforementioned reporting guidelines; 3) did not provide data on guideline adherence; 4) were subsets of the included studies; 5) published abstracts, letters, editorials, or commentaries; and 6) reviews in languages other than English for feasibility and resource purposes.

Search strategy

The search strategy was based on the previously published review⁷ and was updated for this systematic review. We searched four databases (Excerpta Medical Database [EMBASE], MEDLINE, Cumulative Index to Nursing, and Allied Health Literature [CINAHL], and Web of Science) from 1996 (CONSORT inception – first created guideline among all eight included guidelines) to September 30, 2016.

We used the following search terms for each of the four databases: (Systematic reviews OR reviews OR quality of reporting OR completeness of reporting) AND (CONSORT OR STROBE OR QUOROM OR PRISMA OR MOOSE OR STARD OR ARRIVE OR CHEERS) OR adherence. Detailed search terms have been reported in the published protocol.⁹ All stages of search, inclusion, exclusion, and data abstraction were performed independently in duplicate, and agreement was reached through team discussion and consensus.

Outcome measures

The primary outcome was the level of adherence to reporting guidelines and their checklists as reported in the systematic reviews. The secondary outcome included the factors that were associated with improved adherence to guidelines.

Data extraction

A specific data abstraction form was designed to include the following data: 1) general characteristics of the included studies (first author, publication year, country, journal, study field, search time frame, data sources, numbers of included primary studies, and study design), 2) main findings from the included studies, 3) authors' summaries and conclusions, and 4) factors reported to be related to improved guideline reporting adherence. Each assessment of the systematic reviews was conducted in duplicate. Calibration was performed on the data extraction form. If the pair of evaluators was unable to come to a conclusion, a third-party reviewer would have settled the dispute.

Quality evaluation

We used the modified Assessing the Methodological Quality of Systematic Reviews/Overview of Quality Assessment Questionnaire (Assessment of Multiple Systematic Reviews [AMSTAR]/Overview Quality Assessment Questionnaire [OQAQ]), a 10-item scale,⁷ to assess the quality of the

systematic reviews included in this review. We assigned a number out of a maximum of 20 points for each included study. The higher the number assigned, the better the quality of the systematic review.

Data synthesis

We provided a qualitative summary and characteristics of the included studies. We summarized the factors associated with adherence based on the included study results; no quantitative analysis was possible in this review. We also reported the percentage of studies in which the level of adherence to reporting each guideline was suboptimal. This was calculated by dividing the number of studies with this finding by the total number of studies evaluating the guideline.

Results

Our search resulted in a total of 9,123 publications, of which 124 systematic reviews that included 26,819 primary studies were included in this systematic review of reviews. Figure 1 shows the PRISMA flowchart for the included studies.




Figure 1 PRISMA flow diagram.

Abbreviation: PRISMA, Preferred Reporting Items for Systematic Reviews and Meta-Analyses.

The characteristics of the included studies are described in Table 1. The majority of the studies (65% of the total 124 included studies) investigated the adherence to the CONSORT statement as expected since it is the first and oldest guideline. The second most commonly investigated guideline is the PRISMA with 19 studies (15%; Table 1).

The majority of studies used the guideline checklist to evaluate the level of adherence and generated a mean score as summarized in [Table S1](#). [Table S1](#) summarizes the studies' findings by guideline with authors' conclusions for each study. Most studies described the adherence to the different guidelines using the following qualitative descriptors:

deficient, not adequately reported, generally poor, sub-optimal, poor, medium, low, poor to moderate, lack of CONSORT adherence, bad, far from satisfactory, lack of standard reporting, improvement over the years has been minor, weak, quality of the articles varied substantially, insufficient, missed reporting some important factors, deficiencies in reporting, inconsistent, needs to be improved, inadequate, there is a need for improvement in quality of reporting, overall adherence is low.

A summary of the quantitative assessment of adherence to guidelines is presented in Table 2.

The level of adherence to all included reporting guidelines was 87.9% of all guidelines combined showing a need for improvement in reporting. Factors associated with poor adherence to CONSORT guideline included trials with significantly positive results, trials with the categorical outcome, trials conducted in North America compared to Europe, and trials funded by nonindustry source. A summary of factors associated with adherence standards is summarized in Table 3. Several factors were associated with better reporting standards relating to authors, study design, outcome specifications, year of publication (recent years of publications are associated with better reporting standards), journal, funding source, and study/author country.

Factors associated with improved adherence to reporting guidelines

Author factors

The included studies reported that the expertise of the author team, for example, an epidemiologist, improved the quality of reporting the study. In addition, having multiple authors also improved reporting quality.

Study factors

Study design with detailed methods including allocation concealment, randomization, specific outcome measures, sample

size and power calculations, acknowledgment of limitations and sources of bias, larger sample size, registration of clinical trials, pharmacological interventions, and detailed statistical analysis plan were associated with better reporting and adherence to reporting guidelines. Year of publication was also associated with adherence in which the more recently published articles had increased adherence.

Journal factor

Publications in journals endorsing reporting guidelines have better adherence to these guidelines than articles published in journals that do not endorse such guidelines. In addition, journals' impact factor, medical journals, and journals with restriction on the number of words per article also had articles with better reporting standards. Publication in a general medical journal was associated with better reporting quality than a specialty journal.

Ethics and funding factors

Articles that reported ethical approval, participants' consent, and the source of funding were associated with improved adherence to reporting guidelines.

Country of study factors

Geographic location of the study has an impact on the quality of reporting and adherence to reporting guidelines, for example, studies reported from Europe had better reporting standards compared to studies from North America. Studies reported from China had lower adherence to guidelines than elsewhere indicating geographical variations may directly or indirectly impact the level of adherence to reporting guidelines in the medical literature.

Quality assessment of included studies

For each included systematic review, we performed a quality assessment using the modified AMSTAR/OQAQ score. Table 4 provides the total score out of 20 for each study. The scores varied from 9 to 20. The average score for all the included studies is 16.14. The lowest scores were related to items 5 and 6 of the quality assessment related to the availability of the primary studies' characteristics similar to a previously reported study.⁷ Items 5 and 6 were evaluated if there was information on included and excluded studies provided and if the characteristics of included studies provided, respectively.

Discussion

The medical literature is paramount to the progression of the understanding of health and disease and the establishment

Table 1 Characteristics of the included studies

Study	Year	Journal	Country	Statement assessed	Number of studies
Adie ²⁸	2013	<i>Annals of Surgery</i>	Australia	CONSORT	150
Adie et al ²⁹	2015	<i>Annals of Surgery</i>	Australia	PRISMA	150
Agha et al ³⁰	2015	<i>Annals of Plastic Surgery</i>	UK	STROBE	94
Agha et al ³¹	2016	<i>International Journal of Surgery</i>	UK	CONSORT	193
				PRISMA	
				STROBE	
Aguar et al ³²	2014	<i>Annals of Pharmacotherapy</i>	Brazil	PRISMA	7
Aguar et al ³³	2016	<i>Journal of Clinical Pharmacy and Therapeutics</i>	Brazil	CHEERS	8
Al Faleh and Al-Omran ³⁴	2009	<i>BMC Pediatrics</i>	Saudi Arabia	QUOROM	61
Al-Namankany et al ³⁵	2009	<i>International Journal of Pediatric Dentistry</i>	UK	CONSORT	173
Alvarez et al ³⁶	2009	<i>British Journal of Dermatology</i>	France	CONSORT	98
Anttila et al ³⁷	2006	<i>Pediatrics</i>	Finland	CONSORT	15
Areia et al ³⁸	2010	<i>Endoscopy</i>	Portugal	CONSORT	120
Augestad et al ³⁹	2012	<i>Journal of the American Medical Informatics Association</i>	Norway	CONSORT	32
Balasubramanian et al ⁴⁰	2006	<i>Annals of Surgery</i>	UK	CONSORT	69
Bath and Bath ⁴¹	2000	<i>Stroke</i>	UK	CONSORT	114
Bereza et al ⁴²	2008	<i>Annals of Pharmacotherapy</i>	Canada	QUOROM	16
Bian et al ⁴³	2006	<i>Journal of Chinese Integrative Medicine</i>	People's Republic of China	CONSORT	66
Biondi-Zoccai et al ⁴⁴	2006	<i>BMJ</i>	Italy	QUOROM	10
Borg Debono et al ⁴⁵	2012	<i>BMC Anesthesiology</i>	Canada	CONSORT	23
Bousquet et al ⁴⁶	2011	<i>Journal of Allergy and Clinical Immunology</i>	France	CONSORT	94
Bramhall et al ⁴⁷	2015	<i>Inflammatory Bowel Diseases</i>	UK	ARRIVE	58
Cairo et al ⁴⁸	2012	<i>Journal of Clinical Periodontology</i>	Spain	CONSORT	276
Capili et al ⁴⁹	2010	<i>Clinical Journal of Pain</i>	USA	CONSORT	10
Cavadas et al ⁵⁰	2011	<i>International Urogyn J</i>	Portugal	CONSORT	41
Choi et al ⁵¹	2014	<i>Trials</i>	South Korea	CONSORT	29
Chowers et al ⁵²	2009	<i>Journal of Antimicrobial Chemotherapy</i>	Israel	CONSORT	49
Cook et al ⁵³	2011	<i>Medical Education</i>	USA	STROBE	130
Daitch et al ⁵⁴	2016	<i>Journal of Pediatric Gastroenterology and Nutrition</i>	Israel	CONSORT	51
Dasi et al ⁵⁵	2012	<i>Journal of Clinical Pharmacology</i>	Spain	CONSORT	40
Delaney et al ⁵⁶	2010	<i>Transfusion</i>	USA	STROBE,	47
				CONSORT	
DeMauro et al ⁵⁷	2011	<i>Pediatrics</i>	USA	CONSORT	179
de Vries and van Roon ⁵⁸	2010	<i>Archives of Diseases in Childhood</i>	The Netherlands	CONSORT	107
Dias et al ⁵⁹	2006	<i>Human Reproduction</i>	UK	CONSORT	164
Ethgen et al ⁶⁰	2009	<i>BMC Medical Research Methodology</i>	France	CONSORT	132
Eyawo et al ⁶¹	2008	<i>Trials</i>	Canada	CONSORT	47
Fan et al ⁶²	2014	<i>PLoS One</i>	China	CONSORT	21
Farrokhyar et al ⁶³	2007	<i>Canadian Journal of Surgery</i>	Canada	CONSORT	50
Fidalgo et al ⁶⁴	2015	<i>Ophthalmic and Physiological Optics</i>	UK	STARD	58
Fleming et al ⁶⁵	2013	<i>Angle Orthodontist</i>	UK	PRISMA	109
Fontela et al ⁶⁶	2009	<i>PLoS One</i>	Canada	STARD	90
Freeman et al ⁶⁷	2009	<i>European Journal of Obstetrics & Gynecology and Reproductive Biology</i>	UK	STARD	27
Froud et al ⁶⁸	2012	<i>Community Dentistry and Oral Epidemiology</i>	UK	CONSORT	23
Fung et al ⁶⁹	2009	<i>Ophthalmology</i>	USA	CONSORT,	36
				STROBE	
Gagnier et al ⁷⁰	2006	<i>American Journal of Medicine</i>	Canada	CONSORT	206
Gao et al ⁷¹	2015	<i>Trials</i>	China	CONSORT	98
Gianola et al ⁷²	2013	<i>Physical Therapy</i>	Italy	PRISMA	88
Gohari et al ⁷³	2016	<i>Journal of Diabetes and Metabolic Disorders</i>	Iran	CONSORT	185
Gulin et al ⁷⁴	2015	<i>PLoS Neglected Tropical Diseases</i>	Argentina	ARRIVE	83
Halpern et al ⁷⁵	2004	<i>International Journal of Obstetric Anesthesia</i>	Canada	CONSORT	99
Hemels et al ⁷⁶	2004	<i>Current Medical Research and Opinion</i>	France	QUOROM	32

(Continued)

Table 1 (Continued)

Study	Year	Journal	Country	Statement assessed	Number of studies
Herdan et al ⁷⁷	2011	<i>Gynecological Surgery</i>	Germany	CONSORT	37
Huang et al ⁷⁸	2015	<i>Expert Review of Anticancer Therapy</i>	China	CONSORT	40
Hui et al ⁷⁹	2012	<i>Support Care Cancer</i>	USA	CONSORT	44
Junhua et al ⁸⁰	2007	<i>The Journal of Complementary and Alternative Medicine</i>	China	QUOROM	107
Karpouzis and Bonello ⁸¹	2016	<i>Chiropractic and Manual Therapies</i>	Australia	CONSORT	35
Kiehna et al ⁸²	2010	<i>Journal of Neurosurgery</i>	USA	CONSORT	27
Kim et al ⁸³	2014	<i>BMJ Open</i>	South Korea	CONSORT	146
Kober et al ⁸⁴	2006	<i>Journal of the National Cancer Institute</i>	Australia	CONSORT	142
Ladd et al ⁸⁵	2010	<i>Addictive Behaviors</i>	USA	CONSORT	127
Lee et al ⁸⁶	2013	<i>Trauma Acute Care Surgery</i>	UK	CONSORT	83
Lee et al ⁸⁷	2016	<i>JAMA Facial Plastic Surgery</i>	UK	PRISMA	79
Li et al ⁸⁸	2011	<i>Evidence-based Complementary and Alternative Medicine</i>	USA	CONSORT	42
Li et al ⁸⁹	2014	<i>Systematic Reviews</i>	China	PRISMA	487
Li et al ⁹⁰	2014	<i>BMC Complementary and Alternative Medicine</i>	China	CONSORT	6994
Liu et al ⁹¹	2015	<i>PLoS One</i>	China	PRISMA	72
Liu et al ⁹²	2013	<i>Transplant International</i>	UK	CONSORT	290
Liu et al ⁹³	2015	<i>Journal of Evidence-based Medicine</i>	China	CONSORT	76
Liu et al ⁹⁴	2014	<i>PLoS One</i>	China	PRISMA	476
Liu et al ⁹⁵	2016	<i>PLoS One</i>	China	ARRIVE	396
Lu et al ⁹⁶	2015	<i>Archives of Physical Medicine and Rehabilitation</i>	USA	CONSORT	105
Lu et al ⁹⁷	2011	<i>Expert Review of Anticancer Therapy</i>	China	CONSORT	46
Ma et al ⁹⁸	2011	<i>PLoS One</i>	China	PRISMA	369
Ma et al ⁹⁹	2012	<i>The Journal of Alternative and Complementary Medicine</i>	China	PRISMA	88
Marshman and Farid ¹⁰⁰	2010	<i>Community Dental Health</i>	UK	CONSORT	48
McCormick et al ¹⁰¹	2013	<i>Journal of Shoulder and Elbow Surgery</i>	USA	CONSORT	54
Miller et al ¹⁰²	2009	<i>Academic Radiology</i>	Canada	STARD	18
Moberg-Mogren and Nelson ¹⁰³	2006	<i>American Journal of Occupational Therapy</i>	USA	CONSORT	14
Moher et al ¹⁰⁴	2002	<i>BMC Pediatrics</i>	Canada	CONSORT	251
Montané et al ¹⁰⁵	2010	<i>BMC Clinical Pharmacology</i>	Spain	CONSORT	92
Montgomery et al ¹⁰⁶	2011	<i>Trials</i>	UK	CONSORT	76
Nicolau et al ¹⁰⁷	2013	<i>The International Journal of Tuberculosis and Lung Disease</i>	Canada	PRISMA	137
Norton-Mabus and Nelson ¹⁰⁸	2008	<i>OTJR: Occupation, Participation and Health</i>	USA	CONSORT	30
Ntala et al ¹⁰⁹	2013	<i>Primary Care Respiratory Journal</i>	Greece	CONSORT	35
Panic et al ¹¹⁰	2013	<i>PLoS One</i>	Italy	PRISMA	90
Parsons et al ¹¹¹	2011	<i>Journal of Bone and Joint Surgery, British Volume</i>	UK	CONSORT	100
Patel et al ¹¹²	2014	<i>Psychological Medicine</i>	UK	CONSORT	31
Piggott et al ¹¹³	2004	<i>Palliative Medicine</i>	UK	CONSORT	93
Péron et al ¹¹⁴	2012	<i>Journal of the National Cancer Institute</i>	France	CONSORT	357
Peters et al ¹¹⁵	2015	<i>PLoS One</i>	The Netherlands	PRISMA	80
Plint et al ¹¹⁶	2006	<i>Medical Journal of Australia</i>	Canada	CONSORT	8
Prady et al ¹¹⁷	2008	<i>PLoS One</i>	UK	CONSORT	90
Pratoomsoot et al ¹¹⁸	2015	<i>PLoS One</i>	Thailand	CONSORT	71
Rao et al ¹¹⁹	2016	<i>PLoS One</i>	UK	STROBE	37
Rice et al ¹²⁰	2016	<i>Journal of Psychosomatic Research</i>	Canada	PRISMA	21
Rios et al ¹²¹	2008	<i>Journal of Clinical Endocrinology and Metabolism</i>	Canada	CONSORT	89
Rikos et al ¹²²	2016	<i>Multiple Sclerosis and Related Disorders</i>	Greece	CONSORT	102
Schwarz et al ¹²³	2012	<i>Journal of Clinical Periodontology</i>	Germany	ARRIVE	75
Scott et al ¹²⁴	2012	<i>The Pediatric Infectious Disease Journal</i>	Switzerland	CONSORT	70
Shawyer et al ¹²⁵	2015	<i>Journal of Pediatric Surgery</i>	Canada	STROBE	48
Shea et al ¹²⁶	2006	<i>BMC Medical Research Methodology</i>	Canada	QUOROM	53
Shea et al ¹²⁷	2006	<i>The Journal of Rheumatology</i>	The Netherlands	QUOROM	57
Stevely et al ¹²⁸	2015	<i>PLoS One</i>	UK	CONSORT	68

(Continued)

Table 1 (Continued)

Study	Year	Journal	Country	Statement assessed	Number of studies
Strech et al ¹²⁹	2011	<i>Journal of Clinical Psychiatry</i>	Germany	CONSORT	105
Tan et al ¹³⁰	2014	<i>International Journal of Surgery</i>	UK	PRISMA	37
Thabane et al ¹³¹	2007	<i>International Journal of Obesity</i>	Canada	CONSORT	63
Tunis et al ¹³²	2013	<i>Radiology</i>	Canada	PRISMA	130
Turner et al ¹³³	2012	<i>Cochrane Database of Systematic Reviews</i>	Canada	CONSORT	45
Vigna-Taglianti et al ¹³⁴	2006	<i>Annals of Oncology</i>	Italy	QUOROM	80
Walleiser et al ¹³⁵	2011	<i>Journal of Clinical Epidemiology</i>	Switzerland	CONSORT	106
Wang et al ¹³⁶	2007	<i>Clinical Therapeutics</i>	China	CONSORT	7422
Wang et al ¹³⁷	2013	<i>PLoS One</i>	China	CONSORT	27
Wangge et al ¹³⁸	2010	<i>PLoS One</i>	The Netherlands	CONSORT	232
Weingärtner et al ¹³⁹	2016	<i>Expert Review of Clinical Pharmacology</i>	Germany	CONSORT	117
Weir et al ¹⁴⁰	2012	<i>International Journal of Medical Informatics</i>	USA	PRISMA QUOROM	13
Wen et al ¹⁴¹	2008	<i>Journal of Clinical Epidemiology</i>	China	QUOROM	161
Willis and Quigley ¹⁴²	2011	<i>BMC Medical Research Methodology</i>	UK	PRISMA	236
Yao et al ¹⁴³	2014	<i>Eye</i>	UK	CONSORT	65
Zafar et al ¹⁴⁴	2008	<i>Clinical and Experimental Ophthalmology</i>	Pakistan	STARD	76
Zhang ¹⁴⁵	2015	<i>BMJ Open</i>	China	MOOSE	607
Zhao et al ¹⁴⁶	2016	<i>Medicine</i>	China	CONSORT	68
Zheng et al ¹⁴⁷	2016	<i>Open Heart</i>	UK	CONSORT	33
Zhong et al ¹⁴⁸	2011	<i>European Journal of Integrated Medicine</i>	China	CONSORT	153
Zintzaras et al ¹⁴⁹	2010	<i>Clinical Therapeutics</i>	Greece	CONSORT	18
Zintzaras et al ¹⁵⁰	2012	<i>BMC Musculoskeletal Disorders</i>	Greece	STARD	103
Ziogas and Zintzaras ¹⁵¹	2009	<i>Annals of Epidemiology</i>	Greece	CONSORT	261

Abbreviations: ARRIVE, Animal Research: Reporting In Vivo Experiments; BMC, *BioMed central*; BMJ, *British Medical Journal*; CHEERS, Consolidated Health Economic Evaluation Reporting Standards; CONSORT, Consolidated Standards of Reporting Trials; *International Urogyn J*, *International Urogynecology Journal*; JAMA, *The Journal of the American Medical Association*; MOOSE, Meta-analysis of Observational Studies in Epidemiology; OTJR, *Occupational Therapy Journal of Research*; PLoS, Public Library of Science; PRISMA, Preferred Reporting Items for Systematic reviews and Meta-Analyses; QUOROM, Quality of Reporting of Meta-analysis; STARD, Standards for Reporting of Diagnostic Accuracy; STROBE, Strengthening the Reporting of Observational Studies in Epidemiology.

Table 2 Summary of the included studies' conclusions

Type of guideline	Total number of studies	Studies reporting inadequate adherence ^a
CONSORT	81 (three combined studies with both CONSORT and STROBE; one combined study with STROBE, CONSORT, and PRISMA)	71 (88%)
PRISMA	19 (one combined study with both PRISMA and QUOROM; one combined study with STROBE, CONSORT, and PRISMA)	16 (84%)
STROBE	8 (three combined studies with both CONSORT and STROBE; one combined study with STROBE, CONSORT, and PRISMA)	7 (88%)
QUOROM	10 (one combined study with both PRISMA and QUOROM)	5 (50%)
STARD	6	5 (83%)
ARRIVE	4	4 (100%)
CHEERS	1	1 (100%)
MOOSE	1	1 (100%)
All guidelines	124 (distinct studies)	109 (87.9%)

Note: ^aThe number of studies concluding that "some improvements are needed, reporting inadequate, poor, medium, suboptimal, etc."

Abbreviations: ARRIVE, Animal Research: Reporting In Vivo Experiments; CHEERS, Consolidated Health Economic Evaluation Reporting Standards; CONSORT, Consolidated Standards of Reporting Trials; MOOSE, Meta-analysis of Observational Studies in Epidemiology; PRISMA, Preferred Reporting Items for Systematic Reviews and Meta-Analyses; QUOROM, Quality of Reporting of Meta-analysis; STARD, Standards for Reporting of Diagnostic Accuracy; STROBE, Strengthening the Reporting of Observational Studies in Epidemiology.

of priorities and recommendations for prevention, diagnosis, treatment, and measurement of outcomes. To implement research findings, transparent and consistent reporting standards are needed to help make informed decisions. Such

standards have been set by the CONSORT working group and others for the past 2 decades with the aim of improving the reporting standards in biomedical research. It is expected that the introduction of new change to the current practice will

Table 3 Factors associated with reporting quality of articles using the CONSORT guideline

Study	Year	Sample size	Factors associated with adherence (↑↓)
Adie ²³	2013	150	Outcome specification (↑) ^a At least one author with a degree in epidemiology (↑) ^a Length of article in words (↑) ^a Allocation concealment (↑) ^a Random sequence (↑) ^a Power calculation (↑)
Agha et al ³¹	2016	193	Greater details on study design (↑) Detailed outcome definitions and measurements (↑) Indication of how quantitative variables were handled during analyses (↑) Discussion of limits and potential sources of bias (↑)
Al-Namankany et al ³⁵	2009	173	Year of publication (↑)
Alvarez et al ³⁶	2006	98	Pharmaceutical industry funding (↑) ^a Year of publication (↑) ^a Sample size (↑) ^a
Areia et al ³⁸	2010	120	Publication in CONSORT-endorsing journals (↑) Year of publication (↑)
Balasubramanian et al ⁴⁰	2006	69	Number of authors (↑) ^a Multicenter studies (↑) ^a Declared funding sources (↑) ^a Reporting in medical journals (↑) ^a
Bath and Bath ⁴¹	2000	114	Trial quality (↑) ^a Trials with positive outcome (↓) ^a Year of publication (↑) ^a
Borg Debanio et al ⁴⁵	2012	23	Impact factor (↑) Funding reported (↑) Journal adopted CONSORT statement at the time of data collection (↑) Sample size (↑)
Cairo et al ⁴⁸	2012	64	Year of publication (↑) ^a Statistically significant clinical outcomes – positive study results (↓) ^a
Capili et al ⁴⁹	2010	10	Journal requiring the use of CONSORT (↑)
Chowers et al ⁵²	2009	49	Industry-sponsored trials (industry-sponsored vs. nonindustry-sponsored trial) (↑) Year of publication (↑) ^a
de Vries and van Roon ⁵⁸	2010	107	Sponsoring (↑)
DeMauro et al ⁵⁷	2011	179	Time trend (↑) ^a Journal type – general medical journals vs. pediatric journals (↑) ^a
Ethgen et al ⁶⁰	2009	132	Impact factor (↑) ^a Publication in CONSORT-endorsing journals (↑) ^a
Farrokhyar et al ⁶³	2007	50	Sample size (↑) ^a Year of publication (↑) ^a Location of the study (↑) ^a Source of funding (↓) Type of primary outcome in the study (categorical) (↓)
Gao et al ⁷¹	2015	98	Supported by funding (↑) ^a
Herdan et al ⁷⁷	2011	37	Year of publication (↑) ^a
Karpouzis and Bonello ⁸¹	2016	35	Year of publication (↑) ^a Larger sample size (↑) ^a
Kiehna et al ⁸²	2010	27	Publication in CONSORT-endorsing journals (↑) ^a
Kim et al ⁸³	2014	146	Year of publication (↑) ^a
Ladd et al ⁸⁵	2010	127	Year of publication (↑) ^a
Lee et al ⁸⁶	2013	83	Higher impact factor of journal (↑) ^a Journals requiring submission of CONSORT checklist (↑) ^a
Liu et al ⁹²	2013	290	Reporting of funding (↑) Journal endorses CONSORT (↑) ^a Good-quality RCTs (high Jadad scores) (↑) ^a Allocation concealment (↑) ^a Data analysis by randomized group (↑) ^a Sample size > 100 (↑) ^a

(Continued)

Table 3 (Continued)

Study	Year	Sample size	Factors associated with adherence (↑↓)
Liu et al ⁹³	2015	76	Journal adopting CONSORT guidelines (↑) ^a Later publication year (↑) ^a
Lu et al ⁹⁶	2015	105	Year of publication (1976–2001, 2002–2010, 2011–2013) (↑) ^a
McCormick et al ¹⁰¹	2013	54	High Jadad score (↑) ^a
Moberg-Mogren and Nelson ¹⁰³	2006	14	Year of publication (↑) ^a
Montané et al ¹⁰⁵	2010	92	Impact factor (↑) ^a Year of publication (↑) ^a
Montgomery et al ¹⁰⁶	2011	76	Year of publication (↑) ^a
Ntala et al ¹⁰⁹	2013	35	Impact factor (↑) Country with high income (↑) ^a
Péron et al ¹¹⁴	2012	357	Trials with positive results (↓) Year of publication (↑) ^a Impact factor (↑) ^a Geographic region – North American compared to European trials (↓) ^a Sample size (↑)
Plint et al ¹¹⁶	2006	8	Overall consort items (↑) Reporting method of sequence generation (↑) ^a Allocation concealment (↑) ^a
Prady et al ¹¹⁷	2008	90	Standardized page length (↑) Year of publication (↑) ^a
Pratoomsoot et al ¹¹⁸	2015	71	Country of publication (ASEAN ^b vs. plus six) (↑ for some factors for ASEAN; ↑ for some factors for plus six)
Rikos et al ¹²²	2016	102	After the publication of CONSORT (↑) Impact factor (↑) ^a Year of publication (↑) ^a
Rios et al ¹²¹	2008	89	Sample size (↑) ^a Industrial funding (↑) ^a Journal of publication (publication in JCEM) (↑) ^a
Scott et al ¹²⁴	2012	70	Trial registration (↑) Year of publication (↑) Trial size (↑)
Thabane et al ¹³¹	2007	63	Type of intervention (pharmacological intervention vs. non-pharmacological intervention) (↑) ^a Sample sizes (↑) ^a Year of publication (↑) ^a
Turner et al ¹³³	2012	45	Time trend (↑) ^a
Yao et al ¹⁴³	2014	65	Number of authors (↑) Impact factor (↑)
Zhao et al ¹⁴⁶	2016	68	Year of publication (↑) Reporting of funding (↑) Reporting of informed consent form (↑) Reporting of ethical approval (↑)
Zheng et al ¹⁴⁷	2016	33	Number of authors (↑) ^a Number of patients (↑) ^a Impact factor (↑) ^a Time trend (↑) ^a Number of participants (↑) Treatment duration (↑) Reporting of funding (↑)
Zhong et al ¹⁴⁸	2011	153	Non-Chinese reports (compared to those published in mainland China) (↑) ^a Publication in CONSORT-endorsing journals (↑) ^a
Ziogas and Zintzaras ¹⁵¹	2009	261	Year of publication (↑) ^a Impact factor (↑) ^a

Notes: ^aStatistically significant increase/decrease, $p \leq 0.05$; (↑), positively associated with adherence; (↓), negatively associated with adherence. The number of studies concluding that “some improvements are needed, reporting inadequate, poor, medium, suboptimal, etc”. ^bAssociation of Southeast Asian nations, Association of Southeast Asian Nations (ASEAN) plus six groups, which composed of the members of the ASEAN plus Australia, China, India, Japan, New Zealand, and South Korea.

Abbreviations: CONSORT, Consolidated Standards of Reporting Trials; JCEM, *The Journal of Clinical Endocrinology and Metabolism*; RCT, randomized control trial.

Table 4 Reporting quality of the 124 included systematic reviews, assessed by the modified AMSTAR/OQAQ (10 items, score out of 20)

Study	Global score
Adie ²⁸	17
Adie et al ²⁹	18
Agha et al ³⁰	15
Agha et al ³¹	14
Aguiar et al ³²	14
Aguiar et al ³³	19
Al Faleh and Al-Omran ³⁴	16
Al-Namankany et al ³⁵	15
Alvarez et al ³⁶	10
Anttila et al ³⁷	15
Areia et al ³⁸	18
Augestad et al ³⁹	20
Balasubramanian et al ⁴⁰	16
Bath and Bath ⁴¹	16
Bereza et al ⁴²	20
Bian et al ⁴³	15
Biondi-Zoccai et al ⁴⁴	15
Borg Debanò et al ⁴⁵	9
Bousquet et al ⁴⁶	18
Bramhall et al ⁴⁷	10
Cairo et al ⁴⁸	19
Capili et al ⁴⁹	15
Cavadas et al ⁵⁰	17
Choi et al ⁵¹	17
Chowers et al ⁵²	12
Cook et al ⁵³	18
Daitch et al ⁵⁴	17
Dasi et al ⁵⁵	19
Delaney et al ⁵⁶	14
DeMauro et al ⁵⁷	17
de Vries and van Roon ⁵⁸	18
Dias et al ⁵⁹	17
Ethgen et al ⁶⁰	13
Eyawo et al ⁶¹	18
Fan et al ⁶²	18
Farrokhyar et al ⁶³	19
Fidalgo et al ⁶⁴	18
Fleming et al ⁶⁵	15
Fontela et al ⁶⁶	17
Freeman et al ⁶⁷	11
Froud et al ⁶⁸	16
Fung et al ⁶⁹	17
Gagnier et al ⁷⁰	16
Gao et al ⁷¹	13
Gianola et al ⁷²	12
Gohari et al ⁷³	15
Gulin et al ⁷⁴	14
Halpern et al ⁷⁵	14
Hemels et al ⁷⁶	19
Herdan et al ⁷⁷	15
Huang et al ⁷⁸	12
Hui et al ⁷⁹	18
Junhua et al ⁸⁰	13
Karpouzis and Bonello ⁸¹	16

(Continued)

Table 4 (Continued)

Study	Global score
Kiehna et al ⁸²	16
Kim et al ⁸³	16
Kober et al ⁸⁴	17
Ladd et al ⁸⁵	19
Lee et al ⁸⁶	16
Lee et al ⁸⁷	17
Li et al ⁸⁸	18
Li et al ⁸⁹	15
Li et al ⁹⁰	14
Liu et al ⁹¹	19
Liu et al ⁹²	16
Liu et al ⁹³	14
Liu et al ⁹⁴	17
Liu et al ⁹⁵	19
Lu et al ⁹⁶	18
Lu et al ⁹⁷	18
Ma et al ⁹⁸	19
Ma et al ⁹⁹	16
Marshman and Farid ¹⁰⁰	14
McCormick et al ¹⁰¹	16
Miller et al ¹⁰²	17
Moberg-Mogren and Nelson ¹⁰³	16
Moher et al ¹⁰⁴	14
Montané et al ¹⁰⁵	15
Montgomery et al ¹⁰⁶	17
Nicolau et al ¹⁰⁷	16
Norton-Mabus and Nelson ¹⁰⁸	10
Ntala et al ¹⁰⁹	18
Panic et al ¹¹⁰	11
Parsons et al ¹¹¹	17
Patel et al ¹¹²	13
Piggott et al ¹¹³	14
Péron et al ¹¹⁴	15
Peters et al ¹¹⁵	17
Plint et al ¹¹⁶	18
Prady et al ¹¹⁷	19
Pratoomsoot et al ¹¹⁸	15
Rao et al ¹¹⁹	18
Rice et al ¹²⁰	19
Rios et al ¹²¹	20
Rikos et al ¹²²	17
Schwarz et al ¹²³	10
Scott et al ¹²⁴	16
Shawyer et al ¹²⁵	15
Shea et al ¹²⁶	13
Shea et al ¹²⁷	19
Stevely et al ¹²⁸	18
Strech et al ¹²⁹	18
Tan et al ¹³⁰	14
Thabane et al ¹³¹	19
Tunis et al ¹³²	18
Turner et al ¹³³	20
Vigna-Taglianti et al ¹³⁴	15
Walleiser et al ¹³⁵	19
Wang et al ¹³⁶	15
Wang et al ¹³⁷	17

(Continued)

Table 4 (Continued)

Study	Global score
Wangge et al ¹³⁸	12
Weingärtner et al ¹³⁹	17
Weir et al ¹⁴⁰	20
Wen et al ¹⁴¹	18
Willis and Quigley ¹⁴²	20
Yao et al ¹⁴³	16
Zafar et al ¹⁴⁴	16
Zhang ¹⁴⁵	18
Zhao et al ¹⁴⁶	17
Zheng et al ¹⁴⁷	18
Zhong et al ¹⁴⁸	17
Zintzaras et al ¹⁴⁹	18
Zintzaras et al ¹⁵⁰	14
Ziogas and Zintzaras ¹⁵¹	15

Abbreviations: AMSTAR, Assessment of Multiple Systematic Reviews; OQQAQ, Overview Quality Assessment Questionnaire.

take time to adopt and disseminate. However, the uptake of the widely available guidelines has been less than ideal. We define suboptimal and less than ideal as <100%. The whole idea of a systematic review is to have completely transparent methods reported, so everyone can follow and reproduce the results. Inherently, systematic reviews are meant to be a more rigorous study design. This allows them to produce meaningful results than individual studies. Thus, when reviews fail to adhere to reporting guidelines, it calls into question the consistency of their results. Given the weight that systematic reviews have in the scientific community, it is imperative that we hold reviews to a high standard.

Five years ago, we investigated the level of adherence to reporting standards in the medical literature, and we identified 86% of the systematic reviews conducted on the level of adherence to reporting guidelines of the medical literature to be less than ideal.⁷ Since our previous scoping review, many new revisions and updates to reporting guidelines have been introduced. Currently, there are 358 reporting guidelines on the EQUATOR Network website¹⁶ for many study types that are freely available. However, endorsement of reporting guidelines by journals still remains low.

Among all the factors that can improve the reporting quality, such as author factors, study factors, journal factors, ethics and funding factors, and country of study factors, author factors as well as their limitations have been studied in other researches. The author factors were the number of the authors of the publication and the level of expertise in the different research methods. Multiple authorships were shown to be an important determinant of the impact of the research being produced and its likelihood of being cited.¹⁷ The complexity and cost of medical research today requires multiple levels of

expertise in various disciplines as well as accountability and oversight by study team members, institutions, and funding bodies. It is known that the number of authors per article has increased over the past few decades^{18,19} with a concern posed to question the roles of multiple authors and the most senior academics holding senior authorship at the expense of others in the team.²⁰ Other studies have reported that the research produced by teams rather than single authors was impactful and more frequently cited, at least in certain fields.²¹ It is likely that multiple authorships arising from collaborative efforts have advantages of producing good quality impactful research; however, multiple authorships also have limitations and may not be feasible at every setting due to geographical limitations or strict timeline to follow as bringing more authors is time-consuming.²² In this review, we found that having multiple authorships is important to have publications with better adherence to reporting guidelines. However, the role of each author and the hierarchy of authorship should be clarified for successful collaborations and research impact as discussed earlier.

Study factors that improved adherence to reporting guidelines included well-designed, detailed study methods and adequately powered studies. Study results could be altered regarding trial designs, qualities, and methods.²³ Therefore, guidelines such as CONSORT statement that is designed for randomized control trials (RCTs), STROBE guideline for observational studies, and PRISMA guideline for systematic reviews were invented accordingly based on different study designs. RCTs are also considered as the highest level of primary evidence in the clinical practice, and therefore it is vital that these trials are reported according to the expected standards.²⁴

Other factors reported that might improve the level of adherence to reporting guidelines included journals endorsing these guidelines. The Internal Committee of Medical Journal Editors (ICMJE) recognized the importance of reporting guidelines in ensuring study details that are described adequately to be evaluated appropriately and encouraged journals to request these reporting standards from authors.²⁵ The EQUATOR Network has valuable resources and tool kits to assist authors and journal editors to adopt the reporting guidelines and provide case studies of journals endorsing the guidelines. Since journals that endorsed reporting guidelines often ask authors to submit a completed checklist regarding the guidelines, it improves the quality of reporting for those journals endorsing these guidelines. Yet, not all journals currently endorse the guidelines. According to the CONSORT website, there are 585 journals that endorse CONSORT,²⁶

while there are about 30,000 journals indexed in PubMed.²⁷ While not all of these indexed journals publish RCTs, many of them do publish them, but do not adhere to CONSORT guidelines.²⁷

The EQUATOR Network also has tool kits for ethics boards and study sponsors to ensure that the reporting guidelines are considered when these agencies review research submissions for ethical approval or funding requests. It is therefore important that all stakeholders take part in the use and dissemination of the reporting guidelines to enhance the quality of medical research and biomedical literature.

Limitations

The included studies are limited to only eight of the reporting guidelines, and therefore the current study lacks the generalizability to other guidelines that may have a better adherence standard. In addition, there was no comparison between studies to ensure that they are using qualitative descriptors such as “inadequate” or “suboptimal” with the same operational definition. The studies do not provide sufficient information regarding the operationalization of qualitative descriptors to allow us to adequately compare descriptors across studies.

In addition, the study was limited to systematic reviews that present with its own set of limitations. The most notable limitation is the low mean score on the quality assessment since each systematic review follows different reporting guidelines or does not follow guidelines at all and the lack of detailed data on the included studies’ characteristics. Furthermore, a quantitative analysis was not conducted, as not all included studies provided relevant data. Strict inclusion criteria may have allowed a quantitative analysis. However, for the sake of a more representative sample, such criteria were not implemented.

The inclusion of studies in English only is also a limitation to a selected section of the medical literature and did not include other reporting guidelines that may be in use in other languages.

Despite the limited scope of inclusion criteria and quality limitation of the included studies, this review provides an insight into the limited uptake of reporting guidelines and calls for exploring barriers to such uptake. Future studies may include broad surveys of authors, journal editors, funding agencies, ethics boards, and readers to solicit opinions and understanding of the role of reporting guidelines in the medical research and literature.

Conclusion

Current adherence to reporting guidelines in the medical literature is suboptimal. However, there are factors associated with better reporting upon which we can develop strategies

for better reporting. Reporting guidelines are an imperative tool in the endeavor to improve the consistency of reporting in the medical literature. However, the suboptimal uptake and correct usage of reporting guidelines demonstrate the need for further emphasis in the scientific community to encourage the use of reporting guidelines. The responsibility for improving the transparency, quality, and reproducibility of medical literature lies with all stakeholders from the research participants to regulatory authorities and everyone in between including authors, readers, educators, funders, academic and health care institutions, editors, peer reviewers, and guideline developers. Future studies may include broad surveys of authors, journal editors, funding agencies, ethics boards, and readers to solicit opinions and understanding of the role of reporting guidelines in the medical research and literature.

Data sharing statement

Unpublished study data are available upon request.

Author contributions

Contributed to the conception and design of the study, development of data extraction forms, search strategy, analysis of results, manuscript writing, and final review of the manuscript: YJ, NS, IS, CL, HS, and GL. Contributed to the methodological design, critical revision, and final review of the manuscript: MB, LZ, BB, MW, LPFA, IN, AL, LM, MM, YC, GS, MAHL, JDA, and LT. Substantially contributed to the conception and design of the study, critical revision, and final approval of the manuscript: ZS. All the authors read and approved the final manuscript. All the authors consented and approved the manuscript for publication. All authors contributed toward data analysis, drafting and revising the paper and agree to be accountable for all aspects of the work.

Disclosure

The authors report no conflicts of interest in this work.

References

1. NCBI [database on the Internet]. *PUBMED*. 2017. Available from: <https://www.ncbi.nlm.nih.gov/pubmed>. Accessed June 16, 2018.
2. Simera I, Altman DG, Moher D, Schulz KF, Hoey J. Guidelines for reporting health research: the EQUATOR network’s survey of guideline authors. *PLoS Med*. 2008;5(6):e139.
3. Altman DG, Simera I, Hoey J, Moher D, Schulz K. EQUATOR: reporting guidelines for health research. *Lancet*. 2008;371(9619):1149–1150.
4. Schulz KF, Altman DG, Moher D; CONSORT Group. CONSORT 2010 statement: updated guidelines for reporting parallel group randomised trials. *BMJ*. 2010;340:c332.
5. Bernstein AM, Song M, Zhang X, et al. Processed and unprocessed red meat and risk of colorectal cancer: analysis by tumor location and modification by time. *PLoS One*. 2015;10(8):e0135959.
6. Alexander DD, Weed DL, Miller PE, Mohamed MA. Red meat and colorectal cancer: a quantitative update on the state of the epidemiologic science. *J Am Coll Nutr*. 2015;34(6):521–543.

7. Samaan Z, Mbuagbaw L, Kosa D, et al. A systematic scoping review of adherence to reporting guidelines in health care literature. *J Multidisciplinary Healthc.* 2013;6:169–188.
8. Moher D, Liberati A, Tetzlaff J, Altman DG; PRISMA Group. Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. *BMJ.* 2009;339:b2535.
9. Li G, Mbuagbaw L, Samaan Z, et al. State of reporting of primary biomedical research: a scoping review protocol. *BMJ Open.* 2017;7:e014749.
10. von Elm E, Altman DG, Egger M, et al; STROBE Initiative. The strengthening of reporting of observational studies in epidemiology (STROBE) statement: guidelines for reporting observational studies. *Lancet.* 2007;370(9596):1453–1457.
11. Moher D, Cook DJ, Eastwood S, Olkin I, Rennie D, Stroup DF. Improving the quality of reports of meta-analyses of randomised controlled trials: the QUOROM statement. Quality of reporting of meta-analyses. *Lancet.* 1999;354(9193):1896–1900.
12. Bossuyt PM, Reitsma JB, Bruns DE, et al; Standards for Reporting of Diagnostic Accuracy. The STARD statement for reporting studies of diagnostic accuracy: explanation and elaboration. *Clin Chem.* 2003;49(1):7–18.
13. Kilkenny C, Browne WJ, Cuthill IC, Emerson M, Altman DG. Improving bioscience research reporting: the ARRIVE guidelines for reporting animal research. *PLoS Biol.* 2010;8(6):e1000412.
14. Husereau D, Drummond M, Petrou S, et al. Consolidated health economic evaluation reporting standards (CHEERS) statement. *Cost Eff Res Allocation.* 2013;11:6.
15. Stroup DF, Berlin JA, Morton SC, et al. Meta-analysis of observational studies in epidemiology: a proposal for reporting. *JAMA.* 2000;283(15):2008–2012.
16. EQUATOR [home page on the Internet]. Available from: <http://www.equator-network.org>. Accessed June 16, 2018.
17. Thelwall M, Sud P. National, disciplinary and temporal variations in the extent to which articles with more authors have more impact: Evidence from a geometric field normalised citation indicator. *J Inf.* 2016;10(1):48–61.
18. Schrock JB, Kraeutler MJ, McCarty EC. Trends in authorship characteristics in the American journal of sports medicine, 1994 to 2014. *Am J Sports Med.* 2016;44(7):1857–1860.
19. Geminiani A, Ercoli C, Feng C, Caton JG. Bibliometrics study on authorship trends in periodontal literature from 1995 to 2010. *J Periodontol.* 2013;85(5):e136–e143.
20. Drenth JH. Multiple authorship: the contribution of senior authors. *JAMA.* 1998;280(3):219–221.
21. Wuchty S, Jones BF, Uzzi B. The increasing dominance of teams in production of knowledge. *Science.* 2007;316(5827):1036–1039.
22. Bozeman B, Fay D, Slade CP. Research collaboration in universities and academic entrepreneurship: the-state-of-the-art. *J Technol Transf.* 2013;38(1):1–67.
23. Moher D, Pham B, Jones A, et al. Does quality of reports of randomised trials affect estimates of intervention efficacy reported in meta-analyses? *Lancet.* 1998;352(9128):609–613.
24. Atkins D, Best D, Briss PA, et al; GRADE Working Group. Grading quality of evidence and strength of recommendations. *BMJ.* 2004;328(7454):1490.
25. ICMJE. Recommendations for the Conduct, Reporting, Editing, and Publication of Scholarly Work in Medical Journals. *Citeseeer.* 2016; (December):1–17. doi: 10.1080/08941920.2016.1150542
26. CONSORT [webpage on the Internet]. *Consort Endorsers.* 2018. Available from: <http://www.consort-statement.org/about-consort/endorsers>. Accessed June 16, 2018.
27. NIH [webpage on the Internet]. *U.S National Library of Medicine.* 2018. Available from: https://www.nlm.nih.gov/bsd/serfile_addedinfo.html. Accessed June 16, 2018.
28. Adie S. CONSORT compliance in surgical randomized trials: possible solutions. *Ann Surg.* 2013;261(5):4932–4932.
29. Adie S, Ma D, Harris IA, Naylor JM, Craig JC. Quality of conduct and reporting of meta-analyses of surgical interventions. *Ann Surg.* 2015;261(4):685–694.
30. Agha RA, Lee S-Y, Jeong KJL, Fowler AJ, Orgill DP. Reporting quality of observational studies in plastic surgery needs improvement: a systematic review. *Ann Plast Surg.* 2015;76(5):585–589.
31. Agha RA, Fowler AJ, Limb C, et al. Impact of the mandatory implementation of reporting guidelines on reporting quality in a surgical journal: a before and after study. *Int J Surg.* 2016;30:169–172.
32. Aguiar PM, Brito GD, Correr CJ, Lyra DP, Storpirtis S. Exploring the quality of systematic reviews on pharmacist interventions in patients with diabetes: an overview. *Ann Pharmacother.* 2014;48(7):887–896.
33. Aguiar PM, Lima TM, Storpirtis S. Systematic review of the economic evaluations of novel therapeutic agents in multiple myeloma: what is the reporting quality? *J Clin Pharm Ther.* 2016;41(2):189–197.
34. Al Faleh K, Al-Omran M. Reporting and methodologic quality of Cochrane Neonatal review group systematic reviews. *BMC Pediatr.* 2009;9:38.
35. Al-Namankany AA, Ashley P, Moles DR, Parekh S. Assessment of the quality of reporting of randomized clinical trials in paediatric dentistry journals. *Int J Paediatr Dent.* 2009;19(5):318–324.
36. Alvarez F, Meyer N, Gourraud PA, Paul C. CONSORT adoption and quality of reporting of randomized controlled trials: a systematic analysis in two dermatology journals. *Br J Dermatol.* 2009;161(5):1159–1165.
37. Anttila H, Malmivaara A, Kunz R, Autti-Rämö I, Mäkelä M. Quality of reporting of randomized, controlled trials in cerebral palsy. *Pediatrics.* 2006;117(6):2222–2230.
38. Areia M, Soares M, Dinis-Ribeiro M. Quality reporting of endoscopic diagnostic studies in gastrointestinal journals: where do we stand on the use of the STARD and CONSORT statements? *Endoscopy.* 2010;42(2):138–147.
39. Augestad KM, Berntsen G, Lassen K, et al; Study Group of Research Quality in Medical Informatics and Decision Support (SQUID). Standards for reporting randomized controlled trials in medical informatics: a systematic review of CONSORT adherence in RCTs on clinical decision support. *J Am Med Inf Assoc.* 2012;19(1):13–21.
40. Balasubramanian SP, Wiener M, Alshameeri Z, Tiruvoipati R, Elbourne D, Reed MW. Standards of reporting of randomized controlled trials in general surgery: can we do better? *Ann Surg.* 2006;244(5):663–667.
41. Bath FJOVE, Bath PM. Quality of full and final publications reporting acute stroke trials. *Stroke.* 2000;29(10):2203–2210.
42. Bereza BG, Machado M, Einarson TR. Assessing the reporting and scientific quality of meta-analyses of randomized controlled trials of treatments for anxiety disorders. *Ann Pharmacother.* 2008;42:1402–1409.
43. Bian ZX, Moher D, Dagenais S, et al. Improving the quality of randomized controlled trials in Chinese herbal medicine, part IV: applying a revised CONSORT checklist to measure reporting quality. *Zhong Xi Yi Jie He Xue Bao.* 2006;4(3):233–242.
44. Biondi-Zoccai G, Lotrionto M, Abbate A, Testa L. Compliance with QUOROM and quality of reporting of over-lapping meta-analyses on the role of acetylcysteine in the prevention of contrast associated nephropathy. *BMJ.* 2006;332(7535):199–204.
45. Borg Debono V, Zhang S, Ye C, et al. The quality of reporting of RCTs used within a postoperative pain management meta-analysis, using the CONSORT statement. *BMC Anesthesiol.* 2012;12:13.
46. Bousquet PJ, Calderón MA, Demoly P, et al. The consolidated standards of reporting trials (CONSORT) statement applied to allergen-specific immunotherapy with inhalant allergens: A global allergy and asthma European network (GA2LEN) article. *J Allergy Clin Immunol.* 2011;127(1):49–56. doi:10.1016/j.jaci.2010.11.111
47. Bramhall M, Florez-Vargas O, Stevens R, Brass A, Cruickshank S. Quality of methods reporting in animal models of colitis. *Inflamm Bowel Dis.* 2015;21(6):1248–1259.
48. Cairo F, Sanz I, Matesan P, Nieri M, Pagliaro U. Quality of reporting of randomized control trials to implant in dentistry. *J Clin Periodontol.* 2012;39:202–206.

49. Capili B, Anastasi JK, Geiger JN. Adverse event reporting in acupuncture clinical trials focusing on pain. *Clin J Pain*. 2010;26(1):43–48.
50. Cavadas V, Branco F, Carvalho FL, Osório L, Gomes MJ, Silva-Ramos M. The quality of reporting of randomized controlled trials in pelvic organ prolapse. *Int Urogynecol J*. 2011;22(9):1117–1125.
51. Choi J, Jun JH, Kang BK, Kim KH, Lee MS. Endorsement for improving the quality of reports on randomized controlled trials of traditional medicine journals in Korea: a systematic review. *Trials*. 2014;15:429.
52. Chowers MY, Gottesman BS, Leibovici L, Pielmeier U, Andreassen S, Paul M. Reporting of adverse events in randomized controlled trials of highly active antiretroviral therapy: systematic review. *J Antimicrob Chemother*. 2009;64(2):239–250.
53. Cook DA, Levinson AJ, Garside S. Method and reporting quality in health professions education research: a systematic review. *Med Educ*. 2011;45(3):227–238.
54. Daitch V, Babich T, Singer P, Leibovici L. Quality of reporting nutritional randomized controlled trials in patients with cystic fibrosis. *J Pediatr Gastroenterol Nutr*. 2016;63(2):265–269.
55. Dasi F, Navarro-García MM, Jiménez-Heredia M, et al. Evaluation of the quality of publications on randomized clinical trials using the consolidated standards of reporting trials (CONSORT) statement guidelines in a Spanish tertiary hospital. *J Clin Pharmacol*. 2012;52(7):1106–1114.
56. Delaney M, Meyer E, Cserti-Gazdewich C, et al. A systematic assessment of the quality of reporting for platelet transfusion studies. *Transfusion*. 2010;50(10):2135–2144.
57. DeMauro SB, Giaccone A, Kirpalani H, Schmidt B. Quality of reporting of neonatal and infant trials in high-impact journals. *Pediatrics*. 2011;128(3):e639–e644.
58. de Vries TW, van Roon EN. Low quality of reporting adverse drug reactions in paediatric randomised controlled trials. *Arch Dis Child*. 2010;95(12):1023–1026.
59. Dias S, McNamee R, Vail A. Evidence of improving quality of reporting of randomized controlled trials in subfertility. *Hum Reprod*. 2006;21(10):2617–2627.
60. Ethgen M, Boutron L, Steg PG, Roy C, Ravaud P. Quality of reporting internal and external validity data from randomized controlled trials evaluating stents for percutaneous coronary intervention. *BMC Med Res Methodol*. 2009;9:24–24.
61. Eyawo O, Lee C-W, Rachlis B, Mills EJ. Reporting of noninferiority and equivalence randomized trials for major prostaglandins: a systematic survey of the ophthalmology literature. *Trials*. 2008;9:69–69.
62. Fan F-F, Xu Q, Sun Q, Zhao SJ, Wang P, Guo XR. Assessment of the reporting quality of randomized controlled trials on treatment of coronary heart disease with traditional Chinese medicine from the Chinese journal of integrated traditional and Western medicine: a systematic review. *PLoS One*. 2014;9(1):e86360.
63. Farrokhyar F, Chu R, Whitlock R, Thabane L. A systematic review of the quality of publications reporting coronary artery bypass grafting trials. *Can J Surg*. 2007;50(4):266–277.
64. Fidalgo BMR, Crabb DP, Lawrenson JG. Methodology and reporting of diagnostic accuracy studies of automated perimetry in glaucoma: evaluation using a standardised approach. *Ophthalmic Physiol Opt*. 2015;35(3):315–323.
65. Fleming PS, Seehra J, Polychronopoulou A, Fedorowicz Z, Pandis N. A PRISMA assessment of the reporting quality of systematic reviews in orthodontics. *Angle Orthod*. 2013;83(1):158–163.
66. Fontela PS, Pant Pai N, Schiller I, Dendukuri N, Ramsay A, Pai M. Quality and reporting of diagnostic accuracy studies in TB, HIV and malaria: evaluation using QUADAS and STARD standards. *PLoS One*. 2009;4(11):e7753.
67. Freeman K, Szczepura A, Osipenko L. Non-invasive fetal RHD genotyping tests: a systematic review of the quality of reporting of diagnostic accuracy in published studies. *Eur J Obstet Gynecol Reprod Biol*. 2009;142(2):91–98.
68. Froud R, Eldridge S, Diaz Ordaz K, Marinho VCC, Donner A. Quality of cluster randomized controlled trials in oral health: a systematic review of reports published between 2005 and 2009. *Community Dent Oral Epidemiol*. 2012;40(suppl 1):3–14.
69. Fung AE, Palanki R, Bakri SJ, Depperschmidt E, Gibson A. Applying the CONSORT and STROBE statements to evaluate the reporting quality of neovascular age-related macular degeneration studies. *Ophthalmology*. 2009;116(2):286–296.
70. Gagnier JJ, DeMelo J, Boon H, Rochon P, Bombardier C. Quality of reporting of randomized controlled trials of herbal medicine interventions. *Am J Med*. 2006;119(9):800.e1–e11.
71. Gao J, Deng G, Hu Y, et al. Quality of reporting on randomized controlled trials on recurrent spontaneous abortion in China. *Trials*. 2015;16:172–172.
72. Gianola S, Gasparini M, Agostini M, et al. Survey of the reporting characteristics of systematic reviews in rehabilitation. *Phys Ther*. 2013;93(11):1456–1466.
73. Gohari F, Baradaran HR, Tabatabaee M, et al. Quality of reporting randomized controlled trials (RCTs) in diabetes in Iran: a systematic review. *J Diabetes Metab Disord*. 2015;15(1):36.
74. Gulin JEN, Rocco DM, García-Bournissen F. Quality of reporting and adherence to ARRIVE guidelines in animal studies for Chagas disease preclinical drug research: a systematic review. *PLoS Negl Trop Dis*. 2015;9(11):e0004194.
75. Halpern SH, Darani R, Douglas MJ, Wight W, Yee J. Compliance with the CONSORT checklist in obstetric anaesthesia randomised controlled trials. *Int J Obstet Anesth*. 2004;13(4):207–214.
76. Hemels MEH, Vicente C, Sadri H, Masson MJ, Einarson TR. Quality assessment of metaanalyses of RCTs of pharmacotherapy in major depressive disorder. *Curr Med Res Opin*. 2004;20(4):477–484.
77. Herdan A, Roth R, Grass D, et al. Improvement of quality of reporting in randomised controlled trials to prevent hypotension after spinal anaesthesia for caesarean section. *Gynecol Surg*. 2011;8:121–127.
78. Huang D, Jin X, Gao J, et al. Quality evaluation of randomized controlled trials reports of laparoscopy compared with open colorectal resection for colorectal cancer. *Expert Rev Anticancer Ther*. 2015;15(6):727–732.
79. Hui D, Arthur J, Dalal S, Bruera E. Quality of the supportive and palliative oncology literature: A focused analysis on randomized controlled trials. *Support Care Cancer*. 2012;20(8):1779–1785.
80. Junhua Z, Hongcai S, Xiumei G, et al. Methodology and reporting quality of systematic review/meta-analysis of traditional Chinese medicine. *J Altern Complement Med*. 2007;13(8):797–805.
81. Karpouzis F, Bonello R. Quality of reporting of randomised controlled trials in chiropractic using the CONSORT checklist. *Man Ther*. 2016;24:19.
82. Kiehna EN, Starke RM, Pouratian N, Dumont AS. Standards for reporting randomized controlled trials in neurosurgery. *J Neurosurg*. 2010;114(2):280–285.
83. Kim KH, Kang JW, Lee MS, Lee J-D. Assessment of the quality of reporting in randomised controlled trials of acupuncture in the Korean literature using the CONSORT statement and STRICTA guidelines. *BMJ Open*. 2014;4(7):e005068.
84. Kober T, Trelle S, Engert A. Reporting of randomized controlled trials in Hodgkin lymphoma in biomedical journals. *J Natl Cancer Inst*. 2006;98(9):620–625.
85. Ladd BO, McCrady BS, Manuel JK, Campbell W. Improving the quality of reporting alcohol outcome studies: effects of the CONSORT statement. *Addict Behav*. 2010;35:660–666.
86. Lee S-Y, Teoh PJ, Camm CF, Agha RA. Compliance of randomized controlled trials in trauma surgery with the CONSORT statement. *J Trauma Acute Care Surg*. 2013;75(4):562–572.
87. Lee SY, Sagoo H, Whitehurst K, et al. Compliance of systematic reviews in plastic surgery with the PRISMA statement. *JAMA Facial Plast Surg*. 2016;18(2):101–105.

88. Li JY, Zhang YF, Smith GS, et al. Quality of reporting of randomized clinical trials in tai chi interventions—a systematic review. *Evid Based Complement Alternat Med*. 2011;2011:383245.
89. Li J-L, Ge L, Ma JC, et al. Quality of reporting of systematic reviews published in “evidence-based” Chinese journals. *Syst Rev*. 2014;3:58–58.
90. Li J, Liu Z, Chen R, et al. The quality of reports of randomized clinical trials on traditional Chinese medicine treatments: a systematic review of articles indexed in the China National Knowledge Infrastructure database from 2005 to 2012. *BMC Complement Altern Med*. 2014;14:362–362.
91. Liu D, Jin J, Tian J, Yang K. Quality assessment and factor analysis of systematic reviews and meta-analyses of endoscopic ultrasound diagnosis. *PLoS One*. 2015;10:1–13.
92. Liu LQ, Morris PJ, Pengel LHM. Compliance to the CONSORT statement of randomized controlled trials in solid organ transplantation: a 3-year overview. *Transpl Int*. 2013;26:300–306.
93. Liu XT, Zhang X, Wen S, Peng L, Hong Q, Kang D. Impact of the Consolidated Standards of Reporting Trials (CONSORT) checklist on reporting of randomized clinical trials in traditional Chinese medicine. *J Evid Based Med*. 2015;8:192–208.
94. Liu Y, Zhang R, Huang J, et al. Reporting quality of systematic reviews/meta-analyses of acupuncture. *PLoS One*. 2014;9(11):e113172.
95. Liu Y, Zhao X, Mai Y, et al. Adherence to ARRIVE guidelines in Chinese journal reports on neoplasms in animals. *PLoS One*. 2016;11:e0154657.
96. Lu J, Gary KW, Copolillo A, Ward J, Niemeier JP, Lapane KL. Randomized controlled trials in adult traumatic brain injury: a review of compliance to CONSORT statement. *Arch Phys Med Rehabil*. 2015;96:702–714.
97. Lu L, Zeng J, Chen Y. Quality of reporting in randomized controlled trials conducted in China on the treatment of cancer pain. *Expert Rev Anticancer Ther*. 2011;11:871–877.
98. Ma B, Guo J, Qi G, et al. Epidemiology, quality and reporting characteristics of systematic reviews of traditional Chinese medicine interventions published in Chinese journals. *PLoS One*. 2011;6:e20185.
99. Ma B, Qi GQ, Lin XT, Wang T, Chen ZM, Yang KH. Epidemiology, quality, and reporting characteristics of systematic reviews of acupuncture interventions published in Chinese journals. *J Altern Complement Med*. 2012;18:813–817.
100. Marshman Z, Farid F. The quality of reporting of randomised controlled trials in dental public health. *Community Dent Health*. 2010;27:253–256.
101. McCormick F, Cvetanovich GL, Kim JM, et al. An assessment of the quality of rotator cuff randomized controlled trials: utilizing the Jadad score and CONSORT criteria. *J Shoulder Elbow Surg*. 2013;22:1180–1185.
102. Miller E, Roposch A, Uleryk E, Doria AS. Juvenile idiopathic arthritis of peripheral joints. Quality of reporting of diagnostic accuracy of conventional MRI. *Acad Radiol*. 2009;16:739–757.
103. Moberg-Mogren E, Nelson DL. Evaluating the quality of reporting occupational therapy randomized controlled trials by expanding the CONSORT criteria. *Am J Occup Ther*. 2006;60:226–235.
104. Moher D, Sampson M, Campbell K, et al. Assessing the quality of reports of randomized trials in pediatric complementary and alternative medicine. *BMC Pediatr*. 2002;12:20–21.
105. Montané E, Vallano A, Vidal X, Aguilera C, Laporte J-R. Reporting randomised clinical trials of analgesics after traumatic or orthopaedic surgery is inadequate: a systematic review. *BMC Clin Pharmacol*. 2010;10:2–2.
106. Montgomery AA, Astin MP, Peters TJ. Reporting of factorial trials of complex interventions in community settings: a systematic review. *Trials*. 2011;12:179.
107. Nicolau I, Ling D, Tian L, Lienhardt C, Pai M. Methodological and reporting quality of systematic reviews on tuberculosis. *Int J Tuberc Lung Dis*. 2013;17:1160–1169.
108. Norton-Mabus JC, Nelson DL. Reporting of randomized controlled trials in occupational therapy and speech therapy: evaluation using an expansion of the consort statement. *Occup Particip Health*. 2008;28:64–71.
109. Ntala C, Birmipili P, Worth A, Anderson NH, Sheikh A. The quality of reporting of randomised controlled trials in asthma: a systematic review. *Prim Care Respir J*. 2013;22:417–424.
110. Panic N, Leoncini E, De Belvis G, Ricciardi W, Boccia S. Evaluation of the endorsement of the preferred reporting items for systematic reviews and meta-analysis (PRISMA) statement on the quality of published systematic review and meta-analyses. *PLoS One*. 2013;8(12):e83138.
111. Parsons NR, Hiskens R, Price CL, Achten J, Costa ML. A systematic survey of the quality of research reporting in general orthopaedic journals. *J Bone Joint Surg*. 2011;93:1154–1159.
112. Patel MX, Collins S, Hellier J, Bhatia G, Murray RM. The quality of reporting of phase II and III trials for new antipsychotics: a systematic review. *Psychol Med*. 2014;45(3):467–479.
113. Piggott M, McGee H, Feuer D. Has CONSORT improved the reporting of randomized controlled trials in the palliative care literature? A systematic review. *Palliat Med*. 2004;18:32–38.
114. Péron J, Pond GR, Gan HK, et al. Quality of reporting of modern randomized controlled trials in medical oncology: a systematic review. *J Natl Cancer Inst*. 2012;104:982–989.
115. Peters JPM, Hooft L, Grolman W, Stegeman I. Reporting quality of systematic reviews and meta-analyses of otorhinolaryngologic articles based on the PRISMA statement. *PLoS One*. 2015;10:1–11.
116. Plint AC, Moher D, Morrison A, et al. Does the CONSORT checklist improve the quality of reports of randomised controlled trials? A systematic review. *Med J Aust*. 2006;185:263–267.
117. Prady SL, Richmond SJ, Morton VM, MacPherson H. A systematic evaluation of the impact of STRICTA and CONSORT recommendations on quality of reporting for acupuncture trials. *PLoS One*. 2008;3(2):e1577.
118. Pratoomsot C, Sruamsiri R, Dilokthornsakul P, Chaiyakunapruk N. Quality of reporting of randomised controlled trials of herbal interventions in ASEAN plus six countries: a systematic review. *PLoS One*. 2015;10(1):e108681.
119. Rao A, Brück K, Methven S, et al. Quality of reporting and study design of CKD cohort studies assessing mortality in the elderly before and after STROBE: a systematic review. *PLoS One*. 2016;11:1–16.
120. Rice DB, Kloda LA, Shrier I, Thombs BD. Reporting completeness and transparency of meta-analyses of depression screening tool accuracy: a comparison of meta-analyses published before and after the PRISMA statement. *J Psychosom Res*. 2016;87:57–69.
121. Rios LP, Oduyungbo A, Moitri MO, Rahman MO, Thabane L. Quality of reporting of randomized controlled trials in general endocrinology literature. *J Clin Endocrinol Metab*. 2008;93:3810–3816.
122. Rikos D, Dardiotis E, Tsvigoulis G, Zintzaras E, Hadjigeorgiou GM. Reporting quality of randomized-controlled trials in multiple sclerosis from 2000 to 2015, based on CONSORT statement. *Mult Scler Relat Disord*. 2016;9:135–139.
123. Schwarz F, Iglhaut G, Becker J. Quality assessment of reporting of animal studies on pathogenesis and treatment of peri-implant mucositis. *Clin Periodontol*. 2012;39(suppl 12):63–72.
124. Scott P, Ott F, Egger M, Low N. Completeness of reporting in randomized controlled trials of 3 vaccines. *Pediatr Infect Dis J*. 2012;31:1286–1294.
125. Shawyer AC, Pemberton J, Kanters D, Alnaqi AAA, Flageole H. Quality of reporting of the literature on gastrointestinal reflux after repair of esophageal atresia-tracheoesophageal fistula. *J Pediatr Surg*. 2015;50:1099–1103.
126. Shea B, Boers M, Grimshaw JM, Hamel C, Bouter LM. Does updating improve the methodological and reporting quality of systematic reviews? *BMC Med Res Methodol*. 2006;6:1–7.
127. Shea B, Bouter LM, Grimshaw JM, et al. Scope for improvement in the quality of reporting of systematic reviews. From the Cochrane Musculoskeletal Group. *J Rheumatol*. 2006;33:9–15.

128. Stevely A, Dimairo M, Todd S, et al. An investigation of the shortcomings of the CONSORT 2010 statement for the reporting of group sequential randomised controlled trials: a methodological systematic review. *PLoS One*. 2015;10:1–20.
129. Strech D, Soltmann B, Weikert B, Bauer M, Pfennig A. Quality of reporting of randomized controlled trials of pharmacologic treatment of bipolar disorders: a systematic review. *J Clin Psychiatry*. 2011;72:1214–1221.
130. Tan WK, Wigley J, Shantikumar S. The reporting quality of systematic reviews and meta-analyses in vascular surgery needs improvement: a systematic review. *Int J Surg*. 2014;12:1262–1265.
131. Thabane L, Chu R, Cuddy K, Douketis J. What is the quality of reporting in weight loss intervention studies? A systematic review of randomized controlled trials. *Int J Obes*. 2007;31:1554–1559.
132. Tunis AS, McInnes MDF, Hanna R, Esmail K. Association of study quality with completeness of reporting: have completeness of reporting and quality of systematic reviews and meta-analyses in major radiology journals changed since publication of the PRISMA statement? *Radiology*. 2013;269:413–426.
133. Turner L, Shamseer L, Altman DG, et al. Consolidated standards of reporting trials (CONSORT) and the completeness of reporting of randomised controlled trials (RCTs) published in medical journals. *Cochrane Database Syst Rev*. 2012;11:MR000030.
134. Vigna-Taglianti F, Vineis P, Liberati A, Faggiano F. Quality of systematic reviews used in guidelines for oncology practice. *Ann Oncol*. 2006;17:691–701.
135. Walleser S, Hill SR, Bero LA. Characteristics and quality of reporting of cluster randomized trials in children: reporting needs improvement. *J Clin Epidemiol*. 2011;64:1331–1340.
136. Wang G, Mao B, Xiong ZY, et al. The quality of reporting of randomized controlled trials of traditional Chinese medicine: a survey of 13 randomly selected journals from mainland China. *Clin Ther*. 2007;29:1456–1467.
137. Wang P, Xu Q, Sun Q, Fan FF, Guo XR, Guo F. Assessment of the reporting quality of randomized controlled trials on the treatment of diabetes mellitus with traditional Chinese medicine: a systematic review. *PLoS One*. 2013;8(7):e70586.
138. Wangge G, Klungel OH, Roes KC, de Boer A, Hoes AW, Knol MJ. Room for improvement in conducting and reporting non-inferiority randomized controlled trials on drugs: a systematic review. *PLoS One*. 2010;5(10):e13550.
139. Weingärtner V, Dargatz N, Weber C, et al. Patient reported outcomes in randomized controlled cancer trials in advanced disease: a structured literature review. *Expert Rev Clin Pharmacol*. 2016;9:821–829.
140. Weir CR, Stagers N, Laukert T. Reviewing the impact of computerized provider order entry on clinical outcomes: the quality of systematic reviews. *Int J Med Inform*. 2012;81(4):219–231.
141. Wen J, Ren Y, Wang L, et al. The reporting quality of meta-analyses improves: a random sampling study. *J Clin Epidemiol*. 2008;61:770–775.
142. Willis B, Quigley M. The assessment of the quality of reporting of meta-analyses in diagnostic research: a systematic review. *BMC Med Res Methodol*. 2011;11:163–163.
143. Yao AC, Khajuria A, Camm CF, Edison E, Agha R. The reporting quality of parallel randomised controlled trials in ophthalmic surgery in 2011: a systematic review. *Eye*. 2014;28:1341–1349.
144. Zafar A, Khan GI, Siddiqui MA. The quality of reporting of diagnostic accuracy studies in diabetic retinopathy screening: a systematic review. *Clin Experiment Ophthalmol*. 2008;36(6):537–542.
145. Zhang Z-W. Epidemiology, quality, and reporting characteristics of systematic reviews and meta-analyses of observational studies published in Chinese journals. *BMJ Open*. 2015;63:446–455.
146. Zhao X, Zhen Z, Guo J, et al. Assessment of the reporting quality of placebo-controlled randomized trials on the treatment of type 2 diabetes with traditional Chinese medicine in mainland China: a PRISMA-compliant systematic review. *Medicine*. 2016;95(3):e2522.
147. Zheng SL, Chan FT, Maclean E, Jayakumar S, Nabeebaccus AA. Reporting trends of randomized controlled trials in heart failure with preserved ejection fraction: a systematic review. *Open Heart*. 2016;3(2):e000449.
148. Zhong Y, Zhou W, Jiang H, et al. Quality of reporting of two-group parallel randomized controlled clinical trials of multi-herb formulae: a survey of reports indexed in the Science Citation Index Expanded. *Eur J Integr Med*. 2011;3(4):e309–e316.
149. Zintzaras E, Kitsios GD, Papatheanasiou AA, et al. Randomized trials of dopamine agonists in restless legs syndrome: a systematic review, quality assessment, and meta-analysis. *Clin Ther*. 2010;32(2):221–237.
150. Zintzaras E, Papatheanasiou AA, Ziogas DC, Voulgarelis M. The reporting quality of studies investigating the diagnostic accuracy of anti-CCP antibody in rheumatoid arthritis and its impact on diagnostic estimates. *BMC Musculoskelet Disord*. 2012;13:113.
151. Ziogas DC, Zintzaras E. Analysis of the quality of reporting of randomized controlled trials in acute and chronic myeloid leukemia, and myelodysplastic syndromes as governed by the CONSORT statement. *Ann Epidemiol*. 2009;19(7):494–500.

Journal of Multidisciplinary Healthcare

Publish your work in this journal

The Journal of Multidisciplinary Healthcare is an international, peer-reviewed open-access journal that aims to represent and publish research in healthcare areas delivered by practitioners of different disciplines. This includes studies and reviews conducted by multidisciplinary teams as well as research which evaluates the results or conduct of such teams or health

care processes in general. The journal covers a very wide range of areas and welcomes submissions from practitioners at all levels, from all over the world. The manuscript management system is completely online and includes a very quick and fair peer-review system. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <https://www.dovepress.com/journal-of-multidisciplinary-healthcare-journal>

Dovepress