

Does the type of culture medium used influence birthweight of children born after IVF?

Heleen Zandstra¹, Aafke P. A. Van Montfoort², and John C. M. Dumoulin^{2,*}

¹Department of Obstetrics and Gynaecology, GROW School for Oncology and Developmental Biology, Maastricht University Medical Centre, Maastricht, The Netherlands ²Department of Obstetrics and Gynaecology, IVF Laboratory, Maastricht University Medical Center, Maastricht, The Netherlands

*Correspondence address. E-mail: john.dumoulin@mumc.nl

Hum Reprod 2015;**30**:530–542

The authors would like to apologise for a typing error in the results of this article. Ziebe *et al.* (2013) included 304 children in their analysis; GM-CSF n = 163, control n = 141 (and not 204 children as erroneously stated in the article).

: The authors would like to assure readers that this does not affect any
: content of the article.

: Ziebe S, Loft A, Povlsen BB, Erk K, Agerholm I, Aasted M, Gabrielsen A, Hnida
: C, Zobel DP, Munding B *et al.* A randomized clinical trial to evaluate the effect
: of granulocyte-macrophage colony-stimulating factor (GM-CSF) in embryo
: culture medium for *in vitro* fertilization. *Fertil Steril* 2013;**99**:1600–1609.